

A New O-Antigen Gene Cluster Has a Key Role in the Virulence of the *Escherichia coli* Meningitis Clone O45:K1:H7[∇]

Céline Plainvert,¹ Philippe Bidet,¹ Chantal Peigne,¹ Valérie Barbe,² Claudine Médigue,³
Erick Denamur,^{4,5} Edouard Bingen,¹ and Stéphane Bonacorsi^{1*}

Laboratoire d'Études de Génétique Bactérienne dans les Infections de l'Enfant (EA 3105), Université Paris Diderot-Paris 7, Hôpital Robert Debré (APHP), 75019 Paris,¹ Commissariat à l'Énergie Atomique, Direction des Sciences du Vivant, Institut de Génétique, Génomique, 91057 Evry Cedex,² Commissariat à l'Énergie Atomique, Direction des Sciences du Vivant, Institut de Génétique, Atelier de Génétique Comparative, CNRS UMR8030, 91057 Evry Cedex,³ Institut National de la Santé et de la Recherche Médicale U722, 75018 Paris,⁴ and Université Paris Diderot-Paris 7, site Xavier Bichat, 75018 Paris,⁵ France

Received 26 June 2007/Accepted 20 September 2007

A new highly pathogenic clone of *Escherichia coli* meningitis strains harboring the unusual serogroup O45 has recently emerged in France. To gain insight into the pathogenicity of this new clone, we investigated the possible role of antigen O45 in the virulence of strain S88 (O45:K1:H7), representative of this emerging clone. We first showed that the S88 O-antigen gene cluster sequence differs from that of O45 in the reference strain *E. coli* 96-3285, suggesting that the two O45 polysaccharides, while probably sharing a community of epitopes, represent two different antigens. The unique functional organization of the two O-antigen gene clusters and the low DNA sequence homology of the orthologous genes suggest that the two loci originated from a common ancestor and have since undergone multiple recombination events. Phylogenetic analysis based on the flanking gene *gnd* sequences indicates that the S88 antigen O45 (O45_{S88}) gene cluster may have been acquired, at least in part, from another member of the *Enterobacteriaceae*. Mutagenesis of the O45_{S88} antigen gene cluster was used for functional analysis of the loci and revealed the crucial role of the O polysaccharide in S88 virulence in a neonatal rat meningitis model. We also developed a PCR method to specifically identify the O45_{S88} antigen gene cluster. Together, our findings suggest that horizontal acquisition of a new O-antigen gene cluster, at least partly from another species, may have been a key event in the emergence and virulence of the *E. coli* O45:K1:H7 clone in France.

O antigen, the polysaccharide constituent of lipopolysaccharide (LPS), is a major focus of studies of infections caused by extraintestinal pathogenic *Escherichia coli* (ExPEC) strains for two main reasons. First, it can be used for typing studies, which have shown the highly clonal organization of some ExPEC strains, such as neonatal meningitis strains (39). Indeed, the most important globally distributed neonatal meningitis clone is the so-called archetypal clone O18:K1:H7 (1). Second, O antigens may play a major role in the virulence of ExPEC and notably in resistance to serum bactericidal activity; their characterization may thus help to understand the pathophysiology of ExPEC infections (22).

Serotyping combined with molecular methods previously allowed us to detect the emergence in France of a highly virulent meningitis-causing clone closely related to the O18:K1:H7 clone but harboring the unusual serogroup O45, as well as capsular antigen K1 and flagellar antigen H7 (12). O45 antigen has only sporadically been described in ExPEC strains (34, 46) and is absent from most *E. coli* meningitis strains in American and European collections (1, 23), except in Hungary (14).

In contrast, several intestinal pathogenic *E. coli* (InPEC)

strains have been identified as belonging to serogroup O45, such as enterotoxigenic *E. coli* and Shiga toxin-producing *E. coli* (STEC) (9, 41). In addition, *E. coli* O45 strains producing cytotoxic necrotizing factor or expressing the “attaching and effacing” phenotype have been isolated from several diarrheic animals (4, 32). The importance of O45 antigen in InPEC strains led DebRoy et al. to sequence the genomic region involved in O-antigen synthesis in the CDC reference strain 96-3285 (O45:H2) and to develop a specific PCR method to detect *E. coli* O45 (16).

Many PCR tests for *E. coli* serogroups have recently been developed by targeting the genes involved in O-antigen synthesis and clustered in the so-called O-antigen gene cluster. Among these genes, those responsible for O-unit processing—*wzx* and *wzy*, encoding the O-antigen flippase and O-antigen polymerase, respectively—are specific for the O-unit composing the polysaccharide (38). These genes may therefore be ideal targets for specific PCR. Indeed, the classical agglutination reaction with specific antiserum is laborious and expensive, and cross-reactions between serogroups can occur (33).

The emergence of the unusual O45 antigen in ExPEC strains prompted us to analyze the O-antigen gene cluster sequence of S88, a strain representative of the O45:K1:H7 clone causing neonatal meningitis, by comparison with the published sequence of strain 96-3285 (O45:H2) (16). We show that the two O-antigen gene clusters are genetically related but not identical, suggesting that S88 expresses a new O antigen. We also performed a func-

* Corresponding author. Mailing address: Service de Microbiologie, Hôpital Robert Debré, 48 blvd. Sérurier, 75395 Paris cedex 19, France. Phone: 33 1 40 03 23 40. Fax: 33 1 40 03 24 50. E-mail: stephane.bonacorsi@rdb.aphp.fr.

[∇] Published ahead of print on 28 September 2007.

TABLE 1. Oligonucleotide primers for PCR sequencing, mutant construction, and O45 PCR assay

Name ^a	Target	Sequence (5' to 3') ^b	Amplicon size (bp)
gndF	<i>gnd</i>	ATGTCCAAGCAACAGATCGGCGT	1,407
gndR	<i>gnd</i>	TTAATCCAGCCATTCGGTATGGA	
rfb405S88.P0	ORF 2138	GATTCCCTGGTTATTCTCAATGCTCTCGAAGGGAAATCGT GTGTAGGCTGGAGCTGCTT	
rfb397S88.P2	ORF 2130	CACCTGATTATCATAGAAATAGAGTCCAGTTATGGCCCAAC ATATGAATATCCTCCTTAG	
rfb399S88.P0	ORF 2132	TCTGGTGTTTATACAGGGTAAATGTAAGGTTTCATCGCAAT GTGTAGGCTGGAGCTGCTT	
rfb399S88.P2	ORF 2132	GGAAGGTTAGTTAAGGCAGGGGAGCAGGCATAATAACAT CATATGAATATCCTCCTTAG	
C1	<i>cat</i>	TTATACGCAAGGCGACAAGG	
C2	<i>cat</i>	GATCTTCCGTCACAGGTAGG	
rfb405S88.FR1b	ORF 2138	CGTACATACGGTCTTCCAAC	
rfb397S88.FR2b	ORF 2130	CATTAAGCATATCCCGCTCA	
rfb400S88.FR1	ORF 2133	GAGTAGCTTTGTCTTGC	
rfb399S88.FR2	ORF 2132	TATTCCTGCGTATCCTGCTA	
wzxS88b.F	ORF 2134	GTTGCGATAGTCATGTACTG	119
wzxS88b.R	ORF 2134	GCTACAACCCCTCCCCAGAT	
wzyS88b.F	ORF 2132	GGTATCGTTCACATCGCTTA	330
wzyS88b.R	ORF 2132	GAGAAAATACTCGGTTCCGGC	
ColiBglu.1	<i>uidA</i>	TATGAACTGTGCGTCACAGCC	186
ColiBglu.2	<i>uidA</i>	CATCAGCACGTTATCGAATCC	

^a The oligonucleotide primers used for gene recombination are designated by the suffixes P0 and P2; the oligonucleotide primers used to control correct introduction and excision of the *cat* gene are designated by the suffixes FR1 and FR2 and flank the DNA target segment. Oligonucleotide primers used for O45 triplex PCR are shown in bold characters.

^b Boldface characters indicate the 20 nucleotides homologous to the *cat* gene sequence.

tional analysis of the S88 O-antigen gene cluster and developed a specific PCR to detect strains harboring the S88 somatic antigen.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth media. The *E. coli* meningitis strain S88 is representative of the French virulent clone O45:K1:H7 (12). This serotype was kindly determined by three different National Reference Centers in Denmark (Flemming Schultz, Serum Statens Institut, Copenhagen, Denmark), Germany (Lothar Beutin, Robert Koch Institute, Berlin, Germany), and Spain (Jorge Blanco, Lugo, Spain). Strain S88 belongs to phylogenetic group B2 and harbors ribotype B₂ (12). The CDC reference strain for antigen O45, *E. coli* strain 96-3285 (O45:H2), the O-antigen gene cluster of which was recently sequenced, was kindly provided by Chitrita DebRoy (*E. coli* Reference Center, The Pennsylvania State University). This strain was shown in a previously described PCR grouping method to belong to the nonvirulent phylogenetic group B1 (8) and harbors none of the virulence factors usually encountered among meningitis strains (12). The Danish O45 reference strain, strain H61 (O45:K1:H10), which belongs to phylogenetic group A, was purchased from the Staten Serum Institut (Copenhagen, Denmark). Thirty-six meningitis strains and nine urosepsis strains, all belonging to serotype O45:K1:H7 (determined by the Spanish National Reference Center) and to ribotype B₂ (like S88) and sharing the same virulence genotype, were also used (7, 12). A total of 130 reference strains for the different somatic antigens (O antigens 1, 2, 4 to 24, 26 to 30, 32 to 45, 48 to 53, 55, 57, 59 to 61, 63, 64, 68 to 70, 73 to 77, 79, 81, 83 to 86, 89 to 91, 93, 95, 98, 101 to 121, 123, 125 to 129, 134 to 136, 138 to 142, 144 to 152, 154 to 162, and 164) were kindly provided by Patrick Grimont (Institut Pasteur, Paris, France). The 72 *E. coli* reference (ECOR) strains (31) were also used.

The plasmids pKD46, pKD3, and pCP20 were kindly provided by Lionello Bossi (Centre de Genetique Moleculaire, CNRS, Gif sur Yvette, France).

Luria-Bertani (LB) broth and agar were routinely used and were supplemented with chloramphenicol (12 µg/ml) or ampicillin (50 µg/ml) as necessary.

Sequencing of the S88 O-antigen gene cluster. The O-antigen gene cluster in strain S88 was sequenced as part of a whole-genome sequencing project (ColiScope; www.genoscope.cns.fr) at the Genoscope sequencing center (Evry, France). Sequencing and assembly of the S88 genome were performed as previously described (3). Briefly, the complete genome sequence of *E. coli* S88 was determined using the whole-genome shotgun method (10× coverage). Three libraries were made as follows: two plasmid libraries of 3 kb and 10 kb, obtained

by mechanical shearing, were constructed in plasmid pcDNA2.1 (Invitrogen) and in home vector pCNS (pSU18 modified), respectively. One bacterial artificial chromosome library of average insert size of 30 kb was constructed by enzymatic digestion (HindIII) into pBeloBacII (CalTech). The Phred/Phrap/Consed software package (www.phrap.com) was used for sequence assembly and quality assessment. To resolve contigs, sequence finishing was performed by PCR amplification, primer walking, and/or transposition.

The MaGe (magnifying genomes) software program was used for gene annotation and comparative analysis of the S88 genome (43). Using the AMIGene (annotation of microbial genes) (10) program, a total of 4,859 coding sequences were predicted (and assigned a unique identifier prefixed with ECOS88_) and submitted to automatic functional annotation, including synteny computation (that is, conservation of the chromosomal colocalization between pairs of orthologous genes from different genomes). Manual validation of the automatic annotation was performed using the MaGe interface, which allows graphic visualization of the *E. coli* S88 annotations enhanced by a synchronized representation of synteny groups in other genomes chosen for comparisons (43). Protein motifs and domains were identified by using the InterPro databank (2). TMHMM, version 2.0, was used to identify transmembrane domains (26). Sequence data for comparative analyses were obtained from NCBI databases (ftp://ftp.ncbi.nlm.nih.gov). Annotations of the O-antigen gene cluster described in this paper range from ECOS88_2129 (*gnd*) to ECOS88_2139 (*galF*).

Phylogenetic analysis. To gain insight into the evolutionary history of the O45 antigen gene cluster in strains S88 and 96-3285, nucleotide sequences of the internal genes *mmlABC* and the external gene *gnd* from different *E. coli* strains and several other representative gram-negative bacteria were extracted from the GenBank database. The *gnd* sequence of strain 96-3285 was not available and was determined here from a PCR product obtained with the primers indicated in Table 1. The ClustalW program was used to align the sequences (40). Phylogenetic and molecular evolutionary relationships were examined by using the neighbor-joining method implemented with MEGA, version 3.1, software (27). Bootstrap confidence values for each node of the trees were calculated over 100 replicate trees. Phylogenetic analysis was also performed using the maximum parsimony method, also implemented in MEGA, version 3.1, software.

Construction of mutants. S88 mutants were obtained by the method of Datzenko and Wanner (15) as previously described (29). Briefly, this PCR-based method uses the plasmid pKD46, which allows homologous recombination to occur directly with PCR products. This plasmid is a temperature-sensitive replicon that carries the bacteriophage λ Red system under the control of an

TABLE 2. Genes located in the O-antigen gene cluster of *E. coli* S88 serotype O45:K1:H7

ORF	Proposed gene name	Location (nucleotides)	% G+C content	No. of amino acids in gene product	Putative function	Most significant homolog (strain)	% Amino acid identity	% Amino acid similarity
2138	<i>rmlB</i>	2108855–2109937	46.9	1,083	dTDP-glucose-4,6-dehydratase	dTDP-glucose-4,6-dehydratase (<i>E. coli</i> 96-3285)	90	95
2137	<i>rmlC</i>	2108241–2108783	37	543	dTDP-6-deoxy-D-glucose-3,5-epimerase	dTDP-4-dehydrorhamnose-3,5-epimerase (<i>E. coli</i> 96-3285)	73	85
2136	<i>ill</i>	2107363–2108178	34.5	816	dTDP-6-deoxy-L-xylo-4-hexulose reductase	dTDP-6-deoxy-L-xylo-4-hexulose reductase (<i>E. coli</i> O66)	52	68
2135	<i>wbvA</i>	2106363–2107361	30.6	999	Unknown	Hypothetical protein (<i>E. coli</i> 96-3285)	34	51
2134	<i>wzx</i>	2105114–2106376	34.3	1,263	O-antigen flippase	O-antigen flippase (<i>E. coli</i> 96-3285)	23	42
2133	<i>wbvB</i>	2104150–2105127	38.7	978	Glycosyl transferase	Glycosyl transferase (<i>E. coli</i> 96-3285)	38	52
2132	<i>wzy</i>	2103066–2104157	33.7	1,092	O-antigen polymerase	Putative membranous protein (<i>Photobacterium profundum</i>)	34	54
2131	<i>wbvC</i>	2102570–2103100	31.1	531	O-Acetyltransferase	Serine acetyltransferase (<i>E. coli</i> 96-3285)	42	63
2130	<i>rmlA</i>	2101559–2102425	40.8	867	Glucose-1-phosphate thymidyl transferase	Glucose-1-phosphate thymidyl transferase (<i>E. coli</i> 96-3285)	81	88

arabinose-inducible promoter. Once introduced into the bacterium by electroporation, it renders S88 transformable with linear DNA obtained by PCR. The chloramphenicol acetyltransferase (*cat*) gene, carried by plasmid pKD3, was amplified by PCR with primers bearing extensions of 40 nucleotides homologous to the initial and final portions of the DNA target segment. All the primers used in this study are listed in Table 1. Transformation by electroporation of strain S88 expressing bacteriophage λ Red functions with the PCR product yielded recombinants carrying the DNA target fused to the *cat* gene. Correct introduction of the *cat* gene was controlled by PCR with primers (Table 1) flanking the initial and final portions of the DNA target segment and primers homologous to the *cat* gene, as previously described (29). Conservation of the main virulence determinants in the mutants was controlled by multiplex PCR with primers located in the main extraintestinal virulence genes of S88 (*fyuA*, *papC*, *papGII*, *iucC*, and *iroN*) as previously described (11). Antigens O45 and K1 were detected with specific antisera and phage from the Staten Serum Institute (Copenhagen, Denmark).

Analysis of the LPS. The LPS was extracted with denaturing buffer composed as follows (final concentrations): 0.2% sodium dodecyl sulfate (SDS), 1% β -mercaptoethanol, 36% glycerol, 30 mM Tris-HCl (pH 7.4), and 0.001% bromophenol blue. Strains suspended in 1 ml of phosphate-buffered saline at an optical density at 600 nm of 2 were centrifuged at $3,500 \times g$ for 10 min at 25°C, and the pellet was resuspended with 500 μ l of denaturing buffer. The sample was denatured for 5 min at 95°C, and 3 μ l of proteinase K was added after cooling to room temperature. The sample was incubated for 1 h at 55°C with proteinase K and then centrifuged for 30 min at $13,000 \times g$ and 25°C. The LPS-containing supernatants were stored at -20°C. The LPS preparations were separated on 16% SDS-tricine-polyacrylamide gels at 30 V for 30 min and 100 V for 3 h. LPS was then visualized by silver staining as previously described (42).

Animal meningitis model. *E. coli* bacteremia and meningitis were induced in newborn rats as described by Houdouin et al. (21). Briefly, pathogen-free Sprague-Dawley rats were obtained from Charles River Laboratories (L'Abresle, France) at 4 days of age, together with their mothers. At 5 days of age all the pups were inoculated intraperitoneally with ~ 500 CFU of the tested strain in physiological saline. Eighteen hours later, 5 μ l of blood was obtained by tail incision. The animals were then killed, and 5 μ l of cerebrospinal fluid (CSF) was immediately obtained by cisternal puncture. All samples were quantitatively cultured by plating dilutions of blood and cerebrospinal fluid on LB agar.

O-serogroup-specific PCR assay. Genes encoding O-antigen flippase (*wzx*) and O-antigen polymerase (*wzy*) in the O-antigen gene cluster are specific for each O antigen and are suitable targets for serogroup-specific PCR (36). In order to develop a specific PCR assay for identifying *E. coli* strains harboring a somatic antigen identical to that expressed by strains belonging to the O45:K1:H7 meningitis clone, we designed specific primers for *wzx* and *wzy*. Template DNA for the PCR assays was prepared by mixing 2 μ l of bacterial colony formed on LB agar in 500 μ l of sterile water and heating at 100°C for 10 min. Then, the suspension was centrifuged at $11,000 \times g$ for 3 min at 4°C, and the supernatant containing DNA was used for PCR. PCR was performed in a final volume of 50 μ l using a Qiagen Multiplex PCR kit with 5 μ l of template DNA, 0.3 μ M *wzx* primers, and a 0.2 μ M concentration (each) of *wzy* and *uidA* primers. PCR was

performed in an Icyler (Bio-Rad, Marnes la Coquette, France) as follows: denaturation at 95°C for 15 min; 30 cycles of 94°C for 30 s, 55°C for 90 s, and 72°C for 90 s; and final extension at 72°C for 10 min. The PCR products were electrophoresed in 2% standard agarose gels. The gels were stained with ethidium bromide and visualized under UV light. Positive samples were identified from the presence of bands of the expected sizes compared to results obtained with strain S88.

Nucleotide sequence accession number. The DNA sequences of the S88 *E. coli* O-antigen gene cluster have been deposited in the GenBank database under the accession number CU463050.

RESULTS

Functional annotation of the *E. coli* S88 O-antigen gene cluster. Analysis of the DNA sequence of the putative O-antigen gene cluster located between *galF* and *gnd* in strain S88 showed that it contains nine open reading frames (ORFs) with a total length of 8,379 bp, the same transcriptional direction from *galF* to *gnd*, and a low G+C content (30.6 to 46.9%) compared to the *E. coli* core genome (51%). The nine ORFs of the locus were assigned putative functions based on a protein database similarity search and were named using the system proposed by Reeves et al. (36) (Table 2). The most homologous proteins were almost always found in the products of the O-antigen gene cluster of *E. coli* 96-3285, the CDC reference strain for the O45 somatic antigen (Table 2).

Products of ORFs 2130, 2138, and 2137 shared a high level of identity (>70%) with RmlA, RmlB, and RmlC, respectively. ORF 2136, immediately downstream of *rmlC*, putatively encodes a protein sharing 52% identity with the *ill* product in the recently sequenced O66-antigen gene cluster (13). This gene and *rmlABC* are known to be involved in the biosynthesis pathway of 6-deoxy-L-talose, which may therefore compose the oligosaccharide unit of S88 O antigen. The other putative proteins encoded by the five remaining ORFs had identities lower than 50% with available proteins (Table 2). Two genes for S88 O-antigen unit processing, potentially encoding the O-antigen flippase (*wzx*) and the O-antigen polymerase (*wzy*), were presumptively identified. The putative Wzx (ORF 2134) had 12 predicted transmembrane segments, which is the typical number for O-antigen flippase (36). Moreover, it also showed




FIG. 1. Genetic organization and comparison of the O-antigen gene clusters of *E. coli* S88 (upper row) and 96-3285 (lower row). Putative orthologous genes of the O-antigen gene clusters are connected. Values below the ORF numbers are % G+C content; values below the S88 genes are percent amino acid identity/percent amino acid similarity. NH, no homology.

23% identity and 42% similarity to the putative Wzx of *E. coli* 96-3285. The ORF 2132 (putative wzy) had a predicted amino acid sequence corresponding to eight transmembrane segments and a large cytoplasmic loop of 60 amino acids compatible with an O-antigen polymerase function (38). However, this protein had no homology with previously published *E. coli* O-antigen polymerase sequences, and its function is therefore highly hypothetical. Among the three remaining genes, two putative transferases (ORFs 2131 and 2133) were identified; the putative function of the third gene (ORF 2135) could not be assigned (Table 2).


The maps of the O-antigen gene clusters of S88 and *E. coli* 96-3285, based on protein homologies, were highly similar, except for ORF 2135 (Fig. 1). For example, the *rmlABC* and *tll* genes (assuming that *wbhT* corresponds to *tll*) were in the same order. Moreover, *rmlB* and *rmlA* were identically situated at both extremities of each locus, a very unusual disposition for these two genes (38). The only important difference was the presence of four genes located between *rmlB* and *rmlC* in the 96-3285 O-antigen gene cluster but not in S88 (Fig. 1).

Phylogenetic analysis of genes lying within or flanking the O-antigen gene cluster. In order to determine the genetic relatedness of the S88 and 96-3285 O-antigen gene clusters, phylogenetic trees were constructed with GenBank DNA sequences of the *rmlABC* genes from *E. coli* and other genera. These three genes were chosen for their strong similarities at the protein level between *E. coli* strains S88 and 96-3285. Trees obtained with the neighbor-joining method are shown in Fig. 2. Similar topologies were obtained with the maximum parsimony method (data not shown). In the *rmlA* tree (Fig. 2A), a cluster containing only *E. coli* and *Shigella* strains clearly emerged. Of note, all these strains harbored the four *rmlABCD* genes required for L-rhamnose synthesis. All the other *E. coli* *rmlA* genes, including *rmlA* in strains S88 and 96-3285, which are not associated with a complete *rmlABCD* cluster, were distantly related to each other, with low DNA homologies. Indeed, the *rmlA* genes of S88 and 96-3285 were not significantly closer to each other than to the *rmlA* genes of different species such as *Yersinia* spp. (Fig. 2A). In the *rmlB* tree, S88 was again distantly related to the main group of *E. coli* strains, despite a mean identity of 81% with this group. Finally, in the *rmlC* tree, three major groups of genes with very low DNA homologies were distinguished. One of them contained *E. coli* S88, 96-3285, and the O66 reference strain, in which *Vibrio cholerae* and *Aeromonas hydrophila* are nearly equidistant.

As the immediate flanking gene *gnd* is potentially a target

for horizontal transfer of the O-antigen gene cluster (30), we investigated the genetic relationship of this gene between several *E. coli* strains and other *Enterobacteriaceae* (Fig. 2D). Most of the *E. coli* *gnd* sequences, including the *gnd* sequence of strain 96-3285, clustered together, with an average identity of 95%. In contrast, S88 *gnd* was only distantly related to this cluster (mean identity, 85%). As previously described, *gnd* sequences from *E. coli* ECOR-16, ECOR-4, and ECOR-71 were more closely related to *Klebsiella* spp. or *Citrobacter* spp., from which they are thought to have been acquired (30). However, S88 *gnd* could not be linked to any of these species or to other *Enterobacteriaceae* *gnd* sequences available in the GenBank database.

Functional analysis of the O-antigen gene cluster. In order to determine the functional role of the putative O-antigen gene cluster and the putative O-antigen polymerase gene (*wzy*) in *E. coli* strain S88, two mutants were prepared according to the method described by Datsenko and Wanner (29). One mutant, named CelP3, had a deletion of the entire O-antigen gene cluster from the first gene (ORF 2138) to the last (ORF 2130), and the other mutant, named CelP7, had a deletion of the putative *wzy* gene. With both mutants, PCR using primers flanking the initial and final portions of the DNA target segment and primers homologous to the *cat* gene (29) demonstrated the introduction of the *cat* gene instead of the target genes (data not shown). Moreover, multiplex PCR amplification of the main extraintestinal virulence genes and capsular antigen K1 were positive for each mutant, as for the wild-type strain (data not shown). Mutants CelP3 and CelP7 did not agglutinate with O45-specific antiserum, in contrast to strain S88, suggesting the involvement of the two deleted loci in the biosynthesis of the polysaccharide somatic O antigen. We then analyzed the polysaccharide somatic O antigen in the two mutants and the wild-type strain by SDS-polyacrylamide gel electrophoresis (Fig. 3). The S88 polysaccharide somatic O antigen showed a wild-type bimodal distribution of LPS, characterized by a first band of lipid A-core and several more bands that corresponded to O-antigen chain subunits. The CelP3 mutant had a rough phenotype characterized by only one band of lipid A-core and no attached O-antigen chain, demonstrating that the DNA segment located between *galF* and *gnd* is effectively involved in the biosynthesis of the somatic O antigen in strain S88 and therefore corresponds to the O-antigen gene cluster. The CelP7 mutant had a semirough phenotype with only one O-antigen subunit substitution on the core oligosaccharide (Fig. 3). This result indicates that the gene that we presump-






FIG. 3. Visualization of LPS from strains 96-3285 and S88 and mutants by SDS-polyacrylamide gel electrophoresis. Lane 1, 96-3285 (CDC reference strain O45); lane 2, S88 (wild type with long-chain LPS [S LPS]); lane 3, CelP7 (S88 with a deletion of *wzy* and the core replaced by one O unit [SR LPS]); and lane 4, CelP3 (S88 with a deletion of the O-antigen gene cluster and complete loss of O chains [R LPS]).

tively identified as O-antigen polymerase (*wzy*) effectively encodes a polymerase responsible for the assembly of the oligosaccharide subunits.

O somatic antigen has a major role in the virulence of S88. O antigen has been recognized as a potential virulence factor in ExPEC infection. Indeed, somatic antigen O18 is known to play a key role in the virulence of the archetypal O18:K1:H7 meningitis clone. Thus, in order to investigate the pathogenic role of the S88 somatic antigen, we assessed the virulence of the CelP3 and CelP7 mutants, relative to S88, in two different experiments with a neonatal rat meningitis model. For comparison, strain 96-3285 was also assessed in our model. In the first experiment the average bacteremia observed with mutant CelP3 (2.94 ± 0.04 log CFU/ml) was far lower ($P < 0.0001$) than that observed with S88 (6.15 ± 0.89 log CFU/ml) (Table 3). No cases of meningitis were observed with the mutant, whereas five cases occurred with the wild-type strain. Similar results were obtained in the second experiment, in which the average bacteremia observed with mutant CelP7 (2.91 ± 0.04

TABLE 3. Mean bacterial counts in blood and CSF culture positivity in a neonatal rat model of meningitis with S88 and its mutants

Expt. and strain	Description	No. of infected animals	Mean bacteremia (log CFU/ml [SD])	No. of culture-positive CSF specimens
Expt. no. 1				
S88	S88 wild type	15	6.15 (0.89)	5
CelP3	S88 (O-antigen gene cluster with <i>cat</i> gene insertion)	15	2.94 (0.04) ^a	0
Expt. no. 2				
S88	S88 wild type	10	6.23 (1.68)	3
CelP7	S88 (<i>wzy::cat</i>)	10	2.91 (0.01) ^b	0

^a $P = 10^{-9}$ versus S88 bacterial count in experiment 1.

^b $P = 10^{-4}$ versus S88 bacterial count in experiment 2.

log CFU/ml) was lower ($P < 0.0001$) than that observed with S88 (6.23 ± 1.68 log CFU/ml). Again, the only three cases of meningitis occurred with S88. Therefore, each mutant produced a level of bacteremia at least 3 logs lower than the wild-type strain S88. These results suggest that, in this model, the O antigen plays a major role in producing sustained high-level bacteremia. As expected 96-3285 was unable to induce any bacteremia in our model even when the inoculum was increased 100-fold (data not shown).

O-serogroup PCR assay. In order to develop an accurate typing system able to specifically recognize strains harboring the somatic antigen identical to that of strain S88, we designed primers to specifically amplify the *wzx* and *wzy* genes, which are usually O antigen specific. The two genes were amplified simultaneously with the *uidA* gene, which serves as an internal control, resulting in a triplex PCR assay (Fig. 4). First, we assessed the efficiency of our PCR assay with 44 clinical strains of *E. coli* O45:K1:H7 closely related to S88 (12). The 44 strains were PCR positive with both primer pairs. Specificity was then evaluated using as a template DNA extracted from 130 O reference strains, including the reference strains 96-3285 and H61 for O45 somatic antigen, and DNA extracted from the 72 ECOR reference strains. None of the 204 PCR assays was

FIG. 2. Phylogenetic trees generated by the neighbor-joining method for *rmlA*, *rmlB*, *rmlC*, and *gnd* sequences of several representative gram-negative species. Numbers at the branches are bootstrap proportions obtained from 100 replicates. *, *E. coli* or *Shigella* strain harboring a complete *rmlABCD* cluster. Sequences were extracted from the GenBank database, except for the *gnd* sequences of strains 96-3285 and S88. The following *E. coli* serogroups are represented: O1 (strain APEC01; accession number CP000468), O6 (strain CFT073; accession number NC_004431), O7 (strain VW187; accession number AF125322), O9 (strain ECOR-16; accession number M64325), O16 (strain K-12, MG1655; accession number U00096), O18 (strain UTI89; accession number CP000243), O26 (accession number AF529080), O45 (strain 96-3285; accession number AY771223), O52 (strain G1066; accession number AY528413), O66 (accession number DQ069297), O78 (strain ECOR-71; accession number U14461), O91 (strain ECA95; accession number AY035396), O114 (strain G1088; accession number AY573377), and O157 (strain EDL933; accession number NC_002655), and O nontypeable (strain ECOR-4; accession number M64324). Other strains include the following: *Escherichia vulneris* (strain ATCC 33821; accession number U14465), *Shigella boydii* (strain sb227; accession number CP000036), *Shigella dysenteriae* (strain sd197; accession number CP000034), *Shigella flexneri* 2a (strain 301; accession number AE005674), *Salmonella enterica* serovar Typhi (strain CT18; accession number AL627273), *S. enterica* serovar Typhimurium (strain LT2; accession number NC_003197), *Citrobacter koseri* (strain CT19; accession number U14424 and strain CT42; acc. number U14428), *Citrobacter freundii* (strain ATCC 8090; accession number U14466), *Citrobacter amalonaticus* (strain CT28; accession number U14426), *Klebsiella pneumoniae* (strain Chedit *gnd* gene [accession number D21242] and strain KT776 *rml* genes [accession number AF097519]), *Raoultella terrigena* (strain ATCC 33257; accession number AY376146), *Erwinia carotovora* (strain SCRI1043; accession number NC_004547), *Yersinia enterocolitica* serogroup O:3 (strain 6471/76; accession number Z18920), *Vibrio cholerae* (strain S6535; accession number AY239000), *Vibrio fischeri* (strain ES114; accession number CP000020), and *Aeromonas hydrophila* (strain PPD134/91; accession number AF148126).




FIG. 4. O-serogroup PCR assay using *wzx* (119 bp), *wzy* (330 bp), and *uidA* (186 bp) primers. Lane 1, 100-bp molecular weight ladder; lane 2, 96-3285 (CDC reference strain O45); lane 3, H61 (Staten Serum Institute strain O45); lane 4, S88 (wild type); lane 5, clinical strain of *E. coli* O45:K1:H7, closely related to S88.

positive. These results showed the efficiency and specificity of our PCR assay.

DISCUSSION

We recently described the emergence in France of a new highly virulent group B2 *E. coli* neonatal meningitis clone of serotype O45:K1:H7. While capsular antigen K1 and flagellar antigen H7 are common among *E. coli* strains causing neonatal meningitis (1), antigen O45 is rare and, intriguingly, is mainly shared by InPEC strains such as enterotoxigenic *E. coli* and STEC strains (4, 9, 32, 41) which do not usually belong to group B2 (18). One possible explanation for the expression of the same antigen by such different *E. coli* pathotypes belonging to different phylogenetic groups is horizontal transfer of the O-antigen gene cluster between these clones. The transferability of the O-antigen gene cluster has been demonstrated, particularly for the locus encoding the O157 antigen of the worldwide STEC O157:H7 clone (20). Moreover, older and recent phylogenetic analyses (1, 7, 24) showing the existence of common serogroups among distant clonal groups of ExPEC strains suggest that this transferability is shared by many other O-antigen gene clusters. In our study, comparing the published O-antigen gene cluster sequence of the O45 CDC reference strain and the O45-antigen gene cluster sequence obtained through the ongoing S88 sequencing project (www.genoscope.fr), we found that although the two loci shared some similarities, the serogroup identity cannot be explained by simple horizontal transfer of the locus between these two strains.

It is therefore likely that S88 expresses an O polysaccharide related but not identical to O45. The cross-reaction could be due in part to the probable presence of 6-deoxy-L-talose. This sugar is the product of the four genes *rmlABC* and *tll* (38), which we presumptively identified in the O-antigen gene cluster of strain S88. Moreover, 6-deoxy-L-talose is known to be present in O45 and O66 polysaccharides, for which antisera may cross-react (33). Until the structure of the S88 O polysaccharide is elucidated and specific antibodies are available, we propose to name its O serogroup O45_{S88}.

Although not identical, the S88 and 96-3285 O-antigen gene clusters shared a high degree of similarity with respect to the physical map of the loci. *rmlA* and *rmlB* are important in the nucleotide sugar biosynthesis pathways converting glucose-1-phosphate to the dTDP-6-deoxy-D-xylo-4-hexulose intermediate, which is a branch point for several pathways (38). These two genes mainly have two types of disposition. In several *E.*

coli strains and also in *Salmonella* spp. and *Shigella* spp., they are part of the four genes *rmlABCD* clustering at the 5' end of the O-antigen cluster in the order *rmlBDAC*. In other *E. coli* strains and in several *Enterobacteriaceae* and non-*Enterobacteriaceae*, the *rmlAB* genes are present while *rmlCD* genes are absent, and *rmlAB* genes cluster together at the 5' end of the O-antigen gene cluster in the order *rmlBA*. We exhaustively inspected all *rmlAB* genes available in the GenBank database and found no such scattered disposition. Therefore, the separate positions of *rmlA* and *rmlB*, each at one extremity of the locus in strains S88 and 96-3285, is unique. This result and the similar global disposition of the orthologous genes between the two loci, point to a common ancestor for the two O-antigen gene clusters. This common ancestor would have given rise to two different O-antigen gene clusters via several genetic events, such as deletion or acquisition of the four genes lying between *rmlB* and *rmlC* (Fig. 1).

In order to gain insight into the genetic evolution of the O45_{S88} gene cluster, we constructed phylogenetic trees of *rmlABC* DNA sequences. In the three trees, S88 and 96-3285 were distantly related to the main *E. coli* and *Shigella* group. Moreover, the two strains shared a low level of genetic identity, and their *rmlABC* genes did not appear to be more closely related to each other than to the *rmlABC* genes of other species. All these results suggest that although the global organization of the two loci encoding the O-antigen process is similar in S88 and 96-3285, none of the genes was recently exchanged between these strains. In line with the hypothesis postulating a common ancestor for the two loci, the large genetic distance observed between the orthologous genes may be explained by multiple horizontal gene transfers from different species and/or by multiple mutations during a long period of evolution.

Horizontal transfer of O-antigen gene clusters generally involves the flanking conserved genes and, notably, *gnd*, which serves as a target for recombination. Nelson and Selander sequenced *gnd* in several *E. coli* strains and other *Enterobacteriaceae* and found that *gnd* in certain *E. coli* strains had been imported from *Klebsiella* spp. and *Citrobacter* spp. (30). In order to determine whether the O45_{S88} gene cluster was acquired in part from other species, we compared the *gnd* sequences of *E. coli* strains S88 and 96-3285 and of several other *Enterobacteriaceae*. *gnd* in S88 was distantly related to *gnd* in other *E. coli* species but unrelated to the gene in previously identified donor genera such as *Citrobacter* and *Klebsiella*. These results indicate that the O45_{S88} gene cluster may, at least in part (including the *rmlA* gene), have been transferred from another, unidentified species. To confirm this it would be necessary to find a common donor for *gnd* and *rmlA*. In order to obtain more evidence of horizontal transfer, we also analyzed the genetic relationship of *galF*, the opposite flanking gene, and the JUMPStart sequence, a conserved sequence just upstream of the operon encoding many polysaccharides. However, *galF* and the JUMPStart sequences of S88 were strongly homologous to the sequences of other *E. coli* strains (data not shown) and could not, therefore, be used to support our hypothesis.

O polysaccharides contribute to the pathogenicity of ExPEC strains. This virulence factor belongs to the so-called protectin class, which plays an important role in protection against complement-mediated lysis and phagocytosis. Indeed, the O18 an-

tigen plays a key role in virulence of the global meningitis clonal group O18:K1:H7. Pluschke et al., using *E. coli* mutants lacking somatic antigen, showed that polysaccharide O18 is involved in resistance to the classical complement pathway in guinea pig serum (35). In vivo, the same authors as well as Kim et al. demonstrated the crucial role of O18 in sustained high-level bacteremia, which is necessary for blood-brain barrier penetration (25, 35). However, several reports suggest that not all O polysaccharides have identical virulence properties. When exploring the virulence of the avian pathogenic *E. coli* O78:K80:H9, Mellata et al. constructed a mutant lacking antigen O78 and two derivative strains supplemented with antigen O1 or O26 (28). They found that the loss of O78 was associated with lower pathogenicity but that substitution by O1 or O26 did not fully restore the initial virulence. More strikingly, Russo et al. prepared an isogenic O-antigen-deficient mutant from a human blood isolate named CP9 (O4:K54:H5) and unexpectedly observed a slight increase in virulence compared with the parental strain (37). All these results underline the unpredictable nature of O-antigen involvement in ExPEC virulence. We evaluated the role of the somatic antigen in the virulence of the O45:K1:H7 clone in an experimental meningitis model and found that antigen O45 was crucial for the bacteremic step. Although we did not restore O45 polysaccharide expression in our mutants and thereby confirm the involvement of this factor in extraintestinal pathogenicity, we obtained the same results, namely, a complete loss of virulence in our experimental model, with two different mutants in two separate experiments. Thus, the implication of other, uncontrolled genetic events in the loss of virulence of our mutants is very unlikely. Therefore, the new somatic antigen O45_{S88} may in part account for the emergence of the French meningitis clone.

Our work underlined the limits of O serotyping by agglutination, owing to the risk of cross-reactions. Numerous O-antigen gene clusters have been sequenced in recent years, and PCR assays targeting the O-antigen unit processing genes, the O-antigen flippase (*wzx*) and the O-antigen polymerase (*wzy*), have been developed (5, 6, 13, 16, 17, 19, 44, 45). This O-genotyping approach has several advantages, including low cost, rapidity, and detection of strains with the O antigen of interest in a complex microflora. The PCR assay we developed here will allow strains harboring the O45_{S88} antigen to be distinguished from those expressing the reference O45 antigen. This method may also facilitate the search for the ecological niche and the mode of human acquisition of this antigen. Further studies will be necessary to understand the origin of this new somatic antigen, which could serve as a vaccine target for neonatal meningitis prevention. Meanwhile, our PCR test could be used to detect the highly virulent O45:K1:H7 clone in the microflora of neonates or even in the mother's vaginal flora, as is already the case for group B streptococci.

ACKNOWLEDGMENTS

We thank Vanida Vongsouthi for technical assistance with LPS analysis and Olivier Clermont for finishing the sequencing of S88.

This work was supported by Fondation pour la Recherche Médicale grant 20050904666.

REFERENCES

- Achtman, M., A. Mercer, B. Kusecek, A. Pohl, M. Heuzenroeder, W. Aaronson, A. Sutton, and R. P. Silver. 1983. Six widespread bacterial clones among *Escherichia coli* K1 isolates. *Infect. Immun.* **39**:315–335.
- Apweiler, R., T. K. Attwood, A. Bairoch, A. Bateman, E. Birney, M. Biswas, P. Bucher, L. Cerutti, F. Corpet, M. D. Croning, R. Durbin, L. Falquet, W. Fleischmann, J. Gouzy, H. Hermjakob, N. Hulo, I. Jonassen, D. Kahn, A. Kanapin, Y. Karavidopoulou, R. Lopez, B. Marx, N. J. Mulder, T. M. Oinn, M. Pagni, F. Servant, C. J. Sigrist, and E. M. Zdobnov. 2001. The InterPro database, an integrated documentation resource for protein families, domains and functional sites. *Nucleic Acids Res.* **29**:37–40.
- Barbe, V., D. Vallenet, N. Fonknechten, A. Kreimeyer, S. Oztas, L. Labarre, S. Cruveiller, C. Robert, S. Duprat, P. Wincker, L. N. Ornston, J. Weissenbach, P. Marliere, G. N. Cohen, and C. Medigue. 2004. Unique features revealed by the genome sequence of *Acinetobacter* sp. ADP1, a versatile and naturally transformation competent bacterium. *Nucleic Acids Res.* **32**:5766–5779.
- Batissou, I., M. P. Guimond, F. Girard, H. An, C. Zhu, E. Oswald, J. M. Fairbrother, M. Jacques, and J. Harel. 2003. Characterization of the novel factor *paa* involved in the early steps of the adhesion mechanism of attaching and effacing *Escherichia coli*. *Infect. Immun.* **71**:4516–4525.
- Beutin, L., Q. Kong, L. Feng, Q. Wang, G. Krause, L. Leomil, Q. Jin, and L. Wang. 2005. Development of PCR assays targeting the genes involved in synthesis and assembly of the new *Escherichia coli* O174 and O177 O antigens. *J. Clin. Microbiol.* **43**:5143–5149.
- Beutin, L., J. Tao, L. Feng, G. Krause, S. Zimmermann, K. Gleier, Q. Xia, and L. Wang. 2005. Sequence analysis of the *Escherichia coli* O15 antigen gene cluster and development of a PCR assay for rapid detection of intestinal and extraintestinal pathogenic *E. coli* O15 strains. *J. Clin. Microbiol.* **43**:703–710.
- Bidet, P., F. Mahjoub-Messai, J. Blanco, J. Blanco, M. Dehem, Y. Aujard, E. Bingen, and S. Bonacorsi. 2007. Combined multilocus sequence typing and O serotyping distinguishes *Escherichia coli* subtypes associated with infant urosepsis and/or meningitis. *J. Infect. Dis.* **196**:297–303.
- Bidet, P., A. Metais, F. Mahjoub-Messai, L. Durand, M. Dehem, Y. Aujard, E. Bingen, X. Nassif, and S. Bonacorsi. 2007. Detection and identification by PCR of a highly virulent phylogenetic subgroup among extraintestinal pathogenic *Escherichia coli* B2 strains. *Appl. Environ. Microbiol.* **73**:2373–2377.
- Blanco, J., M. Blanco, J. I. Garabal, and E. A. Gonzalez. 1991. Enterotoxins, colonization factors and serotypes of enterotoxigenic *Escherichia coli* from humans and animals. *Microbiologia* **7**:57–73.
- Bocs, S., S. Cruveiller, D. Vallenet, G. Nuel, and C. Medigue. 2003. AMIGene: annotation of microbial genes. *Nucleic Acids Res.* **31**:3723–3726.
- Bonacorsi, S., V. Houdouin, P. Mariani-Kurkdjian, F. Mahjoub-Messai, and E. Bingen. 2006. Comparative prevalence of virulence factors in *Escherichia coli* causing urinary tract infection in male infants with and without bacteremia. *J. Clin. Microbiol.* **44**:1156–1158.
- Bonacorsi, S. P., O. Clermont, V. Houdouin, C. Cordevant, N. Brahimi, A. Marecat, C. Tinsley, X. Nassif, M. Lange, and E. Bingen. 2003. Molecular analysis and experimental virulence of French and North American *Escherichia coli* neonatal meningitis isolates: identification of a new virulent clone. *J. Infect. Dis.* **187**:1895–1906.
- Cheng, J., B. Liu, D. A. Bastin, W. Han, L. Wang, and L. Feng. 2007. Genetic characterization of the *Escherichia coli* O66 antigen and functional identification of its *wzy* gene. *J. Microbiol.* **45**:69–74.
- Czirok, E., M. Herpay, and H. Milch. 1993. Computerized complex typing of *Escherichia coli* strains from different clinical materials. *Acta Microbiol. Hung.* **40**:217–237.
- Datsenko, K. A., and B. L. Wanner. 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* **97**:6640–6645.
- DebRoy, C., P. M. Fratamico, E. Roberts, M. A. Davis, and Y. Liu. 2005. Development of PCR assays targeting genes in O-antigen gene clusters for detection and identification of *Escherichia coli* O45 and O55 serogroups. *Appl. Environ. Microbiol.* **71**:4919–4924.
- DebRoy, C., E. Roberts, J. Kundrat, M. A. Davis, C. E. Briggs, and P. M. Fratamico. 2004. Detection of *Escherichia coli* serogroups O26 and O113 by PCR amplification of the *wzx* and *wzy* genes. *Appl. Environ. Microbiol.* **70**:1830–1832.
- Escobar-Paramo, P., O. Clermont, A. B. Blanc-Potard, H. Bui, C. Le Bouguenec, and E. Denamur. 2004. A specific genetic background is required for acquisition and expression of virulence factors in *Escherichia coli*. *Mol. Biol. Evol.* **21**:1085–1094.
- Feng, L., W. Wang, J. Tao, H. Guo, G. Krause, L. Beutin, and L. Wang. 2004. Identification of *Escherichia coli* O114 O-antigen gene cluster and development of an O114 serogroup-specific PCR assay. *J. Clin. Microbiol.* **42**:3799–3804.
- Feng, P., K. A. Lampel, H. Karch, and T. S. Whittam. 1998. Genotypic and phenotypic changes in the emergence of *Escherichia coli* O157:H7. *J. Infect. Dis.* **177**:1750–1753.

21. Houdouin, V., S. Bonacorsi, N. Brahimi, O. Clermont, X. Nassif, and E. Bingen. 2002. A uropathogenicity island contributes to the pathogenicity of *Escherichia coli* strains that cause neonatal meningitis. *Infect. Immun.* **70**:5865–5869.
22. Johnson, J. R. 2003. Microbial virulence determinants and the pathogenesis of urinary tract infection. *Infect. Dis. Clin. N. Am.* **17**:261–278, viii.
23. Johnson, J. R., E. Oswald, T. T. O'Bryan, M. A. Kuskowski, and L. Spanjaard. 2002. Phylogenetic distribution of virulence-associated genes among *Escherichia coli* isolates associated with neonatal bacterial meningitis in The Netherlands. *J. Infect. Dis.* **185**:774–784.
24. Johnson, J. R., K. L. Owens, C. R. Clabots, S. J. Weissman, and S. B. Cannon. 2006. Phylogenetic relationships among clonal groups of extraintestinal pathogenic *Escherichia coli* as assessed by multi-locus sequence analysis. *Microbes Infect.* **8**:1702–1713.
25. Kim, K. S., H. Itabashi, P. Gemski, J. Sadoff, R. L. Warren, and A. S. Cross. 1992. The K1 capsule is the critical determinant in the development of *Escherichia coli* meningitis in the rat. *J. Clin. Investig.* **90**:897–905.
26. Krogh, A., B. Larsson, G. von Heijne, and E. L. Sonnhammer. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**:567–580.
27. Kumar, S., K. Tamura, and M. Nei. 2004. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* **5**:150–163.
28. Mellata, M., M. Dho-Moulin, C. M. Dozois, R. Curtiss, 3rd, P. K. Brown, P. Arne, A. Bree, C. Desautels, and J. M. Fairbrother. 2003. Role of virulence factors in resistance of avian pathogenic *Escherichia coli* to serum and in pathogenicity. *Infect. Immun.* **71**:536–540.
29. Negre, V. L., S. Bonacorsi, S. Schubert, P. Bidet, X. Nassif, and E. Bingen. 2004. The siderophore receptor IroN, but not the high-pathogenicity island or the hemin receptor ChuA, contributes to the bacteremic step of *Escherichia coli* neonatal meningitis. *Infect. Immun.* **72**:1216–1220.
30. Nelson, K., and R. K. Selander. 1994. Intergeneric transfer and recombination of the 6-phosphogluconate dehydrogenase gene (*gnd*) in enteric bacteria. *Proc. Natl. Acad. Sci. USA* **91**:10227–10231.
31. Ochman, H., and R. K. Selander. 1984. Standard reference strains of *Escherichia coli* from natural populations. *J. Bacteriol.* **157**:690–693.
32. Orden, J. A., J. A. Ruiz-Santa-Quiteria, D. Cid, S. Garcia, and R. de la Fuente. 1999. Prevalence and characteristics of necrotogenic *Escherichia coli* (NTEC) strains isolated from diarrhoeic dairy calves. *Vet. Microbiol.* **66**:265–273.
33. Orskov, I., F. Orskov, B. Jann, and K. Jann. 1977. Serology, chemistry, and genetics of O and K antigens of *Escherichia coli*. *Bacteriol. Rev.* **41**:667–710.
34. Ott, M., L. Bender, G. Blum, M. Schmittroth, M. Achtman, H. Tschape, and J. Hacker. 1991. Virulence patterns and long-range genetic mapping of extraintestinal *Escherichia coli* K1, K5, and K100 isolates: use of pulsed-field gel electrophoresis. *Infect. Immun.* **59**:2664–2672.
35. Pluschke, G., J. Mayden, M. Achtman, and R. P. Levine. 1983. Role of the capsule and the O antigen in resistance of O18:K1 *Escherichia coli* to complement-mediated killing. *Infect. Immun.* **42**:907–913.
36. Reeves, P. R., M. Hobbs, M. A. Valvano, M. Skurnik, C. Whitfield, D. Coplin, N. Kido, J. Klena, D. Maskell, C. R. Raetz, and P. D. Rick. 1996. Bacterial polysaccharide synthesis and gene nomenclature. *Trends Microbiol.* **4**:495–503.
37. Russo, T. A., G. Sharma, C. R. Brown, and A. A. Campagnari. 1995. Loss of the O4 antigen moiety from the lipopolysaccharide of an extraintestinal isolate of *Escherichia coli* has only minor effects on serum sensitivity and virulence in vivo. *Infect. Immun.* **63**:1263–1269.
38. Samuel, G., and P. Reeves. 2003. Biosynthesis of O-antigens: genes and pathways involved in nucleotide sugar precursor synthesis and O-antigen assembly. *Carbohydr. Res.* **338**:2503–2519.
39. Sarff, L. D., G. H. McCracken, M. S. Schiffer, M. P. Glode, J. B. Robbins, I. Orskov, and F. Orskov. 1975. Epidemiology of *Escherichia coli* K1 in healthy and diseased newborns. *Lancet* **1**:1099–1104.
40. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
41. Toth, I., V. Karcagi, B. Nagy, I. Gado, and H. Milch. 1994. Examination of verocytotoxin producing capacity and determination of the presence of Shiga-like toxin genes in human *Escherichia coli* isolates. *Acta Microbiol. Immunol. Hung.* **41**:259–264.
42. Tsai, C. M., and C. E. Frasch. 1982. A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. *Anal. Biochem.* **119**:115–119.
43. Vallenet, D., L. Labarre, Z. Rouy, V. Barbe, S. Bocs, S. Cruveiller, A. Lajus, G. Pascal, C. Scarpelli, and C. Medigue. 2006. MaGe: a microbial genome annotation system supported by synteny results. *Nucleic Acids Res.* **34**:53–65.
44. Wang, L., H. Curd, W. Qu, and P. R. Reeves. 1998. Sequencing of *Escherichia coli* O111 O-antigen gene cluster and identification of O111-specific genes. *J. Clin. Microbiol.* **36**:3182–3187.
45. Wang, L., and P. R. Reeves. 1998. Organization of *Escherichia coli* O157 O antigen gene cluster and identification of its specific genes. *Infect. Immun.* **66**:3545–3551.
46. Wullenweber, M., L. Beutin, S. Zimmermann, and C. Jonas. 1993. Influence of some bacterial and host factors on colonization and invasiveness of *Escherichia coli* K1 in neonatal rats. *Infect. Immun.* **61**:2138–2144.