

Evolutionary Relationships Among γ -Carboxymuconolactone Decarboxylases

WU-KUANG YEH,* DON R. DURHAM,† PAUL FLETCHER,‡ AND L. NICHOLAS ORNSTON

Department of Biology, Yale University, New Haven, Connecticut 06511

Received 17 October 1980/Accepted 19 January 1981

γ -Carboxymuconolactone decarboxylase (EC 4.1.1.44) from *Azotobacter vinelandii* resembled the isofunctional enzymes from *Acinetobacter calcoaceticus* and *Pseudomonas putida*. All three decarboxylases appeared to be hexamers formed by association of identical subunits of about 13,300 daltons. The *A. vinelandii* and *P. putida* decarboxylases cross-reacted immunologically with each other, and the NH₂-terminal amino acid sequences of the enzymes differed in no more than 7 of the first 36 residues. In contrast, the *A. calcoaceticus* decarboxylase did not cross-react with the decarboxylase from *A. vinelandii* or *P. putida*; the NH₂-terminal amino acid sequences of these enzymes diverged about 50% from the NH₂-terminal amino acid sequence of the *A. calcoaceticus* decarboxylase.

Immunological comparisons conducted with protocatechuate oxygenase (EC 1.13.11.3) (2) and γ -carboxymuconolactone decarboxylase (EC 4.1.1.44) (1), enzymes of the protocatechuate branch of the β -ketoacid pathway, indicated that the *Azotobacter* species enzymes are evolutionarily homologous to isofunctional enzymes formed by fluorescent *Pseudomonas* species. As described here and elsewhere (1, 10, 13), isofunctional enzymes formed by members of other bacterial genera appear to be immunologically distant, and these organisms govern expression of the enzymes with induction patterns unlike those shared by *Azotobacter* and *Pseudomonas* species (12). The conclusions drawn from immunological evidence are supported by chemical and physical data presented here. The results show that the γ -carboxymuconolactone decarboxylases of *Azotobacter* and *Pseudomonas* species are closely related to each other and more distantly related to the evolutionarily homologous γ -carboxymuconolactone decarboxylase of *Acinetobacter* species.

MATERIALS AND METHODS

Bacterial strains and growth conditions. *Azotobacter vinelandii* OP was obtained from Paul E. Bishop, North Carolina State University, Raleigh, N.C. Large-scale growth of *A. vinelandii* was accomplished at 30°C in a New Brunswick Fermacell CF-130 fermentor containing 100 liters of modified Burk medium (14). *Pseudomonas putida* PRS 2260 (4) and

Acinetobacter calcoaceticus ADP 152 (17) were described recently. For immunodiffusion studies, the appropriate strains of *A. vinelandii*, *P. putida*, and *A. calcoaceticus* were grown in 500-ml Erlenmeyer flasks containing 150 ml of mineral medium (5, 14) supplemented with 10 mM p -hydroxybenzoate.

Purification of γ -carboxymuconolactone decarboxylase from *A. vinelandii*. All purification procedures were performed between 0 to 4°C. Buffer A was 20 mM Tris-hydrochloride (pH 7.5) containing 25 μ M dithiothreitol; buffer B was 10 mM Na₂HPO₄-KH₂PO₄ (pH 7.0); and buffer C was 20 mM Tris-hydrochloride containing 0.1 M NaCl. Cell pastes with a wet weight of 200 g were suspended in three volumes of buffer A and disrupted by passage through an American Instrument continuous-flow French pressure cell at 12,000 lb/in². Unbroken cells and debris were removed by centrifugation at 40,000 $\times g$ for 20 min. The resultant crude extract (Table 1 step 1) was brought to 30% saturation by the addition of ammonium sulfate followed by centrifugation at 40,000 $\times g$ for 20 min. Ammonium sulfate treatment was repeated on the supernatant fraction until 75% saturation was reached. After centrifugation, the protein pellet was dissolved in buffer A (Table 1, step 2) and dialyzed against three changes of this buffer over 48 h. The dialysis was applied onto a DEAE-cellulose column (5 by 30 cm) previously equilibrated with buffer A. The column was washed with three volumes of the same buffer, after which a continuous linear gradient, constructed from 0 to 0.3 M NaCl in buffer A in a total volume of 6 liters, was applied. Fractions of 15 ml were collected at a flow rate of 60 ml/h, and those containing γ -carboxymuconolactone decarboxylase activity, which eluted between 0.16 to 0.19 M NaCl, were pooled (Table 1, step 3). The DEAE-cellulose eluate was treated with ammonium sulfate, and the protein pellet that precipitated between 45 to 75% saturation was dissolved in buffer B (Table 1, step 4) and dialyzed thoroughly against the same buffer. The dialysate was

† Present address: Department of Microbiology, Medical College of Virginia, Virginia Commonwealth University, Richmond, VA 23298.

‡ Present address: Department of Microbiology, East Carolina University, Greenville, NC 27834.

TABLE 1. Purification of γ -carboxyconolactone decarboxylase from *A. vinelandii*

Step	Volume (ml)	Total activity (U)	Total protein (mg)	Sp act (U/mg)	Yield (%)	Purification (fold)
1. Crude extract	610	34,000	17,670	1.92	100	1.0
2. First 30 to 75% saturated ammonium sulfate fraction	310	30,130	8,290	3.63	89	1.9
3. First DEAE-cellulose eluate	500	32,000	1,575	20.3	94	10.6
4. Second 45 to 75% saturated ammonium sulfate fraction	33	31,350	726	43.2	92	22.5
5. Second DEAE-cellulose eluate	120	26,400	356	74.2	78	38.6
6. Third 40 to 65% saturated ammonium sulfate fraction	7	23,920	248	96.5	70	50.3
7. Sephadex G-200 eluate	32	19,840	30.4	653	58	340
8. Sephadex G-100 eluate	35	12,600	10.0	1,260	37	656
9. Quaternary aminoethyl-Sephadex eluate	12	6,040	4.8	1,258	18	655

applied onto a second DEAE-cellulose column (1.6 by 20 cm) previously equilibrated with buffer B. The column was washed with 2 volumes each of 0.01, 0.05, and 0.1 M sodium potassium phosphate buffer, and then a continuous linear gradient, constructed from 0.1 M sodium potassium phosphate buffer to the same buffer containing 0.3 M NaCl in a total volume of 300 ml, was applied. Fractions of 3 ml were collected at a flow rate of 20 ml/h, and those containing the decarboxylase activity were pooled (Table 1, step 5) and fractionated with ammonium sulfate. The protein pellet that precipitated between 40 to 65% saturation was dissolved in buffer A (Table 1, step 6) and applied onto a Sephadex G-200 column (2.5 by 100 cm) previously equilibrated with buffer A. Fractions of 3 ml were collected at a flow rate of 14 ml/h, and those containing a specific activity of the decarboxylase greater than 600 U/mg were combined (Table 1, step 7) and concentrated with ammonium sulfate (0 to 70% saturation). The resultant pellet was dissolved in buffer A and loaded onto a Sephadex G-100 column (2.5 by 40 cm) previously equilibrated with buffer A. The eluate containing the decarboxylase activity (Table 1, step 8) was concentrated with ammonium sulfate (0 to 70% saturation), dialyzed thoroughly against buffer C, and applied onto a quaternary aminoethyl-Sephadex column (0.9 by 8 cm) previously equilibrated with buffer C. The column was washed with 10 volumes of the same buffer, and then a continuous linear gradient, constructed from 0.1 to 0.4 M NaCl in 20 mM Tris-hydrochloride (pH 7.2) in a total volume of 50 ml, was applied. Fractions of 2 ml were collected at a flow rate of 10 ml/h, and those containing the decarboxylase activity were pooled (Table 1, step 9). The quaternary aminoethyl-Sephadex eluate was stored at 4°C in the presence of ammonium sulfate at 70% saturation.

Polyacrylamide gel electrophoresis. Polyacrylamide gel electrophoresis was performed with 10% disc gels (9). The subunit size of *A. vinelandii* decarboxylase was estimated by sodium dodecyl sulfate-gel electrophoresis (15) with previously described standard proteins (17).

Amino acid analysis. The amino acid composition of *A. vinelandii* decarboxylase was determined by a previously described procedure (4) and analyzed with a computer program (8) for the minimum molecular weight, which is the subunit size, of the enzyme.

NH₂-terminal amino acid sequence determination. Previously published procedures (16) were used for determination of the NH₂-terminal amino acid sequence of *A. vinelandii* decarboxylase.

Serological studies. Antisera against *A. vinelandii*, *P. putida*, and *A. calcoaceticus* decarboxylases were prepared (2), and the method of Stanier et al. (13) was used to examine serological cross-reaction on Ouchterlony double-diffusion plates (7).

Chemicals. Chemicals were described previously (2, 17). Quaternary aminoethyl-Sephadex was obtained from Pharmacia Fine Chemicals.

RESULTS

Purity of *A. vinelandii* decarboxylase. The most purified preparation of *A. vinelandii* decarboxylase (Table 1, step 9) possessed a specific activity of 1,260 U/mg. When subjected to electrophoresis on 10% polyacrylamide gels, the decarboxylase preparation migrated as a major band and a slight minor band (Fig. 1). The minor band may have been an electrophoretically different form of the decarboxylase because, as described below, the preparation was immunologically homogeneous, and the NH₂-terminal amino acid sequence of the enzyme was determined without detectable contamination.

Molecular weight and subunit size determinations. The molecular weight of the *A. vinelandii* decarboxylase, close to the molecular weights of the *A. calcoaceticus* and *P. putida* decarboxylases, was estimated to be 85,500 by gel filtration (Fig. 1). The size of the *A. vinelandii* decarboxylase subunit was determined as 13,300 by sodium dodecyl sulfate-gel electrophoresis and as 13,460 by computer-aided analysis of the amino acid composition of the enzyme (8).

Serological properties. Antisera prepared against the purified *A. vinelandii* decarboxylase formed a single precipitin band when diffused against a crude extract of *A. vinelandii* cells in which the enzyme had been induced (Fig. 2). This band formed a spur with a precipitin band formed by a *P. putida* extract containing the




FIG. 1. Insert depicts a stained 10% polyacrylamide gel on which 50 µg of *A. vinelandii* decarboxylase (Table 1, step 9) had migrated electrophoretically. The graph shows data indicating the molecular weights of decarboxylases determined by filtration on a standardized Bio-Gel agarose A 1.5m column (2.6 by 100 cm).




FIG. 2. Double-diffusion plates showing the immunological cross-reactions of enzymes with *A. vinelandii* decarboxylase. The center wells received 0.2 ml (containing approximately 3 mg of protein) of antiserum against the *A. vinelandii* decarboxylase. The outer wells of the top plate received crude extracts of cells in which the decarboxylase had been induced by growth with *p*-hydroxybenzoate. The outer wells of the bottom plate received the indicated amounts of purified decarboxylases.

decarboxylase (Fig. 2). No precipitin band was formed with extracts of uninduced *A. vinelandii* and *P. putida* cells (not shown) or with *A. calcoaceticus* extract containing the enzyme (Fig. 2). Similar precipitin patterns were formed when purified decarboxylase was substituted for crude extract in the outer wells (Fig. 2). Thus, the *A. vinelandii* decarboxylase appears to be an immunologically homogeneous preparation that cross-reacts strongly with the *P. putida* decarboxylase and not at all with the *A. calcoaceticus* decarboxylase.




FIG. 3. Double-diffusion plates on which purified decarboxylases from different bacterial genera were diffused against antiserum prepared against *P. putida* decarboxylase (center well, top) and against antiserum prepared against *A. calcoaceticus* decarboxylase (center well, bottom).

The conclusion that the *A. vinelandii* and *P. putida* decarboxylases resemble each other more closely than they resemble the *A. calcoaceticus* decarboxylase is fortified by the immunodiffusion patterns shown in Fig. 3. Antisera prepared against the *P. putida* decarboxylase form a precipitin band with the *P. putida* and *A. vinelandii* decarboxylases, but not with the *A. calcoaceticus* decarboxylase (Fig. 3). Antisera prepared against the *A. calcoaceticus* decarboxylase formed a precipitin band with this enzyme, but not with the corresponding decarboxylase from *A. vinelandii* or *P. putida* (Fig. 3).

Amino acid composition. The amino acid composition of the *A. vinelandii* decarboxylase is shown in Table 2. Quantitative comparison of amino acid compositions may be achieved by measurement of SAQ, the sum of the square of the difference in mole fraction of each amino acid that can be readily quantitated in a protein

hydrolysate. In over 5,000 pairwise comparisons, Marchalonis and Weltman (3) found that an SAQ of less than 50 invariably reflected a structural similarity that was revealed by comparison

TABLE 2. Amino acid composition of *A. vinelandii* γ -carboxymuconolactone decarboxylase

Amino acid	Amino acid residues per 13,460 daltons ^a
Asx	11.26
Thr	5.95
Ser	6.10
Glx	15.14
Pro	4.95
Gly	9.86
Ala	13.17
Cys ^b	1.14
Val	9.04
Met	3.05
Ile	6.21
Leu	11.70
Tyr	2.63
Phe	3.66
His	4.12
Lys	4.76
Arg	7.91
Trp ^c	1.18

^a From computer-aided analysis of amino acid composition (8).

^b Estimated as cysteic acid after performic acid oxidation (4).

^c Determined after hydrolysis of a protein sample with 3 N mercaptoethanesulfonic acid (4).

of amino acid sequences. With the *A. vinelandii* decarboxylase as reference we found that SAQs for the amino acid compositions of the *A. calcoaceticus* and *P. putida* decarboxylases were 9 and 24, respectively.

NH₂-terminal amino acid sequence. Quantitative measurements for the first 49 cycles in the NH₂-terminal amino acid sequence determination of the *A. vinelandii* decarboxylase are shown in Table 3.

DISCUSSION

Relationships among γ -carboxymuconolactone decarboxylases from *A. calcoaceticus*, *A. vinelandii*, and *P. putida*. γ -Carboxymuconolactone decarboxylases from the three bacterial genera shared some properties: the enzymes appeared to be hexamers formed by association of identical subunits of about 13,300 daltons (11), and the amino acid compositions of the proteins were similar (17). On the other hand, the specific activity of the *A. calcoaceticus* decarboxylase (140 U/mg) was substantially lower than the specific activities of the *A. vinelandii* and *P. putida* decarboxylases (1,260 and 1,310 U/mg, respectively), and the latter two decarboxylases cross-reacted immunologically with each other, but not with the *A. calcoaceticus* decarboxylase. Thus, it appears that all three decarboxylases were derived from a com-

TABLE 3. Automated sequence analysis of *A. vinelandii* γ -carboxymuconolactone decarboxylase: identification of amino acid residues

Cycle	PTH ^a	Me ₃ Si	BH-ORG ^b	BH-AQU ^c	Residue
1	M (193) ^d	M (181)			Met
2		D (125)	D (73)		Asp
3		E (166)	E (68)		Glu
4			K (37)		Lys
5		E (125)	E (68)		Glu
6			R (4)	R (17)	Arg
7	Y (18)	Y (101)	Y (46)		Tyr
8		D (65)	D (47)		Asp
9	A (131)	A (115)	A (53)		Ala
10	G (129)	G (106)	G (66)		Gly
11	M (66)	M (82)			Met
12		Q (20)	E (49)		Gln
		E (36)			
13	V (208)	V (153)	V (33)		Val
14			R (3)	R (8)	Arg
15			R (3)	R (14)	Arg
16	A (114)	A (104)	A (47)		Ala
17	V (97)	V (75)	V (33)		Val
18			L (35)		Leu
19	G (38)	G (33)	G (26)		Gly
20		D (25)	D (31)		Asp
21	A (72)	A (85)	A (43)		Ala
22				H (7)	His
23	V (230)	V (145)	V (26)		Val
24		D (48)	D (25)		Asp
25			R (2)	R (7)	Arg
26					(Cys or Ser) ^e
27	LI ^f (181)	LI (119)	L (26)		Leu
28			K (10)		Lys
29		D (11)	D (22)		Asn ^g
30	LI (163)	LI (117)	L (25)		Leu
31			T (14)		Thr
32	P (39)		P (11)		Pro
33	F (32)	F (34)	F (14)		Phe
34		D (7)	D (20)		Asn ^g
35		E (46)	E (23)		Glu
36		E (42)	E (25)		Glu
37	F (32)	F (31)	F (11)		Phe
38		Q (5)	E (22)		Gln
		E (26)			
39		E (30)	E (24)		Glu
40	M (9)	M (14)			Met
41	LI (97)	LI (59)	I (11)		Ile
42			T (12)		Thr
43			R (1)	R (3)	Arg
44				H (3)	His
45		A (45)	A (19)		(Ala)
46					(Cys or Ser) ^e
47		G (16)	G (12)		Gly
48		D (10)	D (11)		Asp
49	LI (138)	LI (67)	I (8)		Ile

^a PTH, Phenylthiohydantoin.

^b BH-ORG, Back hydrolysis-extracted organic phase.

^c BH-AQU, Back hydrolysis-remaining aqueous phase.

^d Results indicate the single-letter amino acid designation and (within parentheses) the number of nanomoles recovered.

^e The only two amino acid residues that could not be identified by the procedures used in this sequence determination are cysteine and serine.

^f PTH-leucine and PTH-isoleucine were coeluted by the gas chromatographic technique.

^g PTH-asparagine was distinguished from PTH-asparatic acid by high-pressure liquid chromatography.

mon ancestor; the structural genes for the *A. vinelandii* and *P. putida* decarboxylases seem to have diverged recently relative to their divergence from the *A. calcoaceticus* decarboxylase. These conclusions are strengthened by comparison of the NH₂-terminal amino acid sequences of the decarboxylases. The data allow a three-way comparison of 34 of the first 36 residues. The *A. vinelandii* and *P. putida* sequences were identical in 83% of the compared positions (Fig.

4), whereas comparison of the *A. vinelandii* sequence with the *A. calcoaceticus* sequence reveals an identity of 47% (Fig. 5).

Early events in the evolution of γ -carboxymuconolactone decarboxylases and muconolactone isomerases. γ -Carboxymuconolactone decarboxylases and muconolactone isomerases, enzymes that mediate analogous biochemical reactions, appear to share a common ancestral gene. Alignment of the NH₂-terminal

<u><i>Asotobacter</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	MET	ASP	GLU	LYS	GLU	ARG	TYR	ASP	ALA	GLY	MET	GLN	VAL	ARG	ARG	ALA	VAL	LEU
<u><i>Pseudomonas</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	MET	ASP	GLU	LYS	GLN	ARG	TYR	ASP	ALA	GLY	MET	GLN	VAL	ARG	ARG	ALA	VAL	LEU
<u><i>Asotobacter</i></u> Decarboxylase	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
	GLY	ASP	ALA	HIS	VAL	ASP	ARG	CYS	LEU	LYS	ASN	LEU	THR	PRO	PHE	ASN	GLU	GLU
<u><i>Pseudomonas</i></u> Decarboxylase	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
	GLY	ASP	ALA	HIS	VAL	ASP	ARG	CYS	LEU	GLY	LYS	LEU	ASN	ASP	PHE	ASN	GLY	GLU

FIG. 4. Comparison of the NH₂-terminal amino acid sequences of *A. vinelandii* and *P. putida* γ -carboxymuconolactone decarboxylases. Identical residues are enclosed in boxes.

<u><i>Asotobacter</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	MET	ASP	GLU	LYS	GLU	ARG	TYR	ASP	ALA	GLY	MET	GLN	VAL	ARG	ARG	ALA	VAL	LEU
<u><i>Acinetobacter</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
	MET	ASN	ASP	GLU	GLN	ARG	TYR	LYS	GLN	GLY	LEU	GLU	VAL	ARG	THR	GLU	VAL	LEU
<u><i>Asotobacter</i></u> Decarboxylase	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
	GLY	ASP	ALA	HIS	VAL	ASP	ARG	CYS	LEU	LYS	ASN	LEU	THR	PRO	PHE	ASN	GLU	GLU
<u><i>Acinetobacter</i></u> Decarboxylase	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
	GLY	GLU	LYS	HIS	VAL	ASN	ARG	SER	LEU	GLU	ASN	LEU	ASP	ASP	PHE	ARG	GLN	ASP

FIG. 5. Comparison of the NH₂-terminal amino acid sequences of *A. vinelandii* and *A. calcoaceticus* γ -carboxymuconolactone decarboxylases. Identical residues are enclosed in boxes.

<u><i>Acinetobacter</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	
	MET	ASN	ASP	GLU	-Δ-	GLN	ARG	TYR	LYS	GLN	GLY	LEU	GLU
<u><i>Asotobacter</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	
	MET	-Δ-	ASP	GLU	LYS	GLU	ARG	TYR	ASP	ALA	GLY	MET	GLN
<u><i>Pseudomonas</i></u> Decarboxylase	1	2	3	4	5	6	7	8	9	10	11	12	
	MET	-Δ-	ASP	GLU	LYS	GLN	ARG	TYR	ASP	ALA	GLY	MET	GLN
<u><i>Acinetobacter</i></u> Isomerase	24	25	26	27	28	29	30	31	32	33	34	35	
	LYS	SER	VAL	GLU	LYS	ALA	-Δ-	TYR	SER	GLN	GLU	LEU	GLN
<u><i>Pseudomonas</i></u> Isomerase	24	25	26	27	28	29	30	31	32	33	34	35	
	LYS	ALA	ASP	GLU	LYS	GLU	-Δ-	LEU	ALA	GLN	ARG	LEU	GLN

FIG. 6. A portion of the NH₂-terminal amino acid sequence of γ -carboxymuconolactone decarboxylase appears to be conserved within the primary structure of muconolactone isomerase. Numbers indicate the positions of residues in the primary sequences of the proteins. Boxes enclose residues at positions where identical residues are found in the decarboxylase-isomerase comparison.

amino acids of the enzymes reveals sequence similarity suggesting a low overall homology (17). In addition, the NH₂-terminal amino acid sequence of the decarboxylase appears to be conserved within the primary structure of the muconolactone isomerase: the tetrapeptide extending from residues 2 through 5 in the *A. vinelandii* decarboxylase is represented at residues 26 through 29 in the amino acid sequence of the *P. putida* muconolactone isomerase (Fig. 6). This is consistent with the proposal that, as genes for the β -ketoacid pathway became established, oligonucleotide substitution mutations placed sequences coding for peptides in novel structural contexts (6, 18).

ACKNOWLEDGMENTS

We thank Gary Davis for his expert technical assistance.

This work was supported by grant PCM7724884 from the National Science Foundation and Public Health Service grants GM 21714 and GM 25487 from the National Institutes of Health.

LITERATURE CITED

- Durham, D. R., and L. N. Ornston. 1980. Homologous structural genes and similar induction patterns in *Azotobacter* spp. and *Pseudomonas* spp. *J. Bacteriol.* **143**: 834-840.
- Durham, D. R., L. A. Stirling, L. N. Ornston, and J. J. Perry. 1980. Intergeneric evolutionary homology revealed by the study of protocatechuate 3,4 dioxygenase from *Azotobacter vinelandii*. *Biochemistry* **19**:149-155.
- Marchalonis, J. J., and J. K. Weltman. 1971. Relatedness among proteins: a new method of estimation and its application to immunoglobulins. *Comp. Biochem. Physiol. B* **38**:609-625.
- McCorkle, G. M., W. K. Yeh, P. Fletcher, and L. N. Ornston. 1980. Repetitions in the NH₂-terminal amino acid sequence of β -ketoacid enol-lactone hydrolase from *Pseudomonas putida*. *J. Biol. Chem.* **255**:6335-6341.
- Ornston, L. N., and R. Y. Stanier. 1966. The conversion of catechol and protocatechuate to β -ketoacid by *Pseudomonas putida*. I. *Biochemistry. J. Biol. Chem.* **241**: 3776-3786.
- Ornston, L. N., and W. K. Yeh. 1979. Origins of metabolic diversity: evolutionary divergence by sequence repetition. *Proc. Natl. Acad. Sci. U.S.A.* **76**:3996-4000.
- Ouchterlony, O. 1953. Antigen-antibody reactions in gels; types of reactions in coordinated systems of diffusion. *Acta Pathol. Microbiol. Scand.* **32**:231-240.
- Ozawa, K., and S. Tanaka. 1968. Computer-aided calculation of amino acid composition of proteins. *Anal. Biochem.* **24**:270-280.
- Patel, R. N., R. B. Meagher, and L. N. Ornston. 1974. Relationships among enzymes of the β -ketoacid pathway. IV. Muconolactone isomerase from *Acinetobacter calcoaceticus* and *Pseudomonas putida*. *J. Biol. Chem.* **249**:7410-7419.
- Patel, R. N., and L. N. Ornston. 1976. Immunological comparison of enzymes of the β -ketoacid pathway. *Arch. Microbiol.* **110**:27-36.
- Parke, D. 1979. Structural comparison of γ -carboxymuconolactone decarboxylase and muconolactone isomerase from *Pseudomonas putida*. *Biochim. Biophys. Acta* **578**:145-154.
- Stanier, R. Y., and L. N. Ornston. 1973. The β -ketoacid pathway. *Adv. Microb. Physiol.* **9**:89-151.
- Stanier, R. Y., D. Wachter, C. Gasser, and A. C. Wilson. 1970. Comparative immunological studies of two *Pseudomonas* enzymes. *J. Bacteriol.* **102**:351-360.
- Strandberg, G. W., and P. W. Wilson. 1968. Formation of the nitrogen fixing enzyme system in *Azotobacter vinelandii*. *Can. J. Microbiol.* **14**:25-31.
- Weber, K., and M. Osborn. 1979. The reliability of molecular weight determinations by dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Biol. Chem.* **244**: 4406-4412.
- Yeh, W. K., G. Davis, P. Fletcher, and L. N. Ornston. 1978. Homologous amino acid sequences in enzymes mediating sequential metabolic reactions. *J. Biol. Chem.* **253**:4920-4923.
- Yeh, W. K., P. Fletcher, and L. N. Ornston. 1980. Homologies in the NH₂-terminal amino acid sequences of γ -carboxymuconolactone decarboxylases and muconolactone isomerases. *J. Biol. Chem.* **255**:6347-6354.
- Yeh, W. K., and L. N. Ornston. 1980. Origins of metabolic diversity: substitution of homologous sequences into genes for enzymes with different catalytic activities. *Proc. Natl. Acad. Sci. U.S.A.* **77**:5365-5369.