

Research

Open Access

## AhR transcriptional activity in serum of Inuits across Greenlandic districts

Manhai Long<sup>1</sup>, Bente Deutch<sup>2</sup> and Eva C Bonefeld-Jorgensen\*<sup>1</sup>

Address: <sup>1</sup>Unit of Cellular and Molecular Toxicology, Department of Environmental and Occupational Medicine, Institute of Public Health, University of Aarhus, Denmark and <sup>2</sup>Centre for Arctic Environmental Medicine, Department of Environmental and Occupational Medicine, Institute of Public Health, University of Aarhus, Denmark

Email: Manhai Long - ml@mil.au.dk; Bente Deutch - bd@mil.au.dk; Eva C Bonefeld-Jorgensen\* - ebj@mil.au.dk

\* Corresponding author

Published: 23 October 2007

Received: 13 December 2006

*Environmental Health* 2007, **6**:32 doi:10.1186/1476-069X-6-32

Accepted: 23 October 2007

This article is available from: <http://www.ehjournal.net/content/6/1/32>

© 2007 Long et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Human exposure to lipophilic persistent organic pollutants (POPs) including polychlorinated dibenzo-*p*-dioxins/furans (PCDDs/PCDFs), polychlorinated biphenyls (PCBs) and organochlorine pesticide is ubiquitous. The individual is exposed to a complex mixture of POPs being life-long beginning during critical developmental windows. Exposure to POPs elicits a number of species- and tissue-specific toxic responses, many of which involve the aryl hydrocarbon receptor (AhR). The aim of this study was to compare the actual level of integrated AhR transcriptional activity in the lipophilic serum fraction containing the actual POP mixture among Inuits from different districts in Greenland, and to evaluate whether the AhR transactivity is correlated to the bio-accumulated POPs and/or lifestyle factors.

**Methods:** The study included 357 serum samples from the Greenlandic districts: Nuuk and Sisimiut (South West Coast), Qaanaaq (North Coast) and Tasiilaq (East Coast). The bio-accumulated serum POPs were extracted by ethanol: hexane and clean-up on Florisil columns. Effects of the serum extract on the AhR transactivity was determined using the Hepa 1.12cR mouse hepatoma cell line carrying an AhR-luciferase reporter gene, and the data was evaluated for possible association to the serum levels of 14 PCB congeners, 10 organochlorine pesticide residues and/or lifestyle factors.

**Results:** In total 85% of the Inuit samples elicited agonistic AhR transactivity in a district dependent pattern. The median level of the AhR-TCDD equivalent (AhR-TEQ) of the separate genders was similar in the different districts. For the combined data the order of the median AhR-TEQ was Tasiilaq > Nuuk ≥ Sisimiut > Qaanaaq possibly being related to the different composition of POPs. In overall, the AhR transactivity was inversely correlated to the levels of sum POPs, age and/or intake of marine food.

**Conclusion:** i) We observed that the proportion of dioxin like (DL) compounds in the POP mixture was the dominating factor affecting the level of serum AhR transcriptional activity even at very high level of non DL-PCBs; ii) The inverse association between the integrated serum AhR transactivity and sum of POPs might be explained by the higher level of compounds antagonizing the AhR function probably due to selective POP bioaccumulation in the food chain.

## 1. Background

Being resistant to both biotic and abiotic degradation, most persistent organic pollutants (POPs) bioaccumulate and magnify in animals and humans [1,2]. The polychlorinated dibenzo-*p*-dioxins/furans (PCDDs/PCDFs), polychlorinated biphenyls (PCBs) and organochlorine pesticides are dominating among the POPs. Some POPs e.g. PCBs and persistent organochlorine pesticides, that still are released into the environment due to ongoing use in developing countries and improper storage or disposal in developed countries, can be transported to the Arctic regions by atmospheric and oceanic currents and have been found to bioaccumulate and biomagnify in the Arctic marine food web [3,4]. The traditional diet of the indigenous people in the Arctic largely depends on fish, seabirds and marine mammals being associated with extraordinary high POP exposure [5-10], which may possess a health risk [11]. Since 1994, organochlorines have been measured in fat and plasma samples taken from Inuits of Greenland indicating that the contaminant level in Greenland is very high [5,8-10,12]. It has been documented that exposure to POPs such as dioxins (e.g. 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, TCDD) and dioxin-like (DL) compounds including *non-ortho* and *mono-ortho* PCBs may cause a series of negative effects both in animals and humans including carcinogenicity [13], immunotoxicity and adverse effects on reproductive and neurobehaviour [14]. Most of the toxic and biological effects of dioxins and DL-compounds is mediated mainly through activation of the aryl hydrocarbon receptor (AhR), an intracellular ligand-dependent transcriptional factor expressed in most tissues of mammals [15]. Upon receptor-ligand binding and translocation into the nucleus, the complex with the AhR nuclear translocator (ARNT) binds to the DNA dioxin-responsive elements (DREs), causing induction of gene transcription, for instance, encoding for metabolic enzymes [16].

Since dioxins and DL-compounds exist as complex mixtures of various congeners in environmental and biologic samples, the concept of TEQ (TCDD toxic equivalent) has been introduced to simplify risk assessment and regulatory control [1,17]. The classical TEQs (WHO-TEQ) are calculated by multiplying the concentration of individual PCDDs/PCDFs/PCBs by their respective Toxic Equivalency Factors (TEFs), which correspond to the relative potency of the congener to generate AhR-mediated effects compared to TCDD, the most potent AhR ligand [1,17]. Studies have emphasized that assessment of the toxicological potential of a chemical mixture is much more complex than can be deduced by a calculated TEQ value [7,18,19]. There are several drawbacks using the TEF concept for risk assessment of mixtures of POPs such as expensive and time consuming gas chromatography mass spectrometry (GC-MS) determinations, small concentra-

tions of hardly detectable individual congeners, presence of compounds not routinely measured or unknown substances with AhR affinity, the lack of TEF values for several POPs, and possible antagonistic or synergistic interactions between POPs [2,19-21]. Therefore, there is a need of simple, inexpensive and specific assays that can be used to give an integrated risk assessment of dioxins and DL-compounds. Rapid and sensitive AhR-based screening bioassays are valuable to more fully characterize the ligand-binding specificity of the AhR, to identify AhR ligands and AhR interfering compounds present in the extracts of environmental and biologic samples. A number of bioassays for the determination of TEQ values have been developed [22]. The experts have agreed that the use of *in vitro* bioassays provides an useful tool as a pre-screening method for TEQs in environmental samples [1,23]. The *in vitro* AhR mediated chemical activated luciferase gene expression (CALUX) bioassay is a reporter-gene based cell bioassay using genetically modified cells responding to AhR ligands. The CALUX bioassay has proven to be a quick and sensitive assay to detect the AhR mediated potential of pure chemicals [2,20,21,24], extracts of environmental and biological matrices eliciting the biological integrated net TEQ value (CALUX-TEQ) of complex mixtures as found in sediment, pore water [25], bovine and human milk [19,26], human serum [27] and follicular fluid [28]. Compared to the calculated chemical-derived TEQ (WHO-TEQ), the CALUX based AhR-TEQ might be more biological relevant for the risk assessment of dioxins and DL-compounds [29].

As a part of the human health program of the "Arctic Monitoring and Assessment Program (AMAP)" [30], the aim of the present study was i) to compare the AhR transcriptional activity of the actual mixture of lipophilic POP of serum of subjects from different districts in Greenland and ii) to evaluate whether the AhR transcriptional activity is associated to the profile and level of POPs and/or lifestyle factors.

## 2. Methods

### 2.1. Subjects, sampling and POPs determination

The subjects and sampling methods have been described in detail elsewhere [10]. Briefly, the participants were of Inuit descent from Nuuk, Sisimiut (South West Greenland), Qaanaaq (North West Greenland) and Tasiilaq (East Greenland). The men data of Sisimiut was also a part of the EU project INUENDO. Venous blood samples and questionnaires about demographic and lifestyle factors were collected. The serum was prepared as described [10] and then stored at -80°C until analyzed.

As described [10] plasma was analyzed for POPs including cis-chlordane, trans-chlordane, oxychlordane, *p,p'*-DDE, *p,p'*-DDT, hexachlorobenzene (HCB), beta-Hex-

achlorocyclohexane ( $\beta$ -HCH), mirex, toxaphene 26, toxaphene 50 and 14 PCB congeners (CB28, CB52, CB99, CB101, **CB105**, **CB118**, CB128, CB138, CB153, **CB156**, CB170, CB180, CB183, CB187; in bold are DL-PCB) by gas chromatography (GC). The samples enrolled per district were enough for comparison of the POP levels across districts with a power of 80% in power calculation. Plasma lipids were measured using conventional enzymatic methods and total lipids were calculated as described [31]. The lipid adjusted POP data ( $\mu\text{g}/\text{kg}$  lipid) was accomplished by dividing the wet-weight concentration in plasma ( $\mu\text{g}/\text{l}$ ) by the individual samples lipid concentration and multiplying by 1000 to convert from gram to kilogram [10,31]. Because of high POP intercorrelation [10] (Bonefeld-Jorgensen and Long, submitted), the group-variables were used for evaluation of the data, namely, the sum of 14 PCB congeners ( $\Sigma\text{PCB}_{14}$ ), the sum of pesticides ( $\Sigma$  pesticide) and finally the sum of all the determined POPs ( $\Sigma$  POP).

The fatty acid profiles were determined in plasma phospholipids at the Biology Department, University of Guelph, Canada [32]. The n-3 polyunsaturated fatty acids were reported on the sum of C18:3, n-3, C20:4, n-3, C20:5, n-3, C22:5, n-3 and C22:6, n-3, and the n-6 fatty acids was the sum of C18:2, n-6, C18:3, n-6, C20:2, n-6, C20:3, n-6 and C20:4, n-6 [10]. The ratio between n-3 and n-6 is known to be a strong indicator of marine food intake and thus a good indicator of the relative consumption of traditional versus imported food [8,33].

## 2.2. AhR transcriptional activation assay

The serum was extracted with a mixture of ammonium sulphate/ethanol/hexane (1:1:3) and the lipid extract was concentrated and cleaned up on two Florisil columns to get the actual mixture of POPs [27,31]. The detailed descriptions of dissolving the extract samples and detecting serum AhR transcriptional activity has been described in detail elsewhere [27]. The stably transfected mouse hepatoma cell line Hepa1.12cR cells carrying the AhR-luciferase reporter gene was kindly provided by MS Denison (University of California, Davis, CA, USA). Similar with Ziccardi and co-workers [34], the optimal conditions in our assay for determination of AhR transcriptional activity of samples using Hepa1.12cR cell line were in 100  $\mu\text{l}$  medium containing 10% FCS with  $6 \times 10^4$  cells, final solvent concentration of 0.1–0.4% DMSO and 4 h of exposure. Further validation of our Hepa1.12cR bioassay was obtained by participating in two international programs of inter-laboratory comparisons of dioxin-like compounds (The second round of interlaboratory comparison of dioxin-like compounds in food using bioassays, 2004, Örebro University, Sweden, and Interlaboratory comparison of dioxins and dioxin-like compounds in food using CALUX bioassay, 2006, Scien-

tific Institute of Public Health, Belgium). The limit of detection was 64 fg TEQ/well. The Hepa1.12cR cells were exposed for 4 hours to the serum extract alone (termed AhRag) or to mimic physiological environment in the presence of half maximum effective concentration of TCDD (termed AhRcomp) [27]. The luciferase activity was determined using a luminometer (BMG LUMISTAR) and cell protein was accomplished using a fluorometer (Wallac 1420 Multilabel Counter, Perkin Elmer life science, FIN) as previously described [35]. The luciferase activity of samples was presented as relative light units per microgram protein (RLU/ $\mu\text{g}$  protein) [35]. The average coefficient of variation (CV) between replicates (intra CV) and between day-to-day independent assays (inter CV) of the solvent control was 9% and 17%, respectively and the intra CV of the test samples was 10%.

No cell toxicity on Hepa1.12cR cells was observed after exposure to the serum extract samples determined as described [35].

## 2.3. Calculation and statistical analysis

In the separate assays the activity differences between the triple serum extract determinations and their respective solvent controls [% agonist (samples with agonistic effect), % inhibition (samples with decrease of TCDD induced activity), and % increase (samples with further increase of TCDD induced activity)] were evaluated using the Student t-test with statistical significant level of  $p \leq 0.05$  (Microsoft Excel).

The AhR-TEQs values of serum extract were obtained by interpolation of the AhRag values onto the parallel TCDD dose-response sigmoid Hill curve and only the AhRag values being significantly higher than the solvent control and in the linear range of the TCDD dose-response curve were used to calculate AhR-TEQ [27].

Natural logarithmic transformation of AhR-TEQ/AhR-comp and POP data improved the normality (checked by Q-Q plots) and homogeneity of variance, and the statistical analysis was performed on the ln-transformed data. The comparisons of different variables (POPs, AhR-TEQ and AhRcomp) among the districts were performed with One-way ANOVA test. When ANOVA showed statistical significant differences complementary multiple comparison *ad hoc* tests was performed. Test for equal variances was performed using Levene's test. The least-significant difference (LSD) test was used if the variables showed equal variance; otherwise Dunnett T3 test was used. The comparison of variables between the genders was performed by independent Student t-test.

Bivariate correlations were evaluated by Pearson correlation analyses.

The overall associations between the AhR-mediated activities and POPs across the study groups and/or across genders were assessed by comparing the regression lines for each study group/gender by using multiple regression analysis.

Up to date few studies on AhR transcriptional activity in human serum have been reported [27,36-39], and thus the knowledge is limited about which dietary or other lifestyle determinants might affect serum AhR transcriptional activity. Our hypothesis is that potential determinants of POP bioaccumulation might also be potential determinants for serum AhR transcriptional activity. Guided by the literature [40], age and seafood intake are known determinants affecting the POP serum level, and intake of seabird might as well influence the POP level. The n-3 fatty acids being determinants of seafood intake [8-10], and lifestyle factors such as smoking and BMI might also influence the serum POP level [8]. Using the multiple linear regression model, assessing the relation between the POPs and AhR activities, the impact of potential confounders were evaluated by entering variables together with  $\Sigma$ PCB<sub>14</sub>,  $\Sigma$  pesticide and  $\Sigma$  POP as follows: in the first step, age and then seafood intake (represented by n-3/n-6 ratio) were included in the model, then additionally smoking years and BMI were included in the model. Finally, the bird intake was additionally included in the model. The potential confounder was defined if the regression coefficient ( $\beta$ ) changes more than 10% and when a significant association ( $p \leq 0.05$ ) was obtained after adjustment. Then the calculations were adjusted for the confounders.

The inter-district variations in serum levels of POPs, AhR-TEQ and AhRcomp were also assessed by linear regression models and age was in these models considered as a potential confounder of these variables.

The statistical analysis was performed on SPSS 13.0 (SPSS Inc. Chicago, IL, USA). The statistical significant level was set to  $p = 0.05$ .

### 3. Results

#### 3.1. The demographical and lifestyle factors of study groups

As reported [10], the age of the participants was from 18-77 with the median age of 35 years old. The Nuuk men were significantly older than the other participants, whereas the age range of the remaining male and female participants of all four districts were similar (Table 1). No significant difference was observed for the BMI of the participants from all districts. For the smoking years, Nuuk men and women had the highest median, whereas the other three districts were similar for both sexes (Table 1). The Nuuk and Sisimiut men had the highest and the low-

est median n-3/n-6 ratio, respectively, whereas for women significantly higher n-3/n-6 ratio was observed for Qaanaaq and Tasiilaq. The median seabird intake of the Nuuk men was the highest whilst for the rest of subject's similar median seabird intake was observed (Table 1). Nuuk and Sisimiut participants had higher median consumption of dairy food than that of Qaanaaq and in overall male participants had lower dairy food intake than women (Table 1).

#### 3.2. The plasma level of POPs

For the men, similar median levels of  $\Sigma$  PCB<sub>14</sub>,  $\Sigma$  pesticide and  $\Sigma$  POP were observed among Nuuk, Qaanaaq and Tasiilaq and all were significantly higher than that of Sisimiut with the order of Nuuk  $\geq$  Qaanaaq  $\geq$  Tasiilaq  $>$  Sisimiut. Tasiilaq and Qaanaaq women had significantly higher POPs median levels than that of Nuuk and Sisimiut women (Table 2) as reported [10]. In general, male participants had higher median POP levels than women in all districts. For the combined sex data, Tasiilaq and Qaanaaq had similar POPs median levels which were significantly higher than that of Nuuk and Sisimiut participants (Tasiilaq  $\geq$  Qaanaaq  $>$  Nuuk  $>$  Sisimiut). After adjustment for age, the pattern of regional difference did not change (data not shown).

The dioxin-like PCBs (DL-PCBs) determined in this study were CB105, CB118 and CB156. The Nuuk men had higher median percentage of  $\Sigma$ DL-PCB of the  $\Sigma$  PCB<sub>14</sub> than men from other districts (Nuuk  $>$  Tasiilaq  $\geq$  Qaanaaq  $\geq$  Sisimiut). No significant difference of  $\Sigma$ DL-PCB percentage between women from different districts was observed but a lower trend was observed for Qaanaaq women (Table 2). Women had significantly higher median percentage of DL-PCBs than men. In overall, the median percentage of  $\Sigma$ DL-PCB for the combined data was in the order of Nuuk  $\geq$  Tasiilaq  $\geq$  Sisimiut  $>$  Qaanaaq (Table 2).

#### 3.3. AhR-TEQ and AhRcomp activity

91% of all male and 79% of all female serum samples showed significant agonistic AhR transcriptional activity. The levels of AhR-TEQ differed significantly among the districts for the separate gender (Table 3, Fig. 1A & 1B). The order of median level of AhR-TEQ was Tasiilaq  $\geq$  Sisimiut  $\geq$  Nuuk  $>$  Qaanaaq for men (Fig. 1A), and for women Nuuk  $\geq$  Tasiilaq  $\geq$  Sisimiut  $>$  Qaanaaq (Fig. 1B). Thus for both genders Qaanaaq had a lower AhR-TEQ level than participants from other districts (Table 3).

AhRcomp activity was assessed by co-exposure with serum extract and 60 pM TCDD ( $EC_{50(TCDD)}$ ). The median AhRcomp activity of men was in the order of Qaanaaq  $\geq$  Sisimiut  $\geq$  Tasiilaq  $>$  Nuuk, reflecting the higher percentage of samples elicited inhibition of the TCDD induced AhR activity in the serum samples of Nuuk and Tasiilaq (Table

**Table 1: Basic characteristics of study population**

| Men                                |         | Nuuk      | Sisimiut          | Qaanaaq           | Tasiilaq          | All       |
|------------------------------------|---------|-----------|-------------------|-------------------|-------------------|-----------|
| Age (years)                        | n       | 50        | 52                | 43                | 41                | 186       |
|                                    | median  | 54*       | 30                | 34                | 34                | 37        |
|                                    | mean    | 54        | 31                | 34                | 34                | 39        |
|                                    | min-max | 35–77     | 18–46             | 19–45             | 19–45             | 18–77     |
| BMI (kg/m <sup>2</sup> )           | n       | 46        | 52                | 43                | 41                | 182       |
|                                    | median  | 28        | 26                | 27                | 26                | 26        |
|                                    | mean    | 28        | 27                | 27                | 26                | 27        |
|                                    | min-max | 22–35     | 19–36             | 20–41             | 21–33             | 19–41     |
| Smoking (years)                    | n       | 45        | 52                | 41                | 40                | 178       |
|                                    | median  | 34*       | 12                | 18                | 15                | 18        |
|                                    | mean    | 31        | 12                | 17                | 15                | 19        |
|                                    | min-max | 0–59      | 0–26              | 0–31              | 0–31              | 0–59      |
| n-3/n-6                            | n       | 50        | 37                | 43                | 41                | 171       |
|                                    | median  | 0.61*     | 0.22 <sup>▲</sup> | 0.40              | 0.40              | 0.40      |
|                                    | mean    | 0.74      | 0.29              | 0.48              | 0.45              | 0.50      |
|                                    | min-max | 0.20–1.93 | 0.10–0.73         | 0.09–1.45         | 0.12–1.03         | 0.09–1.93 |
| Seabird intake (per month)         | n       | 47        | 49                | 41                | na                | 137       |
|                                    | median  | 2.0*      | 1.0               | 1.0               | na                | 2.0       |
|                                    | mean    | 5.5       | 2.6               | 2.6               | na                | 3.6       |
|                                    | min-max | 0–20      | 0–28              | 0–20              | na                | 0–28      |
| Dairy food consumption (per month) | n       | 48        | 51                | 43                | na                | 142       |
|                                    | median  | 41        | 51                | 32* <sup>▲</sup>  | na                | 41        |
|                                    | mean    | 39        | 49                | 30                | na                | 40        |
|                                    | min-max | 2–84      | 8–84              | 3–58              | na                | 2–84      |
| Women                              |         |           |                   |                   |                   |           |
| Age (years)                        | n       | 45        | 42                | 36                | 48                | 171       |
|                                    | median  | 38        | 31                | 34                | 29                | 34        |
|                                    | mean    | 36        | 33                | 33                | 30                | 33        |
|                                    | min-max | 19–45     | 18–44             | 18–44             | 19–45             | 18–45     |
| BMI (kg/m <sup>2</sup> )           | n       | 43        | 42                | 36                | 48                | 169       |
|                                    | median  | 26        | 26                | 24                | 25                | 25        |
|                                    | mean    | 26        | 27                | 26                | 26                | 26        |
|                                    | min-max | 18–32     | 19–47             | 18–34             | 17–40             | 17–47     |
| Smoking (years)                    | n       | 42        | 42                | 34                | 48                | 166       |
|                                    | median  | 20        | 12                | 16                | 12                | 14        |
|                                    | mean    | 16        | 13                | 16                | 13                | 14        |
|                                    | min-max | 0–33      | 0–27              | 0–29              | 0–31              | 0–33      |
| n-3/n-6                            | n       | 45        | 38                | 36                | 48                | 167       |
|                                    | median  | 0.25      | 0.29              | 0.40 <sup>#</sup> | 0.41 <sup>#</sup> | 0.33      |
|                                    | mean    | 0.29      | 0.32              | 0.48              | 0.46              | 0.38      |
|                                    | min-max | 0.12–0.72 | 0.14–0.82         | 0.15–1.53         | 0.18–1.0          | 0.12–1.53 |
| Seabird intake (per month)         | n       | 44        | 41                | 35                | na                | 120       |
|                                    | median  | 1.0       | 1.0               | 1.0               | na                | 1.0       |
|                                    | mean    | 1.4       | 1.3               | 3.4               | na                | 1.9       |
|                                    | min-max | 0–8.0     | 0–2.0             | 1.0–28            | na                | 0–28      |
| Dairy food consumption (per month) | n       | 45        | 42                | 35                | na                | 122       |
|                                    | median  | 56        | 57                | 37                | na                | 49        |
|                                    | mean    | 53        | 49                | 36                | na                | 47        |
|                                    | min-max | 2–84      | 0–84              | 0–84              | na                | 0–84      |
| Men + Women                        |         |           |                   |                   |                   |           |
| Age (years)                        | n       | 95        | 94                | 79                | 89                | 357       |
|                                    | median  | 44        | 31                | 34                | 32                | 35        |
|                                    | mean    | 46        | 31                | 33                | 32                | 36        |
|                                    | min-max | 19–77     | 18–46             | 18–45             | 19–45             | 18–77     |
| BMI (kg/m <sup>2</sup> )           | n       | 89        | 94                | 79                | 89                | 351       |
|                                    | median  | 26        | 26                | 26                | 25                | 26        |
|                                    | mean    | 27        | 27                | 27                | 26                | 26        |
|                                    | min-max | 18–35     | 19–47             | 18–41             | 17–40             | 17–47     |
| Smoking (years)                    | n       | 87        | 94                | 75                | 88                | 344       |

**Table 1: Basic characteristics of study population (Continued)**

|                                    |         |           |           |           |           |           |
|------------------------------------|---------|-----------|-----------|-----------|-----------|-----------|
|                                    | median  | 23        | 12        | 17        | 13        | 16        |
|                                    | mean    | 23        | 12        | 16        | 14        | 16        |
|                                    | min-max | 0-59      | 0-27      | 0-31      | 0-31      | 0-59      |
| n-3/n-6                            | n       | 95        | 75        | 79        | 89        | 338       |
|                                    | median  | 0.36      | 0.25      | 0.40      | 0.40      | 0.36      |
|                                    | mean    | 0.51      | 0.30      | 0.48      | 0.45      | 0.44      |
|                                    | min-max | 0.12-1.93 | 0.10-0.82 | 0.09-1.53 | 0.12-1.03 | 0.09-1.93 |
| Seabird intake (per month)         | n       | 91        | 90        | 76        | na        | 257       |
|                                    | median  | 2.0       | 1.0       | 1.0       | na        | 1.0       |
|                                    | mean    | 3.5       | 2.0       | 3.0       | na        | 2.8       |
|                                    | min-max | 0-20      | 0-28      | 0-28      | na        | 0-28      |
| Dairy food consumption (per month) | n       | 93        | 93        | 78        | na        | 264       |
|                                    | median  | 48        | 51        | 36        | na        | 43        |
|                                    | mean    | 46        | 49        | 33        | na        | 43        |
|                                    | min-max | 2-84      | 0-84      | 0-84      | na        | 0-84      |

na: not available.

\*:  $p < 0.05$  versus other districts; ▲:  $p < 0.05$  versus Nuuk and Sisimiut men; #:  $p < 0.05$  versus Nuuk and Sisimiut women; ♣:  $p < 0.05$  versus Nuuk and Sisimiut men and all women.

3, Fig. 1C). Higher percentage of serum samples of men from Qaanaaq and Sisimiut elicited a further increase of the TCDD induced AhR activity (Table 3, Fig. 1C). The serum samples of women from the four districts had similar AhRcomp level; mainly eliciting a further increase of the TCDD induced AhR activity (Table 3, Fig. 1D).

No significant difference of the AhR-TEQ level was found between the genders in the separate district (Table 3). In the Nuuk study group, men had a lower level of AhRcomp than women, whereas similar AhRcomp levels were found for the two genders for Sisimiut, Qaanaaq and Tasiilaq participants (Table 3).

Adjustment for age did not change the regional and gender difference of AhR-TEQ and AhRcomp (data not shown).

### 3.4. The association between AhR-TEQ, AhRcomp and POPs

#### 3.4.1. The correlations between the single POPs

Pearson correlation analyses showed that most PCB congeners were significantly and mutually intercorrelated ( $r > 0.82$ ;  $p < 0.001$ ), whereas CB 28, CB52, CB128 and CB101 had a relatively lower correlation coefficients to other PCB congeners ( $r = 0.29-0.82$ ;  $p < 0.05$ ). Also the determined pesticides were intercorrelated ( $r > 0.75 - 0.99$ ). The grouping variables ( $\Sigma$ PCB<sub>14</sub> and  $\Sigma$  pesticide) intercorrelated with  $r = 0.97$  and the sum of determined POPs as  $\Sigma$  POP was therefore acceptable.

#### 3.4.2. The correlations between AhR-TEQ, AhRcomp and POPs

##### 3.4.2.1. Multiple regressions of AhR-TEQ and AhRcomp on POPs across the study groups and genders

Multiple regression analysis of the combined districts and genders data was performed to analyze for homogeneity/heterogeneity of the associations between POPs and AhR-

TEQ/AhRcomp (see Additional file 1). The analyses indicated that the associations of POPs and AhR-TEQ were allowed to be evaluated in each district for the combined gender, whereas the AhRcomp activity and POPs association must be evaluated in each single district for men and for women in combined districts.

However, in order to obtain the overall trend of the studied groups and a better statistical power, the data based on the combined genders of the single districts for POPs and AhRcomp and the combined genders across the districts were also evaluated for the correlation of POPs and AhR-TEQ/AhRcomp.

##### 3.4.2.2. Correlations between AhR-TEQ, AhRcomp and POPs

An inverse correlation between AhR-TEQ and POPs was observed for the combined men data and the total combined data (Table 4). Few significant correlations between AhR-TEQ and POPs were found for the single districts for the two sexes. Significantly inverse correlations between AhR-TEQ and the POP variables were observed for both the Qaanaaq women and the combined Qaanaaq participants before and after adjustment for age and/or n-3/n-6 (Table 4).

As shown in Table 5, the total combined data and the combined Tasiilaq data showed an inverse relationship between AhRcomp and POPs before and after adjustment for age and n-3/n-6. The significantly inverse correlations of AhRcomp and POPs observed for all men data disappeared upon adjustment for age and/or n-3/n-6 ratio (Table 5). Also for the combined Nuuk data, negative correlations of AhRcomp and POPs were observed, which disappeared upon adjustment for age plus n-3/n-6 ratio (Table 5). For the two sexes in the single districts, few significant correlations between AhRcomp and POPs were observed but Tasiilaq women showing an inverse correla-

**Table 2: The plasma POP levels in the subjects of the different district**

| Men                               |         | Nuuk              | Sisimiut  | Qaanaaq           | Tasiilaq          | All        |
|-----------------------------------|---------|-------------------|-----------|-------------------|-------------------|------------|
| Σ PCB <sub>14</sub> (µg/kg lipid) | n       | 38                | 51        | 41                | 37                | 167        |
|                                   | median  | 2682              | 540.9*    | 1854              | 1890              | 1526       |
|                                   | mean    | 3094              | 752.1     | 2699              | 2079              | 2057       |
|                                   | min-max | 490-11113         | 132-2366  | 617-7352          | 318-4527          | 132-11113  |
| Σ pesticide (µg/kg lipid)         | n       | 38                | 51        | 41                | 41                | 171        |
|                                   | median  | 3401              | 737.8*    | 2630              | 2242              | 2126       |
|                                   | mean    | 4034              | 1012      | 3824              | 2372              | 2684       |
|                                   | min-max | 405-19118         | 141-3542  | 617-14366         | 194-7224          | 141-19118  |
| Σ POP (µg/kg lipid)               | n       | 38                | 51        | 41                | 37                | 167        |
|                                   | median  | 6322              | 1353*     | 4411              | 4067              | 3551       |
|                                   | mean    | 7127              | 1765      | 6524              | 4249              | 4692       |
|                                   | min-max | 896-30231         | 285-5908  | 1234-19807        | 499-11260         | 285-30231  |
| ΣDL-PCB/Σ PCB <sub>14</sub> (%)   | n       | 38                | 51        | 41                | 37                | 167        |
|                                   | median  | 8.42 <sup>▲</sup> | 7.07      | 7.53              | 7.51              | 7.70       |
|                                   | mean    | 8.36              | 7.41      | 7.44              | 7.73              | 7.71       |
|                                   | min-max | 5.0-11.0          | 4.0-11.0  | 4.0-11.0          | 6.0-11.0          | 4.0-11.0   |
| <b>Women</b>                      |         |                   |           |                   |                   |            |
| Σ PCB <sub>14</sub> (µg/kg lipid) | n       | 44                | 42        | 34                | 47                | 167        |
|                                   | median  | 295.7             | 346       | 1188 <sup>▲</sup> | 1660 <sup>▲</sup> | 635.5      |
|                                   | mean    | 492.4             | 459.7     | 1406              | 1809              | 1041       |
|                                   | min-max | 97.5-2428         | 67.2-1384 | 99-5980           | 112-6701          | 67.2-6701  |
| Σ pesticide (µg/kg lipid)         | n       | 44                | 42        | 34                | 48                | 168        |
|                                   | median  | 481.6             | 496.3     | 1671 <sup>▲</sup> | 2044 <sup>▲</sup> | 860        |
|                                   | mean    | 678               | 700       | 2018              | 2397              | 1446       |
|                                   | min-max | 83.6-2881         | 75.6-3014 | 77.9-7529         | 153-6995          | 85.5-7529  |
| Σ POP (µg/kg lipid)               | n       | 44                | 42        | 34                | 47                | 167        |
|                                   | median  | 748               | 823       | 2847 <sup>▲</sup> | 3797 <sup>▲</sup> | 1469       |
|                                   | mean    | 1171              | 1159      | 3424              | 4169              | 2480       |
|                                   | min-max | 181-5309          | 143-4337  | 177-12432         | 256-12829         | 143-12829  |
| ΣDL-PCB/Σ PCB <sub>14</sub> (%)   | n       | 44                | 42        | 34                | 47                | 167        |
|                                   | median  | 9.01              | 9.26      | 8.45              | 8.96              | 9.00       |
|                                   | mean    | 9.02              | 9.40      | 8.78              | 9.02              | 9.10       |
|                                   | min-max | 4.0-13.0          | 6.0-15.0  | 6.0-13.0          | 6.0-12.0          | 4.0-15.0   |
| <b>Men + Women</b>                |         |                   |           |                   |                   |            |
| Σ PCB <sub>14</sub> (µg/kg lipid) | n       | 82                | 93        | 75                | 84                | 334        |
|                                   | median  | 912.4             | 495.3     | 1379 <sup>#</sup> | 1806 <sup>#</sup> | 1068       |
|                                   | mean    | 1698              | 620.0     | 2113              | 1928              | 1549       |
|                                   | min-max | 97.5-11113        | 67.2-2366 | 99-7352           | 112-6701          | 67.2-11113 |
| Σ pesticide (µg/kg lipid)         | n       | 82                | 93        | 75                | 89                | 339        |
|                                   | median  | 1336              | 685.3     | 2156 <sup>#</sup> | 2208 <sup>#</sup> | 1327       |
|                                   | mean    | 2234              | 871       | 3006              | 2386              | 2071       |
|                                   | min-max | 83.6-19118        | 75.6-3542 | 77.9-14366        | 153-7224          | 85.5-19118 |
| Σ POP (µg/kg lipid)               | n       | 82                | 93        | 75                | 84                | 334        |
|                                   | median  | 2322              | 1244      | 3437 <sup>#</sup> | 4020 <sup>#</sup> | 2420       |
|                                   | mean    | 3931              | 1491      | 5119              | 4206              | 3597       |
|                                   | min-max | 181-30231         | 143-5908  | 177-19807         | 256-12829         | 143-30231  |
| ΣDL-PCB/Σ PCB <sub>14</sub> (%)   | n       | 82                | 93        | 75                | 84                | 334        |
|                                   | median  | 8.79              | 7.92      | 7.88              | 8.40              | 8.30       |
|                                   | mean    | 8.71              | 8.31      | 8.05              | 8.45              | 8.39       |
|                                   | min-max | 4.0-13.0          | 4.0-15.0  | 4.0-13.0          | 6.0-12.0          | 4.0-15.0   |

Σ PCB<sub>14</sub>: sum of 14 PCB congeners; Σ pesticide: sum of 10 pesticides; Σ POP: Σ PCB<sub>14</sub> + Σ pesticide; ΣDL-PCBs: sum of dioxin-like PCBs (PCB105, PCB118, PCB156).

\*: p < 0.05, versus men from other districts; <sup>▲</sup>: p < 0.05, versus Nuuk and Sisimiut women;

<sup>#</sup>: p < 0.05, versus Nuuk and Sisimiut subjects; <sup>▲</sup>: p < 0.05, versus men from other districts.

**Table 3: The serum levels of AhR transcriptional activity of subjects of the different district**

| Men                                   |            | Nuuk        | Sisimiut    | Qaanaaq     | Tasiilaq    | All         |
|---------------------------------------|------------|-------------|-------------|-------------|-------------|-------------|
| AhR-TEQ <sup>1</sup> (pg/g lipid)     | n          | 21          | 42          | 38          | 38          | 139         |
|                                       | median     | 166         | 189         | 110*        | 240         | 153         |
|                                       | mean       | 206         | 247         | 118         | 296         | 219         |
|                                       | min-max    | 95 – 1019   | 26 – 1202   | 58 – 271    | 98 – 1452   | 26 – 1452   |
|                                       | % agonist  | 79          | 88          | 100         | 97          | 91          |
| AhRcomp <sup>2</sup> (RLU/μg protein) | n          | 38          | 51          | 43          | 38          | 170         |
|                                       | median     | 0.78*       | 1.21        | 1.35        | 1.04        | 1.15        |
|                                       | mean       | 0.82        | 1.24        | 1.38        | 1.12        | 1.15        |
|                                       | min-max    | 0.38 – 1.18 | 0.59 – 2.46 | 0.77 – 2.17 | 0.31 – 3.24 | 0.31 – 3.24 |
|                                       | % increase | 0           | 44          | 54          | 29          | 33          |
| % inhibition                          | 32         | 4           | 2           | 32          | 16          |             |
| <b>Women</b>                          |            |             |             |             |             |             |
| AhR-TEQ <sup>1</sup> (pg/g lipid)     | n          | 12          | 32          | 32          | 44          | 120         |
|                                       | median     | 235         | 188         | 122*        | 228         | 167         |
|                                       | mean       | 326         | 221         | 131         | 332         | 248         |
|                                       | min-max    | 115 – 1055  | 93 – 1032   | 63 – 259    | 100 – 1987  | 63 – 1987   |
|                                       | % agonist  | 64          | 64          | 100         | 91          | 79          |
| AhRcomp <sup>2</sup> (RLU/μg protein) | n          | 45          | 42          | 36          | 44          | 167         |
|                                       | median     | 1.41        | 1.33        | 1.31        | 1.28        | 1.32        |
|                                       | mean       | 1.50        | 1.38        | 1.33        | 1.92        | 1.54        |
|                                       | min-max    | 0.9 – 4.54  | 0.94 – 3.88 | 0.74 – 1.87 | 0.25 – 15   | 0.25 – 15   |
|                                       | % increase | 49          | 64          | 58          | 46          | 54          |
| % inhibition                          | 0          | 0           | 11          | 21          | 8           |             |
| <b>Men + Women</b>                    |            |             |             |             |             |             |
| AhR-TEQ <sup>1</sup> (pg/g lipid)     | n          | 33          | 74          | 70          | 82          | 259         |
|                                       | median     | 188         | 189         | 115         | 239         | 161         |
|                                       | mean       | 250         | 236         | 124         | 315         | 232         |
|                                       | min-max    | 94 – 1055   | 26 – 1202   | 58 – 271    | 98 – 1987   | 26 – 1987   |
|                                       | % agonist  | 71          | 76          | 100         | 94          | 85          |
| AhRcomp <sup>2</sup> (RLU/μg protein) | n          | 83          | 93          | 79          | 82          | 337         |
|                                       | median     | 1.08        | 1.26        | 1.33        | 1.18        | 1.23        |
|                                       | mean       | 1.19        | 1.30        | 1.36        | 1.55        | 1.35        |
|                                       | min-max    | 0.38 – 4.54 | 0.59 – 3.88 | 0.74 – 2.17 | 0.25 – 15   | 0.25 – 15   |
|                                       | % increase | 27          | 53          | 56          | 38          | 43          |
| % inhibition                          | 15         | 2           | 6           | 26          | 12          |             |

<sup>1</sup>AhR-TEQ (TCDD equivalents): The AhR-TEQ of samples eliciting significantly agonistic activity was calculated by interpolation to the TCDD dose-response curve using the Sigmaplot program, given as pg/g serum lipid. The % agonistic indicates the % of samples eliciting a significant increase in AhR activity compared to the solvent control. <sup>2</sup>AhRcomp: AhR competitive activity of serum extract + 60 pM TCDD (EC<sub>50</sub>) given as RLU/μg protein. EC<sub>50</sub> solvent control = 1 RLU/μg protein; % increase and % inhibition indicates the % of samples responding with a further increase or decrease of the EC<sub>50</sub>(TCDD) induced activity, respectively. \*: p < 0.05 versus subjects from other districts.

tion upon adjustment for age and/or n-3/n-6. In overall, negative association between AhRcomp and POP data was found but Sisimiut men where AhRcomp activity was positively related to the determined ΣPCB<sub>14</sub> upon adjustment for age and n-3/n-6.

Further adjustment for BMI, smoking year and bird intake did not change the pattern of associations of POPs and AhR-TEQ, AhRcomp (data not shown).

Scattered and few significant correlations of AhR-TEQ/AhRcomp and the single PCB and pesticides and/or Σ DL-PCBs were observed (data not shown). Since the congeners of PCBs were highly intercorrelated and the AhR-TEQ represents the integrated activity of the POP mixture, it is


hard to assess which of the compounds having the highest impact on the AhR-TEQ/AhRcomp level.

**3.5. Correlations between lifestyle factors, POPs and AhR transcriptional activity**

**3.5.1. POPs**

As found earlier [10], for the combined data the levels of ΣPCB<sub>14</sub>, Σpesticide and ΣPOPs were positively associated with age (r > 0.449, p < 0.001) and seafood intake represented by n-3/n-6 ratio (r > 0.614, p < 0.001), and age was positively correlated to seafood intake (r = 0.472, p < 0.001).





**Figure 1**  
**The serum AhR transcriptional activity.** A) The AhR-TEQ in men from different districts; B) The AhR-TEQ in women from different districts; C) The AhRcomp activity in men from different districts and D) The AhRcomp activity in women from different districts. The reference line of the control is given as dash line (C, D). The values in the figure are percentage of sample with agonistic (A, B) and increase or inhibition of TCDD induced AhR transcriptional activity (C, D).

**Table 4: Multivariate linear regression analysis of AhR-TEQ and POPs**

|                          |                       | Nuuk |       |      | Sisimiut |      |      | Qaanaaq   |             |                | Tasiilaq |      |     | All        |             |                |
|--------------------------|-----------------------|------|-------|------|----------|------|------|-----------|-------------|----------------|----------|------|-----|------------|-------------|----------------|
| Men <sup>a</sup>         |                       | n    | β     | p    | n        | β    | p    | n         | β           | p              | n        | β    | p   | n          | β           | p              |
| Σ PCB <sub>14</sub>      | raw <sup>1</sup>      | 21   | -.01  | .97  | 41       | -.24 | .13  | 37        | -.14        | .41            | 33       | -.06 | .72 | <b>135</b> | <b>-.21</b> | <b>.01</b>     |
|                          | +age                  | 21   | .05   | .86  | 41       | -.09 | .64  | 37        | .05         | .80            | 33       | .03  | .87 | 135        | -.16        | .11            |
|                          | +n-3/n-6 <sup>2</sup> | 21   | -.01  | .98  | 29       | -.25 | .25  | 37        | -.30        | .23            | 33       | -.11 | .54 | 123        | -.22        | .07            |
|                          | +age+n-3/n-6          | 21   | .07   | .87  | 29       | -.06 | .81  | 37        | -.14        | .60            | 33       | .01  | .98 | 123        | -.16        | .21            |
| Σ pesticide              | raw                   | 21   | .08   | .72  | 41       | -.27 | .087 | 37        | -.20        | .23            | 37       | -.13 | .46 | <b>139</b> | <b>-.25</b> | <b>&lt;.01</b> |
|                          | +age                  | 21   | .15   | .57  | 41       | -.14 | .44  | 37        | -.01        | .98            | 37       | -.05 | .78 | <b>139</b> | <b>-.21</b> | <b>.04</b>     |
|                          | +n-3/n-6 <sup>2</sup> | 21   | .16   | .61  | 29       | -.34 | .13  | 37        | -.49        | .06            | 37       | -.20 | .23 | <b>127</b> | <b>-.37</b> | <b>&lt;.01</b> |
|                          | +age+n-3/n-6          | 21   | .22   | .53  | 29       | -.20 | .39  | 37        | -.30        | .32            | 37       | -.09 | .60 | <b>127</b> | <b>-.31</b> | <b>.01</b>     |
| Σ POP                    | raw <sup>1</sup>      | 21   | .04   | .86  | 41       | -.26 | .10  | 37        | -.19        | .24            | 33       | -.11 | .56 | <b>135</b> | <b>-.16</b> | <b>.01</b>     |
|                          | +age                  | 21   | .11   | .69  | 41       | -.12 | .50  | 37        | -.01        | .96            | 33       | -.02 | .93 | 135        | -.20        | .06            |
|                          | +n-3/n-6 <sup>2</sup> | 21   | .09   | .78  | 29       | -.30 | .17  | 37        | -.48        | .07            | 33       | -.15 | .40 | <b>123</b> | <b>-.29</b> | <b>.02</b>     |
|                          | +age+n-3/n-6          | 21   | .16   | .66  | 29       | -.15 | .54  | 37        | -.29        | .31            | 33       | -.04 | .82 | 123        | -.23        | .06            |
| Women <sup>b</sup>       |                       |      |       |      |          |      |      |           |             |                |          |      |     |            |             |                |
| Σ PCB <sub>14</sub>      | raw <sup>1</sup>      | 12   | .06   | .85  | 32       | .23  | .20  | <b>32</b> | <b>-.46</b> | <b>.01</b>     | 43       | -.09 | .55 | 119        | -.04        | .70            |
|                          | +age                  | 12   | -.01  | .99  | 32       | .25  | .24  | <b>32</b> | <b>-.40</b> | <b>.05</b>     | 43       | -.21 | .35 | 119        | -.002       | .99            |
|                          | +n-3/n-6 <sup>2</sup> | 12   | .58   | .19  | 28       | .17  | .43  | 32        | -.32        | .18            | 43       | -.13 | .50 | 115        | .06         | .62            |
|                          | +age+n-3/n-6          | 12   | .53   | .31  | 28       | .20  | .38  | 32        | -.25        | .36            | 43       | -.23 | .35 | 115        | .08         | .53            |
| Σ pesticide              | raw <sup>1</sup>      | 12   | .11   | .74  | 32       | .25  | .16  | <b>32</b> | <b>-.53</b> | <b>&lt;.01</b> | 44       | -.11 | .46 | 120        | -.06        | .52            |
|                          | +age                  | 12   | .06   | .89  | 32       | .26  | .19  | <b>32</b> | <b>-.50</b> | <b>.01</b>     | 44       | -.28 | .20 | 120        | -.03        | .74            |
|                          | +n-3/n-6 <sup>2</sup> | 12   | .69   | .12  | 28       | .19  | .39  | <b>32</b> | <b>-.51</b> | <b>.04</b>     | 44       | -.17 | .35 | 116        | .01         | .97            |
|                          | +age+n-3/n-6          | 12   | .65   | .20  | 28       | .20  | .37  | 32        | -.47        | .09            | 44       | -.31 | .19 | 116        | .02         | .89            |
| Σ POP                    | raw <sup>1</sup>      | 12   | .09   | .79  | 32       | .24  | .18  | <b>32</b> | <b>-.51</b> | <b>&lt;.01</b> | 43       | -.11 | .49 | 119        | -.05        | .57            |
|                          | +age                  | 12   | .03   | .95  | 32       | .25  | .22  | <b>32</b> | <b>-.48</b> | <b>.02</b>     | 43       | -.24 | .28 | 119        | -.02        | .83            |
|                          | +n-3/n-6 <sup>2</sup> | 12   | .66   | .14  | 28       | .17  | .42  | 32        | -.47        | .07            | 43       | -.15 | .43 | 115        | .04         | .77            |
|                          | +age+n-3/n-6          | 12   | .62   | .23  | 28       | .20  | .39  | 32        | -.42        | .14            | 43       | -.27 | .27 | 115        | .05         | .67            |
| Men + Women <sup>c</sup> |                       |      |       |      |          |      |      |           |             |                |          |      |     |            |             |                |
| Σ PCB <sub>14</sub>      | raw <sup>1</sup>      | 33   | -.19  | .29  | 74       | -.05 | .65  | <b>70</b> | <b>-.32</b> | <b>.01</b>     | 77       | -.10 | .40 | <b>254</b> | <b>-.15</b> | <b>.02</b>     |
|                          | +age                  | 33   | -.09  | .79  | 74       | .03  | .83  | <b>70</b> | <b>-.22</b> | <b>.01</b>     | 77       | -.07 | .65 | 254        | -.10        | .18            |
|                          | +n-3/n-6 <sup>2</sup> | 33   | .12   | .72  | 58       | -.15 | .30  | <b>70</b> | <b>-.34</b> | <b>.02</b>     | 77       | -.15 | .24 | 238        | -.16        | .06            |
|                          | +age+n-3/n-6          | 33   | .24   | .60  | 58       | -.09 | .53  | 70        | -.26        | .09            | 77       | -.11 | .49 | 238        | -.12        | .17            |
| Σ pesticide              | raw <sup>1</sup>      | 33   | -.13  | .47  | 74       | -.07 | .58  | <b>70</b> | <b>-.38</b> | <b>&lt;.01</b> | 82       | -.12 | .28 | <b>254</b> | <b>-.18</b> | <b>&lt;.01</b> |
|                          | +age                  | 33   | .06   | .84  | 74       | .01  | .94  | <b>70</b> | <b>-.30</b> | <b>.03</b>     | 82       | -.12 | .38 | 254        | -.13        | .07            |
|                          | +n-3/n-6 <sup>2</sup> | 33   | .29   | .37  | 58       | -.16 | .29  | <b>70</b> | <b>-.47</b> | <b>&lt;.01</b> | 82       | -.21 | .09 | <b>238</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                          | +age+n-3/n-6          | 33   | .41   | .29  | 58       | -.12 | .44  | 70        | -.40        | .02            | 82       | -.17 | .22 | <b>238</b> | <b>-.21</b> | <b>.02</b>     |
| Σ POP                    | raw <sup>1</sup>      | 33   | -.16  | .38  | 74       | -.06 | .59  | <b>70</b> | <b>-.37</b> | <b>&lt;.01</b> | 77       | -.12 | .30 | <b>254</b> | <b>-.17</b> | <b>.01</b>     |
|                          | +age                  | 33   | -.001 | .997 | 74       | .01  | .92  | <b>70</b> | <b>-.29</b> | <b>.04</b>     | 77       | -.10 | .49 | 254        | -.12        | .09            |
|                          | +n-3/n-6 <sup>2</sup> | 33   | .23   | .50  | 58       | .16  | .28  | <b>70</b> | <b>-.45</b> | <b>&lt;.01</b> | 77       | -.18 | .17 | <b>238</b> | <b>-.20</b> | <b>.02</b>     |
|                          | +age+n-3/n-6          | 33   | .36   | .395 | 58       | -.11 | .46  | 70        | -.31        | .02            | 77       | -.14 | .36 | 238        | -.16        | .07            |

The analyses were performed on ln transformed data with AhR-TEQ as dependent and POPs as independent variables before and upon adjusted for the potential confounders.

<sup>1</sup>: non-adjusted data.

<sup>2</sup>: ratio of n-3 to n-6 fatty acids in serum.

<sup>3</sup>: Combining districts is not allowed according to the multiple regression analyses.

<sup>b</sup>: Combining districts is not allowed according to the multiple regression analyses.

<sup>c</sup>: According to the multiple regression analyses, combining sex for each district is allowed while combining sexes for all districts is not allowed.

For Σ PCB<sub>14</sub>, Σ pesticide and Σ POP, AhR-TEQ, see the legend of Table 2 and Table 3

**3.5.2. AhR-TEQ**

As shown in Table 6, scattered and few correlations between AhR-TEQ and lifestyle factors were found in the single districts and the separate genders, most of which involved the age, and/or the n-3/n-6 ratio. A significantly negative correlation between age and AhR-TEQ was observed for Sisimiut men and the combined men data. For Sisimiut women, a positive correlation between BMI

and AhR-TEQ and for Qaanaaq women a negative correlation of the n-3/n-6 ratio and AhR-TEQ were observed. Significantly negative correlation between age and AhR-TEQ was also found for the combined Qaanaaq data and the total combined data. For the combined Nuuk, Sisimiut and Qaanaaq data a significantly, positive correlation between AhR-TEQ and consumption of dairy food was found (Table 6), suggesting that dairy products may

**Table 5: Multivariate linear regression analysis of AhRcomp and POPs**

|                     |  | Nuuk      |             |               | Sisimiut  |            |            | Qaanaaq |       |      | Tasiilaq  |             |               | All        |             |                |
|---------------------|--|-----------|-------------|---------------|-----------|------------|------------|---------|-------|------|-----------|-------------|---------------|------------|-------------|----------------|
|                     | Men <sup>a</sup>                             | n         | β           | p             | n         | β          | p          | n       | β     | p    | n         | β           | p             | n          | β           | p              |
| Σ PCB <sub>14</sub> | raw <sup>1</sup>                             | 38        | -.11        | .52           | 51        | -.13       | .36        | 41      | -.07  | .68  | 34        | -.18        | .31           | <b>164</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | 38        | -.16        | .47           | 51        | -.03       | .87        | 41      | -.001 | 1.00 | 34        | -.29        | .15           | 164        | -.05        | .56            |
|                     | +n-3/n-6 <sup>2</sup>                        | 38        | -.24        | .37           | 37        | .29        | .12        | 41      | .08   | .68  | 34        | -.20        | .27           | 150        | -.01        | .95            |
|                     | +age+n-3/n-6                                 | 38        | -.30        | .34           | <b>37</b> | <b>.42</b> | <b>.04</b> | 41      | .11   | .61  | 34        | -.29        | .15           | 150        | .10         | .35            |
| Σ pesticide         | raw <sup>1</sup>                             | 38        | -.09        | .59           | 51        | -.20       | .17        | 41      | -.12  | .46  | 38        | -.27        | .11           | <b>168</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | 38        | -.10        | .59           | 51        | -.12       | .45        | 41      | -.07  | .75  | 38        | -.36        | .05           | 168        | -.06        | .53            |
|                     | +n-3/n-6 <sup>2</sup>                        | 38        | -.16        | .52           | 37        | .22        | .27        | 41      | .01   | .97  | 38        | -.29        | .10           | 154        | -.04        | .73            |
|                     | +age+n-3/n-6                                 | 38        | -.16        | .52           | 37        | .29        | .17        | 41      | .04   | .87  | 38        | -.36        | .06           | 154        | .07         | .49            |
| Σ POP               | raw <sup>1</sup>                             | 38        | -.10        | .55           | 51        | -.18       | .22        | 41      | -.10  | .53  | 34        | -.19        | .28           | <b>164</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | 38        | -.13        | .53           | 51        | -.09       | .57        | 41      | -.04  | .84  | 34        | -.30        | .13           | 164        | -.06        | .53            |
|                     | +n-3/n-6 <sup>2</sup>                        | 38        | -.20        | .43           | 37        | .25        | .20        | 41      | .04   | .86  | 34        | -.21        | .25           | 150        | -.02        | .87            |
|                     | +age+n-3/n-6                                 | 38        | -.22        | .43           | 37        | .34        | .10        | 41      | .07   | .76  | 34        | -.30        | .13           | 150        | .09         | .46            |
| Σ PCB <sub>14</sub> | Women <sup>b</sup><br>raw <sup>1</sup>       | 44        | .08         | .60           | 42        | .17        | .28        | 34      | -.31  | .07  | 43        | -.18        | .24           | 163        | -.09        | .24            |
|                     | +age   | 44        | .16         | .37           | 42        | .09        | .59        | 34      | -.15  | .46  | <b>43</b> | <b>-.51</b> | <b>.02</b>    | 163        | -.11        | .17            |
|                     | +n-3/n-6 <sup>2</sup>                        | 44        | .18         | .25           | 38        | .09        | .62        | 34      | -.49  | .06  | 43        | -.27        | .14           | 159        | -.11        | .25            |
|                     | +age+n-3/n-6                                 | 44        | .25         | .14           | 38        | .07        | .71        | 34      | -.33  | .25  | <b>43</b> | <b>-.56</b> | <b>.02</b>    | 159        | -.13        | .21            |
| Σ pesticide         | raw <sup>1</sup>                             | 44        | .07         | .67           | 42        | .09        | .57        | 34      | -.25  | .15  | 44        | -.21        | .18           | 164        | -.11        | .18            |
|                     | +age   | 44        | .11         | .49           | 42        | .          | .995       | 34      | -.07  | .72  | <b>44</b> | <b>-.55</b> | <b>.01</b>    | 164        | -.13        | .13            |
|                     | +n-3/n-6 <sup>2</sup>                        | 44        | .18         | .26           | 38        | -.03       | .85        | 34      | -.37  | .17  | 44        | -.30        | .10           | 160        | -.14        | .18            |
|                     | +age+n-3/n-6                                 | 44        | .22         | .19           | 38        | -.06       | .76        | 34      | -.19  | .52  | <b>44</b> | <b>-.59</b> | <b>.01</b>    | 160        | -.15        | .15            |
| Σ POP               | raw <sup>1</sup>                             | 44        | .08         | .62           | 42        | .12        | .45        | 34      | -.27  | .12  | 43        | -.195       | .21           | 163        | -.10        | .21            |
|                     | +age   | 44        | .14         | .42           | 42        | .03        | .85        | 34      | -.10  | .64  | <b>43</b> | <b>-.54</b> | <b>.01</b>    | 163        | -.12        | .15            |
|                     | +n-3/n-6 <sup>2</sup>                        | 44        | .19         | .24           | 38        | .01        | .97        | 34      | -.42  | .12  | 43        | -.29        | .12           | 159        | -.13        | .21            |
|                     | +age+n-3/n-6                                 | 44        | .24         | .15           | 38        | -.02       | .94        | 34      | -.24  | .42  | <b>43</b> | <b>-.59</b> | <b>.01</b>    | 159        | -.14        | .17            |
| Σ PCB <sub>14</sub> | Men + Women <sup>c</sup><br>raw <sup>1</sup> | <b>82</b> | <b>-.55</b> | <b>&lt;.0</b> | 93        | -.05       | .64        | 75      | -.14  | .22  | 77        | -.22        | .06           | <b>327</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | <b>82</b> | <b>-.35</b> | <b>.02</b>    | 93        | -.07       | .58        | 75      | -.04  | .76  | <b>77</b> | <b>-.39</b> | <b>.01</b>    | <b>327</b> | <b>-.19</b> | <b>&lt;.01</b> |
|                     | +n-3/n-6 <sup>2</sup>                        | <b>82</b> | <b>-.39</b> | <b>.01</b>    | 75        | -.06       | .66        | 75      | -.07  | .62  | <b>77</b> | <b>-.27</b> | <b>.03</b>    | <b>309</b> | <b>-.21</b> | <b>&lt;.01</b> |
|                     | +age+n-3/n-6                                 | <b>82</b> | <b>-.21</b> | <b>.25</b>    | 75        | -.08       | .55        | 75      | -.001 | 1.00 | <b>77</b> | <b>-.41</b> | <b>.01</b>    | <b>309</b> | <b>-.18</b> | <b>.02</b>     |
| Σ pesticide         | raw <sup>1</sup>                             | <b>82</b> | <b>-.53</b> | <b>&lt;.0</b> | 93        | -.10       | .34        | 75      | -.15  | .20  | <b>82</b> | <b>-.23</b> | <b>.04</b>    | <b>332</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | <b>82</b> | <b>-.30</b> | <b>.03</b>    | 93        | -.13       | .27        | 75      | -.04  | .80  | <b>82</b> | <b>-.38</b> | <b>.01</b>    | <b>332</b> | <b>-.18</b> | <b>&lt;.01</b> |
|                     | +n-3/n-6 <sup>2</sup>                        | <b>82</b> | <b>-.34</b> | <b>.02</b>    | 75        | -.12       | .35        | 75      | -.08  | .61  | <b>82</b> | <b>-.28</b> | <b>.02</b>    | <b>314</b> | <b>-.21</b> | <b>.01</b>     |
|                     | +age+n-3/n-6                                 | <b>82</b> | <b>-.17</b> | <b>.28</b>    | 75        | -.14       | .29        | 75      | .01   | .93  | <b>82</b> | <b>-.40</b> | <b>&lt;.0</b> | <b>314</b> | <b>-.18</b> | <b>.02</b>     |
| Σ POP               | raw <sup>1</sup>                             | <b>82</b> | <b>-.54</b> | <b>&lt;.0</b> | 93        | -.09       | .42        | 75      | -.15  | .21  | <b>77</b> | <b>-.22</b> | <b>.05</b>    | <b>327</b> | <b>-.24</b> | <b>&lt;.01</b> |
|                     | +age   | <b>82</b> | <b>-.32</b> | <b>.03</b>    | 93        | -.11       | .35        | 75      | -.04  | .80  | <b>77</b> | <b>-.40</b> | <b>.02</b>    | <b>327</b> | <b>-.19</b> | <b>&lt;.01</b> |
|                     | +n-3/n-6 <sup>2</sup>                        | <b>82</b> | <b>-.37</b> | <b>.01</b>    | 75        | -.10       | .43        | 75      | -.07  | .62  | <b>77</b> | <b>-.27</b> | <b>.03</b>    | <b>309</b> | <b>-.21</b> | <b>&lt;.01</b> |
|                     | +age+n-3/n-6                                 | <b>82</b> | <b>-.19</b> | <b>.28</b>    | 75        | -.13       | .35        | 75      | .011  | .95  | <b>77</b> | <b>-.41</b> | <b>.01</b>    | <b>309</b> | <b>-.18</b> | <b>.02</b>     |

The analyses were performed on ln transformed data with AhRcomp as dependent and POPs as independent variables before and upon adjusted for the potential confounders.

<sup>1</sup>: non-adjusted data.

<sup>2</sup>: ratio of n-3 to n-6 fatty acids in serum.

<sup>a</sup>: Combining districts is not allowed according to the multiple regression analyses.

<sup>b</sup>: Combining districts is allowed according to the multiple regression analyses.

<sup>c</sup>: Combining districts and sex is not allowed according to the multiple regression analyses.

For Σ PCB<sub>14</sub>, Σ pesticide and Σ POP and AhRcomp, see the legend of Table 2 and Table 3.

be one source of DL-compounds and/or AhR activating compounds for the Inuits living in West Greenland.

### 3.5.3. AhRcomp activity

The AhRcomp data was also in overall inversely related to age and/or the n3/n6 ratio, but also the seabird intake seemed to be a determinant. A negative correlation for

AhRcomp to seabird intake was observed for the Qaanaaq men and all men data (Table 6). Moreover, the AhRcomp of all men was found to be inversely correlated to age, n-3/n-6 ratio and smoking years. Negative correlations between AhRcomp and the n-3/n-6 ratio and age and smoking years were found for the women of Nuuk and Qaanaaq, respectively. For the combined sex data of Nuuk subjects, negative correlations between AhRcomp and age, n-3/n-6 ratio and seabird intake were found (Table 6). By combining all data (districts and sexes), AhRcomp was found to be negatively correlated to age and n3/n6 ratio (Table 6). Besides, seabird intake was found to be negatively correlated to AhRcomp for all men data, the

total combined data of Nuuk, Sisimiut and Qaanaaq groups (Table 6).

**4. Discussion**

In the present study we compared the *ex vivo* net integrated AhR transcriptional activity of the lipophilic serum POP mixture of Inuits among different Greenlandic districts and evaluated whether the serum AhR transcriptional activity was related to the bioaccumulated POPs and/or lifestyle factors. The serum levels of POPs, AhR-TEQ and AhRcomp activity differed among the two sexes living in different Greenlandic districts (Northwest-Qaanaaq, South West-Nuuk and Sisimiut and East-Tasiilaq). Also the association of the AhR transcriptional activ-

**Table 6: The Pearson correlation coefficients of serum AhR-TEQ/AhRcomp and life style factors**

|             | Men        |             |            |            |             |                | Women     |             |               |           |             |            | Men + Women |             |            |            |             |                |
|-------------|------------|-------------|------------|------------|-------------|----------------|-----------|-------------|---------------|-----------|-------------|------------|-------------|-------------|------------|------------|-------------|----------------|
|             | AhR-TEQ    |             |            | AhRcomp    |             |                | AhR-TEQ   |             |               | AhRcomp   |             |            | AhR-TEQ     |             |            | AhRcomp    |             |                |
| Nuuk        | n          | r           | p          | n          | r           | p              | n         | r           | p             | n         | r           | p          | n           | r           | p          | n          | r           | p              |
| age         | 20         | -.08        | .74        | 37         | -.02        | .91            | 11        | .12         | .72           | 44        | -.13        | .41        | 32          | -.19        | .28        | <b>82</b>  | <b>-.53</b> | <b>&lt;.01</b> |
| n-3/n-6     | 20         | -.01        | .97        | 37         | -.02        | .91            | 11        | -.32        | .32           | <b>44</b> | <b>-.30</b> | <b>.05</b> | 32          | -.26        | .15        | <b>82</b>  | <b>-.50</b> | <b>&lt;.01</b> |
| BMI         | 19         | .40         | .08        | 35         | .24         | .15            | 11        | .07         | .84           | 42        | -.19        | .22        | 31          | .13         | .48        | 78         | -.17        | .14            |
| smoking     | 19         | .04         | .86        | 32         | .12         | .52            | 8         | .56         | .15           | 34        | .22         | .21        | 27          | .13         | .53        | 66         | -.22        | .08            |
| seabird     | 19         | .001        | 1.0        | 34         | .01         | .94            | 8         | -.47        | .20           | 37        | -.09        | .56        | 28          | -.19        | .32        | <b>72</b>  | <b>-.43</b> | <b>&lt;.01</b> |
| diary food  | 20         | -.26        | .25        | 36         | -.12        | .48            | 11        | .34         | .29           | 44        | .014        | .93        | 32          | .03         | .85        | 81         | .19         | .09            |
| Sisimiut    |            |             |            |            |             |                |           |             |               |           |             |            |             |             |            |            |             |                |
| age         | <b>41</b>  | <b>-.32</b> | <b>.04</b> | 50         | -.21        | .14            | 31        | .08         | .65           | 41        | .21         | .18        | 73          | -.18        | .12        | 92         | .01         | .91            |
| n-3/n-6     | 29         | -.06        | .75        | 36         | -.31        | .06            | 27        | .21         | .28           | 37        | .29         | .07        | 57          | -.10        | .48        | 74         | .08         | .55            |
| BMI         | 41         | -.05        | .76        | 50         | -.16        | .28            | <b>31</b> | <b>.52</b>  | <b>&lt;.0</b> | 41        | -.18        | .27        | 73          | .19         | .10        | 92         | -.16        | .14            |
| smoking     | <b>35</b>  | <b>-.36</b> | <b>.03</b> | 43         | -.12        | .44            | 29        | .11         | .56           | 33        | .01         | .94        | 64          | -.15        | .22        | 76         | -.03        | .78            |
| seabird     | 38         | -.01        | .97        | 46         | -.12        | .43            | 28        | -.17        | .38           | 38        | -.05        | .74        | 67          | -.05        | .72        | 85         | -.12        | .28            |
| diary food  | 40         | .15         | .37        | 49         | .06         | .69            | 30        | -.003       | .99           | 40        | .23         | .15        | 71          | .09         | .44        | 90         | .14         | .18            |
| Qaanaaq     |            |             |            |            |             |                |           |             |               |           |             |            |             |             |            |            |             |                |
| age         | 37         | -.28        | .08        | 42         | -.11        | .48            | 31        | -.33        | .07           | <b>35</b> | <b>-.34</b> | <b>.05</b> | <b>69</b>   | <b>-.30</b> | <b>.01</b> | 78         | -.21        | .66            |
| n-3/n-6     | 37         | -.002       | .99        | 42         | -.16        | .30            | <b>31</b> | <b>-.42</b> | <b>.02</b>    | 35        | -.06        | .74        | 69          | -.17        | .15        | 78         | -.11        | .31            |
| BMI         | 37         | -.02        | .49        | 42         | -.08        | .60            | 31        | -.11        | .54           | 35        | .04         | .83        | 69          | -.13        | .28        | 78         | -.02        | .87            |
| smoking     | 33         | -.12        | .49        | 37         | -.12        | .49            | 29        | -.23        | .22           | <b>33</b> | <b>-.35</b> | <b>.05</b> | 62          | -.20        | .12        | 70         | -.21        | .08            |
| seabird     | 35         | -.19        | .27        | <b>39</b>  | <b>-.33</b> | <b>.04</b>     | 30        | -.05        | .81           | 34        | -.23        | .19        | 66          | -.01        | .37        | 74         | -.28        | .17            |
| diary food  | 37         | -.02        | .91        | 42         | .15         | .34            | 29        | -.01        | .96           | 35        | .14         | .42        | 67          | .001        | .99        | 76         | .14         | .22            |
| Tasiilaq    |            |             |            |            |             |                |           |             |               |           |             |            |             |             |            |            |             |                |
| age         | 37         | -.19        | .25        | 37         | .05         | .76            | 43        | .04         | .81           | 43        | .10         | .53        | 81          | -.07        | .54        | 81         | .03         | .76            |
| n-3/n-6     | 37         | .26         | .12        | 37         | .01         | .95            | 43        | .12         | .90           | 43        | .02         | .91        | 81          | .12         | .28        | 81         | .007        | .95            |
| BMI         | 37         | -.002       | .91        | 37         | -.002       | .99            | 43        | -.27        | .08           | 43        | -.08        | .60        | 81          | -.19        | .10        | 81         | -.07        | .52            |
| smoking     | 33         | .01         | .97        | 33         | -.003       | .99            | 41        | .06         | .72           | 41        | .05         | .75        | 74          | .03         | .79        | 74         | .02         | .90            |
| seabird     | na         | na          | na         | na         | na          | na             | na        | na          | na            | na        | na          | na         | na          | na          | na         | na         | na          | na             |
| diary food  | na         | na          | na         | na         | na          | na             | na        | na          | na            | na        | na          | na         | na          | na          | na         | na         | na          | na             |
| All         |            |             |            |            |             |                |           |             |               |           |             |            |             |             |            |            |             |                |
| age         | <b>138</b> | <b>-.19</b> | <b>.03</b> | <b>169</b> | <b>-.35</b> | <b>&lt;.01</b> | 119       | -.09        | .34           | 166       | .03         | .66        | <b>258</b>  | <b>-.16</b> | <b>.01</b> | <b>336</b> | <b>-.20</b> | <b>&lt;.01</b> |
| n-3/n-6     | 126        | .10         | .27        | <b>155</b> | <b>-.20</b> | <b>.01</b>     | 115       | -.12        | .19           | 162       | -.04        | .59        | 242         | -.02        | .81        | <b>318</b> | <b>-.16</b> | <b>.01</b>     |
| BMI         | 137        | -.02        | .86        | 167        | -.01        | .77            | 119       | .01         | .87           | 164       | -.02        | .23        | 257         | -.01        | .93        | 332        | -.09        | .10            |
| smoking     | 120        | -.17        | .07        | <b>145</b> | <b>-.23</b> | <b>.01</b>     | 107       | -.01        | .96           | 141       | .04         | .68        | 227         | -.11        | .11        | 286        | -.11        | .06            |
| seabird*    | 94         | -.07        | .47        | <b>121</b> | <b>-.35</b> | <b>&lt;.01</b> | 68        | -.20        | .10           | 111       | -.13        | .16        | 163         | -.12        | .12        | <b>233</b> | <b>-.33</b> | <b>&lt;.01</b> |
| diary food* | 99         | .13         | .18        | 129        | .02         | .79            | 72        | .21         | .08           | 119       | .13         | .15        | <b>172</b>  | <b>.17</b>  | <b>.03</b> | 249        | .11         | .08            |

na: not available

\*: For the combined data of Nuuk, Sisimiut and Qaanaaq.

ity to the determined POPs differed between the districts and genders. We observed for the combined data an inverse association between AhR-TEQ and POPs where the Qaanaaq women had main influence. The negative association of AhRcomp activity to POPs was mainly contributed by Nuuk and Tasiilaq subjects.

As reported [10], the POPs levels of Tasiilaq and Qaanaaq participants (including men and women) were in general higher than that of Nuuk and Sisimiut subjects, which is consistent with previous reports that inhabitants of East Greenland had higher POP levels than those in the West Greenland [6,8,41] and in accordance with POP levels in the marine biota of North West Greenland and East Greenland being higher than that of the Central West Coast [41]. A greater prevalence of western lifestyle habits (consumption of imported food) was reported for Nuuk and Sisimiut subjects [10,12], which can explain the lower POP burden in Sisimiut and Nuuk subjects compared to Qaanaaq and Tasiilaq. However, in the present study, Nuuk men elicited higher POP levels than men from other districts, which may be due to their higher age and high intake of seafood and seabird [10,42].

The pattern of regional differences of serum AhR-TEQ levels did not follow the district-distribution of POPs levels. The AhR-TEQ level of Nuuk men having the highest POP level was lower than that of Sisimiut men with the lowest POPs level, whereas Qaanaaq women with relatively higher POPs level elicited the lowest AhR-TEQ level. However, the regional differences of AhR-TEQ seemed to mimic the district variation of the ratio of the  $\Sigma$ DL-PCB to the  $\Sigma$ PCB<sub>14</sub>. For both sexes, the DL-PCB proportion of Qaanaaq subjects was in general lower than that of Nuuk, Tasiilaq and Sisimiut (men: Nuuk > Tasiilaq  $\geq$  Qaanaaq  $\geq$  Sisimiut; women: Sisimiut  $\geq$  Nuuk  $\geq$  Tasiilaq  $\geq$  Qaanaaq) and consistently lower AhR-TEQ level was observed for Qaanaaq men and women (men: Tasiilaq  $\geq$  Sisimiut  $\geq$  Nuuk > Qaanaaq; women: Nuuk  $\geq$  Tasiilaq  $\geq$  Sisimiut > Qaanaaq). Similarly, male participants in all districts had higher POP burden but lower DL-PCB proportion and accordingly a tendency of lower AhR-TEQ level compared to female was observed in the present study. Moreover, we also found that individuals with higher  $\Sigma$ DL-PCB/ $\Sigma$ PCB<sub>14</sub> ratio tend to have higher AhR-TEQ levels while those with higher  $\Sigma$  POP have significantly lower AhR-TEQ than individuals with lower  $\Sigma$  POP (data not shown). Our study further support the previous report that DL-PCBs were major contributors of the total WHO-TEQ concentration in the plasma of Arctic Inuits [43]. Therefore, the total POPs level may not be a suitable proxy of the DL-compounds levels and thus the composition of the POPs in the body must be taken into account for the assessment of the net AhR transcriptional activity of serum samples [27].

Ayotte and coworkers reported that the average plasma WHO-TEQ level of Inuit adults living in Arctic Quebec was 184 pg/g lipid [43], being compatible with the serum AhR-TEQ level of Greenlandic Inuits determined in the present study (161 pg/g lipid). Previously, we reported a lower AhR-TEQ level for Greenlandic Inuit males (197 pg/g lipid) than for European men from Sweden, Warsaw and Kharkiv with the average serum AhR-TEQ level of 317 pg/g lipid [27]. The present study further supports that Greenlandic Inuits have lower level of AhR-TEQ than European individuals. The lower level of AhR-TEQ in Inuits compared to Europeans might be due to lower level of DL-compounds and lipophilic AhR activating compounds as well as higher level of AhR inhibiting POPs in the Inuit samples in addition to the report of lower level of PCDD/Fs in the Arctic region versus Europe and America [44]. Furthermore, re-evaluation of our previous study [27] by stratifying on the Inuits and European Caucasians showed a significantly positive correlation of POP proxy marker CB-153 and AhR-TEQ in European Caucasians ( $r_s = 0.20$ ,  $p = 0.001$ ), combined with their higher AhR-TEQ level, suggesting European Caucasians were more close to DL-compounds sources.

Knowing that the measured net integrated AhR induced activity of the lipophilic serum extract are in the linear range (Bonefeld-Jorgensen and Long, submitted), and running a TCDD dose-response measurement in parallel, we suggest that the chemical mixture in serum extract can further increase or inhibit the TCDD induced AhR mediated activity (termed as AhRcomp). In the present study, the AhRcomp differed between districts and between sexes. Compared to Qaanaaq and Sisimiut men, Nuuk and Tasiilaq men had the lowest AhRcomp median level being in accordance with the higher frequency of samples eliciting inhibition of the TCDD induced AhR activity, suggesting the presence of compounds having the potency to compete with TCDD for the AhR activation. The higher incidence of further increase of the TCDD induced effect observed for Sisimiut and Qaanaaq men as well as women from all districts suggest the presence of compounds interacting with TCDD for AhR mediated effects. The differences in the incidence of further increase and inhibition of TCDD induced activity might be caused by the different POP composition of Inuits from the different districts and for genders also differences in diet intake might have an impact on the AhR function [8,10].

Humans are exposed to a combination of several xenobiotics. It is likely that the interaction between these contaminants, additivity and/or synergy and/or antagonism, will be reflected in the final toxic effect. *In vivo* and *in vitro* studies have reported that PCB mixtures and some individual PCB congeners antagonized TCDD effects [45,46], and Arochlor 1254 was reported to inhibit TCDD induced

7-ethoxyresorufin O-deethylase (EROD) activity [47]. A mixture of PCBs, based on the congeners identified in human milk, significantly decreased TCDD induced rat liver EROD activity [45]. For the POPs determined in the serum extract in this study, only 3 DL-PCBs (CB105, CB118 and CB156) were shown to have agonistic AhR potentials [1,17,23], while other PCBs such as PCB153, PCB 52, PCB128 have the potential to antagonize the AhR pathway [20,48] and HCB was also reported to inhibit the TCDD induced activity [49]. Chlordane was reported to decrease the basal and TCDD induced CYP 1A1 promoter activity in MCF-7 cells [50]. Furthermore, most of the determined pesticides and PCBs in this study were reported as xenoestrogens (p,p'-DDT, p,p'-DDE, toxaphene, chlordane) [51-54] or antiestrogens (CB138, CB153, CB180) [55,56]. Previous studies showed that o,p'-DDT and other xenoestrogenic pesticides such as endosulfan and dieldrin significantly decreased the basal and TCDD induced AhR mediated gene expression [50,57]. Moreover, high concentrations of weak agonists can lead to antagonism [21,58]. Also, there may be other forms of antagonism when there are large concentrations of other compounds relative to high affinity ligands. Non-DL-PCBs can both antagonize and/or synergize the effects of DL-compounds, it depends on the response, the dose, and the concentrations [48,59]. As humans are exposed to a complex mixture and many PCBs can act as AhR antagonists most likely in concert when present together, this could actually influence the overall effect of DL-compounds by decreasing the net serum AhR transcriptional activity. Since the serum AhR transcriptional activity determined in the present study is the net integrated effect of the AhR agonists and antagonists of the constituents in the mixture, the negative correlation of AhR-TEQ, AhR-comp and POPs/PCBs/pesticides observed might be due to a higher concentration of compounds having antagonistic activity on AhR transactivity.

The biomagnification of PCBs in the aquatic food-chain is congener-specific. For most marine mammals, non-DL-PCBs are observed more resistant to biodegradation than DL-PCBs [60] and a previous study reported that the concentration of CB77, CB126 and CB169 showed no obvious biomagnification although the concentration of non-DL-PCBs such as CB138 increased from plankton to piscivores to herring gulls [61]. The lack of biomagnification of CB77 was attributed to its rapid elimination by aquatic species [62] and the CB105 and CB118 are more easily metabolized congeners than non-DL-PCBs such as CB153 [63]. Owing to this selective biotransformation during the bioaccumulation in the food chain [10], the concentration of some DL-PCBs might be reduced, whereas non-DL-PCBs accumulate, causing the variation of the composition of bioaccumulated PCBs.

In overall, the serum AhR-TEQ/AhRcomp negatively correlated to age and n-3/n-6 ratio, being in accordance to the frequency of marine food intake and POPs bioaccumulation over time. For the separate districts and genders a scattered correlation of AhR-TEQ/AhRcomp and lifestyle factors was observed. This non-consistent pattern might be explained by relative low statistical power for the separate gender and/or districts and needs further confirmation.

The inverse correlation of AhR-TEQ to seafood intake given by the n-3/n-6 ratio, suggests that the AhR transcriptional activity might follow the intake of n-6 fatty acids possibly with the origin from non-marine diet being supported by our previous report (Bonefeld-Jorgensen and Long, submitted). Moreover, it was reported that the important source of DL-compounds for the population of the western world is dairy food intake [64-66]. In the present study, we observed that the consumption of dairy food was higher for the Nuuk and Sisimiut participants than that of Qaanaaq and a positive correlation between dairy food intake and serum AhR-TEQ level was observed for the combined Nuuk and Sisimiut data and the combined Nuuk, Sisimiut and Qaanaaq data, respectively. Being more westernized, the import of western food and intake of milk products was reported to be higher for the subjects living in Nuuk and Sisimiut, while the intake of local Greenlandic products was relatively higher in Qaanaaq [10]. Hence, the dairy food might be an important source of DL-compounds for the Inuits living in bigger cities such as Nuuk and Sisimiut, relating to the higher AhR-TEQ level observed for Nuuk and Sisimiut compared to Qaanaaq. Moreover, we observed that POPs increase significantly with age for Inuits in this study as reported [10] as well as for Inuits and Europeans in an earlier study [67]. Evaluation of our previous data [27] showed that in Europeans the AhR-TEQ positively correlated to age ( $r = 0.12$ ,  $p = 0.05$ ), whereas for Inuits negative correlation was observed ( $r = -0.3$ ,  $p = 0.01$ ), suggesting different POP profiles resulted in different outcome on the AhR transcriptional activity.

Although the correlations found in the present study were weak, the chance finding may be excluded because the significant correlations were higher than 5%.

In the present study, the AhR transcriptional activation of serum extract was measured as an integrated net AhR-TEQ value without providing information on the levels of individual congeners. This bioassay responds to environmental contaminants acting through the AhR-pathway and allows the detection of chemicals that are not included in the chemical analytical method based on the standard TEF scheme. However, as discussed previously [27], upon serum extraction and clean-up the polycyclic aromatic

hydrocarbons (PAHs) and other plant dietary AhR agonists are not expected to contribute to our determined AhR transcriptional activity (personal communication, Pierre Dumas, Quebec, Canada). The CALUX bioassay including AhR transcription activation assay takes into account all possible interactions (synergistic, additive and/or antagonistic interaction) between congeners while chemical analytical method like GC-MS focuses on a selected numbers of compounds (i.e. those documented to produce AhR-dependent toxicity) and only assumes the additive interaction of known detectable compounds. Therefore, the AhR-TEQ value derived from CALUX bioassays (AhR transcription activation bioassay) and chemical methods (WHO-TEQ) can not be expected to be equal. In most of the cases, higher AhR-TEQ value are usually obtained with AhR-CALUX bioassays compared to the calculated WHO-TEQ derived from chemical method [18,68]. However, the AhR-TEQ and WHO-TEQ values derived from AhR transcriptional activation CALUX bioassays and chemical analyses, respectively, are correlated [68] and these two methods are complimentary where the AhR transcription activation CALUX bioassay provides an overall biological response of the mixture, whereas the chemical analysis provides the concentration of specific compounds. Thus the key application of AhR transcriptional activation CALUX bioassay is screening and prioritization of samples [28] and is a valuable tool for comparison of different level of AhR transcriptional activity including DL-activity across various population.

## 5. Conclusion

The POPs and serum AhR transcriptional activity differed among Greenlandic districts, suggesting that the difference of serum AhR transcriptional activity depends on the composition of bioaccumulated POPs. The observed inverse correlation of POPs and AhR-TEQ or AhRcomp data suggests the presence of compounds with antagonistic impact on the AhR signalling pathway, probably due to the selective PCB bioaccumulation in the food chain. Further study is needed to elucidate this preliminary conclusion.

## Abbreviations

AhR aryl hydrocarbon receptor

AhRag agonistic AhR activity

AhRcomp competitive AhR activity

CALUX Chemical activated luciferase gene expression

DDT 2, 2-bis (p-chlorophenyl)-1, 1, 1-trichloroethane

DL dioxin-like

GC-MS gas chromatography – mass spectrometry

HCB hexachlorobenzene

β-HCH Hexachlorocyclohexane

PAHs polycyclic aromatic hydrocarbons

PCDDs/PCDFs polychlorinated dibenzo-p-dioxins/furans

PCBs polychlorinated biphenyls

POPs persistent organic pollutants

TCDD 2,3,7,8-tetrachlorodibenzo-p-dioxin

TEFs Toxic Equivalency Factors

TEQs TCDD toxic equivalents

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

ML and EBJ were responsible for design of the AhR transcriptional activity study and preparation of the manuscript; ML performed the mechanistic work and data evaluation. BD has provided the epidemiological data concerning POPs, questionnaires and fatty acids. All authors have read and approved the final manuscript.

## Additional material

### Additional file 1

*Multiple regressions of the combined study groups. The data provided represents the statistical analysis of the homogeneity/heterogeneity of the associations between POPs and AhR-TEQ/AhRcomp among the districts and genders.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1476-069X-6-32-S1.pdf>]

## Acknowledgements

We thank technical assistant Birgitte S Andersen and Inger Sørensen for their excellent skills in the laboratory work and all CMT group members for scientific support. We thank Jens C Hansen for comments on the manuscript and Henning S Pedersen for sampling. This study was supported by DANCEA, The Danish Environmental Ministry as a part of the "AMAP Human Health Effects Monitoring programme", and by the EU commission (INUENDO project).

## References

- Van den Berg M, Birnbaum L, Bosveld AT, Brunstrom B, Cook P, Feeley M, Giesy JP, Hanberg A, Hasegawa R, Kennedy SW, Kubiak T, Larsen JC, van Leeuwen FX, Liem AK, Nolt C, Peterson RE, Poellinger L, Safe S, Schrenk D, Tillitt D, Tysklind M, Younes M, Waern F, Zacharewski T: **Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife.** *Environ Health Perspect* 1998, **106(12)**:775-792.
- Safe SH: **Polychlorinated biphenyls (PCBs): environmental impact, biochemical and toxic responses, and implications for risk assessment.** *Crit Rev Toxicol* 1994, **24(2)**:87-149.
- Vorkamp K, Riget F, Glasius M, Pecseli M, Lebeuf M, Muir D: **Chlorobenzenes, chlorinated pesticides, coplanar chlorobiphenyls and other organochlorine compounds in Greenland biota.** *Sci Total Environ* 2004, **331(1-3)**:157-175.
- Barrie LA, Gregor D, Hargrave B, Lake R, Muir D, Shearer R, Tracey B, Bidleman T: **Arctic contaminants: sources, occurrence and pathways.** *Sci Total Environ* 1992, **122(1-2)**:1-74.
- Deutch B, Hansen JC: **High human plasma levels of organochlorine compounds in Greenland. Regional differences and lifestyle effects.** *Dan Med Bull* 2000, **47(2)**:132-137.
- Van Oostdam JC, Dewailly E, Gilman A, Hansen JC, Odland JO, Chashchin V, Berner J, Butler-Walker J, Lagerkvist BJ, Olafsdottir K, Soininen L, Bjerregard P, Klopov V, Weber JP: **Circumpolar maternal blood contaminant survey, 1994-1997 organochlorine compounds.** *Sci Total Environ* 2004, **330(1-3)**:55-70.
- Bonefeld Jorgensen EC, Ayotte P: **Toxicological properties of persistent organic pollutants and related health effects of concern for the arctic populations.** *AMAP Assessment 2002: Human Health in the Arctic 2003, chapter 6*:57-74.
- Deutch B, Pedersen HS, Hansen JC: **Dietary composition in Greenland 2000, plasma fatty acids and persistent organic pollutants.** *Sci Total Environ* 2004, **331(1-3)**:177-188.
- Deutch B, Dyerberg J, Pedersen HS, Asmund G, Moller P, Hansen JC: **Dietary composition and contaminants in north Greenland, in the 1970s and 2004.** *Sci Total Environ* 2006, **370**:372-381.
- Deutch B, Pedersen HS, Asmund G, Hansen JC: **Contaminants, diet, plasma fatty acids and smoking in Greenland 1999-2005.** *Sci Total Environ* 2007, **372(2-3)**:486-496.
- Kinloch D, Kuhnlein H, Muir DC: **Inuit foods and diet: a preliminary assessment of benefits and risks.** *Sci Total Environ* 1992, **122(1-2)**:247-278.
- Dewailly E, Mulvad G, Pedersen HS, Ayotte P, Demers A, Weber JP, Hansen JC: **Concentration of organochlorines in human brain, liver, and adipose tissue autopsy samples from Greenland.** *Environ Health Perspect* 1999, **107(10)**:823-828.
- Steenland K, Bertazzi P, Baccarelli A, Kogevinas M: **Dioxin revisited: developments since the 1997 IARC classification of dioxin as a human carcinogen.** *Environ Health Perspect* 2004, **112(13)**:1265-1268.
- Lindstrom G, Hooper K, Petreas M, Stephens R, Gilman A: **Workshop on perinatal exposure to dioxin-like compounds. I. Summary.** *Environ Health Perspect* 1995, **103 Suppl 2**:135-142.
- Rowlands JC, Gustafsson JA: **Aryl hydrocarbon receptor-mediated signal transduction.** *Crit Rev Toxicol* 1997, **27(2)**:109-134.
- Safe S, Krishnan V: **Cellular and molecular biology of aryl hydrocarbon (Ah) receptor-mediated gene expression.** *Arch Toxicol Suppl* 1995, **17**:99-115.
- Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, Fiedler H, Hakansson H, Hanberg A, Haws L, Rose M, Safe S, Schrenk D, Tohyama C, Tritscher A, Tuomisto J, Tysklind M, Walker N, Peterson RE: **The 2005 World Health Organization reevaluation of human and Mammalian toxic equivalency factors for dioxins and dioxin-like compounds.** *Toxicol Sci* 2006, **93(2)**:223-241.
- Van Overmeire I, Clark, G.C., Brown, D.J., Chu, M.D., Cooke, W. M., Denison, M.S., Baeyene, W., Srebrnik S., Goeyens, L.: **Trace contamination with dioxin-like chemicals: evaluation of bioassay-based TEQ determination for hazard assessment and regulatory responses.** *Environmental Science & Policy* 2001, **4(6)**:345-357.
- Laier P, Cederberg, T, Larsen, J.C., Vinggaard, A.M.: **Applicability of the CALUX bioassay for screening of dioxin levels in human milk samples.** *Food Additives and Contaminants* 2003, **20(6)**:583-595.
- Aarts JM, Denison MS, Cox MA, Schalk MA, Garrison PM, Tullis K, de Haan LH, Brouwer A: **Species-specific antagonism of Ah receptor action by 2,2',5,5'-tetrachloro- and 2,2',3,3',4,4'-hexachlorobiphenyl.** *Eur J Pharmacol* 1995, **293(4)**:463-474.
- Long M, Laier P, Vinggaard AM, Andersen HR, Lynggaard J, Bonefeld-Jorgensen EC: **Effects of currently used pesticides in the AhR-CALUX assay: comparison between the human TV101L and the rat H4IIE cell line.** *Toxicology* 2003, **194(1-2)**:77-93.
- Behnisch PA, Hosoe K, Sakai S: **Bioanalytical screening methods for dioxins and dioxin-like compounds: a review of bioassay/biomarker technology.** *Environ Int* 2001, **27(5)**:413-439.
- van den Berg M, Peterson RE, Schrenk D: **Human risk assessment and TEFs.** *Food Addit Contam* 2000, **17(4)**:347-358.
- Garrison PM, Tullis K, Aarts JM, Brouwer A, Giesy JP, Denison MS: **Species-specific recombinant cell lines as bioassay systems for the detection of 2,3,7,8-tetrachlorodibenzo-p-dioxin-like chemicals.** *Fundam Appl Toxicol* 1996, **30(2)**:194-203.
- Murk AJ, Legler J, Denison MS, Giesy JP, van de Guchte C, Brouwer A: **Chemical-activated luciferase gene expression (CALUX): a novel in vitro bioassay for Ah receptor active compounds in sediments and pore water.** *Fundam Appl Toxicol* 1996, **33(1)**:149-160.
- Aarts JMMJG, Cenijn PH, Blankvoort BMG, Murk A, Brouwer A, Bovee TFH, Traag WA, Hoogenboom LAP, Patandin S, Weisglas-Kuperus N, Sauer PJJ, Denison MS: **Application of the chemical-activated luciferase expression (CALUX) bioassay for quantification of dioxin-like compound in small samples of human milk and blood plasma.** *Organohal Compds* 1996, **27**:285-290.
- Long M, Andersen BS, Lindh CH, Hagmar L, Giwercman A, Manicardi GC, Bizzaro D, Spano M, Toft G, Pedersen HS, Zvezday V, Bonde JP, Bonefeld-Jorgensen EC: **Dioxin-like activities in serum across European and Inuit populations.** *Environ Health* 2006, **5**:14.
- Windal I, Denison MS, Birnbaum LS, Van Wouwe N, Baeyens W, Goeyens L: **Chemically activated luciferase gene expression (CALUX) cell bioassay analysis for the estimation of dioxin-like activity: critical parameters of the CALUX procedure that impact assay results.** *Environ Sci Technol* 2005, **39(19)**:7357-7364.
- Brouwer A, Ahlborg UG, Van den Berg M, Birnbaum LS, Boersma ER, Bosveld B, Denison MS, Gray LE, Hagmar L, Holene E, et al.: **Functional aspects of developmental toxicity of polyhalogenated aromatic hydrocarbons in experimental animals and human infants.** *Eur J Pharmacol* 1995, **293(1)**:1-40.
- AMAP: **Arctic Monitoring and Assessment Programme.** <http://www.amap.no>.
- Butler Walker J, Seddon L, McMullen E, Houseman J, Tofflemire K, Corriveau A, Weber JP, Mills C, Smith S, Van Oostdam J: **Organochlorine levels in maternal and umbilical cord blood plasma in Arctic Canada.** *Sci Total Environ* 2003, **302(1-3)**:27-52.
- Deutch B, Pedersen HS, Jorgensen EC, Hansen JC: **Smoking as a determinant of high organochlorine levels in Greenland.** *Arch Environ Health* 2003, **58(1)**:30-36.
- Tjonneland A, Overvad K, Thorling E, Ewertz M: **Adipose tissue fatty acids as biomarkers of dietary exposure in Danish men and women.** *Am J Clin Nutr* 1993, **57(5)**:629-633.
- Ziccardi MH, Gardner IA, Denison MS: **Development and modification of a recombinant cell bioassay to directly detect halogenated and polycyclic aromatic hydrocarbons in serum.** *Toxicol Sci* 2000, **54(1)**:183-193.
- Bonefeld-Jorgensen EC, Grunfeld HT, Gjermansen IM: **Effect of pesticides on estrogen receptor transactivation in vitro: a comparison of stable transfected MVLN and transiently transfected MCF-7 cells.** *Mol Cell Endocrinol* 2005, **244(1-2)**:20-30.
- Covaci A, Koppen G, Van Cleuvenbergen R, Schepens P, Winneke G, van Larebeke N, Nelen V, Vlietinck R, Schoeters G: **Persistent organochlorine pollutants in human serum of 50-65 years old women in the Flanders Environmental and Health Study (FLEHS). Part 2: Correlations among PCBs, PCDD/PCDFs and the use of predictive markers.** *Chemosphere* 2002, **48(8)**:827-832.
- Pauwels A, Cenijn PH, Schepens PJ, Brouwer A: **Comparison of chemical-activated luciferase gene expression bioassay and gas chromatography for PCB determination in human serum and follicular fluid.** *Environ Health Perspect* 2000, **108(6)**:553-557.



38. Warner M, Eskenazi B, Patterson DG, Clark G, Turner WE, Bon-signore L, Mocarelli P, Gerthoux PM: **Dioxin-Like TEQ of women from the Seveso, Italy area by ID-HRGC/HRMS and CALUX.** *J Expo Anal Environ Epidemiol* 2004.
39. Ayotte P, Dewailly E, Lambert GH, Perkins SL, Poon R, Feeley M, Larochelle C, Pereg D: **Biomarker measurements in a coastal fish-eating population environmentally exposed to organochlorines.** *Environ Health Perspect* 2005, **113(10)**:1318-1324.
40. James RA, Hertz-Picciotto I, Willman E, Keller JA, Charles MJ: **Determinants of serum polychlorinated biphenyls and organochlorine pesticides measured in women from the child health and development study cohort, 1963-1967.** *Environ Health Perspect* 2002, **110(7)**:617-624.
41. Riget F, Dietz R, Vorkamp K, Johansen P, Muir D: **Levels and spatial and temporal trends of contaminants in Greenland biota: an updated review.** *Sci Total Environ* 2004, **331(1-3)**:29-52.
42. Johansen P, Muir D, Asmund G, Riget F: **Human exposure to contaminants in the traditional Greenland diet.** *Sci Total Environ* 2004, **331(1-3)**:189-206.
43. Ayotte P, Dewailly E, Ryan JJ, Bruneau S, Lebel G: **PCBs and dioxin-like compounds in plasma of adult Inuit living in Nunavik (Arctic Quebec).** *Chemosphere* 1997, **34(5-7)**:1459-1468.
44. Schlabach M, Biseth A, Gundersen H: **Sampling and measurement of PCDD/PCDF and non-ortho PCB in Arctic air at Ny-Ålesund, Spitzbergen.** *Organohal Compds* 1996, **28**:325-239.
45. Chu I, Lecavalier P, Hakansson H, Yagminas A, Valli VE, Poon P, Feeley M: **Mixture effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin and polychlorinated biphenyl congeners in rats.** *Chemosphere* 2001, **43(4-7)**:807-814.
46. Chen G, Bunce NJ: **Interaction between halogenated aromatic compounds in the Ah receptor signal transduction pathway.** *Environ Toxicol* 2004, **19(5)**:480-489.
47. Bannister R, Davis D, Zacharewski T, Tizard I, Safe S: **Aroclor 1254 as a 2,3,7,8-tetrachlorodibenzo-p-dioxin antagonist: effects on enzyme induction and immunotoxicity.** *Toxicology* 1987, **46(1)**:29-42.
48. Suh J, Kang JS, Yang KH, Kaminski NE: **Antagonism of aryl hydrocarbon receptor-dependent induction of CYP1A1 and inhibition of IgM expression by di-ortho-substituted polychlorinated biphenyls.** *Toxicol Appl Pharmacol* 2003, **187(1)**:11-21.
49. Schroiens C, Windal I, Goeyens L, Baeyens W: **Study of the interference problems of dioxin-like chemicals with the bio-analytical method CALUX.** *Talanta* 2004, **63**:1261-1268.
50. Coumoul X, Diry M, Robillot C, Barouki R: **Differential regulation of cytochrome P450 1A1 and 1B1 by a combination of dioxin and pesticides in the breast tumor cell line MCF-7.** *Cancer Res* 2001, **61(10)**:3942-3948.
51. Bonefeld Jorgensen EC, Autrup H, Hansen JC: **Effect of toxaphene on estrogen receptor functions in human breast cancer cells.** *Carcinogenesis* 1997, **18(8)**:1651-1654.
52. Bonefeld-Jorgensen EC: **The Human Health Effect Programme in Greenland, a review.** *Sci Total Environ* 2004, **331(1-3)**:215-231.
53. Welch RM, Levin W, Conney AH: **Estrogenic action of DDT and its analogs.** *Toxicol Appl Pharmacol* 1969, **14(2)**:358-367.
54. Cassidy RA, Vorhees CV, Minnema DJ, Hastings L: **The effects of chlordane exposure during pre- and postnatal periods at environmentally relevant levels on sex steroid-mediated behaviors and functions in the rat.** *Toxicol Appl Pharmacol* 1994, **126(2)**:326-337.
55. Bonefeld-Jorgensen EC, Andersen HR, Rasmussen TH, Vinggaard AM: **Effect of highly bioaccumulated polychlorinated biphenyl congeners on estrogen and androgen receptor activity.** *Toxicology* 2001, **158(3)**:141-153.
56. Rasmussen TH, Nielsen F, Andersen HR, Nielsen JB, Weihe P, Grandjean P: **Assessment of xenoestrogenic exposure by a biomarker approach: application of the E-Screen bioassay to determine estrogenic response of serum extracts.** *Environ Health* 2003, **2(1)**:12.
57. Jeong HG, Kim JY: **Effects of o,p'-DDT on the 2,3,7,8-tetrachlorodibenzo-p-dioxin-inducible CYP1A1 expression in murine Hepa-1c1c7 cells.** *Food Chem Toxicol* 2002, **40(11)**:1685-1692.
58. Bonefeld-Jorgensen EC, Long M, M.V. H, Vinggaard AM: **Endocrine disrupting potentials of bisphenol A, bisphenol A dimethylacrylate, 4-n-nonylphenol and 4-n-octylphenol in vitro-new data and a brief review.** *Environ Health Perspect* 2007, in press: <http://www.ehponline.org/docs/admin/newest.html>.
59. van Birgelen AP, Fase KM, van der Kolk J, Poiger H, Brouwer A, Seinen W, van den Berg M: **Synergistic effect of 2,2',4,4',5,5'-hexachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-p-dioxin on hepatic porphyrin levels in the rat.** *Environ Health Perspect* 1996, **104(5)**:550-557.
60. Boon JP, van der Meer J, Allchin CR, Law RJ, Klungsoyr J, Leonards PE, Spliid H, Storr-Hansen E, McKenzie C, Wells DE: **Concentration-dependent changes of PCB patterns in fish-eating mammals: structural evidence for induction of cytochrome P450.** *Arch Environ Contam Toxicol* 1997, **33(3)**:298-311.
61. IPCS: **Concise International Chemical Assessment Document 55: Polychlorinated biphenyls: human health aspects.** <http://www.inchem.org/documents/cicads/cicads/cicad55.htm> ;2003.
62. Koslowski SE, Metcalfe CD, Lazar R, Haffner GD: **The distribution of 42 PCBs, including three coplanar congeners, in the food web of the western basin of Lake Erie.** *Journal of Great Lakes Research* 1994, **20**:260-270.
63. Demers A, Ayotte P, Brisson J, Dodin S, Robert J, Dewailly E: **Plasma concentrations of polychlorinated biphenyls and the risk of breast cancer: a congener-specific analysis.** *Am J Epidemiol* 2002, **155(7)**:629-635.
64. Baars AJ, Bakker MI, Baumann RA, Boon PE, Freijer JJ, Hoogenboom LA, Hoogerbrugge R, van Klaveren JD, Liem AK, Traag WA, de Vries J: **Dioxins, dioxin-like PCBs and non-dioxin-like PCBs in foodstuffs: occurrence and dietary intake in The Netherlands.** *Toxicol Lett* 2004, **151(1)**:51-61.
65. Llobet JM, Domingo JL, Bocio A, Casas C, Teixido A, Muller L: **Human exposure to dioxins through the diet in Catalonia, Spain: carcinogenic and non-carcinogenic risk.** *Chemosphere* 2003, **50(9)**:1193-1200.
66. Taioli E, Marabelli R, Scortichini G, Migliorati G, Pedotti P, Cigliano A, Caporale V: **Human exposure to dioxins through diet in Italy.** *Chemosphere* 2005, **61(11)**:1672-1676.
67. Jonsson BA, Rylander L, Lindh C, Rignell-Hydbom A, Giwercman A, Toft G, Pedersen HS, Ludwicki JK, Goralczyk K, Zvezday V, Spano M, Bizzaro D, Bonefeld-Jorgensen EC, Manicardi GC, Bonde JP, Hagmar L: **Inter-population variations in concentrations, determinants of and correlations between 2,2',4,4',5,5'-hexachlorobiphenyl (CB-153) and 1,1-dichloro-2,2-bis(p-chlorophenyl)-ethylene (p,p'-DDE): a cross-sectional study of 3161 men and women from Inuit and European populations.** *Environ Health* 2005, **4**:27.
68. Van Wouwe N, Windal I, Vanderperren H, Eppe G, Xhrouet C, Mas-sart AC, Debacker N, Sasse A, Baeyens W, De Pauw E, Sartor F, Van Oyen H, Goeyens L: **Validation of the CALUX bioassay for PCDD/F analyses in human blood plasma and comparison with GC-HRMS.** *Talanta* 2004, **63**:1157-1167.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

