

IMMUNOLOGICAL UNRESPONSIVENESS TO  
THYMUS-INDEPENDENT ANTIGENS:  
TWO FUNDAMENTALLY DIFFERENT GENETIC  
MECHANISMS OF  
B-CELL UNRESPONSIVENESS TO DEXTRAN\*

BY CARMEN FERNANDEZ AND GÖRAN MÖLLER

*(From the Department of Immunobiology, Karolinska Institutet, Wallenberglaboratory, Lilla  
Frescati, 104 05 Stockholm 50, Sweden)*

It is well established that the ability to respond to certain antigens is under genetic control and several different types of control mechanisms are known. The genetic control of responses to thymus-dependent (TD) antigens usually affects IgG synthesis and two separate types have been observed. The immune response to certain antigens is under control of antigen-specific immune response genes, which are localized in the chromosome region determining the major histocompatibility antigens and they exert their effect on thymus-derived lymphocytes (1). These genes do not code for the variable region of the immunoglobulin receptors on the B cells and a normal response can be induced provided adequate T-cell help is given. A second type of genetic control to TD antigens is not antigen specific, only effects the quantity of the immune response, and is not controlled by histocompatibility-linked genes (2).

The immune response to thymus-independent (TI) antigens can also be under genetic control and also in this case two mechanisms, which primarily affect IgM synthesis, can be distinguished. One of these mechanisms concerns the ability of B cells to become activated by the polyclonal B-cell activating (PBA) properties inherent to TI antigens. In one case (C3H/HeJ), the gene(s) determine(s) the presence of the nonimmunoglobulin receptors on B cells for lipopolysaccharide, which are responsible for B-cell activation. The genes are codominantly expressed (3, 4) and localized on chromosome 4 in the mouse.<sup>2</sup> They do not code for the variable part of the immunoglobulin receptors. Another nonresponding strain to certain TI antigens (CBA/N) lacks PBA receptors or receptor-bearing cells capable of reacting to, for example, Ficoll, and cannot mount an immune response to any hapten coupled to this TI carrier (5). The gene is localized on the X chromosome. A second type of genetic control of the response to TI antigens concerns the  $\alpha$ -1-3 epitope of dextran (Dx) (6). The gene(s) determine(s) low immune re-

\* Supported by grants from the Swedish Cancer Society and Ministerio de Educación y Ciencia Espanol.

<sup>1</sup> Abbreviations used in this paper: BSS, balanced salt solution; Con A, concanavalin A; Dx, dextran; FITC, fluorescein isothiocyanate; HRC, horse erythrocytes; NNP, (4-hydroxy-3,5-dinitrophenyl)acetyl; PBA, polyclonal B-cell activator; PFC, plaque-forming cells; PHA, phytohemagglutinin; RBC, erythrocytes; SRBC, sheep erythrocytes; TD, thymus dependent; TI, thymus independent.

<sup>2</sup> J. Watson, K. Kelly, M. Largen, and B. A. Taylor. The genetic mapping of a defective LPS response gene in C3H/HeJ mice. Manuscript submitted for publication.

sponsiveness rather than absence of a response and at least one of the genes is closely linked to allotypes of the heavy chain locus (7).

We have found that different mouse strains vary in their ability to produce antibodies to the  $\alpha$ -1-6 epitope of Dx B512, and in certain strains a large proportion of mice are total nonresponders (8). Since Dx activates B cells in the absence of helper cells, such as T cells and macrophages (9), and since suppressor T cells do not affect this response (10), only two fundamental mechanisms can account for unresponsiveness, specifically: (a) lack of expression of V genes coding for antibodies against the  $\alpha$ -1-6 epitope and (b) lack of triggering receptors on the B cells possessing immunoglobulin receptors for the epitope. We will show that both types of unresponsiveness can occur in different mouse strains.

### Materials and Methods

Mice of the following strains were used in the present study: A/Sn, A.CA, A.SW, BALB/c, CBA, CBA/N, C3H/Tif, C3H/HeJ, C57BL, A.TH, A.TL, 4R, 2R, B10.5M, and certain hybrids among these strains.

*Antigens and Polyclonal Activators.* Native Dx from *Lactobacillus mesenteroides* B512 (average mol wt  $5-40 \times 10^6$  daltons), was obtained from ICN Pharmaceuticals Inc., Cleveland, Ohio. Other Dx preparations were obtained from Pharmacia, Uppsala, Sweden.

Native fluorescein isothiocyanate (FITC)-Dx was synthesized from native Dx B512 by reacting it with FITC, and was provided by Dr. van de Belder, Pharmacia. The final conjugation ratio was one molecule of FITC for every 200 glucose residues.

Lipopolysaccharide from *Escherichia coli* 055:B5 was prepared from phenol water extraction provided by professor T. Holme (Department of Bacteriology, Karolinska Institutet).

Concanavalin A (ConA) and phytohemagglutinin (PHA) were obtained from Pharmacia, Uppsala, Sweden, and Wellcome Reagents Limited, Beckenham, England, respectively.

*Preparation of Lymphocytes.* Spleens were removed and teased with forceps in ice-cold balanced salt solution (BSS). After brief sedimentation, the cells in the supernate were washed three times in 50 ml of cold BSS and subsequently suspended in culture medium to the desired cell concentration. Cellular viability was determined in a hemocytometer after staining the damaged cells with 0.02% trypan blue.

*Assay of Antibody Synthesis.* Anti- $\alpha$ -1-6 plaque-forming cells (PFC) were detected by a direct PFC assay with sheep erythrocytes (SRBC) with stearyl Dx B512 with a mol wt of 70,000 daltons, as described before by Howard et al. (11).

*Anti-FITC-PFC.* The coupling of FITC to erythrocytes (RBC) has been described before (12). For detection of high, medium, and low affinity PFC the following concentrations of FITC in carbonate bicarbonate buffer pH 9.2 were used: 0.05, (of 0.1) 0.5, and 5 mg/ml.

*Anti-NNP PFC.* RBC were coupled with (4-hydroxy-3,5-dinitrophenyl)acetyl (NNP) as described by Pasanen and Mäkelä (13). These cells were used to detect polyclonal antibody synthesis.

*Medium.* The medium used in most of the experiments was Eagle's Minimum Essential Medium in Earle's solution, supplemented with glutamine, non-essential amino acids, and pyruvate, and containing 100 IU of penicillin and 100  $\mu$ g of streptomycin/ml, as described by Mishell and Dutton (15). The medium was further buffered by 10 mM of HEPES and the pH adjusted to 7.2. All these reagents were obtained from Flow Laboratories, Irvine, Scotland. Most experiments were carried out in serum-free medium (14), except where specifically indicated in the figure legends.

*Induction of Polyclonal Antibody Synthesis.* Spleen cells were cultured serum-free in 3-cm diameter plastic Petri dishes or tubes (NUNC, Roskilde, Denmark) with a cell concentration of  $10^7$  cells/ml in 1 ml medium (15) set up in triplicate unless otherwise stated. The cultures were incubated at 37°C in plastic boxes filled with a mixture of 10% CO<sub>2</sub>, 83% N<sub>2</sub>, and 7% O<sub>2</sub>.

TABLE I  
*Immune Response to the  $\alpha$ -1-6 Epitope in 3-4 Mo Old Mice of Different Strains after Immunization with 2  $\mu$ g Native Dx 5 Days Earlier*

Strain	No. of mice	PFC/spleen against	
		SRBC	$\alpha$ -1-6*
CBA	16	90 $\pm$ 18	33,340 $\pm$ 4,900
C57BL	8	330 $\pm$ 100	46,440 $\pm$ 6,900
A.CA	8	178 $\pm$ 52	2,778 $\pm$ 1,679
A	17	105 $\pm$ 34	2,786 $\pm$ 695
A.TH	11	11 $\pm$ 9	1,036 $\pm$ 770
A.TL	9	33 $\pm$ 15	788 $\pm$ 377
CBA/N	6	2 $\pm$ 1	5 $\pm$ 3

\* The PFC response against  $\alpha$ -1-6 was tested against SRBC coated with stearyl Dx (40  $\mu$ g in 10 ml) to detect PFC of medium affinity.

### Results

*Strain Variability in the Response to the  $\alpha$ -1-6 Epitope.* The immune response to native Dx B512 ( $\alpha$ -1-6) was tested in 14 inbred strains and was found to be very strain dependent. High responder strains, such as C57BL and CBA gave 400-1,000 PFC/ $10^6$  cells 5 days after immunization with the optimal dose 1-2  $\mu$ g/mouse. In contrast, A, A.CA, A.SW, A.TH, A.TL, and CBA/N mice were found to be low or nonresponders to the  $\alpha$ -1-6 epitope (8). Over 200 A.CA were tested and about 90% of them only gave 0-10 PFC/ $10^6$  cells, which, at best, can be classified as an extremely low response. Other mouse strains, such as BALB/c were intermediate responders, exhibiting PFC values between the extremes reported above. Occasionally, A.CA mice exhibited a high response in terms of the number of PFC. This response was of very low affinity and the causes for this will be the topic of a separate report. A variable of importance in determining the level of the response was the age of the mice; young mice of all strains were nonresponders. However, when comparisons were made with mice of the same age (3-4 mo) the strain differences still existed (Table I). Since the mechanism of low responsiveness was found to be different with strains A, A.CA, A.SW, A.TH, and A.TL as compared with strain CBA/N, they will be dealt with separately.

*Mechanism of Unresponsiveness in Strains on A Background.* The low responsiveness of strains on A background could not be attributed to differences in dose requirements, affinity, or kinetics of the immune response, since A.CA mice failed to respond to the  $\alpha$ -1-6 epitope over a wide range of Dx or FITC Dx concentrations and at any time after immunization (Table II). Nonresponsiveness also affected the immune response to Dx conjugated to the TD carrier horse erythrocytes (HRC) (Table III) even in the presence of PHA as a polyclonal T-cell activator. A.CA mice did not produce anti- $\alpha$ -1-6 antibodies of noncomplement fixing classes, since sera from A.CA mice did not agglutinate Dx-coated RBC. The incorporation of native Dx with Freund's adjuvant did not make the mice responsive.

*Low Responsiveness is Not Due to Suppressor Cells.* The existence of

TABLE II  
*Dose Dependence, Kinetics, and Affinity of the Immune Response to Native Dx or Dx T 2000\* in High and Low Responder Strains*

Strain	Antigen	Dose ( $\mu\text{g}/\text{mouse}$ )	Day of test	PFC/ $10^6$ against $\alpha$ -1-6 by using the following indicator $\ddagger$ cells			
				4	40	400	
CBA	Native Dx	1	4	228	289	268	
				174	541	289	
				54	92	72	
		10	6	1	131	339	—
					100	108	133
					100	—	92
		10	15	1	64	183	143
					29	208	248
					100		
A.CA	Native Dx	1	4	1	2	1	
				3	2	6	
				4	4	8	
		10	6	1	2	4	1
					5	2	13
					1	2	1
		10	15	1	1	1	3
					1	10	10
					100	1	6
CBA	T 2000	1	4	192	217	129	
				202	327	199	
				47	127	107	
		10	6	1	270	237	210
					151	239	150
					230	—	180
		10	15	1	113	223	185
					105	327	646
					17	446	757
A.CA	T 2000	1	4	5	—	3	
				0	0	0	
				6	4	4	
		10	6	1	0	1	0
					2	1	0
					1	4	2
		10	15	1	2	28	41
					1	0	1
					1	28	31

\* Mol wt =  $2 \times 10^6$ .

$\ddagger$  SRBC coated with 4, 40, or 400  $\mu\text{g}/10$  ml of stearyl Dx during sensitization.

suppressor T cells capable of specifically suppressing an immune response has been reported in several different systems (16). Suppressor T cells were not involved here, since thymectomized, lethally irradiated A.CA mice that had been repopulated with syngeneic anti-theta serum-treated bone marrow failed to respond to native Dx (reference 8 and not shown).

It still seemed possible that the environment of A.CA mice contained suppressor elements for the anti- $\alpha$ -1-6 responding cells. To test this possibility,

TABLE III  
*Nonresponder Strains to  $\alpha$ -1-6 Cannot Respond to Dx Conjugated to TD RBC*

Strain	Antigen‡	Mitogen	PFC/spleen against	
			$\alpha$ -1-6*	HRC
A.CA	Dx 4-HRC	—	0	42,800
	Dx 40-HRC	—	50	48,000
	Dx 400-HRC	—	0	35,600
CBA $\times$ C57BL	Dx 4-HRC	—	200	180,800
	Dx 40-HRC	—	550	87,600
	Dx 400-HRC	—	4,550	144,000
A.CA	Dx 4000-HRC	—	0	23,600
	Dx 4000-HRC	PHA 100 $\mu$ g	400	40,800
	Dx 4000-HRC	PHA 200 $\mu$ g	50	33,500
	—	—	0	150
CBA $\times$ C57BL	Dx 4000-HRC	—	1,800	48,000
	Dx 4000-HRC	PHA 100 $\mu$ g	12,300	132,800
	Dx 4000-HRC	PHA 200 $\mu$ g	6,100	154,000
	—	—	0	275

\* The PFC response was tested against SRBC coated with stearyl Dx 40-SRBC (40  $\mu$ g stearyl Dx per 10 ml). The background to SRBC was deducted.

‡ Four different conjugates of stearyl Dx-HRC were used as antigen. The figures indicate the concentration in  $\mu$ g/10 ml of stearyl Dx used for conjugation.

A.CA mice were lethally irradiated and repopulated with spleen cells from a high responder strain (B10.5M or CBA) and immunized at the time of repopulation with native Dx and as a control with HRC. In addition, spleen cells from A.CA mice were transferred into lethally irradiated high responder mice (B10.5M or CBA). The results (Table IV) showed that spleen cells from high responder strains responded well to native Dx in the A.CA environment, whereas spleen cells from A.CA mice failed to respond even in the high responder strains. Both groups responded to HRC. Thus, it appears unlikely that any type of specific suppressor influence can account for the lack of immune reactivity against the  $\alpha$ -1-6 epitope in A.CA mice.

*Dx is an Efficient Carrier in Dx Low Responsive Mice.* Lack of an immune response to TI antigens could be theoretically caused by two fundamental mechanisms (since suppressive effects have been excluded) specifically: (a) lack of PBA receptors on the responding B cells, which therefore cannot be activated after binding the antigens or (b) absence of Ig receptors competent to passively focus the antigen to the responding B cells. In the latter case the cells cannot produce antibodies, even if given the correct triggering signal via the PBA receptors.

The possibility that A.CA mice lack the PBA receptor for Dx was tested in two ways. First, it was investigated if Dx could induce polyclonal activation in these mice, as determined by induction of antibodies to unrelated antigens. Second, it was studied if Dx could act as a carrier for an unrelated hapten in a dextran nonresponder strain. Since activation of B cells is caused by signals

TABLE IV  
*Immune Response to  $\alpha$ -1-6 and HRC in Irradiated Recipients Reconstituted with Spleen Cells from High and Low Responder Strains*

Donor	Irradiated recipients	PFC/10 <sup>6</sup> spleen cells against		
		$\alpha$ -1-6	HRC	NNP
CBA	CBA	165	320	95
ACA	CBA	14	84	76
ACA	ACA	0	62	218
CBA	ACA	139	268	116
5M	ACA	1,544	2,500	734
ACA	ACA	6	340	179
ACA	5M	2	183	127

\* The irradiated recipients (750 R) were given  $30 \times 10^6$  spleen cell i.v. and were immediately immunized with 1  $\mu$ g native Dx together with 0.2 ml of a 10% suspension of HRC. The PFC against the immunizing antigens as well as against SRBC and the hapten NNP were determined 4 days (CBA) or 5 days (5M) later.

given to the cells by polyclonal B-cell activating properties of the TI antigen, Dx should not be able to work as a carrier for unrelated haptens; if the mechanism of unresponsiveness was lack of PBA receptors for Dx. The results were clear in both systems and showed that Dx was as potent PBA also in A.CA mice. Thus, Dx induced polyclonal antibody synthesis in A.CA mice in vivo or in vitro as it did in high responder strains (Figs. 1 and 2).

When the hapten FITC was conjugated to Dx and the conjugate used to immunize A.CA mice as well as high responder strains, it was found that the conjugation of FITC to Dx did not make the A.CA mice respond to  $\alpha$ -1-6 (Fig. 3), in agreement with the findings that conjugation of Dx to HRC did not make A.CA mice competent to respond to the  $\alpha$ -1-6 epitope. However, A.CA mice responded strongly to the FITC epitope and were equal in this respect to the  $\alpha$ -1-6 high responder strains (Fig. 3). Thus, Dx can work as an efficient carrier for unrelated haptens, indicating that A.CA mice possess PBA receptors for Dx.

*Polyclonal Activation of Cells from Low Responder Mice Does Not Induce Anti- $\alpha$ -1-6 Antibodies.* The second alternative given above (lack of V gene expression against the  $\alpha$ -1-6 epitope) can be tested by the use of polyclonal B-cell activators, since these substances are competent to reveal the V gene repertoire of the B cells that are susceptible to activation by a particular PBA. This approach has been used successfully to study the presence of immunocompetent cells against the tolerogen in specifically tolerant animals (17) and even to induce autoantibody formation in resting B lymphocytes (18).

Consequently, polyclonal concentrations of the PBA lipopolysaccharide (LPS) were injected into A.CA mice, (or added to such cells in culture) and as a control also injected into strains that were high responders to the  $\alpha$ -1-6 epitope (or added to such cells in vitro). The groups were tested at various time intervals against different antigens (SRBC, HRC, NNP, FITC, and  $\alpha$ -1-6 of high and medium affinity). The results showed that LPS was competent to induce antibody synthesis against all epitopes studied in the high responder strains, but A.CA and A.TH mice consistently failed to express anti- $\alpha$ -1-6 PFC,




FIG. 1. Induction of polyclonal antibody synthesis by 2 mg native Dx in spleen cells from A.CA and A x C57BL mice.  $10^7$  spleen cells were cultivated in serum-free medium in the presence or absence of native dx for 2 days and thereafter assayed against FITC-SRBC by using a high epitope density. Bg indicates background in nonimmunized animals.




FIG. 2. Induction of polyclonal antibody synthesis by different concentrations of native Dx in vitro by using spleen cells from A.CA (■) and C57BL (□) mice. The response was tested after 2 days in culture by using the culture system described in Fig. 1 against NNP-coated target cells.

even though a PFC response could easily be detected against all other antigens used (Fig. 4 and Table V).

Analogous experiments carried out in vivo with strains A and CBA mice gave similar results (Fig. 5).

*CBA/N Mice Lack the Major Part of the B-cell Population Responding to the PBA Property of Dx.* A similar analysis was carried out with CBA/N mice to determine the mechanism of unresponsiveness to  $\alpha$ -1-6. The results were

## B-CELL UNRESPONSIVENESS TO DEXTRAN




FIG. 3. Immune response against  $\alpha$ -1-6 and FITC in A  $\times$  CBA F<sub>1</sub> and A.CA mice after immunization i.v. with 100  $\mu$ g/mouse of native FITC-Dx B512. The response was determined at day 5 by using the Jerne plaque assay as modified for the detection of PF Cells against  $\alpha$ -1-6. For detection of anti- $\alpha$ -1-6 PFC, SRBC were conjugated with 40  $\mu$ g of stearyl Dx in 10 ml of a 5% suspension of SRBC. Anti-FITC PFC were detected with SRBC conjugated with a solution of 5 mg/ml of FITC.




FIG. 4. Induction of polyclonal antibody synthesis in vivo by injection of 100  $\mu$ g of LPS i.p. into B10.5M and A.CA mice. The spleens were taken out after 2 and 4 days and tested for the number of PFC against  $\alpha$ -1-6 and the hapten NNP. The response against  $\alpha$ -1-6 was undetectable in A.CA mice.

completely different from those obtained with strains on A background. Thus, native Dx induced a very small polyclonal antibody response against the haptens FITC, NNP, or heterologous RBC (Fig. 6). Also the response to LPS was impaired, but was stronger than the response to Dx.

When CBA/N mice (or CBA  $\times$  DBA F<sub>1</sub> males) were immunized with FITC-Dx they neither produced PFC nor  $\alpha$ -1-6 against FITC. However, they gave a



TABLE V  
*Polyclonally Activating Concentrations of LPS Fail to Induce Anti- $\alpha$ -1-6 PFC in Cells from Low Responder Strains In Vitro\**

Strain	Treatment of cultures <sup>‡</sup>	PFC/10 <sup>6</sup> cells against the indicated targets at day			
		3		6	
C57BL	—	$\alpha$ -1-6 <sup>§</sup>	FITC <sup>  </sup>	$\alpha$ -1-6	FITC
	LPS	0	55	0	100
A.CA	—	380	2,042	600	5,000
	LPS	10	70	0	67
A.TH	—	0	1,410	16	1,766
	LPS	30	50	0	17
		0	880	0	1,700

\* The lymphocytes were cultivated in 2 ml medium containing 5% human A serum and  $5 \times 10^{-9}$  M 2-mercaptoethanol at a cell concentration of  $3 \times 10^5$  cells/ml.

<sup>‡</sup> 100  $\mu$ g/ml of LPS.

<sup>§</sup> Stearoyl Dx at a concentration of 40  $\mu$ g/10 ml was used to sensitize SRC as targets in the PFC assay.

<sup>||</sup> 5 mg/ml of FITC was used to sensitize SRBC as targets in the PFC assay.




FIG. 5. Induction of polyclonal antibody synthesis in vivo by the injection of 100  $\mu$ g LPS into A and CBA mice. Their PFC response against  $\alpha$ -1-6 and NNP was followed up to day 6. A mice (■) responded only to NNP (·····) and not to  $\alpha$ -1-6 (—), whereas CBA mice ( $\Delta$ ) responded to both epitopes.

definite, although small response against FITC when the hapten was conjugated to TD HRC, indicating that the mice possess V genes for the FITC epitope (Table VI). Thus, the cause for the unresponsiveness to FITC-Dx in this case appears to be the lack of the major part of the B-cell population having PBA receptors for Dx, in analogy with the findings that these mice lack triggering receptors for other PBA, such as Ficoll. CBA/N mice are analogous in this way




FIG. 6. Polyclonal activation by LPS and native Dx of spleen cells from CBA/N x DBA male and female mice. Half of the mice were untreated, whereas the other half were immunized 5 days earlier with 100  $\mu$ g of native FITC-Dx and were tested for PFC response against FITC. Only the immunized females developed an immune response to FITC, the males were totally unresponsive. The spleen cells were cultivated in RPMI medium containing 5% fetal calf serum and  $5 \times 10^{-5}$  M 2-mercaptoethanol by using  $2 \times 10^6$  cells/ml and 0.5 ml per culture tube. LPS (50  $\mu$ g/ml) and native Dx (4 mg/ml) were added to the indicated cultures. The cells were harvested on day 3 and tested against NNP-SRBC. Bg indicates background in nonactivated cultures.

TABLE VI  
Immune Response of CBA/N and BALB/c Mice to Native FITC Dx and FITC-HRC\*

Strain	Antigen	SRBC	HRC	PFC/spleen against $\alpha$ -1-6 $\ddagger$		FITC $\S$	
				40	400	0.2	5
CBA/N	-	10	20	0	0	120	-
	FITC-Dx 100 $\mu$ g	5 $\pm$ 5	NT	10 $\pm$ 10	0	45 $\pm$ 45	155 $\pm$ 63
	FITC-HRC	35 $\pm$ 17	3,625 $\pm$ 880	NT	NT	1,690 $\pm$ 630	2,615 $\pm$ 660
BALB/c¶	-	25	50	25	425	175	1,250
	FITC-Dx 100 $\mu$ g	116 $\pm$ 116	NT	1,933 $\pm$ 1,137	1,816 $\pm$ 1,238	30,233 $\pm$ 13,700	46,066 $\pm$ 20,500
	FITC-HRC	200 $\pm$ 57	93,000 $\pm$ 30,324	NT	NT	35,666 $\pm$ 9,700	50,530 $\pm$ 13,300

\* The mice were immunized with 100  $\mu$ g native FITC-Dx i.v. and with 0.2 ml of a 10% suspension of FITC-HRC labeled with 0.5 mg/ml of FITC.

$\ddagger$  40 and 400  $\mu$ g/10 ml stearoyl Dx were used to coat SRBC.

$\S$  0.2 and 5 mg/ml FITC were used to coat SRBC.

|| Four mice/group.

¶ Three mice/group.

to the LPS nonresponder C3H/HeJ, lacking PBA receptors for LPS, but the defect is apparently more pronounced and affects cells with receptors for several different PBA and the genetic localization of the defect is different (it is located on the X chromosome).

Finally it was tested if CBA/N mice could produce anti- $\alpha$ -1-6 PFC after immunization with the TD conjugate Edistine-Dx (a plant protein conjugated

to Dx and provided by Dr. Richter, Pharmacia, Uppsala, Sweden). However, no detectable response was obtained.

### Discussion

The findings in this paper illustrate two fundamentally different causes of immunological unresponsiveness at the B-cell level to the same epitope on Dx. Strains on A background were nonresponders because they failed to express V genes coding for antibodies against  $\alpha$ -1-6. In contrast, strain CBA/N failed to respond to Dx or FITC-Dx, because the cell population having PBA receptors for Dx was severely depleted. This strain is therefore analogous to C3H/HeJ which cannot respond to LPS or any hapten coupled to LPS, because there is a selective loss of PBA receptors for LPS. CBA/N not only has defective B cells capable of responding to the PBA property of Dx, as shown here, but also cells responding to Ficoll, pneumococcal polysaccharide SIII, and polyacrylamide. It has been claimed that this strain responds normally to haptened LPS, indicating that it possesses B cells with PBA receptors for LPS (5). CBA/N responded, although poorly, to the PBA property of LPS. Although C3H/HeJ and CBA/N principally exhibit analogous defects, there are marked differences between the two strains. The defect in C3H/HeJ is selective for LPS, and is determined by one gene localized on the fourth chromosome, whereas CBA/N lacks receptors or B-cell populations for several different PBA's and the defect is X-linked. It could not be formally established whether CBA/N mice possess V genes for  $\alpha$ -1-6, since they did not respond to TD Dx conjugates.

The fact that A mice did not respond to Dx, although they have cells with PBA receptors for Dx, is a new type of immunological unresponsiveness. It is a genetically determined (unpublished observations) inability to respond, which could not be ascribed to lack of helper T cells, presence of suppressor T cells, or any suppressive influence of the environment of the mice that did not allow the expression of an anti- $\alpha$ -1-6 response. The unresponsive state exhibited stringent immunological specificity, since only the  $\alpha$ -1-6 epitope was affected, whereas other epitopes conjugated to Dx induced normal immune responses. Since it was not possible to convert nonresponders into responders by a variety of procedures that engage different helper cells, such as conjugating Dx to TD HRC, injecting it with polyclonal T-cell activators such as PHA and Con A, or incorporating it into Freund's adjuvant, it is unlikely that the defect can be ascribed to lack of helper cells. Furthermore, several different PBA's such as LPS and PPD, which could activate polyclonal antibody synthesis to all other antigenic determinants failed to do so against  $\alpha$ -1-6 in the nonresponder strains, whereas expected anti- $\alpha$ -1-6 responses occurred with similarly-treated cells from high responder strains. It follows from these findings that B cells which can be activated by helper T cells, LPS, PPD, and Dx cannot express Ig molecules with the  $\alpha$ -1-6 variable region. Since these polyclonal activators probably affect a major part of the B-cell population (LPS alone activates about 30% of all B cells and these are different from those activated by PPD and Dx) it follows that most if not all B cells do not express the gene responsible for the formation of the combining site for  $\alpha$ -1-6.

The lack of detectable anti- $\alpha$ -1-6 antibody-producing cells could be due to two

main reasons. It is possible that an  $\alpha$ -1-6 clone or several such clones are present in the nonresponders, but the number of cells in the clones is too small to be detected. Alternatively, lymphocytes from these animals either lack V genes coding for antibodies against  $\alpha$ -1-6 or the cells cannot selectively express these V genes. Although the possibility that the clone size is too small to allow detection may be used to explain the *in vitro* data with PBA, it is highly unlikely that it is applicable for the *in vivo* experiments, since the amplification of the response was very pronounced in responder strains, which presumably have very few clones because of the homogeneity of the affinity of the antibodies produced. Usually the number of PFC against  $\alpha$ -1-6 was 500–2,000 times the background 5 days after immunization. The exponential increase of the immune response during the first 5 days is predominantly due to division of initially activated precursor cells rather than to recruitment of new precursor cells. Actually it is possible to detect one precursor cell in adoptive transfer experiments and there is no reason why this would not also be possible against  $\alpha$ -1-6 *in vivo* with the marked clonal expansion that occurs in high responders. There is no reason why precursor cells could not multiply in nonresponder strains; since no suppressor cells have been detected, there is not a need for helper cells, and other epitopes coupled to Dx induce a normal anti-hapten response.

The only tenable conclusion is that strains on A background fail to express Ig receptors and secrete antibodies directed against the  $\alpha$ -1-6 epitope in the majority of the individuals. The mechanism by which a few individuals secrete antibodies against this epitope will be discussed below. It seems highly unlikely that some control mechanism would exist that selectively prevents the synthesis from messenger RNA of antibodies directed only against one particular epitope. It is more plausible that the defect is at the DNA level. Two mechanisms could account for the genetically determined state of unresponsiveness to Dx at the DNA level, namely lack of V genes coding for antibodies against  $\alpha$ -1-6 or lack of V-C gene translocation for this particular specificity. Neither alternative has precedence, but both are possible. The possibility that different V genes would translocate at different rates (or not at all) is compatible with our findings (unpublished observations) that high responder strains to the  $\alpha$ -1-6 epitope do not express this antibody during the 1st mo of life, even when the cells are polyclonally activated, for example, by LPS, whereas they express antibodies of all other specificities. This suggests that the V-C translocation can occur at different times for different V genes. It is possible that the V gene coding for anti- $\alpha$ -1-6 actually exists in strains on A background, but that it never, or rarely, translocates to a C gene. Conclusive experiments on this point require experiments at the DNA level.

A certain proportion of mice on strain A background produced antibodies against  $\alpha$ -1-6, in most cases to a very small degree. Usually the PFC differed markedly from those in high responder strains, because they were unclear and of very low affinity, since they were detected easily and in high numbers only with target cells coated with a high epitope density. This is in contrast to the PFC obtained in high responder strains, which were of uniform affinity; no difference in number or clarity was observed with target cells of different epitope density, suggesting a mono- or pauciclonal response. Mice lacking a

particular V gene may occasionally make use of another V gene, which may have mutated so as to cause the appearance of an antibody crossreacting with the epitope for which a V gene is lacking. This may very well be the case with the small proportion of animals responding to the  $\alpha$ -1-6 epitope in strains of A background. Occasionally, the PFC of low responder mice were undistinguishable from those of high responder strains. This could be caused by a successful V-C translocation also in strains on A background, provided that a lack of this type of translocation is the mechanisms for unresponsiveness.

The possibility that mice on A background do not express Ig receptors, because they have been tolerized by exposure to bacterial products containing  $\alpha$ -1-6 is highly unlikely. Thus, we have previously shown (19) that immunological tolerance to an epitope on Dx only affects a subpopulation of B cells with Ig receptors for the epitope. This subpopulation is characterized by having PBA receptors for Dx as well as Ig receptors for the epitope. Thus, tolerance is not due to clonal deletion and many specific B cells in tolerant mice can be activated by different PBA's to synthesize antibodies against the tolerogen. Similarly, we have shown (unpublished observations) that lymphocytes from Dx tolerant after treatment with dextranase can be activated by LPS, but not with Dx to the synthesis of antibodies against the tolerogen. Finally, Dx-tolerant mice injected with dextranase can make specific high affinity antibodies to Dx after immunization with dextran coupled to a variety of TD and TI carriers, but never after immunization with Dx alone. These findings show that only the B cells which have both Ig and PBA receptors for Dx are irreversibly tolerized, other specific B cells remain unaffected and can be activated by immunization with dx coupled to other carriers. Since we have performed analogous experiments here (Tables III and V) and failed to activate anti-Dx antibodies in A.CA mice, the possibility that A.CA mice are nonresponders because they are tolerant can be ruled out.

A detailed study on the genetics of unresponsiveness to the  $\alpha$ -1-6 epitope will be given elsewhere, but it is clear that the genes responsible are not linked to H-2 region since A, A.CA, A.SW, A.TL, and A.TH all have differences in different parts of the H-2 complex including the I region and they are all low responders. These strains have the same genetic background except H-2 indicating that the genes are not present in the major histocompatibility complex.

### Summary

The immune response of mice to the  $\alpha$ -1-6 epitope of dextran (Dx) B512 was found to be under genetic control. The congenic mouse strains A, A.CA, A.SW, A.TH, and A.TL exhibited a specific defect in their response to  $\alpha$ -1-6. Also strain CBA/N was unresponsive to  $\alpha$ -1-6, but the mechanism of unresponsiveness was found to be different.

Unresponsiveness to  $\alpha$ -1-6 in congenic A strains was not due to suppressor cells. Although these strains failed to respond to the  $\alpha$ -1-6 epitope, they responded strongly to the hapten Fluorescein isothiocyanate (FITC) conjugated to Dx, indicating that the Dx can function as an efficient carrier in these strains. Dx was a potent polyclonal B-cell activator in congenic A strains as well as in high responder strains. Polyclonally-activating concentrations of

lipopolysaccharide (LPS) failed to induce the synthesis of anti- $\alpha$ -1-6 antibodies in congenic A strains, although antibodies of all other specificities studied were produced. However, in high responder strains, LPS induced the synthesis of anti- $\alpha$ -1-6 antibodies. It was concluded that congenic A strains do not express V genes coding for antibodies against  $\alpha$ -1-6.

In contrast, strain CBA/N failed to respond to both the  $\alpha$ -1-6 and FITC epitope on Dx, whereas they could respond to FITC conjugated to horse erythrocytes. Dx induced a very small, if any, polyclonal antibody response in B cells from CBA/N mice or male CBA/N  $\times$  DBA hybrids, whereas Dx was a very potent polyclonal B-cell activator in female hybrids. It is concluded that CBA/N mice are nonresponders to Dx or haptenated Dx, because the cell population that can respond to the polyclonal B-cell activating properties of Dx is severely depleted.

The technical assistance of Susanne Bergstedt is gratefully acknowledged.

Received for publication 6 May 1977.

### References

1. Benacerraf, B. and H. O. McDevitt. 1972. Histocompatibility linked immune response genes. *Science (Wash. D. C.)*. 175:273.
2. Feingold, N., J. Feingold, D. Mouton, Y. Bouthilier, C. Stiffel, and G. Biozzi. 1976. Polygenic regulation of antibody synthesis to sheep erythrocytes in the mouse: a genetic analysis. *Eur. J. Immunol.* 6:43.
3. Coutinho, A., G. Möller, and E. Gronowicz. 1975. Genetical control of B-cell responses. IV. Inheritance of the unresponsiveness to lipopolysaccharides. *J. Exp. Med.* 142:253.
4. Watson, J., and R. Riblet. 1974. Genetical control of responses to bacterial lipopolysaccharides in mice. I. Evidence for a single gene that influences mitogenic and immunogenic responses to lipopolysaccharides. *J. Exp. Med.* 140:1147.
5. Mosier, D. E., I. Scher, H. Ruhl, P. L. Cohen, I. Zitran, and W. E. Paul. 1976. Activation of normal and defective B lymphocytes by thymus-independent antigen. In *Mitogens in Immunobiology*. J. J. Oppenheim and D. L. Rosenstreich, editors. Academic Press, Inc., New York. 313.
6. Blomberg, B., W. R. Geckeler, and M. Weigert. 1972. Genetics of the antibody response to dextran in mice. *Science (Wash. D. C.)*. 177:178.
7. Riblet, R., B. Blomberg, M. Weigert, R. Lieberman, B. A. Taylor, and M. Potter. 1975. Genetics of mouse antibodies. I. Linkage of the dextran response locus,  $V_H$ -DEX to allotype. *Eur. J. Immunol.* 5:775.
8. Fernandez, C., and G. Möller. The immune response against two epitopes on the same thymus-independent polysaccharide carrier. I. Role of epitope density in carrier dependent immunity and tolerance. *Immunology*. 33:59.
9. Coutinho, A., G. Möller, and W. Richter. 1974. Molecular basis of B cell activation. I. Mitogenicity of native and substituted dextrans. *Scand. J. Immunol.* 3:321.
10. Howard, J. G., B. M. Courtenay, and C. Hale. 1976. Lack of effect of thymectomy on spontaneous recovery from tolerance to levan. *Eur. J. Immunol.* 6:838.
11. Howard, J. G., G. Vicari, and B. M. Courtenay. 1975. Influence of molecular structure on the tolerogenicity of bacterial dextrans. I. The  $\alpha$  1-6 linked epitope of dextran B512. *Immunology*. 29:585.
12. Möller, G. 1974. Effect of B-cell mitogens on lymphocyte subpopulations possessing C'3 and Fc receptors. *J. Exp. Med.* 139:969.

13. Pasanen, V. J., and O. Mäkelä. 1969. Effect on the number of haptens coupled to each erythrocyte on haemolytic plaque-formation. *Immunology*. 16:399.
14. Coutinho, A., and G. Möller. 1973. In vitro induction of specific immune responses in the absence of serum: requirements for non-specific T or B cell mitogens. *Eur. J. Immunol.* 3:531.
15. Mishell, R. I. and R. W. Dutton. 1967. Immunization of dissociated spleen cell cultures from normal mice. *J. Exp. Med.* 126:423.
16. Möller, G. 1975. Suppressor T lymphocytes. *Transplant. Rev.* 26.
17. Möller, G., E. Gronowicz, U. Persson, A. Coutinho, E. Möller, L. Hammarström, and E. Smith. 1976. Spleen cells from animals tolerant to a thymus-dependent antigen can be activated by lipopolysaccharide to synthesize antibodies against the tolerogen. *J. Exp. Med.* 143:1429.
18. Primi, D., E. Smith, L. Hammarström, and G. Möller. Sera from LPS injected mice exhibit complement-dependent cytotoxicity against syngeneic and autologous spleen cells. *Cell. Immunol.* 39:252.
19. Fernandez, C., and G. Möller. 1977. Induction of immunological tolerance requires that the B cells can respond to the polyclonal B-cell-activating properties of the thymus-independent antigens. *J. Exp. Med.* 146:308.