

Isolation, cDNA Cloning, and Overexpression of a 33-kD Cell Surface Glycoprotein that Binds to the Globular "Heads" of C1q

By Berhane Ghebrehiwet,*† Boon-Leong Lim,§
Ellinor I. B. Peerschke,† Anthony C. Willis,§ and K. B. M. Reid§

From the Departments of *Medicine and †Pathology, State University of New York, Stony Brook, New York 11794-8161; and the §Medical Research Council Immunochimistry Unit, Department of Biochemistry, Oxford University, Oxford OX1 3QU, UK

Summary

This work describes the functional characterization, cDNA cloning, and expression of a novel cell surface protein. This protein designated gC1q-R, was first isolated from Raji cells and was found to bind to the globular "heads" of C1q molecules, at physiological ionic strength, and also to inhibit complement-mediated lysis of sheep erythrocytes by human serum. The NH₂-terminal amino acid sequence of the first 24 residues of the C1q-binding protein was determined and this information allowed the synthesis of two degenerate polymerase chain reaction primers for use in the preparation of a probe in the screening of a B cell cDNA library. The cDNA isolated, using this probe, was found to encode a pre-pro protein of 282 residues. The NH₂ terminus of the protein isolated from Raji cells started at residue 74 of the predicted pre-pro sequence. The cDNA sequence shows that the purified protein has three potential N-glycosylation residues and is a highly charged, acidic molecule. Hence, its binding to C1q may be primarily but not exclusively due to ionic interactions. The "mature" protein, corresponding to amino acid residues 74–282 of the predicted pre-pro sequence, was overexpressed in *Escherichia coli* and was purified to homogeneity. This recombinant protein was also able to inhibit the complement-mediated lysis of sheep erythrocytes by human serum and was shown to be a tetramer by gel filtration in nondissociating conditions. Northern blot and RT-PCR studies showed that the C1q-binding protein is expressed at high levels in Raji and Daudi cell lines, at moderate levels in U937, Molt-4, and HepG2 cell lines, and at a very low level in the HL60 cell line. However, it is not expressed in the K562 cell line. Comparison of gC1q-R NH₂-terminal sequence with that of the receptor for the collagen-like domain of C1q (cC1q-R) showed no similarity. Furthermore, antibodies to gC1q-R or an 18-amino acid residue-long NH₂-terminal synthetic gC1q-R peptide did not cross-react with antibodies to cC1q-R. Anti-gC1q-R immunoblotted a 33-kD Raji cell membrane protein, whereas anti cC1q-R recognized a molecule of ~60 kD. The NH₂-terminal sequence of gC1q-R appears to be displayed extracellularly since anti-gC1q-R peptide reacted with surface molecules on lymphocytes, polymorphonuclear leukocytes, and platelets, as assessed by flow cytometric and confocal laser scanning microscopic analyses. In addition, all or part of the gC1q binding domain may reside within the 24 amino acid stretch of the NH₂-terminal sequence of gC1q-R since the 18 amino acid residue long-synthetic peptide corresponding to this region inhibited serum C1q hemolytic activity. The data presented in this report suggest that there are at least two types of C1q-R which appear to be expressed on the same type of cells and these receptors individually or in concert may contribute to the diversity of C1q-mediated responses.

C1q binds to a variety of cells such as B cells, monocytes, macrophages, neutrophils, eosinophils, fibroblasts, platelets, endothelial and smooth muscle cells (1–9). The biological responses elicited by C1q are also as diverse as the cell types that express the receptor (1). It enhances FcR- and CR1-mediated phagocytosis in monocytes and macrophages (10,

11), stimulates Ig production by B cells (12, 13), activates platelets to express α IIb/ β ₃ integrins, P-selectin, and procoagulant activity (14), and plays a role in the activation of tumor cytotoxicity of macrophages (15, 16).

Most of the above actions of C1q are considered to be mediated by a C1q receptor that binds to the collagen-like regions

of C1q (1). This C1q receptor is a single chain, acidic glycoprotein with a molecular mass of ~ 60 kD (1). However, this receptor has a relatively low affinity to the collagen-like regions of C1q and binds at a very low ionic strength (10 mM potassium phosphate buffer) (17). In 1989, a high affinity receptor, which binds to the globular heads of C1q, was reported to be present, along with the low affinity receptor, on human diploid fibroblasts (18), thereby indicating that at least two types of C1q receptors might be involved to impart the vast array of C1q-mediated biologic functions.

In this paper and in a previous report (19), a cell surface protein, which has a high affinity for the C1q globular heads, has been characterized. By the use of NH₂-terminal protein sequence information and PCR techniques, a full-length cDNA clone was isolated and the primary sequence of this protein has been predicted. Northern blot and RT-PCR studies were carried out to demonstrate the different levels of expression of this protein in different cell lines. Moreover, a recombinant form of this protein has been produced in *Escherichia coli* and shown to be able to inhibit complement-mediated lysis of sheep erythrocytes by human serum.

Materials and Methods

Chemicals and Reagents

The following chemicals and reagents were purchased from the sources indicated: FCS (Hyclone Laboratories, Logan, UT); RPMI 1640, 100 \times antibiotic-antimycotic mixture (GIBCO BRL, Gaithersburg, MD); Emulphogene BC 720 (polyoxyethylene-10 tridecyl ether), FITC-conjugated goat anti-rabbit IgG, KLH, DMSO (Sigma Chemical Co., St. Louis, MO); NHS-LC-biotin (Sulfosuccinimidyl-6-[biotinamido] Hexanoate), ImmunoPure A/G IgG purification kit, ImmunoPure horseradish peroxidase-conjugated goat anti-rabbit IgG, 4-chloro-1-naphthol, Tween 20 (Pierce, Rockford, IL); Con A-Sepharose PD-10 columns (Pharmacia, Biotech, Inc., Uppsala, Sweden); Mono-Q, TSKgel DEAE-NPR columns (TosoHaas, Montgomeryville, PA); ¹²⁵I-Na (Amersham, Aylesbury, UK); and polyvinylidene difluoride (PVDF) membranes, (Immobilon-P; Millipore Corp., Bedford, MA).

Cultured Cells

Raji cells were used to prepare membrane proteins and were grown in RPMI 1640 containing 10% (vol/vol) heat-inactivated (3 h, 56°C) FCS and 1% (vol/vol) of a 100 \times antibiotic-antimycotic mixture that consisted of 10,000 U/ml penicillin G, 25 μ g/ml amphotericin B, and 100,000 μ g/ml streptomycin sulfate and maintained in an atmosphere of 95% air and 5% CO₂. When large quantities of Raji cells were required, the cells were grown in 2-liter roller bottles as described earlier (20). Other cell lines such as Daudi, U937, Molt-4, and HL60 were grown under similar conditions.

Purified Proteins

Human C1q used in these studies was purified by the method of Reid (21). For comparison, C1q was also purified from a pool of freshly obtained sera according to the procedures of Yonemasu and Stroud (22) followed by purification on Con A-Sepharose 4B as described (23).

The collagen "stalks" of C1q (cC1q) were prepared by pepsin digestion of the purified C1q molecule following the procedure

of Reid (24) and the globular "heads" of C1q were prepared by collagenase digestion as described (25).

Radioiodination

Except for C1q, which was labeled by the method of Bolton and Hunter (26), all other proteins were radiolabeled by the Iodogen method (27) with 1 mCi of Na¹²⁵I as described elsewhere (3, 28). After labeling, the free ¹²⁵I was removed by gel filtration using PD-10 (Sephadex G-25) columns equilibrated with the appropriate buffer system in which the labeled protein was to be kept.

Purification of C1q-R

The purification of C1q-R in these studies used a combination of two published methods (2, 3) with some minor modifications. These methods were: (a) affinity purification on C1q-Sepharose CL4B, (b) ion-exchange separation on FPLC using a Mono-Q column, and (c) HPLC purification using a TSKgel DEAE-NPR column as described (3, 28).

C1q-Sepharose Affinity Chromatography. Raji cell membranes were prepared from 20-liter cultures (4×10^{10} cells) as described in detail earlier (2, 20). Half of the solubilized membrane solution was then applied to a column packed with 3 ml of C1q-Sepharose CL4B (3 mg C1q/ml Sepharose) equilibrated with 10 mM sodium phosphate, pH 7.4, containing 20 mM NaCl, 0.1% Emulphogene BC720, and a cocktail of enzyme inhibitors (2, 20). After washing (5 \times column volume), the bound proteins were eluted with the same buffer containing 1 M NaCl. The eluted material was dialyzed against equilibrating buffer, and a 0.5-ml aliquot was radiolabeled and then tested for C1q-binding activity by a solid phase radioligand binding assay described below. Preliminary studies showed that the eluted material contained most of the C1q-binding activity.

FPLC on Mono-Q Column. The C1q-binding material eluted from the C1q-Sepharose CL4B column was mixed with 100 μ l ($\sim 10^6$ cpm) of the radiolabeled sample and applied to a 1-ml Mono-Q column (9, 17). The column was washed until cpm/ml ≤ 100 , and then the bound proteins were eluted with a linear NaCl concentration gradient (0–600 mM NaCl). C1q-containing fractions were identified by solid phase binding assay (see below), pooled, and concentrated.

HPLC on DEAE Column. The concentrated, C1q-binding fraction from the Mono-Q column was dialyzed against 50 mM Tris, pH 8.0, and then subjected to HPLC using a 1-ml TSK gel DEAE-NPR column as described (9, 17). The bound proteins were eluted using first a 0–500-mM NaCl gradient in equilibrating buffer (10 min) and then a 500 mM–1 M gradient (3 min).

Solid Phase Binding Assay

Solid phase radioligand assays were performed using ¹²⁵I-labeled C1q-R or C1q-R-enriched material by a previously described method (20, 29). Briefly, 50 μ l, 5 μ g/ml each (15 mM Na₂CO₃, 35 mM NaHCO₃, pH 9.6) of C1q, cC1q, or gC1q were applied to duplicate wells of flat-bottomed strip plate-8 (Costar Corp., Cambridge, MA) and incubated for 2 h at 37°C. Wells coated with either 1%, BSA, 100 μ g/ml IgG, or 100 μ g/ml properdin, served as irrelevant or positively charged protein controls. After incubation, the wells were washed (3 \times 300 μ l) with TBST (50 mM Tris-HCl, pH 7.5, containing 150 mM NaCl and 0.05% Tween 20). Then 50 μ l of ¹²⁵I-labeled C1q-R in TBB (20 mM Tris-HCl, pH 7.5, containing 100 μ g/ml BSA) containing a fixed or increasing amounts of counts per minute were added and further incubated

(1 h, 37°C). Next, the unbound radioactivity was removed, the wells washed (5 × 300 μl) with TBB, dried by tapping over a stack of filter papers, and the individual wells counted in a γ-counter.

SDS-PAGE and Western Blot Analysis

SDS-PAGE analysis was performed on 1.5-mm-thick slab gels according to the method of Laemmli (30). Samples were either run unreduced or reduced and alkylated by boiling for 5 min in the presence of 0.1 M dithiothreitol and 0.2 M iodoacetamide. After running, the gels were stained with Coomassie Brilliant Blue, destained and dried. Gels containing radiolabeled samples were exposed to Kodak X-OMAT AR film at -80°C and then analyzed by autoradiography. Samples for Western blot analysis (31) were first run on SDS-PAGE, then electrotransferred to polyvinylidene difluoride (PVDF) nitrocellulose membranes, blocked with 5% nonfat milk containing TBST, and the bound proteins probed with predetermined dilutions of rabbit antibodies in the above buffer. The bound antibodies were visualized by horseradish peroxidase-conjugated goat anti-rabbit IgG followed by reaction with 4-chloro-1-naphthol substrate.

NH₂-terminal Amino Acid Sequencing and Peptide Synthesis

The NH₂-terminal sequence information of gC1q-R was generated using a protein sequencer and analyzer (models 470A and 120A; Applied Biosystems, Inc., Foster City, CA) as described elsewhere (28, 32). An 18-amino acid residue-long synthetic peptide (gC1q-R₁₈) derived from the NH₂-terminal sequence of gC1q-R was synthesized, on a peptide synthesizer (model 430A; Applied Biosystems, Inc.) at the Center for Analysis and Synthesis of Macromolecules (Department of Medicine, State University of New York, Stony Brook, NY). The peptide was then purified by gel filtration on HPLC.

Antibody Production

Antibody to gC1q-R was prepared in rabbits according to the following schedule. Purified gC1q-R (10–20 μg) in 200 μl PBS was mixed with 200 μl each of complete and IFA and injected at three sites: one subcutaneous and two intramuscular. Booster injection was given as above 2 wk later and, thereafter, injections were made every 2 wk in IFA until a total of four such injections were made. The rabbits were then rested for 4–6 wk before a final injection was given and the rabbits bled and the serum collected and tested by ELISA against purified gC1q-R.

Antibody to gC1q-R peptide was generated by injection of 500 μg of KLH-conjugated peptide following the immunization protocol described above. Conjugation of gC1q-R peptide to KLH was accomplished by the glutaraldehyde method as described (33). The IgG fraction from each antiserum was prepared using the ImmunoPure A/G IgG purification kit following the manufacturer's recommendations (Pierce). Antipeptide antibodies were further purified by affinity purification on KLH-Sepharose CL4B.

Hemolytic Assays

The effect of gC1q-R₁₈, KLH-gC1q-R₁₈, or the soluble, recombinant form of gC1q-R (rgC1q-R) was assessed by incubating various amounts of these molecules with 10 μl normal human serum (NHS)¹ for 60 min at room temperature in a total volume of 100

μl GVB (Veronal-buffered saline, pH 7.4, containing 0.15 mM CaCl₂, 1 mM MgCl₂, and 0.1% gelatin). At the end of the incubation, the volume of the reaction mixtures was brought to 950 μl GVB, and 50 μl of EA (10⁸/ml, sheep erythrocytes [E] sensitized with IgM anti-sheep E [A]) was added and further incubated for 60 min at 37°C. The reaction was then stopped, immediately centrifuged, and free hemoglobin in the supernatant assessed spectrophotometrically at 412 nm.

Confocal Laser Scanning Microscopy

PBL, monocytes, and PMN were purified by a combination of Ficoll-Hypaque sedimentation (to separate PMNs from mononuclear cells) and plastic adherence in autologous serum (to separate monocytes from lymphocytes), as described (34). For confocal laser scanning microscopic (CLSM) analyses, the cells were suspended at 10⁶ cells/0.3 ml GVB (veronal-buffered saline, pH 7.4, containing 0.1% gelatin, 0.02% NaN₃, and 5 μg/ml cytochalasin B), preincubated for 30 min at 4°C with a 1:5 dilution of C1q-depleted and heat-inactivated (56°C, 30 min) human serum to block Fc-R sites, and then incubated (1 h, 37°C or 4°C) with 50 μl of a 1 mg/ml each of either IgG anti gC1q-R or IgG from preimmune rabbit serum. After incubation, the cells were washed (2 × 2 ml) with cold GVB by centrifugation (800 g, 4°C), the pellet resuspended in 0.3 ml GVB and further incubated (1 h, 37°C or 4°C) with an appropriate dilution of FITC-conjugated goat anti-rabbit antibody. The cells were then washed (3 × 2 ml) in GVB resuspended in original volume and analyzed by CLSM.

cDNA Cloning and Overexpression of gC1q-R

Generation of the First Strand cDNA from Raji Cell Total RNA by AMV Reverse Transcriptase. After purification of the C1q-binding protein, the sequence of the 24 NH₂-terminal amino acids was determined by automated protein sequencing (Table I). Based on this peptide, two degenerate PCR primers (BH-S and BH-3) (Table 2) were designed and synthesized on a DNA synthesizer (model 381A; Applied Biosystems, Cheshire, UK). The antisense primer (BH-3) was used as the primer for the first strand cDNA synthesis from total RNA. All the components in the reverse transcription reaction were from a kit (cDNA Synthesis System Plus; Amersham International, Amersham, Bucks, UK), except for total RNA and primer. In a 10-μl reaction: 3.0 μl H₂O; 2.0 μl 5 × first strand reaction buffer; 0.5 μl sodium pyrophosphate solution; 0.5 μl RNase inhibitor; 1.0 μl deoxynucleoside triphosphate mix; 1.0 μl BH-3 primer (1 μg/ml), 2.0 μl Raji cell total RNA (10 μg/μl); and 10 U AMV reverse transcriptase were mixed together. After incubation at 42°C for 40 min, the tube was stored at -20°C for subsequent PCR amplification.

Table 1. Comparison of the NH₂-terminal Sequence of Raji cC1q-R and gC1q-R

| Receptor type | NH ₂ -terminal sequence |
|---------------|------------------------------------|
| cC1q-R | EPAVYFKEQFLDGDG |
| gC1q-R | LHTDGDKAFVDFLSDE I <u>KEE</u> RKIQ |

The underlined sequence in gC1q-R was used for the construction of two degenerate oligonucleotide probes as well as the generation of a synthetic peptide (gC1q-R₁₈).

¹ Abbreviations used in this paper: CLSM, confocal laser scanning microscopy; IPTG, isopropyl β-D-thiogalactopyranoside; NHS, normal human serum.

Table 2. Sequences of PCR/Library Screening Oligonucleotides

| Name | Sequence (5' → 3') | Direction | Position |
|------|---------------------------------|-----------|--------------|
| BH-S | 5'-ACIGAYGGIGAYAARGCITTT | Sense | a.a. 76-82 |
| BH-3 | 5'-YTGIAATYTTICKYTCYTC | Antisense | a.a. 92-97 |
| KS-1 | 5'-CAGGAATTCCTGAGTGATGAAATTAAGG | Sense | ntd. 331-352 |
| KA-1 | 5'-TGGAATTCATCACTCAGGAAATCAACA | Antisense | ntd. 324-343 |
| KS-2 | 5'-AACCCCTCGCAAGGGCAGAA | Sense | ntd. 521-539 |
| BM-1 | 5'-GTAGGATTTGTTCACTGGCCA | Antisense | ntd. 967-988 |
| BN-1 | 5'-GCCATGGCTCTGCACACCGACGGAGAC | Sense | ntd. 297-315 |
| F11 | 5'-ACTCCTGGAGCCCGTCAGTAT | | λgt11 primer |
| R11 | 5'-GACCAACTGGTAATGGTAGCGAC | | λgt11 primer |

Added restriction sites are underlined: EcoRI(KS-1 and KA-1); NcoI(BN-1).
a.a., amino acid; I, Inosine; ntd., nucleotide; R, A/G; Y, C/T.

Generation of the First Exact Match Oligonucleotide Probe by PCR Purification. The first strand cDNA generated as described above was used as template and the two degenerate, inosine-containing oligonucleotides (BH-S and BH-3) were used as primers in the PCR reaction. A PCR reaction was performed with 30 cycles in a DNA thermal cycler (Perkin-Elmer/Cetus, Norwalk, CT) using a cycle of denaturation for 0.5 min at 94°C, annealing for 0.5 min at 48°C, and an extension at 72°C for 0.5 min. The reaction mixture (50 μl) contained 25 pmol of each primer, 0.25 mM dNTP, 1.25 U Taq polymerase (Promega, Southampton, UK) with the manufacturer's buffer system, together with 1.0 μl first strand cDNA. The PCR product (5 μl) was electrophoresed on a 4% (wt/vol) agarose gel (Nusieve; FMC Bioproducts Europe, Vallensbaek, Denmark). The expected 66-bp product was obtained.

Subcloning and DNA Sequencing. After electrophoresis, the 66-bp fragment was cut out from the gel and extracted from the gel slice by centrifugation through glass wool (35). The DNA fragment (6 μl of eluant) was bluntend-ligated to the HincII site of pBluescript SK (0.1 μg) using T4 DNA ligase (1 U, Amersham) in a 10-μl reaction volume overnight at room temperature. The *E. coli* strain XL-Blue 1 was used as the host in the transformation of cells with the ligation mixture. After transformation, the cells were plated on LAT plates (LB plates with 100 μg/ml ampicillin and 10 μg/ml tetracycline), which had been spread with X-Gal and isopropyl-β-D-thiogalactopyranoside (IPTG) several hours before the cells were inoculated. White colonies were screened by the PCR using the T7 and T3 primers and conditions as used previously. Colonies that yielded the 246-bp PCR product (a 180-bp fragment was generated if there is no insert) were selected and the plasmids were purified by alkaline lysis method from a 10-ml miniprep culture. Double-stranded DNA sequencing was performed on two purified plasmids by the dideoxynucleotide chain termination procedure using T7 polymerase (Pharmacia).

Screening of the λgt 11 Human B Cell cDNA Library. Two oligonucleotide probes (KS-1 and KA-1, Table 2) were synthesized based on the DNA sequence of the clones generated by PCR. An EcoRI site was added at the 5' end of each probe. These oligonucleotides were endlabeled with [³²P]ATP using T4 kinase and were used to screen the library (~10⁶ plaques, Clonetech, Palo Alto, CA) separately, with KS-1 being used for first screening and KA-1 for second screening. Positive plaques were picked up and the phages were resuspended in 1 ml SM buffer (50 mM Tris-HCl, 10 mM

NaCl, 10 mM MgSO₄, and 0.01% gelatin). Screening of the positive plaques was performed by PCR reactions with primers KS-1 and the primers F11 or R11 in 25-μl reaction volumes. F11 (left arm) and R11 (right arm) were vector-specific primers. Cultures (10 ml) were grown from four positive plaques and λ DNA was isolated by λ mini kit (Qiagen Hybaid Ltd., Middlesex, UK). EcoRI inserts were released by digestion, with EcoRI purified and subcloned into pBluescript for sequence analysis. Three cDNA clones of 1.0-1.1 kb were sequenced by the dideoxynucleotide chain termination procedure using T7 polymerase.

Subcloning the cDNA Insert Encoding the gC1q-R into Plasmid pGex-2T. First, a NcoI site was created in the pGex-2T plasmid (36). pGex-2T plasmid was digested by SmaI, desphosphorylated by alkaline phosphatase, and ligated to a NcoI linker (10 mer) by T4 DNA ligase. Plasmid (pGex-2TN) with the NcoI site inserted was selected, cultured, and purified by alkaline lysis method. Second, in order to create a NcoI site on the gC1q-R protein insert for subcloning, a PCR primer (BN-1) was synthesized. A 50-μl PCR reaction was carried out using BN-1 and T7P (a pBluescript primer) as primers and pBluescript containing the gC1q-R cDNA insert as a template. PFU DNA polymerase (Promega) was used to replace Taq polymerase so as to secure a high fidelity. The amplified PCR product was then precipitated and digested by NcoI and BamHI. The digested insert was purified by gel and subcloned into the NcoI site and BamHI site of pGex-2TN. The plasmid that contained the right insert was isolated (pGex-2TNH) and transformed into *E. coli* strain NM554.

Overexpression of Clq-binding Protein (Residues 74-282 of the Amino Acid Sequence Predicted from the cDNA Clone) in *E. coli* Employing pGex-2T Vector. A colony carrying pGex-2TNH was inoculated to a 100-ml LA (LB plus 50 mg/liter Ampicillin) and grown overnight at 37°C. The next morning, the 100-ml culture was inoculated into 900 ml LA. After 1 h at 37°C, protein expression was induced by 0.1 mM IPTG, the culture allowed to grow for 7 h at 37°C, and the cell harvested by centrifugation at 5,000 g for 15 min. For every liter of culture, 15 ml MTPBSN (150 mM NaCl, 16 mM Na₂HPO₄, 4 mM NaH₂PO₄, 0.05% NaN₃, 0.1% Triton X-100, pH 7.3) was used to resuspend the cell pellet. The cells were lysed by sonication (3 × 1 min) and the supernatant generated by centrifugation at 12,000 g for 15 min contained soluble proteins, including GST-gC1q-R fusion product. Glutathione agarose beads (5 ml, sulphur linkage; Sigma Chemical Co.) were pre-

pared by preswelling in MTPBSN, then washing twice in the same buffer. The supernatant was allowed to pass through a column (10 mm × 150 mm) of the beads and the eluant was discarded. The column was washed by 50 ml MTPBSN to elute nonbinding proteins. The fusion protein was eluted by competition with free glutathione using 20 ml 50 mM Tris-HCl, pH 8.0, containing 5 mM reduced glutathione (Sigma Chemical Co.) (final pH 7.5, freshly prepared). The eluent (20 ml) was made 150 mM with respect to NaCl and 2.5 mM with CaCl₂ before thrombin digestion. The concentration of the fusion protein at this stage was estimated to be 1.6 mg/ml. Thrombin was added to a concentration of 3.2 μg/ml and the digestion was allowed to proceed at room temperature for 2 h on a rotating platform. Complete digestion was achieved, as judged by SDS-PAGE carried out after reduction of disulfide bonds. A Mono-Q column (Pharmacia) was employed to purify the recombinant protein. The pH of the sample was adjusted by adding one-twentieth the volume of 1 M Tris, pH 8.6. The sample was applied to a Mono-Q column (1 ml) preequilibrated in buffer A (20 mM Tris, pH 8.5, 0.05% sodium azide) at a flow rate of 0.5 ml/min. A large peak which was not retained on the column, was collected and then the column was washed with buffer A containing 0.4 M NaCl until the base line returned to zero. Another peak was eluted by buffer A containing 0.6 M NaCl. The two peaks were run on a 15% (wt/vol) SDS-PAGE under reducing conditions and the band giving the expected sequence of the gC1q-R protein was determined, by NH₂-terminal peptide sequencing to be in the peak eluted with buffer A containing 0.6 M NaCl.

Gel Filtration Chromatography. The recombinant protein and the markers were dialyzed into the gel filtration running buffer (50 mM Tris-HCl, 150 mM NaCl, 0.05% (wt/vol) Na₂S₂O₅, pH 7.5) and then loaded onto a Superose 12 column, which was connected to a FPLC system (Pharmacia). The molecular size markers employed were bovine IgG (160 kD), OVA (43 kD), soybean trypsin inhibitor (20.3 kD), and cytochrome C (12.4 kD).

Purification of Total RNA and Northern Blot Analysis. Total RNA (10 μg) from each sample was electrophoresed on a formaldehyde-containing 1% (wt/vol) agarose gel. After electrophoresis, the RNA was transferred to a Hybond-N membrane (Amersham) by capillary blotting and fixed onto the membrane by UV cross-linking (XL-1500 UV crosslinker; Spectronics Corporation, AMS, Burford, UK). A high specific activity, single strand DNA probe was generated by using the PCR method, with a 1.0-kb cDNA clone as a template and the antisense primer, BM-1, using 30 cycles of denaturation for 0.5 min at 94°C, annealing for 0.5 min at 53°C, and an extension at 72°C for 0.5 min. The reaction mixture (50 μl) contained 25 pmol of the primer, 0.2 mM cold dATP, dTTP, and dGTP, 0.625 U Taq polymerase (Promega) with the manufacturer's buffer system, together with 2 μl α-[³²P]dCTP (3,000 Ci/mmol, 10 mCi/ml).

After labeling, the probe was purified on a nick column (Pharmacia) and added directly to the formamide containing prehybridization buffer. The membrane was hybridized with the probe overnight at 42°C. After hybridization, the filter was washed three times with 2× SSC, 0.1% SDS at room temperature, and then washed once in 0.1× SSC, 0.1% SDS at 65°C for 15 min. The filter was exposed to x-ray film for 7 d at -70°C.

RT-PCR. First strand cDNA from various cell lines was generated from total RNAs and AMV reverse transcriptase as discussed above, except that oligo dT was used as primer. The first strand cDNAs were used as templates and the two oligonucleotides (KS-2 and BM-1) were used as primers in the PCR reaction. PCR reactions were performed with 30 cycles of denaturation for 0.5 min at 94°C, annealing for 0.5 min at 53°C, and an extension at 72°C

for 45 s. The reaction mixture (50 μl) contained 25 pmol of each primer, 0.25 mM dNTP, 1.25 U Taq polymerase with the manufacturer's buffer system, together with 1.0 μl first strand cDNA. The PCR product (5 μl) was electrophoresed on a 2% (wt/vol) NuSieve agarose gel.

Results

Purification and Characterization of gC1q-R from Raji Cells. The purification of gC1q-R used sequentially, C1q-Sepharose CL-4B affinity chromatography (2), followed by FPLC on Mono-Q and then HPLC using a column of TSK gel DEAE-NPR (28, 32). When the C1q-binding fraction from the FPLC Mono-Q column was subjected to purification on HPLC, the profile depicted in Fig. 1 was obtained. Fractions eluting between 9 and 18 min were coated onto microtiter plates and tested for their ability to bind ¹²⁵I-C1q by the solid phase radioligand binding assay. Three major regions of activity were observed with the strongest C1q binding noted with fractions corresponding to peak 4. The position where the RO-SS/A-associated calreticulin is expected to elute is marked in Fig. 1 and is close to the position of cC1q-R which eluted at ~450 mM NaCl (peak 1). Both of these molecules possess C1q-binding activity. To assess the structure and composition of the molecules contained within this region of C1q-binding activity, 250-μl samples were taken from each region and analyzed by SDS-PAGE and autoradiography. As shown in the composite autoradiogram (*inset*, Fig. 1), peak 1 (*inset A*) contained a band of ~60 kD (non-reduced), which corresponds to the molecular weight of cC1q-R; peak 2 (*inset B*) contained a 60-kD band and an additional 80-kD band; peak 3 (*inset C*) contained predominantly the 80-kD band (nonreduced) whereas peak 4 (*inset D*) consisted of a single band that migrated with an apparent molecular mass of 33 kD. Solid phase binding studies performed on the 33-kD molecule showed that it binds to C1q very strongly in a dose-dependent and saturable manner (Fig. 2). Although the binding of this molecule to C1q is enhanced by low ionic strength conditions, significant binding (70%, *n* = 3) does occur even at physiologic ionic strength. Furthermore, whereas the binding of cC1q-R and calreticulin to whole C1q was indistinguishable from each other, the binding of the 33-kD molecule was in general, two to three times higher than either the cC1q-R or calreticulin (Fig. 2). In addition, both the 60- and 80-kD species in peak 2 (Fig. 1) appeared to bind to C1q as assessed by autoradiographic analysis of these proteins after elution from C1q-coated microtiter plates (data not shown).

The 33-kD Molecule Binds to Globular Heads of C1q. To identify the domain of C1q to which the 33-kD molecule binds, solid phase binding studies were performed by binding ¹²⁵I-33 kD to microtiter plate wells coated with C1q, cC1q, gC1q, or BSA as described in Materials and Methods. As shown in Fig. 3, the 33-kD molecule binds with high affinity to C1q and gC1q but not to cC1q, suggesting that it is a unique molecule with preferential binding to the globular heads of C1q. Because its elution position on the HPLC




Figure 1. HPLC profile of C1q-R purification. Solubilized Raji cell membranes were sequentially subjected to purification on C1q-Sepharose CL-4B and FPLC Mono-Q. Fractions containing C1q-binding activity were pooled and purified on HPLC using TSK gel DEAE-NPR column and proteins were eluted using concentration gradients of 0–500 mM NaCl (10 min) and 500 mM–1 M (3 min). The positions of cC1q-R which eluted at ~450 mM NaCl and gC1q-R which eluted at about 700 mM NaCl are indicated. (Inset) SDS-PAGE analysis of HPLC peaks. Samples from major peaks (1–4) of HPLC were radiolabeled and analyzed by a 10% SDS-PAGE (with [+], or without [–] reduction) and autoradiography. Lanes: (A) HPLC peak 1, contains cC1q-R; (B) HPLC peak 2; (C) peak 3; and (D) HPLC peak 4, shows a single chain 33-kD gC1q-R.

suggests this molecule to be highly acidic, an experiment was performed to demonstrate that the binding of this molecule to C1q, a basic protein, was not charge dependent. To this end, two very basic proteins, human IgG and properdin, were included in the binding assay and the binding of ^{125}I -gC1q-R to these molecules compared with that of C1q and gC1q. The result of a representative experiment ($n = 3$) is depicted in Fig. 4 and shows that ^{125}I -gC1q-R binds to C1q and gC1q but not to BSA, IgG, or properdin. Furthermore, gC1q-R binds to C1q whose collagen stalks have been occupied by specific anti-stalks IgG (anti-cC1q), whereas no binding was observed to IgG anti-cC1q alone (Fig. 4).

The NH₂-terminal Sequence of gC1q-R Is Not Similar to that of cC1q-R. To determine if the NH₂-terminal amino acid sequence of gC1q-R shows any relationship to that of cC1q-R, both molecules (i.e., peaks 1 and 4, Fig. 1) were

subjected to NH₂-terminal amino acid sequence analyses. The results presented in Table 1 show that the two molecules do not appear to be related. Furthermore, within the 24 amino acid residues of the 33-kD gC1q-R there are four aspartic acid and three glutamine acid residues suggestive of its acidic nature.

The NH₂-terminal Sequence of gC1q-R Contains gC1q-Binding Activity. In an effort to raise antipeptide antibody for affinity purification and immunochemical studies, an 18 amino acid residue-long synthetic peptide (gC1q-R₁₈) of gC1q-R in Table 1 (underlined), was synthesized and purified on HPLC. Since a preliminary study had indicated that the peptide binds to C1q (data not shown) it prompted the question: Does this peptide inhibit the hemolytic activity of C1q in serum? To address this question, various concentrations of either gC1q-R₁₈ or KLH-conjugated peptide were incubated




Figure 2. Binding of ^{125}I gC1q-R to C1q, and influence of ionic strength or metal ions on binding. Binding of radiolabeled gC1q-R to increasing concentrations of C1q was done in PB or PBS (150 mM NaCl) with or without 0.15 mM CaCl_2 and 1 mM MgCl_2 . For comparison, the binding of radiolabeled cC1q-R and the RO-SS/A-associated calreticulin are included.




Figure 3. Solid phase binding assay. Radiolabeled gC1q-R was incubated (2 h, 37°C) with increasing concentrations of immobilized Clq, cClq, or BSA in PBS, pH 7.5. After incubation, individual wells were washed five times and counted. Each data point represents a mean of three separate experiments run in quadruplicate.

with $10\ \mu\text{l}$ of NHS or GVB for 1 h at 37°C in a total volume of $100\ \mu\text{g}$ GVB. After incubation, the residual hemolytic activity was determined as described in Materials and Methods. Both the KLH-peptide and the gC1q-R₁₈ (data not shown) inhibited the hemolytic activity of NHS. Neither KLH alone nor irrelevant peptides of similar length and charge as the gC1q-R₁₈ had such inhibitory activity.

Anti-gC1q-R and Anti-peptide Antibodies Immunoblot a 33-kD Membrane Molecule. Polyclonal antibodies raised against the 33-kD molecule and its 18 amino acid-NH₂-terminal peptide were used in immunoblotting studies. These results shown in Fig. 5 demonstrate that both antibodies recognize the same membrane molecule which migrates with an approximate molecular mass of 33 kD.

CLSM Analysis Shows the 33-kD to Be Expressed on the Surface of Many Peripheral Blood Cells. To ascertain that gC1q-R was a surface molecule, peripheral blood mononuclear leukocytes were purified by Ficoll-Hypaque centrifugation as described




Figure 4. Binding of gC1q-R to C1q and gC1q is specific. Microtiter wells were coated with 500 ng each of either C1q or gC1q; 1,000 ng each of properdin, rabbit anti-cC1q IgG, or a preincubated mixture of C1q (500 ng) and anti-cC1q IgG (1,000 ng) to form cC1q-anti-cC1q immune complex. After washing three times and blocking with 1% BSA-TBST, ^{125}I gC1q-R was added, incubated (2 h, 37°C), and analyzed as described above. The mean percent binding of three experiments run in duplicate is indicated above each bar.

in Materials and Methods and then analyzed by incubation with either IgG of preimmune rabbit, anti-33-kD IgG or anti-peptide IgG. The bound antibodies were detected by incubation with FITC-conjugated F(ab')₂ goat anti-rabbit antibody. Approximately 90–95% of the mononuclear cell population stained with various intensities with anti gC1q-R antibodies. A representative experiment ($n = 4$) is shown in Fig. 6 and is presented as 0.5 μm optical sections of a representative stained cell.

Isolation of a cDNA Clone Coding for the C1q-binding Protein. Since the C1q-binding protein was isolated from the Raji cell line, total RNA from Raji cell was used as a template for first strand cDNA synthesis and subsequent PCR amplification, which employed primers BH-S and BH-3 (Table 2). A DNA fragment of the expected size (66 bp) was generated by PCR. This fragment was subcloned and three clones were sequenced. Based on the sequences, two exact match primers (KS-1 and KA-1) were synthesized for use in the screening of a λ gt 11 library. Three positive clones were isolated from the human B cell library and all were found to contain the same cDNA sequence (Fig. 7). The longest clone was 1,139 bp long whereas the other two clones encompassed nucleotides 106–1113, found in the longest clone. The derived amino acid sequence (Fig. 7) confirms the protein sequencing data (amino acid sequence 74–97) and indicates that




Figure 5. Western blot analysis of Raji gC1q-R. Solubilized Raji cell membrane proteins were analyzed by SDS-PAGE and Western blot analysis using anti-gC1q-R (lane 1), anti-peptide (lane 2), and anti-peptide preabsorbed with KLH-Sepharose CL-4B (lane 3). A blot with rabbit IgG from preimmune serum was negative (data not shown).

the C1q-binding protein is synthesized as a pre-pro protein of 282 residues. The mature protein which starts at residue 74 is predicted from cDNA studies to be preceded by a 60 residue-long hydrophobic stretch containing five cysteines. This in turn is predicted to be preceded by a 13 residue-long leader peptide (37). The predicted molecular weight of the 209 residue-long protein is 24.3 kD and contains three potential *N*-glycosylation sites at residues 114 (Asn-Gly-Thr), 136 (Asn-Asn-Ser), and 223 (Asn-Tyr-Thr).

Northern Blot Analysis. Total RNA from various cell lines (38) was analyzed and is shown in Fig. 8. After stringent washing (0.1 \times SSC, 0.1% SDS, 65°C, 15 min) and exposure for 7 d, strong signals were seen in both B cell lines (Daudi and Raji, Fig. 8), whereas weaker signals could be seen in HepG2 and Molt-4 cell lines. No signal was seen in K562 and only a smear could be seen in HL60. In another Northern blot carried out using the same conditions (data not shown), the U937 cell line (monocyte cell line) yielded a weaker signal than the Raji cell line but a comparable signal to that obtained with HepG2 and Molt-4 cell lines. The size of the mRNA for the C1q-binding protein is \sim 1.5–1.6 kb.

RT-PCR. Using primers BM-1 and KS-2, a PCR fragment of 468 bp should be generated from first strand cDNAs of positive cell lines. The PCR products were run into a 2% (wt/vol) NuSieve gel and the results are shown in Fig. 9. In the positive control, plasmid containing the C1q-binding protein cDNA was used as template in the PCR reaction, whereas there is no first strand cDNA template in the negative control. There are strong signals in HepG2, Molt-4, Daudi, Raji, and U937 cells, but none in K562 cells. There is a weak signal from HL60, showing that it has a low level of the C1q-binding protein mRNA which is not detectable by Northern blotting.

Production of Recombinant Protein and its Properties. The





Figure 6. Confocal laser scanning microscopic imaging (CLSM). Mononuclear leukocytes were incubated (30 min, 37°C) in GVB containing 0.02% NaN₃ with IgG preimmune rabbit serum, IgG anti-gC1q-R, or anti-gC1q-R peptide. The bound antibodies were detected by goat FITC-IgG anti-rabbit and analyzed by CLSM. Approximately 90–95% of the cells stained with both antibodies. The image in the top panel (B) is a 0.5 μm section of a representative cell stained with anti-gC1q-R and (A) is the transmitted light image of the section in (B), whereas (C) represents a composite image of (A) and (B). The bottom panel is a composite of a series of 0.5-μm sections taken from the center of the cell (“equator”) upwards to the “pole” (top left to right and bottom left to right).

```

ccggcggcgcctc#99tgcggggggcgcctaggcctgggtgtcctcttgcatctgcacgtgttcgacgtgtttccggatgctgcctcctg
M L P L I 90
5 L R C V P R V L G S S V A G L R A A A P A S P F R O I L O P
CTGCGCTGCTGGCCGCGTGTGCTGGCCCTCCGCTGCCGCGCTCCGCGCTGCCCGCCGCGCTGCGCTTCCGCGCTCTCGAAGCCG 180
35 A P R L C T R P F G L I S V R A G S E R R P G L L R P R G P
GCACCCCGCTGTGCACCCGGCCCTTCGGCTCTCAGCTGCGCGAGGTTCCGAGCGGGCGCGCGCTCTCCGCGCTCCGGGACCC 270
65 C A C G C G C G S L N T D R A F V D F I S D R I K E E R
TGGCCCTGTGGCTGTGGCTGGGCTGCTGCACACCGACGGAGCAAGCTTTTGTGATTTCCCTGAGTGATGAAATTAAGGGGAAAGA 360
95 K I Q K H K T L P K M S G G W E L E L N G T E A K L V R K V
AAATTCAGAGGCATAAAACCCCTCCCTAAGATGTCTGGAGGTTGGGAGCTGGAACGTAATGGGACAGAGCGAAATTAAGTGGGAAAGTT 450
125 A G E K I T V T F N I N N S I P P T F D G E E P S Q G Q K
GCCGGGAAAAAATCACGGTCACTTTCAACATTAACACAGCATCCACCAACATTTGATGTTGAGGAGGAAACCCCTCCGAAGGCGAGAAG 540
155 V E E O E P E L T S T P N F V V E V I K N D D G K K A L V L
GTTGAAGAAGAGGCGCTGAACCTGACATCACTCCCAATTTCTGGTTGAAGTTATAAAGAATGATGATGGCAAGAGGCCCTTGTGTTG 630
185 D C H Y P E D E V G Q E D E A E S D I F S I R E V S F Q S T
GACTGTCAATTATCCAGAGGATGAGGTTGACAGAAAGACGAGGCGTGAAGTGAACATCTTCTATCAGGGAAGTTAGCTTTCAGTCCA 720
215 G G E S E M K D T N Y T L N T D S L D M A L Y D H L H D F L A
GGCGAGTCTGAATGGAAGGATACATAATTATCACTCAACACAGATTCCTGGACTGGGCCCTATATGACACACTTAAGTATTCCTGGCC 810
245 D R G V D N T F A D E L V E L S T A L E H O E Y I T F L E D
GACCGAGGGGTGGACACACTTTTCAGATGAGCTGGTGGAGCTGACACAGCCCTGGAGCACAGGAGTACATTAATTTCTTGAAGAC 900
275 L K S F V K S Q *
CTCAGAGTTTTGTCAGAGCCAGTAgcagcagcagatgctgaaagccatggttcctcagggcggcttggccagtgcaacaatcctactc 990
tgaagctagacatgctgttgaatgcttatcatcctaatatcatgggggaaanataccaanetttaatttatatgcttgcgtctcat1080
ttatttatcattttctctgcaaaactatattttctagatttttctataacatgatag

```

Figure 7. cDNA and derived amino acid sequences of the C1q-binding protein. Nucleotide sequences are numbered (right) and amino acid sequences are numbered (left). Amino acid residue number one is the first methio-

glutathione-S-transferase (GST)-gC1q-R fusion protein was purified as described in Materials and Methods on a GST affinity column followed by digestion of fusion protein with thrombin and passage over a Mono-Q column. The various stages in the purification procedures were assessed by 15% (wt/vol) SDS-PAGE and the major bands identified by NH₂-terminal protein sequencing. The protein band identified by NH₂-terminal sequencing to be glutathione-S-transferase, had a molecular mass of 26 kD. The protein band shown to be the C1q-binding protein, had an apparent molecular

nine of the putative translated sequence. The derived amino acid sequence confirms the NH₂-terminal protein sequencing data obtained from the mature protein isolated from Raji cells (amino acid sequence residues 75-97, underlined). (*) The stop codon tag at nucleotides 924-926; (double underline) the three potential N-glycosylation sites. These sequence data are available from EMBL/GenBank/DDBJ under accession number X75913.




Figure 8. Northern blot analysis. Total RNA (10 μ g) from various cell lines were analyzed. The samples were HepG2 cells (lane 1), Molt-4 cells (lane 2); K562 cells (lane 3); Daudi cells (lane 4), Raji cells (lane 5), blank (lane 6), and KL-60 cells (lane 7). The 18S and 28S rRNA were taken as size markers. The size of the signal is \sim 1.55 b.

mass of 27-28 kD on SDS-PAGE (even though the molecular weight calculated from the amino acid sequence is only 24.3 kD). In gel filtration studies using a Superose 12 column, the recombinant C1q-binding protein was eluted at a position corresponding to 97.2 kD which is consistent with it being a tetramer of a 24.3-kD polypeptide chain.

Recombinant gC1q-R (rgC1q-R) Inhibits C1q Hemolytic Activity. The recombinant, soluble form of gC1q-R generated by expression in *E. coli* was used to see its effect on C1q hemolytic activity in serum by incubating various concentrations of the recombinant molecule with NHS as described above.




Figure 9. RT-PCR. Lanes 1 and 11 contain molecular weight standards. The samples are: HepG2 cells (lane 2); Molt-4 cells (lane 3); K562 cells (lane 4); Daudi cells (lane 5); Raji cells (lane 6); U937 cells (lane 7); HL60 cells (lane 8); negative control (lane 9); and positive control (lane 10); respectively. A PCR fragment of 468 bp could be seen in positive cell lines.




Figure 10. Inhibition of serum hemolytic activity by recombinant gC1q-R (rgC1q-R). Various concentrations of rgC1q-R were incubated with NHS as described in the legend to Fig. 5 and assessed for complement inhibitory activity.

As shown in Fig. 10, the rgC1q-R was capable of inhibiting the hemolytic activity of C1q in serum in a dose-dependent manner. To verify the specificity of this inhibition, rgC1q-R was first preincubated with C1q before addition to C1q-depleted serum. Whereas, addition of C1q preincubated with rgC1q-R did not reconstitute the hemolytic activity of C1q-depleted serum, the addition of C1q alone did, suggesting that addition of rgC1q-R to serum indeed prevents the binding of C1q to immune complexes (data not shown). The gC1q-R molecule does not, however, activate complement as assessed by C4 titration (data not shown).

Discussion

The existence of two distinct types of C1q receptors had been suggested earlier by Bordin et al. (6, 18) who showed that human diploid fibroblasts express a low affinity C1q-R for the collagen-like domain and a small subpopulation expressing high affinity C1q-R which binds to the globular region of C1q. The present report describes the purification, functional characterization, cDNA cloning, and overexpression of this molecule henceforth referred to as the gC1q-R.

The gC1q-R molecule was isolated to homogeneity using the same purification procedures for cC1q-R and employed a combination of two previously published methods (2, 28, 32). Functional and immunochemical characterization of the gC1q-R molecule revealed that: (a) it is a unique protein that does not share cross-reactive epitopes with cC1q-R; (b) it binds with high affinity to the globular heads of C1q even when the collagen tail of C1q is saturated by anti-cC1q IgG; (c) both the purified and recombinant forms of the molecule in-

hibit the hemolytic activity of C1q in serum and this inhibitory activity may reside within the 24 amino acid residues of the NH₂ terminus; and (d) anti-gC1q-R antibodies blot a 33-kD molecule from Raji cells as well as other types of cells. Furthermore, CLSM analyses of peripheral blood monocytes and lymphocytes (Fig. 6) and other blood cells (8) with anti gC1q-R antibodies show that gC1q-R is expressed on the surface of these cells, although the degree of expression was found to differ from cell type to cell type. These results indicate that gC1q-R is coexpressed with cC1q-R and is likely to be ubiquitously distributed (1).

The cDNA for the C1q-binding protein encodes a pre-pro protein of 282 residues, as determined from the first methionine, whereas NH₂-terminal sequence of the mature protein isolated from Raji cell begins with residue 74 (Leu) (Fig. 7). It is uncertain if this is due to proteolytic cleavage during protein purification or if this protein has an unusually long signal peptide of 73 residues. It seems probable that residues 1-13, or 1-7, form the signal peptide of the C1q-binding protein since they give scores of 5.5 and 4.5, respectively, in the SIGCLEAVE command of the GCG program (39). The mature protein (residues 74-282) is highly charged and is very acidic, with a calculated pI of 4.15. Out of 209 residues in the mature protein, there are 28 glutamic acid, 20 aspartic acid, 16 lysine, 5 histidine, and 4 arginine residues. In contrast, the first 73 residue-long stretch of the pre-pro protein does not have any glutamic acid, aspartic acid, lysine, or histidine residues, but does have 11 arginine residues. The mature protein has only one cystine and thus should not have any intra-chain disulfide bonding.

Since the mature protein contains many charged residues distributed all through the sequence, it is unlikely that it contains a transmembrane domain. Rather, the NH₂-terminal 73 residue-long stretch, which is too long for a signal peptide, is more likely to contain a transmembrane and a cytoplasmic domain, in addition to a signal peptide. The mature protein purified from the Raji cell line has an apparent molecular mass of 33 kD in SDS-PAGE, much bigger than the calculated molecular weight from the primary sequence (24 kD). This could be explained partly by three potential *N*-glycosylation sites and partly by its highly acidic nature. For instance, although the recombinant protein has a calculated molecular mass of 24.3 kD and should be smaller than its fusion partner, glutathione-S-transferase (26 kD), it had a larger apparent molecular weight (~27.5 kD) than GST. On the other hand, the recombinant protein behaved as a tetramer of the 24.3-kD polypeptide chain on gel filtration in nondissociating conditions, i.e., as a protein of 97.2 kD.

Northern blot studies and RT-PCR showed that this protein was expressed in a variety of different types of cell lines. Daudi and Raji cells, which are both B cell lines, express a higher amount of mRNA than the other cell lines. The protein is also expressed in the U937 (a monocyte cell line), Molt-4 (a T cell line) and HepG2 cell lines (a hepatocyte cell line), but at a lower level than in the B cell lines. It was shown by RT-PCR that HL-60 (a cell line derived from human promyelocytic leukemia, which is able to differentiate into neutrophil) expresses a low level of C1q-binding protein

mRNA which was not detected by the Northern blotting study. However, both Northern blotting and RT-PCR confirmed that there is no expression in K562 cell line (a cell line derived from human myelogenous leukaemia, which is highly undifferentiated and of the granulocytic series). Furthermore, in previous studies (8, 19), Western blot analyses showed that this protein is also expressed on eosinophils and neutrophils. On the other hand, a protein, apparently identical to the C1q-binding protein described in this paper, has been purified from HeLa cell line (a fibroblast cell line), and was initially identified as being a pre-mRNA splicing factor, designated as p32 or SF2 (40, 41). However, the claims that it was a pre-mRNA splicing factor, were withdrawn by the same group in their subsequent paper (42). The cDNA clone encoding p32 (41) lacked the first 90 nucleotides seen at the 5' end of the cDNA clone encoding the C1q-binding protein (Fig. 7) and, therefore, it was concluded that p32 lacked an initiating methionine and a signal peptide. However, both these features are seen in the amino acid sequence predicted for the C1q-binding protein (Fig. 7) and are consistent with this protein being secreted from the cell and found on the cell membrane.

Whether this novel molecule plays a primary or auxiliary role within the broad scheme of C1q-C1q-R interaction cannot be ascertained from the present studies. However, the existence of a multi-receptor system may explain in part the participation of C1q-R in such a vast array of cellular responses. It is shown that gC1q-R is able to bind to the globular heads of C1q, even when it is associated with the C1r₂-C1s₂ complex. The binding of gC1q-R to C1 complexes can inhibit the binding of C1 to IgM sensitized erythrocyte and thus block the lysis of the erythrocyte by complement. Of course, under in vivo conditions, the relative ratio of gC1q-R and C1 is not the same as in vitro experiments and, it is possible that C1 is able to bind to immune complexes, activate complement, and cross-link the immune complexes to the C1q-binding protein on effector cell surfaces, like B cell, T cell, monocytes, etc. The binding may result in various biological responses. That gC1q-R participates in a variety of biological responses is evident from a number of experiments. Aggregated C1q-mediated platelet activation which leads to the induction of α_{IIb}/β_3 integrins and expression of procoagulant activity (14) can be inhibited by either preincubation of the platelets with anti gC1q-R antibody or, preexposure of the aggregated C1q with gC1q-R₁₈ (43). In addition, the gC1q-R and cC1q-R, both of which are expressed on eosinophils, have been shown recently to mediate chemotaxis and chemokinesis (8). These results clearly demonstrate that both receptors alone or in conjunction with other cell surface molecules are involved in various C1q-mediated cellular responses. Since the C1q molecule is postulated to circulate in plasma with its globular heads potentially available for binding, it is not clear how a gC1q-R-bearing cell avoids binding to C1q. One possible hypothesis is that the conformational change of C1q followed by binding to cC1q-R must take place before the gC1q-R is engaged to induce secondary messages.

The authors are grateful to Drs. C. Milner and A. Law for total RNAs, Mr. T. Gascoyne for the synthesis of oligonucleotides, David Colflesh for confocal laser scanning microscopy, and Mrs. Maureen Veprek and Miss A. Marsland for excellent secretarial assistance.

This work was supported by grants CA-41047 from the National Cancer Institute and 900696 from the American Heart Association (National Center), by a National Institutes of Health-Fogarty Senior International Fellowship (FO6-TW01732) awarded to B. Ghebrehwet, and by grants 900994 (E. I. B. Peerschke and B. Ghebrehwet) from the American Heart Association-Wyeth Ayerst Laboratories and HL28183 from the National Heart, Lung and Blood Institute awarded to E. I. B. Peerschke. Part of this work was performed during the sabbatical leaves of B. Ghebrehwet and E. I. B. Peerschke at the MRC Immunochemistry Unit, Oxford University (UK). B. L. Lim is supported by the Croucher Foundation.

Address correspondence to Dr. B. Ghebrehwet, Department of Medicine, State University of New York, HSC, T-16, Room 040, Stony Brook, NY 11794-8161.

Received for publication 13 October 1993 and in revised form 3 March 1994.

References

1. Ghebrehwet, B. 1989. Functions associated with the C1q receptor. *Behring Inst. Mitt.* 84:204.
2. Ghebrehwet, B., L. Silvestri, and C. McDevitt. 1984. Identification of the Raji cell membrane-derived C1q inhibitor as a receptor for human C1q. Purification and immunochemical characterization. *J. Exp. Med.* 160:1375.
3. Malhotra, R., and R.B. Sim. 1989. Chemical and hydrodynamic characterization of the human leucocyte receptor for complement subcomponent C1q. *Biochem. J.* 262:625.
4. Peerschke, E.I.B., and B. Ghebrehwet. 1987. Human blood platelets possess specific binding sites for C1q. *J. Immunol.* 138:1537.
5. Tenner, A.J., and N.R. Cooper. 1981. Identification of cell types in human peripheral blood that bind C1q. *J. Immunol.* 126:1174.
6. Bordin, S., W.P. Kolb, and R.C. Page. 1983. C1q receptors on cultured human gingival fibroblasts: analysis of binding properties. *J. Immunol.* 130:1871.
7. Daha, M.R., A.M. Miltenburg, P.S. Hiemstra, N. Klar Mohamad, L.A. Van Es, and V.W. Van Hinsbergh. 1988. The complement subcomponent C1q mediates binding of immune complexes and aggregates to endothelial cells in vitro. *Eur. J. Immunol.* 18:783.
8. Kuna, P., M. Iyer, E.I.B. Peerschke, A.P. Kaplan, K.B.M. Reid, and B. Ghebrehwet. 1993. Eosinophil C1q-R induces chemokinesis and/or chemotaxis. *Mol. Immunol.* 30:26. (Abstr.)
9. Bordin, S., M. Smith, B. Ghebrehwet, D. Oda, and R.C. Page. 1992. Smooth muscle and epithelial cells express specific binding sites for the C1q component of complement. *Clin. Immunol. Immunopathol.* 63:51.
10. Bobak, D.A., M.M. Frank, and A.J. Tenner. 1988. C1q acts synergistically with phorbol dibutyrate to activate CR1-mediated phagocytosis by human mononuclear phagocytes. *Eur. J. Immunol.* 18:2001.
11. Bobak, D.A., T.A. Gaither, M.M. Frank, and A.J. Tenner. 1987. Modulation of FcR function by complement: subcomponent C1q enhances the phagocytosis of IgG-opsonized targets by human monocytes and culture-derived macrophages. *J. Immunol.* 138:1150.
12. Daha, M.R., N. Klar, R. Hoekzema, and L.A. Van Es. 1990. Enhanced Ig production by human peripheral lymphocytes induced by aggregated C1q. *J. Immunol.* 144:1227.
13. Young, K.R., J.L. Ambrus, Jr., A. Malbran, A.S. Fauci, and A.J. Tenner. 1991. Complement subcomponent C1q stimulates Ig production by human B lymphocytes. *J. Immunol.* 146:3356.
14. Peerschke, E.I.B., K.B.M. Reid, and B. Ghebrehwet. 1993. Platelet activation by C1q results in the induction of α_{IIb}/β_3 integrins (GPIIb-IIIa) and the expression of P-selectin and procoagulant activity. *J. Exp. Med.* 178:579.
15. Leu, R.W., A.Q. Zhou, B.J. Shannon, and M.J. Herriott. 1990. Inhibitors of C1q biosynthesis suppress activation of murine macrophages for both antibody-independent and antibody-dependent tumor cytotoxicity. *J. Immunol.* 144:2281.
16. Leu, R.W., A.Q. Zhou, J.A. Rummage, M.J. Kennedy, and B.J. Shannon. 1989. Exogenous C1q reconstitutes resident but not inflammatory mouse peritoneal macrophages for Fc receptor-dependent cellular cytotoxicity and phagocytosis. Relationship to endogenous C1q availability. *J. Immunol.* 143:3250.
17. Malhotra, R., S. Thiel, K.B.M. Reid, and R.B. Sim. 1990. Human leukocyte C1q receptor binds other soluble proteins with collagen domains. *J. Exp. Med.* 172:955.
18. Bordin, S., and R.C. Page. 1989. Detection of a high-affinity binding site for the globular head regions of the C1q complement protein on a human diploid fibroblast subtype. *Mol. Immunol.* 26:677.
19. Ghebrehwet, B., B.L. Lim, E.I.B. Peerschke, Y. Hong, A.C. Willis, and K.B.M. Reid. 1993. Isolation and molecular characterization of a 33 kDa cell surface protein which binds to the globular "heads" of C1q. *Mol. Immunol.* 30:12. (Abstr.)
20. Ghebrehwet, B., E.I.B. Peerschke, Y. Hong, P. Munoz, and P.D. Gorevic. 1992. Short amino acid sequences derived from C1q receptor (C1q-R) show homology with alpha chains of fibronectin and vitronectin receptors and collagen type IV. *J. Leukocyte Biol.* 51:546.
21. Reid, K.B.M. 1981. Preparation of human C1q, a subcomponent of the first component of the classical pathway of complement. *Methods Enzymol.* 80:16.
22. Yonemasu, K., and R.M. Stroud. 1971. C1q: rapid purification method for preparation of monospecific antisera for biochemical studies. *J. Immunol.* 106:304.
23. Conradie, J.D., J.E. Volanakis, and R.M. Stroud. 1975. Evidence for a serum inhibitor of C1q. *Immunochemistry.* 12:967.
24. Reid, K.B.M. 1976. Isolation by partial pepsin digestion of the three collagen-like regions present in subcomponent C1q of the first component of human complement. *Biochem. J.* 155:5.

25. Paques, E.P., R. Huber, and H. Preiss. 1979. Isolation of the globular region of the subcomponent of complement. *Hoppe-Seyler's Z. Physiol. Chem.* 360:177.
26. Bolton, A.E., and W.M. Hunter. 1973. The labelling of proteins to high specific radioactivities by conjugation to a ¹²⁵I-containing acylating agent. Application to the radioimmunoassay. *Biochem. J.* 138:259.
27. Fraker, P.J., and J.C. Speck, Jr. 1978. Protein and cell membrane iodination with a sparingly soluble chloramide 1,3,4,6-tetra chloro 3a, 6a diphenylglycoluril. *Biochem. Biophys. Res. Commun.* 80:849.
28. Peerschke, E.I.B., R. Malhotra, B. Ghebrehiwet, K.B.M. Reid, A.C. Willis, and R.B. Sim. 1993. Isolation of a human endothelial C1q receptor (C1q-R). *J. Leukocyte Biol.* 53:179.
29. Ghebrehiwet, B., S. Bossone, A. Erdei, and K.B.M. Reid. 1988. Reversible biotinylation of C1q with a cleavable bioinyl derivative-application in C1q receptor (C1q-R) purification. *J. Immunol. Methods.* 110:251.
30. Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.)* 227:680.
31. Towbin, H., T. Staehelin, and J. Gordon. 1979. Electrophoretic transfer from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA.* 76:4350.
32. Malhotra, R., A.C. Willis, J.-C. Jensenius, J. Jackson, and R.B. Sim. 1993. Structure and homology of human C1q receptor (collectin receptor). *Immunology.* 78:341.
33. Stähli, C., T. Staehelin, and V. Miggiano. 1983. Spleen cell analysis and optimal immunization for high frequency production of specific hybridomas. *Methods Enzymol.* 92:26.
34. Böyum, A. 1968. Separation of lymphocytes and erythrocytes by centrifugation. *Scand. J. Lab. Invest.* 97:77.
35. Heery, D.M., F. Gannon, and R. Powell. 1990. A simple method for subcloning DNA fragments from gel slices. *Trends Genet.* 6:173.
36. Smith, D.B., and K.S. Johnson. 1988. Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-Transferase. *Gene (Amst.)* 67:31.
37. Kyte, J., and R.R. Doolittle. 1982. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 257:105.
38. European Collection of Animal Cell Cultures Catalogue. 4th Edition. 1992. PHLS Center for Applied Microbiological Research, Salisbury, UK.
39. Program Manual for the GCG Package, Version 7. April 1991. Genetics Computer Inc., Madison, WI.
40. Krainer, A.R., G.C. Conway, and D. Kozak. 1990. Purification and characterization of pre-mRNA splicing factor SF2 from HeLa cells. *Genes & Dev.* 4:1158.
41. Krainer, A.R., A. Mayeda, D. Kozak, and G. Binns. 1991. Functional expression of cloned human splicing factor SF2: homology to RNA-binding proteins, U1 70K, and Drosophila sequencing regulators. *Cell.* 66:383.
42. Mayeda, A., A.M. Zahler, A.R. Krainer, and M.B. Roth. 1992. Two members of a conserved family of nuclear phosphoproteins are involved in pre-mRNA splicing. *Proc. Natl. Acad. Sci. USA.* 89:1301.
43. Peerschke, E.I.B., K.B.M. Reid, and B. Ghebrehiwet. 1993. Platelet activation by C1q multimers: involvement of two distinct receptors. *Mol. Immunol.* 30:42. (Abstr.)