

Research article

Open Access

## Antitumor activity and mechanisms of action of total glycosides from aerial part of *Cimicifuga dahurica* targeted against hepatoma

Ze Tian<sup>1,2</sup>, Jianyong Si<sup>2</sup>, Qi Chang<sup>2</sup>, Liang Zhou<sup>2</sup>, Shilin Chen<sup>2</sup>, Peigen Xiao<sup>2</sup> and Erxi Wu\*<sup>1</sup>

Address: <sup>1</sup>Children's Hospital Informatics Program at Harvard-MIT Division of Health Sciences and Technology, Children's Hospital Boston, Harvard Medical School, Boston, Massachusetts 02115, USA and <sup>2</sup>Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences, Peking Union Medical College, Beijing 100094, China

Email: Ze Tian - ze.tian@childrens.harvard.edu; Jianyong Si - jysi@implad.ac.cn; Qi Chang - qchang@implad.ac.cn; Liang Zhou - zhouliang\_phar@hotmail.com; Shilin Chen - slchen@implad.ac.cn; Peigen Xiao - xiaopg@public.bta.net.cn; Erxi Wu\* - erxi.wu@childrens.harvard.edu

\* Corresponding author

Published: 31 December 2007

Received: 18 June 2007

BMC Cancer 2007, 7:237 doi:10.1186/1471-2407-7-237

Accepted: 31 December 2007

This article is available from: <http://www.biomedcentral.com/1471-2407/7/237>

© 2007 Tian et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Medicinal plant is a main source of cancer drug development. Some of the cycloartane triterpenoids isolated from the aerial part of *Cimicifuga dahurica* showed cytotoxicity in several cancer cell lines. It is of great interest to examine the antiproliferative activity and mechanisms of total triterpenoid glycosides of *C. dahurica* and therefore might eventually be useful in the prevention or treatment of Hepatoma.

**Methods:** The total glycosides from the aerial part (TGA) was extracted and its cytotoxicity was evaluated in HepG2 cells and primary cultured normal mouse hepatocytes by an MTT assay. Morphology observation, Annexin V-FITC/PI staining, cell cycle analysis and western blot were used to further elucidate the cytotoxic mechanism of TGA. Implanted mouse H<sub>22</sub> hepatoma model was used to demonstrate the tumor growth inhibitory activity of TGA *in vivo*.

**Results:** The IC<sub>50</sub> values of TGA in HepG2 and primary cultured normal mouse hepatocytes were 21 and 105 µg/ml, respectively. TGA induced G<sub>0</sub>/G<sub>1</sub> cell cycle arrest at lower concentration (25 µg/ml), and triggered G<sub>2</sub>/M arrest and apoptosis at higher concentrations (50 and 100 µg/ml respectively). An increase in the ratio of Bax/Bcl-2 was implicated in TGA-induced apoptosis. In addition, TGA inhibited the growth of the implanted mouse H<sub>22</sub> tumor in a dose-dependent manner.

**Conclusion:** TGA may potentially find use as a new therapy for the treatment of hepatoma.

### Background

Hepatocellular carcinoma (HCC) is the fifth most common tumor worldwide, and the incidence of HCC has been rising over the past few decades in some areas such as Europe, USA and far eastern Asian countries [1].

Despite advances in diagnosis and standard therapies such as surgery, radiation, and chemotherapy, HCC remains a formidable challenge for clinical therapy [2-5]. In the search for new cancer therapeutics with low toxic-

ity, traditional Chinese medicines are promising candidates.

The dried rhizomes of *Cimicifuga dahurica* (Turcz) Maxim (Ranunculaceae) have been used as cooling, detoxification, antipyretic and analgesic agents for the treatment of some types of headaches and toothaches in Chinese folk medicine and were included in the Chinese Pharmacopoeia [6]. The rhizomes are traditionally the portion of the plant used for medicinal purposes in *Cimicifuga* species, however the aerial part of the plant is usually discarded. Previous phytochemical studies demonstrated that both the rhizomes and the aerial part of the species are rich in cycloartane triterpenoids [7-10]. Some biological activities of total glycosides of rhizomes of *C. dahurica* (TGR) have been investigated by earlier studies of our group. It was reported that TGR could reduce the production of Simian Immunodeficiency Virus (SIV) by inhibition of PHA stimulated <sup>3</sup>H-TdR transportation in lymph cells as well as suppression of the Sister Chromatid Exchange frequency induced by mitomycin C in human peripheral lymphocytes [11,12]. Nevertheless, there are still few reports on the bioactivity of the aerial part of *C. dahurica*. Our recent study has demonstrated cytotoxicity of TGA and three cycloartanes 23, 24 and 25-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside isolated from the aerial part of *C. dahurica* against several cancerous cell lines. These three compounds showed similar effects and induced apoptosis and G<sub>2</sub>/M cell cycle arrest in hepatoma HepG2 and leukemia HL-60 cell lines. Down regulated expression of cdc2 and COX-2 contributed to the apoptosis and cell cycle arrest in HepG2 cells [13]. However, the cytotoxic mechanism and *in vivo* anti-tumor activity of TGA is still unknown.

In the current study, we investigated the anti-tumor activity and the underlying mechanism of TGA both *in vitro* and *in vivo*. Our findings show the novel anticancer activity of TGA and this may provide a new approach to the hepatoma therapy.

## Methods

### Extraction of triterpene components from aerial part of *C. dahurica*

The aerial part of *Cimicifuga dahurica* (Turcz) Maxim (synonyms: *Actinospora dahurica* Turczaninow ex Fischer & C. A. Meyer, Index Sem. Hort. Petrop. 1: 21. 1835; *Actaea dahurica* (Turczaninow ex Fischer & C. A. Meyer) Turczaninow ex Fischer & C. A. Meyer) was collected in Maojingba, Kalaqin Qi, Inner Mongolia Autonomous Region, China, in September 1999, and was identified by Prof. Ruile Pan of the Institute of Medicinal Plant Development, Chinese Academy of Medical Sciences and Peking Union Medical College. A voucher specimen has been deposited in the Herbarium of the Institute (XA99-09).

The powdered aerial part of the plant (14.5 kg) was extracted exhaustively with 10 folds volume of 80% ethanol under refluxing for three times, one hour each time. Following combination and filtering, the solvent was evaporated under vacuum to obtain the crude extract (2.0 kg). Then the crude extract was mixed with siliceous earth and eluted with ethyl acetate. Removal of the solvent in vacuo, the TGA was (210 g) obtained.

### Determination of total content of triterpenes

Twenty nine triterpene glycosides (Table 1) including 25-anhydrocimigenol-3-O- $\beta$ -D-xylopyranoside, 23-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside, 24-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside, and 25-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside, cimigenol xylopyranoside together with ferulic acid and isoferulic acid were isolated from TGA. The total triterpene glycosides content in TGA was estimated by a colorimetric method as we described previously [14], with slight modifications. A 50- $\mu$ l aliquot of TGA methanol solution (1.50 mg/ml) was diluted with 1 ml water and then applied to a Waters Oasis HLB cartridge, which was preconditioned by rinsing with 1 ml methanol and followed by 1 ml water. The cartridge was washed with 2 ml water to remove carbohydrate compounds for interference, and then the triterpenes were eluted with 2 ml methanol from the cartridge to a clean glass tube. After drying by a gentle stream of nitrogen, a 0.2-ml aliquot of 5% vanillin acetic acid (w/v) and a 0.8-ml of aliquot of perchloric acid were added to the residue in the tube. Then the tube was kept in a 80°C water bath for 15 min. After cooling with water, the absorbance of the mixture was determined at 544 nm. The assay was conducted in triplicates. The total triterpene glycosides content was 72.67%  $\pm$  2.03 expressed as cimigenol xylopyranoside equivalents. Usually the total triterpene glycosides content in *C. racemosa* is calculated as 27-deoxyactein equivalent [15]; However since 27-deoxyactein was not isolated from TGA, cimigenol xylopyranoside, one of the main components in TGA and many other *Cimicifuga* plants was used as a standard.

### Cell culture and drug treatment

HepG2 (ATCC, Rockville, MD) cells were maintained in RPMI 1640 containing 10% FBS (Gibco, BRL, Carlsbad, CA), 2 mg/ml sodium bicarbonate, 100  $\mu$ g/ml penicillin sodium salt and 100  $\mu$ g/ml streptomycin sulfate. Cells were grown to 70% confluence, trypsinized with 0.25% trypsin-2 mM EDTA, and plated in 96 well plates. Mouse hepatocytes were isolated from normal CD-1 (ICR) mice (Beijing Vital Laboratory Animal Technology, Beijing, China) with enzymatic perfusion technique as we described previously [13]. The viability of the mouse hepatocytes, tested with Trypan blue was about 80%. In all experiments, cells were grown in RPMI-1640 medium with 10% FBS for 24 h prior to treatment.

**Table 1: Triterpene constituents from *C. dahurica* Thurez Maxim**

1	Cimilactone A [12 $\beta$ -acetoxy-3 $\beta$ - $\beta$ -D-xylopyrano-syloxy-24, 25, 26, 27-tetranor-9,19-cyclolanost-16, 23 -lactone]
2	Cimilactone B [12 $\beta$ -acetoxy-3 $\beta$ - $\beta$ -D-xylopyranosyloxy-24, 25, 26, 27-tetranor-9,19-cyclolanost-7-ene - 16, 23-lactone]
3	Cimidahuside C [12 $\beta$ -acetoxy- 15-oxo-shengmanol-3-O- $\beta$ -D-xylopyranoside]
4	Cimidahuside D [12 $\beta$ -acetoxy- 15-oxo-7, 8-didehydroshengmanol- 3-O- $\beta$ -D-xylopyranoside]
5	Cimidahuside E [(20R, 24R)-24, 25-epoxy-3 $\beta$ -( $\beta$ -D-xylopyranosyloxy)-9,19-cyclolanost-7-ene-16, 23-dione]
6	Cimidahuside F [(20R, 24R)-24, 25-epoxy-15a-hydroxy-3 $\beta$ -( $\beta$ -D-xylopyranosyloxy)-9,19-cyclolanost-7-ene-16,23-dione]
7	Cimidahuside G [(23R,24S)-15- oxo-16-enol-9,19-cyclolanostane-3-O- $\beta$ -D-xylopyranoside]
8	Cimidahuside H [(23R,24S)-15- oxo-16-enol-9, 19-cyclolanostane -7-ene-3-O- $\beta$ -D-xylopyranoside]
9	Cimidahuside I [(23R, 24S)- 12 $\beta$ -acetoxy-15-oxo-16-enol-9,19-cyclolanostane-3-O- $\beta$ -D-xylopyranoside]
10	Cimidahuside J [(23R,24S)- 12 $\beta$ -acetoxy-15-oxo-16-enol-9, 19-cyclolanostane-7-ene-3-O- $\beta$ -D-xylopyranosid e]
14	(20R, 24R)-11 $\beta$ ,24,25-trihydroxy -3- $\beta$ -( $\beta$ -D-xylopyranosyloxy)- 9,19-cyclolanost-7-ene-16,23- dione
15	25-anhydrocimigenol-3-O- $\beta$ -D-xylopyranoside
16	24-epi-7,8-didehydrocimigenol -3-O- $\beta$ -D-xylopyranoside
17	cimigenol-3-O- $\beta$ -D- xylopyranoside
18	7,8-didehydrocimigenol-3-O- $\beta$ -D-xylopyranoside
19	25-O-methylcimigenol-3-O- $\beta$ -D- xylopyranoside
20	15a-hydroxycimicidol-3-O- $\beta$ -D-xylopyranoside
21	7 $\beta$ -hydroxycimigenol-3-O- $\beta$ -D-xylopyranoside
22	12 $\beta$ -hydroxycimigenol-3-O- $\beta$ -D-xylopyranoside
23	24-O-acetyl-7, 8-didehydrocimigenol-3-O- $\beta$ -D-xylopyranoside
24	24-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside
25	25-O-methyl-24-O-acetylcimigenol-3-O- $\beta$ -D-xylopyranoside
26	12 $\beta$ -O-acetylcimiaceroside A
27	12 $\beta$ -O-acetylcimiaceroside B
28	cimiaceroside A
29	cimiaceroside B

TGA was dissolved in DMSO at a concentration of 250 mg/ml, then diluted in tissue culture medium and filtered before use. The final concentration of DMSO (0.1%) did not affect the cell viability.

#### Cytotoxicity assay

$1.5 \times 10^4$  HepG2 cells and  $8 \times 10^3$  mouse hepatocytes were seeded in 96 well plates and treated with TGA or vehicle (0.1% DMSO) at various concentrations and incubated for 48 h, followed by MTT (3-[4, 5-dimethylthiazol-2-yl]-2, 5-diphenyltetrazolium bromide) assay [16]. Briefly,  $IC_{50}$  of the TGA in HepG2 cells and normal mouse hepatocytes were derived from the dose-response curves.

#### Morphology observation in HepG2 cells

AO/EB (acridine orange/ethidium bromide) fluoresce staining method was used to observe the apoptosis morphological changes [17]. Briefly, HepG2 cells were cultured in 3.5 cm dishes and treated with TGA at concentration of 50  $\mu$ g/ml for 0, 12, 24 and 48 h respectively. After treatment, all the cultures were incubated at 37°C, 5% CO<sub>2</sub> for the indicated time. Photographs were taken under an inverted Leica fluorescence 40  $\times$  10 microscope after staining.

#### Annexin V-FITC/PI assay

Apoptosis was quantified by detecting surface exposure of phosphatidylserine in apoptotic cells using Annexin V-

FITC/PI (propidium iodide) apoptosis detection kit (BD Biosciences Clontech). Cells were seeded in 3.5 cm dishes in 1 ml medium and incubated with TGA at the dose of 25, 50 and 100  $\mu$ g/ml for 24 h, respectively. The adherent and floating cells were combined and treated according to the manufacturer's instruction and measured with FITC/PI staining using flow cytometry (Becton Dickinson, San Jose, CA). Apoptotic cells (annexin V+PI-) were differentiated from necrotic cells (annexin V+PI+, including apoptotic cells at late stage).

#### Cell cycle analysis

HepG2 cells were treated with TGA at different concentrations (25, 50 and 100  $\mu$ g/ml for 48 h) and time points (at 50  $\mu$ g/ml for 12, 24 and 48 h). Then cells were collected and fixed in 70% cold ethanol (-20°C) overnight. After washing twice with PBS, cells were resuspended in PBS. RNase A (0.5 mg/ml) and PI (2.5  $\mu$ g/ml) were added to the fixed cells for 30 min. The DNA content of cells was then analyzed with a FACSCalibur instrument (Becton Dickinson, San Jose, CA).

#### Western blotting

After treatments, cells were washed three times with ice-cold PBS and lysed with lysis buffer (50 mM Tris-HCl, pH 7.4, 10 mM EDTA, 1% Triton X-100, 26% urea, and 1 tablet/10 ml protease inhibitor cocktail tablets). Sticky DNA was removed from lysates with a sterile toothpick. The

protein concentration of the supernatant was determined by the Bradford method. The lysates were subjected to electrophoresis on a 10 % SDS-polyacrylamide gel and then transferred to a nitrocellulose membrane [18]. The nitrocellulose membrane was then incubated with mouse monoclonal anti-Bcl-2 and anti-Bax antibody (Santa Cruz Biotechnology, Santa Cruz, CA; sc-509 and sc-7480). Mouse monoclonal  $\beta$ -actin (Lab Vision, Fremont, CA) was used as an internal control. Secondary antibody to IgG conjugated to horseradish peroxidase was used. The blots were probed with the ECL Western blot detection system according to the manufacturer's instructions. The ratio of Bax/Bcl-2 was analyzed by pImage tool.

#### Antitumor evaluation on implanted mouse $H_{22}$ cells

Male CD-1 (ICR) mice (Beijing Vital Laboratory Animal Technology, Beijing, China), weighing 20–22 g, were used for implantation of hepatoma  $H_{22}$  cells (s.c.), which was maintained by weekly (i.p.) passage in CD-1 (ICR) mice. Ascites (0.2 ml of 1:6 dilution) from tumor-bearing mice 7 days after tumor inoculation were implanted (s.c.) into the armpit region of mice. Ten mice each were treated with either TGA (200, 100 and 50 mg/kg b.w., i.g.) or vehicle, once a day for 10 days, 24 h after tumor inoculation. Cyclophosphamide (15 mg/kg b.w., i.p.) was used as a positive control. The tumor inhibition rate (TIR %) was calculated as we described previously [19]. All the animal procedures were conducted in compliance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals.

#### Statistics

One-way ANOVA was used and followed by Dunnett's test.  $p < 0.05$  was considered significant.


## Results

#### Cytotoxic activity

The cytotoxicity of TGA was evaluated by an MTT assay and the  $IC_{50}$  values were derived from the dose-response curves (Fig 1). A 48 h exposure to TGA decreased the proliferation of HepG2 cells in a concentration-dependent manner with an  $IC_{50}$  at 21  $\mu$ g/ml; however, the effect on normal mouse hepatocytes showed bi-directional property: the proliferation was inhibited at higher concentrations, but promoted at lower concentrations, with an  $IC_{50}$  at 105  $\mu$ g/ml. This indicates that TGA may possess relative selective cytotoxicity to hepatoma cancer cells.

#### Induction of apoptosis in HepG2 cells by TGA

Individual apoptosis in the cell population in HepG2 cells treated by TGA was studied by fluorescence staining method. Morphological alteration, such as chromatin aggregation, nuclear and cytoplasmic condensation, and partition of cytoplasm and nucleus into membrane-bound vesicles (apoptotic bodies) were observed in




**Figure 1**  
**Cytotoxic activity of TGA in HepG2 cells and normal mouse hepatocytes.** The cells were treated with vehicle or TGA from 7.8125  $\mu$ g/ml to 250  $\mu$ g/ml. Figure shown represents one of three independent experiments.

HepG2 cells treated by TGA at 50  $\mu$ g/ml for 12, 24 and 48 h (Fig 2).

To further confirm that TGA induces HepG2 cell apoptosis, cells were stained with Annexin V-FITC and PI, and then subsequently analyzed by flow cytometry. This assay is based on the translocation of phosphatidylserine from the inner leaflet of the plasma membrane to the cell surface in the early apoptotic cells. HepG2 cells were treated with TGA at 25, 50 and 100  $\mu$ g/ml for 24 h. The dual parameter fluorescent dot plots showed the viable cell population in the lower left quadrant (annexin V<sup>-</sup>PI<sup>-</sup>), the cells at the early apoptosis are in the lower right quadrant (annexin V<sup>+</sup>PI<sup>-</sup>), and the ones at the late apoptosis are in the upper right quadrant (annexin V<sup>+</sup>PI<sup>+</sup>). As indicated in Fig. 3, in untreated cells, 0.29% of cells were Annexin V-positive/PI-negative, whereas 1.02% of cells were Annexin V/PI double positive. After treatment with TGA at 25, 50 and 100  $\mu$ g/ml for 24 h, the corresponding quantities were 0.75 and 1.76%; 9.05 and 0.93%; 13.23 and 6.09% respectively.

#### Effect of TGA on cell cycle distribution in HepG2 cells

The ability of a substance to affect specific phases of the cell cycle may provide clues to its mechanism of action. To determine the effects of TGA on the cell cycle, HepG2 cells were treated with TGA at different concentrations (25, 50 and 100  $\mu$ g/ml) and time points (at 50  $\mu$ g/ml for 12, 24 and 48 h) respectively. The cells were then stained with PI and analyzed DNA content by flow cytometry. After exposure to 25  $\mu$ g/ml of the TGA, there was an increase of cells in  $G_0/G_1$  when compared to the DMSO solvent control and a concomitant decrease of cells in S and  $G_2/M$  phases. After treatment with 50 and 100  $\mu$ g/ml of TGA, there was



**Figure 2**  
**Morphological changes of HepG2 cell line in response to TGA at 50 µg/ml for different time points.** 1–4, HepG2-cells treated with TGA at 50 µg/ml for 0, 12, 24 and 48 h, respectively.

a decrease of cells in  $G_0/G_1$  and an increase of cells in  $G_2/M$ . Moreover, the sub- $G_1$  apoptotic peak was induced by TGA in dose- and time- dependent manners (Fig 4). This indicates that TGA contains more than one component with the more active or abundant component inducing  $G_0/G_1$  arrest and the less active component inducing  $G_2/M$  arrest and/or individual component(s) in TGA exert different effects at different concentrations.


#### **Upregulation of Bax/Bcl-2 ratio**

The raise of the ratio of Bax/Bcl-2 is of benefit to apoptosis. In light of our study, following treatment with TGA at 50 µg/ml, pro-apoptotic protein Bax expression was up regulated in a time-dependent manner; the anti-apoptotic protein Bcl-2 was down regulated at 12 h and 24 h time points, but slightly up regulated at 48 h time point. At all

events, the ratio of Bax/Bcl-2 was increased during all time points compared with control. Although the ratio of Bax/Bcl-2 at 48 h time point was decreased in some degree than 24 h time point, it was still far higher than that of control (Fig 5). The reason of this might be due to more apoptosis at 48 h in HepG2 cells negatively feedback to inhibit the ratio of Bax/Bcl-2 via enhancing the expression of anti-apoptotic protein.

#### **Tumor growth inhibition of implanted $H_{22}$ cells by TGA**

After tumor implantation for 24 h, administration of TGA (200, 100 or 50 mg/kg b.w., i.g) and cyclophosphamide (15 mg/kg B.W., i.p) once a day for 10 days, could significantly suppress the growth of  $H_{22}$  tumor and TGA at 200 mg/kg was more effective than the lower dosages. One-way ANOVA followed by Dunnett's test was used for sta-




**Figure 3**  
**Flow cytometric analysis of HepG2 cells treated by TGA for 24 h.** 1, Control; 2–4, HepG2 cells treatment with 25, 50 and 100 µg/ml TGA for 24 h.

tistic analysis and significant differences were found for treatment groups *vs.* control (Table 2). In addition, marked body weight loss was observed in cyclophosphamide-treated group compared to the control group, whereas only slight body weight loss was observed in TGA-treated groups. This implies that TGA might be a promising antitumor agent with low toxicity.

**Discussion**


A major complication of chemotherapy is toxicity to normal cells, which is due to the inability of drugs to differentiate between normal and malignant cells. This often impacts the efficacy of the treatment and even makes it impossible to cure the patients. One of the requisite of cancer chemopreventive agent is elimination of damaged or malignant cell through cell cycle inhibition or induction of apoptosis without or with less toxicity in normal cells [20,21].



**Figure 4**  
**Cell cycle distributions of HepG2 cells treated with TGA at different times and dosages.** Cells were stained with PI and analyzed by flow cytometry. 1–3, HepG2 control for 12, 24 and 48 h respectively; 4–6, HepG2 cells treated with TGA at 50 µg/ml for 12, 24 and 48 h respectively. 7–8, HepG2 cells treated with TGA at 25 and 100 µg/ml for 48 h. Figures shown are one of three representative experiments.

First we investigated the cytotoxicity of TGA in HepG2 cells and primary cultured normal mouse hepatocytes. The primary cultured mouse hepatocytes were chosen as normal cells to seek selective hepatoma cytotoxic agents, because these primary cultured cells closely resemble normal cells *in vivo*. Our results indicate that TGA has relatively selective cytotoxicity to hepatoma cells based on the higher IC<sub>50</sub> value in the primary cultured normal hepato-

cytes than that of carcinoma HepG2 cells. The relative selective cytotoxicity of TGA in HepG2 cells may be due to some of the relative selective cytotoxic components 23-, 24- and 25-O-acetylcimigenol-3-O-β-D-xylopyranoside, 25-anhydrolcimigenol-3-O-β-D-xylopyranoside and hepatoprotective constituent cimigenol xylopyranoside in it [13,22,23].



**Figure 5**  
**Regulation of Bax and Bcl-2 protein expression on HepG2 cells by TGA.** Cellular lysate protein (50 µg/lane) was loaded on a 10% SDS-polyacrylamide gel, electrophoresed, and subsequently transferred onto nitrocellulose. Immunoblots were detected with antibody specific for Bcl-2 and Bax. Lysates were from HepG2 cells treated with 50 µg/ml TGA for 0, 12, 24 and 48 h, respectively. The ratio of Bax/Bcl-2 was analyzed by plmage.

Cell proliferation is governed by the cell cycle, which is the target of many anti-cancer agents. Previous studies have demonstrated that extracts and some constituents of rhizomes of *C. racemosa*, the same genus as *C. dahurica*, possess cytotoxic activity against estrogen receptor positive (MCF-7) and estrogen receptor negative (MDA-MB231 and MDA-MB-453) human breast carcinoma cell lines by induction of cell cycle arrest and apoptosis; furthermore, glycosidic fraction could induce G<sub>0</sub>/G<sub>1</sub> cell cycle arrest when tested at 30 µg/ml and G<sub>2</sub>/M arrest when tested at 60 µg/ml in MCF7 cells [15,24]. In addition, it was found that actein and a fraction of black cohosh potentiated antiproliferative effects of chemotherapy agents on human breast cancer cells in more recent research [25]. In light of our study, TGA could induce G<sub>0</sub>/G<sub>1</sub> cell cycle arrest at lower concentration (25 µg/ml) and G<sub>2</sub>/M arrest at higher concentration (50 and 100 µg/ml). This suggests that TGA contains more than one component with the more active or abundant component inducing G<sub>0</sub>/G<sub>1</sub> arrest and the less active component inducing G<sub>2</sub>/M arrest. Active components either for G<sub>0</sub>/G<sub>1</sub> or G<sub>2</sub>/M cell cycle arrests have been detected in TGA by our previ-

ous studies. 23, 24 and 25-O-acetylcimigenol-3-O-β-D-xylopyranoside, isolated from TGA could induce G<sub>2</sub>/M arrest [13]; while 25-anhydrolcimigenol-3-O-β-D-xylopyranoside, which exists in TGA, could induce G<sub>0</sub>/G<sub>1</sub> arrest [26]. There might be some other potent G<sub>0</sub>/G<sub>1</sub> active components undiscovered.

Apoptosis is a tightly regulated process, which involves changes in the expression of a distinct set of genes [27,28]. Two of the major genes responsible for regulating mitochondrial apoptosis pathway are antiapoptotic Bcl-2 and proapoptotic bax [29-31]. In particular, Bax can homodimerize with itself and heterodimerize with Bcl-2 or Bcl-xL. It appears that Bax homodimers activates apoptosis while heterodimers inhibits the process [32]. Moreover, an elevated intracellular ratio of Bax to Bcl-2 occurs during increased apoptotic cell death [33]. In our study, pronounced apoptotic cells were found in HepG2 cells treated with TGA by fluorescence staining and flow cytometric analysis. Moreover, further study showed that enhanced ratio of Bax/Bcl2 at all time points contributed to TGA induced apoptosis. The attenuation of ratio of

**Table 2: Tumor growth inhibitory effect of TGA on H<sub>22</sub> cells (mean ± SD, n = 10)**

Samples	Dosage (mg/kg)	Tumor weight (g)	Growth inhibition %
Control	-	3.28 ± 1.27	
Cyclophosphamide	15	0.93 ± 0.45 <sup>a</sup>	71.67
TGA	200	1.64 ± 0.76 <sup>b</sup>	49.92
TGA	100	1.99 ± 0.82 <sup>c</sup>	39.30
TGA	50	2.06 ± 1.30 <sup>c</sup>	36.98

<sup>a</sup>p < 0.001, <sup>b</sup>p < 0.01, <sup>c</sup>p < 0.05 vs control.



Bax/Bcl-2 at 48 h time point than that of 24 h might be the way of self-protection for cell survival. More apoptosis at 48 h might in turn, attenuate the increased ratio of Bax/Bcl-2 by negative feedback.

### Conclusion

In conclusion, for the first time, the potential anticancer activity and the underlying mechanisms of TGA against hepatoma were investigated in this study. TGA exhibited relative cytotoxicity to HepG2 cells *in vitro* and inhibited growth of H<sub>22</sub> tumor *in vivo*. The results of this study suggest that TGA might be a promising anti-hepatoma agent. Apoptosis and cell cycle arrest could be attributed, in part to its proliferating inhibition, and alteration of ratio of Bax/Bcl-2 might be one of possible mechanisms of TGA inducing apoptosis.

### Competing interests

The author(s) declare that they have no competing interests.

### Authors' contributions

ZT was mainly responsible for the experimental design, the bioactivity assays and the manuscript writing; JS and LZ did all the phytochemical work; QC did triterpene content determination and wrote the draft of that part; SC, PX and EW participated in the experimental design, technique and financial support, and manuscript writing. All authors have contributed to this work, read and approved the final manuscript.

### Acknowledgements

This work is supported by the National Natural Science Foundation of China (30470195) and National Institutes of Health training grant 2 Ta5 LM 07092-1 I. We also thank Prof. Isaac S. Kohane (Children's Hospital Boston, Harvard Medical School) for his support.

### References

- Marrero JA: **Hepatocellular carcinoma.** *Curr Opin Gastroenterol* 2006, **22**:248-253.
- Okita K: **Clinical Aspects of Hepatocellular Carcinoma in Japan.** *Intern Med* 2006, **45**:229-233.
- Bosch X, Ribes J, Borrás J: **Epidemiology of primary liver cancer.** *Semin Liver Dis* 1999, **19**:271-285.
- Zhu AX: **Hepatocellular carcinoma: are we making progress?** *Cancer Invest* 2003, **21**:418-428.
- Obi S, Yoshida H, Toune R, Unuma T, Kanda M, Sato S, Tateishi R, Teratani T, Shiina S, Omata M: **Combination therapy of intraarterial 5-fluorouracil and systemic interferon-alpha for advanced hepatocellular carcinoma with portal venous invasion.** *Cancer* 2006, **106**(9):1990-1997.
- Pharmacopoeia Commission of the People's Republic of China.** In *The Pharmacopoeia of the People's Republic of China* Chemical Industry Publishing House: Beijing; 2000:55.
- Li CJ, Chen DH, Xiao PG: **[Chemical constituents of traditional Chinese drug "sheng-ma" (*Cimicifuga dahurica*)].** *Yao Xue Xue Bao* 1993, **28**(10):777-781. [Article in Chinese]
- Li CJ, Chen DH, Xiao PG: **Chemical Constituents of Traditional Chinese Drug Sheng-ma (*Cimicifuga Dahurica*) III Structures of Cimisine C and Cimisine D.** *Acta Chimica Sinica* 1994, **52**:722-726.
- Liu Y, Chen DH, Si JY, Tu GZ, An DG: **Two new cyclolanstanol xylosides from the aerial parts of *Cimicifuga dahurica*.** *J Nat Prod* 2002, **65**:1486-1488.
- Ye WC, Zhang JW, CheU CT, Ye T, Zhao SX: **New Cycloartane Glycosides from *Cimicifuga dahurica*.** *Planta Med* 1999, **65**:770-772.
- Lin X, Cai YF, Xiao PG: **The effect of *cimicifuga dahurica* saponins on SCE frequency induced by MMC in peripheral lymphocytes of human.** *Aibian Jibian Tubian* 1994, **6**:30-3.
- Lin X, Cai YF, Xiao PG: **Inhibition of SIV *in vitro* by *cimicifuga dahurica* and its action mechanism.** *Hua Xi Yao Xue Za Zhi* 1994, **9**:221-4.
- Tian Z, Yang MS, Huang F, Li KG, Si JY, Shi L, Xiao PG: **Cytotoxicity of cyclartane triterpenoids from *cimicifuga dahurica*.** *Cancer Lett* 2005, **226**:65-75.
- Chang Q, Chen DH, Si JY, Shen LG: **Determination of total triterpene glycosides content in *Siraitia grosvenorii*.** *Zhongguo Zhong Yao Za Zhi* 1995, **20**:554-5.
- Hostanska K, Nisslein T, Freudenstein J, Reichling J, Saller R: ***Cimicifuga racemosa* extract inhibits proliferation of estrogen receptor-positive and negative human breast carcinoma cell lines by induction of apoptosis.** *Breast Cancer Res Treat* 2004, **84**:151-160.
- Carmichael J, DeGraff WC, Gazdar AF: **Evaluation of a tetrazolium-based semiautomated colorimetric assay: assessment of chemosensitivity.** *Cancer Res* 1987, **47**:936-942.
- Chan HL, Liu HQ, Tzeng BC, You YS, Peng SM, Yang M, Che CM: **Syntheses of ruthenium (II) quinonediimine complexes of cyclam and characterization of their DNA-binding activities and cytotoxicity.** *Inorg Chem* 2002, **41**:3161-3171.
- Tian Z, Xu LJ, Chen SB, Zhou L, Yang MS, Chen SL, Xiao PG, Wu E: **Cytotoxic activity of Schisandrolid and Isoschisandrolid acids involves induction of apoptosis.** *Chemotherapy* 2007, **53**:257-262.
- Tian Z, Lin G, Zheng RX, Yang MS, Xiao PG: **Anti-hepatoma activity and mechanisms of components isolated from *Aralia decaisneana*.** *World J Gastroenterol* 2006, **12**:874-9.
- Stolarska M, Mlynarski W, Zalewska-Szewczyk B, Bodalski J: **Cyto-protective effect of amifostine in the treatment of childhood neoplastic diseases--a clinical study including the pharmacoeconomic analysis.** *Pharmacol Rep* 2006, **58**(1):30-34.
- Srivastava JK, Gupta S: **Tocotrienol-rich fraction of palm oil induces cell cycle arrest and apoptosis selectively in human prostate cancer cells.** *Biochem Biophys Res Commun* 2006, **346**:447-453.
- Tian Z, Pan RL, Si JY, Xiao PG: **Cytotoxicity of cycloartane triterpenoids from aerial part of *Cimicifuga foetida*.** *Fitoterapia* 2006, **77**:39-42.
- Yamahara J, Kobayashi M, Kimura H: **Biologically active principles of crude drugs. The effect of *Cimicifugae Rhizoma* and constituents in preventive action on the carbon tetrachloride-induced liver disorder in mice.** *Shoyakugaku Zasshi* 1985, **39**:80-84.
- Einbond LS, Shimizu M, Xiao D, Nuntanakorn P, Lim JT, Suzui M: **Growth inhibitory activity of extracts and purified components of black cohosh on human breast cancer cells.** *Breast Cancer Res Treat* 2004, **83**:221-231.
- Einbond LS, Shimizu M, Nuntanakorn P, Seter C, Cheng R, Jiang B, Kronenberg F, Kennelly EJ, Weinstein IB: **Actein and a fraction of black cohosh potentiate antiproliferative effects of chemotherapy agents on human breast cancer cells.** *Planta Med* 2006, **72**:1200-1206.
- Tian Z, Zhou L, Huang F, Chen SB, Xiao PG, Yang M, Wu E: **Anti-cancer activity and mechanisms of 25-anhydrocimigenol-3-O-beta-D-xylopyranoside isolated from *Souliea vaginata* on hepatomas.** *Anti-cancer drugs* 2006, **17**:545-551.
- Cummings MC, Winterford CM, Walker NI: **Apoptosis.** *Am J Surg Pathol* 1997, **21**:88-101.
- Tong XH, Lin SG, Fujii M, Hou DX: **Molecular mechanisms of echinocystic acid-induced apoptosis in HepG2 cells.** *Biochem Biophys Res Commun* 2004, **321**:539-546.
- Zhong LT, Sarafian T, Kane DJ, Charles AC, Mah SP, Edwards RH, Bredesen DE: **Bcl-2 inhibits death of central neural cell induced by multiple agents.** *Proc Natl Acad Sci* 1993, **90**:4533-4537.
- Bruce-Keller AJ, Begley JG, Fu W, Butterfield DA, Bredesen DE, Hutchins JB, Hensley K, Mattson MP: **Bcl-2 protects isolated**

**plasma and mitochondrial membranes against lipid peroxidation induced by hydrogen peroxide and amyloid beta-peptide.** *J Neurochem* 1998, **70**:31-39.

31. Wolter KG, Hsu YT, Smith CL, Nechushtan A, Xi XG, Youle RJ: **Movement of Bax from the cytosol to mitochondria during apoptosis.** *J Cell Biol* 1997, **139**:1281-1292.
32. Adams JM, Cory S: **The Bcl-2 protein family arbiters of cell survival.** *Science* 1998, **281**:1322-1326.
33. Zha H, Reed JC: **Heterodimerization-independent functions of cell death regulatory proteins Bax and Bcl-2 in yeast and mammalian cells.** *J Biol Chem* 1997, **272**:31482-31488.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2407/7/237/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

