

Zahradníková, A., I. Zahradník, I. Györke, and S. Györke
The Journal of General Physiology. Volume 114, No. 6, December 1999. 787–798.

Page 789

Due to a typesetting error, the symbols for rate constants in Table II were printed incorrectly. The corrected table appears below:

TABLE II
Rate Constants of RyR Models Used for Simulation of Channel Activity

Rate constant	Model 1Ca	Model 2Ca	Model 3Ca	Model 4Ca	Model 5Ca	Unit
k_{on}^*	1.0×10^3	9.2×10^2	8.2×10^2	7.1×10^2	7.1×10^2	$\mu\text{M}^{-1} \cdot \text{s}^{-1}$
k_{off}^*	1.0×10^5	1.4×10^4	5.5×10^3	3.0×10^3	2.0×10^3	s^{-1}
$k_{Cn O1}$			1.0×10^4			s^{-1}
$k_{Cn O2}$			1.0×10^0			s^{-1}
$k_{O1 Cn}$			5.0×10^2			s^{-1}
$k_{O2 Cn}$			5.0×10^{-1}			s^{-1}
$k_{O1 Cn+1}$			2.0×10^0			s^{-1}
$k_{Cn+1 O1}$			6.7×10^{-1}			s^{-1}
$k_{O2 Cn+1}$			3.0×10^3			s^{-1}
$k_{Cn+1 O2}$			1.0×10^2			s^{-1}
$k_{Cn+1 I}$			5.0×10^{-1}			s^{-1}
$k_{I Cn+1}$			1.5×10^0			s^{-1}

*In Model 1Ca, $k_{RC1} = k_{on}$; $k_{C1R} = k_{off}$. In Model 2Ca–Model 5Ca, n independent subunits bind Ca^{2+} with the rate constants k_{on} and k_{off} . The rate constants in $R \leftrightarrow C1 \leftrightarrow \dots \leftrightarrow Cn \dots$ are then: $k_{RC1} = n \times k_{on}$; $k_{C1C2} = (n - 1) \times k_{on}$; \dots ; $k_{Cn-1Cn} = k_{on}$; $k_{C1R} = k_{off}$; $k_{C2C1} = 2 \times k_{off}$; \dots ; $k_{CnCn-1} = n \times k_{off}$.