



Published in final edited form as:  
*Health Psychol.* 2007 January ; 26(1): 121–125.

## Mechanisms by which Childhood Personality Traits Influence Adult Health Status:

### Educational Attainment and Healthy Behaviors

**Sarah E. Hampson,**  
*University of Surrey and Oregon Research Institute, Honolulu*

**Lewis R. Goldberg,**  
*Oregon Research Institute, Honolulu*

**Thomas M. Vogt,** and  
*Kaiser Permanente Center for Health Research Hawaii, Honolulu*

**Joan P. Dubanoski**  
*Kaiser Permanente Center for Health Research Hawaii, Honolulu*

### Abstract

**Objective**—To test a lifespan health-behavior model in which educational attainment and health behaviors (eating habits, smoking, and physical activity) were hypothesized as mechanisms to account for relations between teacher ratings of childhood personality traits and self-reported health status at midlife.

**Design**—The model was tested on 1,054 members of the Hawaii Personality and Health cohort, which is a population-based cohort participating in a longitudinal study of personality and health spanning 40 years from childhood to midlife.

**Outcome**—Self-reported health status as a latent construct indicated by general health, functional status, and body mass index.

**Results**—Childhood Agreeableness, Conscientiousness, and Intellect/Imagination influenced adult health status indirectly through educational attainment, healthy eating habits, and smoking. Several direct effects of childhood traits on health behaviors and health status were also observed.

**Conclusion**—The model extends past associations found between personality traits and health behaviors or health status by identifying a life-course pathway based on the health-behavior model through which early childhood traits influence adult health status. The additional direct effects of personality traits indicate that health-behavior mechanisms may not provide a complete account of relations between personality and health.

### Keywords

childhood Big Five; lifespan health-behavior model; longitudinal study; health behaviors; educational attainment

---

Despite the steady accumulation of evidence that personality traits prospectively predict adult health outcomes, including mortality, there are few studies of the mechanisms by which they do so (Friedman, 2000). Investigating these mechanisms will identify life-course pathways

reflecting the interplay of variables internal and external to the individual that give rise to health status at any particular stage of life. In this study, we tested a life-course pathway by which childhood personality traits influence adult health status at midlife using data from the Hawaii Personality and Health cohort.

The broad traits comprising the Big Five or five-factor model of personality have each been associated with longevity. Introversion (e.g., Wilson, Mendes de Leon, Bienias, Evans, & Bennett, 2004), low Conscientiousness (e.g., Friedman et al., 1993), low Agreeableness (e.g., Miller, Smith, Turner, Guijarro, & Hallet, 1996), Neuroticism, (e.g., Almada et al., 1991), and low intelligence (related although not directly comparable to Intellect/Imagination or Openness; e.g., Whalley & Deary, 2001) have all been identified as mortality risk factors. The lifespan health behavior model provides one theoretical account of the mechanisms by which personality traits affect morbidity and mortality: personality influences on health status are mediated by patterns of health-enhancing and health-damaging behaviors over the life course (Smith & Spiro, 2002). Consistent with this model, there is evidence of associations between all of the Big Five traits and self-reported health practices (e.g., Booth-Kewley & Vickers, 1994; Lemos-Giraldez & Fidalgo-Aliste, 1997), although Conscientiousness has received the most attention. Low Conscientiousness has been associated with numerous behavioral risk factors, including unhealthy eating, tobacco use, and lower levels of physical activity (Bogg & Roberts, 2004). Furthermore, health behaviors such as these are associated with morbidity and mortality (e.g., Thompson, Edelsberg, Colditz, Bird, & Oster, 1999; U.S. DHHS, 2000).

There are few opportunities to test lifespan health-behavior models within the same sample, one exception being the Terman Life Cycle study. Friedman et al. (1993) found support for smoking and alcohol use as partial mediators of the effects of childhood Conscientiousness on mortality. The Hawaii cohort enables the study of life-course pathways spanning four decades from personality in childhood to health status at midlife in a multiethnic population-based sample. In a previous study of this cohort, childhood personality traits predicted adult smoking, alcohol use, body mass index (BMI), and self-rated health (Hampson, Goldberg, Vogt, & Dubanoski, in press). That study also demonstrated partial mediation of the effects of Conscientiousness on self-rated health by smoking and BMI. In the present study, we tested a more complete lifespan health-behavior model involving all of the Big Five traits and two intervening mechanisms connecting childhood personality to adult health status over the life course: educational attainment and health behaviors.

Educational attainment is associated with longevity (Adler, Marmot, McEwen, & Stewart, 1999; Elo & Preston, 1996), and a healthy lifestyle may be one of the numerous benefits that education confers (Yen & Moss, 1999). Lower educational attainment is associated with less healthy eating habits (e.g., Kristal, Hedderson, Patterson, & Neuhauser, 2001), greater risk of cigarette smoking (e.g., Bergen & Caporaso, 1999), and less leisure-time physical activity (He & Baker, 2005). Educational attainment characterizes a person's life pathway in a single, summary variable. It is typically achieved in the early adult years and remains stable, so there is no question of reverse causality in its association with later outcomes. For the Hawaii cohort, in the absence of any data collection between elementary school and midlife, educational attainment provides a proxy for numerous intervening variables during those unobserved years. Moreover, childhood personality traits are reliable predictors of educational attainment both in this cohort and other longitudinal samples (Digman, 1989; Shiner, Masten, & Roberts, 2003). The health behaviors included in the model were the three key dimensions of a healthy lifestyle: healthy eating, tobacco use, and physical activity. Adult health status was measured by three indices. Self-rated general health is a remarkably reliable predictor of mortality (Idler & Benyamini, 1997), and better self-rated health is associated with higher levels of Extraversion and Agreeableness (Goodwin & Engstrom, 2002) and lower levels of Neuroticism or Negative Affect (Costa & McCrae, 1987). Functional status, as an indicator of quality of

life, is regarded as a major component of good health (Ware, & Sherbourne, 1992), and obesity (BMI) has numerous health-damaging consequences including increased risk for cardiovascular disease, cancer, and diabetes (Thompson et al., 1999).

It was hypothesized that a model depicting a life-course pathway from childhood personality traits to health status at midlife, with educational status and health behaviors serving as intervening variables, would prove a good fit to the data. In view of gender differences in the associations between personality traits and longevity (Friedman et al. 1993), the fit of the model was compared for men versus women.

## Method

### Participants

Of the 2,338 members of the original cohort for whom first and second names were available, 1,755 (75%) were found in the first phase of our location efforts.<sup>1</sup> Of those, 1,057 (60% of those located) completed the follow-up questionnaire from which the data analyzed here were derived. These participants are representative of the original children in terms of gender (47% women) and scores (gender-normed) on the Big Five personality factors (Hampson et al., 2001). They ranged in age from 41- 50 years (mean age = 45 years, SD = 2 years) at the time that they completed the follow-up survey. The adult sample for the present analysis was composed of 372 (35%) Japanese Americans, 225 (21%) Native Hawaiian or part-Hawaiian Americans, 194 (18%) European Americans, and 266 (25%) of Filipino, Chinese, Okinawan, Korean, or other Pacific Island ancestry. Three participants failed to complete the measures of eating habits, resulting in complete data for 1,054.

### Measures

Teacher assessments of childhood personality traits were obtained between 1959 and 1967 when the children were in grades 1, 2, 5, or 6. Self-reports of educational attainment, and health behaviors, and health status were obtained between 1999 and 2000 when most participants were aged between 41-50 years.

**Child personality traits**—Teachers rank-ordered all the students in their classroom from lowest to highest on each of 36 to 63 personality attributes, derived from those used by Cattell and Coan (1957), using a fixed 9-step quasi-normal distribution. Definitions developed from focus groups with teachers were provided along with each trait (e.g., “Persevering: Keeps at his (her) work until it is completed; sees a job through despite difficulties, painstaking and thorough”). The five-factor structure was recovered from analyses of selected marker traits and from analyses of all the traits included in the assessments (for details, see Goldberg, 2001). For the present analyses, factor scores on five orthogonal childhood personality factors were derived for each participant based on the analyses of all the traits available. (In contrast to some of our other reports about this cohort, these child factor scores were not separately normed for boys and girls.)

**Educational attainment**—Participants responded to the question “What is the highest grade or year of school you completed?” by selecting one of nine levels: 1 = “Eighth grade or less,” 2 = “Junior high or intermediate school,” 3 = “Some high school,” 4 = “High school graduate or GED certificate,” 5 = “Some technical school,” 6 = “Technical or nursing school graduate,” 7 = “Some college or community college,” 8 = “College graduate,” or 9 = “Postgraduate or professional degree.”

---

<sup>1</sup>The findings reported here are from the first wave of adult recruitment. To date, we have now located 1,978 and recruited 1,241 members of the original cohort.

**Eating habits**—Typical dietary patterns were assessed by a 22-item version of the Food Habits Questionnaire (Kristal, Shattuck, Henry, & Fowler, 1990) and by a 24-item modified Hawaii Food Frequency Questionnaire, both of which have been shown to correlate with more objective measures, including food records and cholesterol levels. Items assessed the frequency with which a wide variety of foods were consumed, including ones typical for various Hawaiian cultural groups (e.g., rice, spam, poi). Exploratory factor analysis of all these items was used to identify three oblique factors accounting for 29% of the variance. Three eating habits scales were formed using items with loadings higher than 0.40 on the target factor. The fiber scale measured consumption of fruit, vegetables, and low-fat dairy products (15 items,  $\alpha = .83$ ); the high-fat scale measured consumption of meat and high-fat, high-carbohydrate foods (13 items,  $\alpha = .80$ ); and the food-preparation scale measured healthy practices such as removing skin from chicken before cooking (7 items,  $\alpha = .69$ ).

**Smoking history**—Questions on smoking history were modeled on those used in national health interview surveys. Participants were asked if they had ever smoked at least 100 cigarettes in their lives. If they reported that they were current smokers, they were asked how many cigarettes they smoked on an average day. From these questions a scale assessing extent of smoking was constructed where 0 = “Never smoked,” 1 = “Ex-smoker,” 2 = “Smokes less than half a pack a day,” and 3 = “Smokes half a pack a day or more.”

**Physical activity**—A physical activity variable was constructed from the Godin questionnaire indicating the total amount of exercise over the past week, with strenuous, moderate, or mild activities weighted differently (higher scores indicated more physical activity). Godin and Shepherd (1985) reported two-week and one-month test-retest correlations of .68 and .55, respectively, and correlations ranging between 0.32 (activity monitors) and 0.56 (maximum oxygen consumption) with more objective criteria.

**Health status**—Self-assessed general health status was assessed by a widely used item from the SF-36 Medical Outcomes Health Survey (Ware & Sherbourne, 1992): “Compared to others of your same age and gender, would you say that in general your health is (1) Poor, (2) Fair, (3) Good, (4) Very Good, or (5) Excellent?” Functional status was assessed by five items from the SF-36 that assessed bodily pain, physical role functioning, work functioning, emotional role functioning, and social role functioning during the past four weeks; standard scores on each of these five variables were combined to form a composite functional status scale, with higher scores indicating better functional status ( $\alpha = .88$ ). Participants reported their height and weight, which were converted to BMI (weight in kilograms divided by height in meters squared).

## Statistical Analysis

The hypothesized life-course pathway was tested using structural equation modeling conducted with Mplus, version 3, using standard maximum likelihood estimation (Muthén & Muthén, 1998-2004). The childhood Big-Five factor scores were treated as exogenous variables. Healthy eating was measured as a latent variable with three indicators (fiber, high fat, and food preparation). In preliminary analyses, healthy eating, smoking, and physical activity did not form a latent construct so they were included individually in the model but allowed to correlate. Health status was a latent construct with three indicators: self-assessed general health status, functional status, and BMI. Correlated error was allowed among indicators of the same construct but not across constructs. Indices of fit of the model to the data included the Root Mean Square Error of Approximation (RMSEA), the Comparative Fit Index (CFI), and the Tucker-Lewis Index (TLI). The significance of parameter estimates was assessed by evaluating the critical ratio of the unstandardized estimate to the standard error. The chi-square difference test was used to compare the overall fit of two competing models. Multiple-group analyses

were conducted to test the effects of gender. Starting with the final model, with the parameter estimates constrained to be equal between the genders, estimates were sequentially freed if freeing them significantly improved the fit of the model.

## Results

Table 1 presents the means and standard deviations for all variables. The largest adult gender difference was on childhood conscientiousness. Girls in this sample were viewed as almost half a standard deviation above the mean, whereas boys were somewhat below the mean on this trait. Women had healthier eating habits, and smoked and exercised less than men.

### Model Testing

In the final model (Figure 1) with all non-significant paths removed, childhood Agreeableness, Conscientiousness, and Intellect/Imagination were positively related to educational attainment. Higher educational attainment predicted healthy eating and less smoking but, contrary to expectation, *less* physical activity. Healthy eating, less smoking, and more physical activity predicted better health status. Conscientiousness and educational attainment were also direct predictors of health status. In addition, Extraversion had a direct positive association with smoking and physical activity, and Agreeableness had a direct negative association with smoking. The fit of the final model was satisfactory (RMSEA = 0.040, 90% CI = .032, .048, CFI = .939; TLI = .908). The fit of the model without the direct path from childhood Conscientiousness to adult health status (RMSEA = .042, 90% CI = .034, .050, CFI = .932; TLI = .900) was significantly worse than the fit of the final model including this direct path ( $\chi^2$  difference = 10.13,  $df = 1$ ,  $p < .01$ ). The fit of the model without the direct path from educational attainment to adult health status (RMSEA = .047, 90% CI = .039, .055, CFI = .915; TLI = .875) was significantly worse than the final model including this direct path ( $\chi^2$  difference = 32.65,  $df = 1$ ,  $p < .001$ ). Multiple-group analysis showed that only one path in the model differed significantly between genders. Women (but not men) who were less Agreeable as children were more likely to smoke ( $-.21$ ,  $p < .001$ ).

## Discussion

A life-course pathway was supported. Children whom teachers regarded as more Agreeable, Conscientious, and Intellectual/Imaginative went on to do better educationally, and also to have healthier eating habits, to smoke less, and to report better health status at midlife. However, adult health status was also influenced by effects not involving this pathway. Conscientiousness had a direct effect, and educational attainment had a direct effect not involving health behaviors. Moreover, there were also direct effects of personality traits on health behavior that were not mediated by educational status (Extraversion on smoking and physical activity, and Agreeableness on smoking). Despite men and women differing in levels of all three health behaviors, the same model (with the exception of just one path) applied equally well to them both.

This test of a lifespan model was limited by the measures available for the Hawaii cohort. The measures were all by self-report, and they were obtained concurrently in the same questionnaire, whereas the model postulates that health behaviors over a lifetime contribute to health status. However, given the difficulty of permanently changing eating and exercise habits (Katz, 2005), it is likely that reports of typical and recent health habits reflected more than a current fad. The smoking measure did not indicate amount of smoking since initiation (e.g., pack years), which would have been a more accurate measure of lifetime exposure. The measure of physical activity failed to distinguish between work-related and leisure-time activity over a 7-day period, which may account for the negative association between physical activity and educational attainment (He & Baker, 2005).

Despite these limitations, the model suggests that educational attainment and health-behavior mechanisms (as studied here) explain some, but not all, of the influence of childhood personality on adult health status. The wider implication is that additional mechanisms are necessary. The main alternative to health-behavior models of personality and health are biological ones that postulate personality pathways through physiological mechanisms (Wiebe & Smith, 1997). Emotional Stability (Neuroticism) did not enter the present model, suggesting that its effects may be better explained by a biological model. The combination of direct and indirect effects of Conscientiousness found both here and by Friedman et al. (1993) suggests the intriguing possibility that this trait, and perhaps others, may influence health through a combination of health-behavior and other mechanisms. Further tests of personality and health models using a lifespan approach will be necessary to more fully determine the multiple mechanisms by which childhood traits have their enduring effects on health.


### Acknowledgements

This research was supported by grants from the National Institute on Aging (AG20048) and the National Institute of Mental Health (MH55600). The authors are indebted to John M. (Jack) Digman (1923-1998) for obtaining the childhood teacher assessments. We gratefully acknowledge Judy A. Andrews and William F. Chaplin for their comments on an earlier version of this manuscript. We thank Maureen Barkley for her assistance with the data analyses, and Chris Arthun for his help with manuscript preparation.

### References

- Alamada SJ, Zonderman AB, Shekelle RB, Dyer AR, Daviglus ML, Costa PT, et al. Neuroticism, cynicism, and risk of death in middle-aged men: The Western Electric Study. *Psychosomatic Medicine* 1991;53:165–175. [PubMed: 2031070]
- Adler, NE.; Marmot, M.; McEwen, B.; Stewart, J., editors. *Socioeconomic status and health in industrialized nations: Social, psychological, and biological pathways*. New York Academy of Sciences; New York: 1999.
- Bergen AW, Caporaso N. Cigarette smoking. *Journal of the National Cancer Institute* 1999;91(16):1365–1375. [PubMed: 10451441]
- Bogg T, Roberts BW. Conscientiousness and health-related behaviors: A meta-analysis of the leading behavioral contributors to mortality. *Psychological Bulletin* 2004;130:887–919. [PubMed: 15535742]
- Booth-Kewley S, Vickers RR Jr. Associations between major domains of personality and health behavior. *Journal of Personality* 1994;62:281–298. [PubMed: 7965560]
- Cattell RB, Coan RA. Child personality structure as revealed in teachers' ratings. *Journal of Clinical Psychology* 1957;13:315–327. [PubMed: 13463129]
- Costa PT Jr, McCrae RR. Neuroticism, somatic complaints, and disease: Is the bark worse than the bite? *Journal of Personality* 1987;55:299–316. [PubMed: 3612472]
- Digman JM. Five robust trait dimensions: Development, stability, and utility. *Journal of Personality* 1989;57:195–214. [PubMed: 2671337]
- Elo IT, Preston SA. Educational differentials in mortality: United States, 1979-1985. *Social Science and Medicine* 1996;42:47–57. [PubMed: 8745107]
- Friedman HS. Long-term relations of personality and health: Dynamisms, mechanisms, and tropisms. *Journal of Personality* 2000;68:1089–1107. [PubMed: 11130733]
- Friedman HS, Tucker JS, Tomlinson-Keasy C, Schwartz JE, Wingard DL, Criqui MH. Does childhood personality predict longevity? *Journal of Personality and Social Psychology* 1993;65:176–185. [PubMed: 8355139]
- Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Canadian Journal of Applied Sports Science* 1985;10:141–146.
- Goldberg LR. Analysis of Digman's child-personality data: Derivation of Big-Five factor scores from each of six samples. *Journal of Personality* 2001;69:709–743. [PubMed: 11575511]
- Goodwin R, Engstrom G. Personality and the perception of health in the general population. *Psychological Medicine* 2002;32:325–332. [PubMed: 11866326]

- Hampson SE, Goldberg SE, Vogt TM, Dubanoski JP. Forty Years on: Teachers' assessments of children's personality traits predict self-reported health behaviors and outcomes at midlife. *Health Psychology*. (in press)
- Hampson SE, Dubanoski JP, Hamada W, Marsella AJ, Matsukawa J, Suarez E, Goldberg LR. Where are they now? Locating former elementary-school students after nearly 40 years for a longitudinal study of personality and health. *Journal of Research in Personality* 2001;35:375–387.
- He XZ, Baker DW. Differences in leisure-time, household, and work-related physical activity by race, ethnicity, and education. *Journal of General Internal Medicine* 2005;20:259–266. [PubMed: 15836530]
- Idler EL, Benyamini Y. Self-rated health and mortality: A review of twenty-seven community studies. *Journal of Health and Social Behavior* 1997;38:21–37. [PubMed: 9097506]
- Katz DL. Competing claims for weight loss. *Annual Review of Public Health* 2005;26:61–88.
- Kristal AR, Hedderston MM, Patterson RE, Neuhauser ML. Predictors of self-initiated, healthful dietary change. *Journal of the American Dietetic Association* 2001;101:762–766. [PubMed: 11478472]
- Kristal AR, Shattuck AL, Henry HJ, Fowler AS. Rapid assessment of dietary intake of fat, fiber, and saturated fat: Validity of an instrument suitable for community intervention research and nutritional surveillance. *American Journal of Health Promotion* 1990;4:288–295.
- Lemos-Giráldez S, Fidalgo-Aliste AM. Personality dispositions and health related habits and attitudes: A cross-sectional study. *European Journal of Personality* 1997;11:197–209.
- Miller TQ, Smith TW, Turner CW, Guijarro ML, Hallet AJ. A meta-analytic review of research on hostility and physical health. *Psychological Bulletin* 1996;119:322–348. [PubMed: 8851276]
- Muthén, LK.; Muthén, BO. *Mplus user's guide*. 3rd ed.. Muthén & Muthén; Los Angeles, CA: 19982004.
- Shiner RL, Masten AS, Roberts JM. Childhood personality foreshadows adult personality and life outcomes two decades later. *Journal of Personality* 2003;71:1145–1170. [PubMed: 14633061]
- Smith TW, Spiro A III. Personality, health, and aging: Prolegomenon for the next generation. *Journal of Research in Personality* 2002;36:363–394.
- Thompson D, Edelsberg J, Colditz GA, Bird AP, Oster G. Lifetime health and economic consequences of obesity. *Archives of Internal Medicine* 1999;159:2177–83. [PubMed: 10527295]
- U. S. DHHS. *Reducing tobacco use: A report of the Surgeon General*. U. S. Department of Health and Human Services; Atlanta, GA: 2000.
- Ware JJ, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Medical Care* 1992;30:473–483. [PubMed: 1593914]
- Weibe, JS.; Smith, TW. Personality and health: Progress and problems in psychosomatics. In: Hogan, R.; Johnson, JA.; Briggs, SR., editors. *Handbook of Personality Psychology*. Academic Press; San Diego, CA: 1997. p. 891-918.
- Whalley LJ, Deary IJ. The longitudinal cohort study of childhood IQ and survival up to age 76. *British Medical Journal* 2001;322:819–824. [PubMed: 11290633]
- Wilson RS, Mendes de Leon CF, Bienias JL, Evans DA, Bennett DA. Personality and mortality in old age. *Journal of Gerontology* 2004;59B:110–116.
- Yen, IH.; Moss, N. Unbundling education: A critical discussion of what education confers and how it lowers risk for disease and death. In: Adler, NE.; Marmot, M.; McEwen, B.; Stewart, J., editors. *Socioeconomic status and health in industrialized nations: Social, psychological, and biological pathways*. New York Academy of Sciences; New York: 1999. p. 350-351.



**Figure 1.** Structural model for the entire sample of direct and indirect influences of childhood personality traits on adult health status through educational attainment and health behaviors. Only significant paths are included and standardized path coefficients are shown. SRH = self-rated health, FS = function status, BMI = body mass index.  
 \*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$ ,  $^\dagger$  parameters fixed at 1.



**Table 1**  
Means and Standard Deviations on all Variables for the Entire Sample (N = 1,054) and Separately for Men (n = 561) and Women (n = 493)

	Total Sample		Men		Women		Gender Difference p value
	Mean	SD	Mean	SD	Mean	SD	
Age (in years)	45.00	2.00	45.00	2.00	45.00	2.02	ns
Educational Attainment	6.74	1.87	6.60	1.92	6.89	1.80	<.01
Fiber	.00	.55	-.09	.52	.10	.56	<.001
High Fat	.00	.55	.08	.57	-.09	.52	<.001
Food Preparation	.00	.60	-.07	.59	.08	.59	<.001
Smoking	.97	1.13	1.08	1.18	.83	1.05	<.001
Physical Activity	40.39	29.52	43.88	30.90	36.44	27.39	<.001
Self-rated Health	3.39	.96	3.34	.94	3.45	.97	ns
Functional Status	.00	.69	.04	.64	-.05	.73	<.05
Body Mass Index	27.20	5.00	28.12	5.46	26.14	6.34	<.001
Childhood Personality							
Extraversion	-.00	1.02	.12	1.03	-.14	.98	<.001
Agreeableness	.02	.97	-.05	.98	.09	.98	<.05
Conscientiousness	.07	1.01	-.23	1.01	.41	.89	<.001
Emotional Stability	.02	1.00	.10	.97	-.08	1.03	<.01
Intellect/Imagination	.04	1.00	.05	1.03	.02	.96	ns

*Note.* Extraversion, Agreeableness, Conscientiousness, Emotional Stability, Intellect/Imagination, Fiber, High Fat, Food Preparation, and Functional Status were measured in standard scores.