

# First line chemotherapy with planned sequential administration of gemcitabine followed by docetaxel in elderly advanced non-small-cell lung cancer patients: a multicenter phase II study

**C Tibaldi<sup>\*,1</sup>, E Vasile<sup>1</sup>, A Antonuzzo<sup>2</sup>, R Di Marsico<sup>3</sup>, A Fabbri<sup>4</sup>, F Innocenti<sup>4</sup>, G Tartarelli<sup>5</sup>, D Amoroso<sup>5</sup>, M Andreuccetti<sup>1</sup>, M Lo Dico<sup>1</sup> and A Falcone<sup>1,6,7</sup>**

<sup>1</sup>Division of Oncology, Department of Oncology, UO Oncologia Medica, Presidio Ospedaliero, Azienda USL-6 of Livorno Viale Alfieri 36, Livorno 57100, Italy; <sup>2</sup>Department of Oncology Civil Hospital, Via Forlanini 22, Piombino 57025, Italy; <sup>3</sup>Department of Oncology Civil Hospital, Via O. Marrucci 66, Cecina 57023, Italy; <sup>4</sup>Division of Pneumology Civil Hospital, Viale Matteotti 35, Pistoia 51100, Italy; <sup>5</sup>Division of Oncology, Department of Oncology Civil Hospital, Via Aurelia 335, Viareggio 55041, Italy; <sup>6</sup>Department of Oncology, University of Pisa Via Roma 67, Pisa 56100, Italy; <sup>7</sup>Istituto Toscano Tumori (ITT) Via Taddeo Alderotti 26/N, Firenze 50139, Italy

This multicenter phase II study evaluated, in chemo-naïve patients with stage IIIB–IV NSCLC, age  $\geq 70$  and with a performance status 0–2, the activity, efficacy and tolerability of planned sequential administration of gemcitabine 1200 mg m<sup>-2</sup> on days 1 and 8 every 3 weeks for three courses followed by three cycles of docetaxel 37.5 mg m<sup>-2</sup> on days 1 and 8 every 3 weeks, provided there was no evidence of disease progression. A total of 56 patients entered the study. According to intention-to-treat analysis, the objective response rate was 16.0% (95% CI 7.6–28.3%); 23 patients (41.0%) had stable disease and 24 patients (43%) had progressive disease. Five patients who had a stable disease after three courses of gemcitabine obtained a conversion to partial response by docetaxel. Median time to progression was 4.8 months (95% CI 3.6–6.0 months) and median duration of survival was 8.0 months (95% CI 5.6–10.5 months). The 1-year survival rate was 34%. No grade 4 haematological toxicity was observed and grade 3 neutropenia and thrombocytopenia were reported in 5.4 and 3.6% of the patients, respectively. Grade 3/4 mucositis and grade 3 diarrhoea, both occurred in 3.6% of the patients and grade 3 asthenia was observed in 9% of patients. One patient reported a grade 4 skin toxicity. No treatment-related deaths occurred. Sequential gemcitabine and docetaxel is a well-tolerated and effective regimen in elderly advanced NSCLC patients.

*British Journal of Cancer* (2008) **98**, 558–563. doi:10.1038/sj.bjc.6604187 www.bjcancer.com

Published online 22 January 2008

© 2008 Cancer Research UK

**Keywords:** gemcitabine; docetaxel; advanced NSCLC; elderly patients

Lung cancer is the leading cause of cancer-related deaths in Western countries. Non-small-cell lung cancer, (NSCLC) accounts for 87% of primary lung cancers and approximately two-thirds of NSCLC patients are in an advanced stage at diagnosis (Govindan *et al*, 2006). Most malignancies, including lung cancer, occur more commonly in the elderly, with almost 50% diagnosed in patients aged  $\geq 70$  years (Havlik *et al*, 1994).

Although third-generation doublets, platinum-based regimens, represent the gold standard in the treatment of advanced NSCLC patients who have good performance status (PS), most elderly advanced patients may be more susceptible to the toxic side effects of platinum-containing combinations due to higher rates of comorbid illness and the age-related impairment of organ function so that single-agent chemotherapy, such as gemcitabine, is considered a valid therapeutic choice (Gridelli *et al*, 2003; Pfister *et al*, 2004). Moreover, the large randomised phase III MILES-trial concluded that the non-platinum containing regimen vinorelbine

plus gemcitabine did not provide a survival benefit over single-agent vinorelbine or gemcitabine, and that the two-drugs combination was more toxic than single-agent therapy (Gridelli *et al*, 2003).

Among the last generation drugs tested in NSCLC, docetaxel seems very promising. This drug has shown single-agent efficacy as a second-line treatment: in patients pre-treated with platinum-based chemotherapy as a first-line therapy, single-agent docetaxel proved to have a survival advantage over best supportive care alone (Shepherd *et al*, 2000), and in comparison to vinorelbine or ifosfamide alone (Fossella *et al*, 2000).

This drug also showed activity as a single agent in untreated elderly advanced NSCLC patients enrolled in the West Japan Thoracic Oncology Group Phase III Trial (WJTOG 9904), comparing docetaxel 60 mg m<sup>-2</sup> to vinorelbine 25 mg m<sup>-2</sup> on days 1 and 8 every 3 weeks. Indeed a statistically significant advantage in terms of progression-free survival (5.5 vs 3.1 months), response rate (22.7 vs 9.9%), and improvement of disease-related symptoms was observed in favour of docetaxel; docetaxel had a non-significantly prolonged median overall survival (14.4 vs 9.9 months) (Kudoh *et al*, 2006). In the first line setting, docetaxel demonstrated both an interesting activity and efficacy whether

\*Correspondence: Dr C Tibaldi; E-mail: tiby@katamail.com

Received 2 July 2007; revised 6 December 2007; accepted 12 December 2007; published online 22 January 2008

combined with a platinum compound (Fossella *et al*, 2003) or with gemcitabine. In particular, among non-platinum-based chemotherapy regimens, the combination docetaxel–gemcitabine is considered one of the most promising. In fact, a randomised phase II trial and two randomised phase III trials compared this association with last generation platinum-containing doublets, docetaxel–cisplatin (Georgoulas *et al*, 2001) and cisplatin–vinorelbine (Georgoulas *et al*, 2005; Pujol *et al*, 2005) yielding similar activity and efficacy. Therefore, the combination docetaxel–gemcitabine could be considered a regimen to test in elderly subjects, as tolerability, especially haematological toxicity, can be problematic (Georgoulas *et al*, 2005; Pujol *et al*, 2005; Manegold *et al*, 2006).

A possible approach to reduce the toxicity of combination regimens consists of administering the same drugs in a sequential manner. This strategy may allow the administration of full dose single agents sequentially without compromising efficacy, while reducing potential toxicity expected with concurrent administration. Preclinical models (Norton and Simon, 1977; Day, 1986) as well as recent clinical trial in NSCLC suggest a benefit for the sequential administration of chemotherapy agents. A randomised phase III trial on advanced NSCLC (Manegold *et al*, 2006), having clinically relevant haematological toxicity (defined as thrombocytopenia with platelets transfusions, anaemia with RBC-transfusions or febrile neutropenia) as a primary end point, compared a concomitant administration of gemcitabine 1000 mg m<sup>-2</sup> on days 1 and 8 and docetaxel 75 mg m<sup>-2</sup> on day 1 every three weeks for 6 courses vs their sequential administration (3 courses of gemcitabine followed by 3 courses of docetaxel at the same dosages). Clinically relevant haematological toxicity occurred less frequently in the sequential arm and the quality of life also improved with this approach.

Three randomised phase III trials (Gridelli *et al*, 2004; Schuette *et al*, 2005; Camps *et al*, 2006) have compared, in second line advanced NSCLC, docetaxel at the dose of 75 mg m<sup>-2</sup> every 3 weeks, considered the standard of care, against a weekly schedule, indicating a similar efficacy but a significantly less severe toxicity in terms of leukopenia, neutropenia and febrile neutropenia with weekly docetaxel.

We carried out a multicenter phase II study to evaluate the activity, efficacy and tolerability, of a sequential regimen consisting of three courses of gemcitabine followed by three courses of docetaxel provided there was no evidence of disease progression. The schedule chosen for docetaxel was 37.5 mg m<sup>-2</sup> on days 1 and 8 every 3 weeks according to our previous study carried out in elderly patients in a second-line setting (Tibaldi *et al*, 2006; Gridelli *et al*, 2007b).

## PATIENTS AND METHODS

### Patient selection criteria

Chemotherapy-naïve patients with histologically or cytologically confirmed NSCLC, aged ≥70 years, measurable disease, and clinical stage IIIB (cytologically positive pleural effusion or metastatic supraclavicular lymph nodes) or stage IV disease were eligible if they also met the following criteria: Eastern Cooperative Oncology Group PS ≤2, life expectancy >3 months, adequate bone marrow reserve (leukocyte count ≥4.0 × 10<sup>9</sup> l<sup>-1</sup>, platelet count ≥100 × 10<sup>9</sup> l<sup>-1</sup>), adequate hepatic function (bilirubin level ≤1.5 mg dl<sup>-1</sup>) and renal function (creatinine level ≤1.5 mg dl<sup>-1</sup>). Prior radiotherapy was allowed provided that the irradiated area was not the only source of measurable disease and that radiation therapy had been completed 7 days before chemotherapy was initiated.

Patients were excluded for the presence of active infections, concomitant malignancy, or a second primary malignancy, recent

myocardial infarction, unstable angina, congestive heart failure, dementia, symptomatic brain metastases. A written informed consent was obtained from each patient before enrolment. The protocol was approved by each local ethics committee of every institution participating to the study and the trial was conducted according to the Helsinki declaration of the World Medical Association.

### Treatment

The chemotherapy regimen consisted of gemcitabine (Gemzar<sup>®</sup>, Eli Lilly and Company, Indianapolis, Indiana, USA) 1200 mg m<sup>-2</sup> in 250 ml of normal saline administered intravenously over 30 min on days 1 and 8 every 3 weeks for 3 courses followed by, docetaxel (Taxotere<sup>®</sup>, Aventis Pharma, Antony Cedex, France) 37.5 mg m<sup>-2</sup> in 500 ml of normal saline administered intravenously over 60 min on days 1 and 8 every 3 weeks for 3 courses, provided there was no evidence of disease progression; premedication consisted of 20 mg dexamethasone i.v. and 5-hydroxytryptamine-3 receptor antagonists as antiemetic prophylaxis. Patients whose white blood cell count, neutrophil count and platelet count were greater than 3.5 × 10<sup>9</sup>, 1.5 × 10<sup>9</sup> and 100 × 10<sup>9</sup> l<sup>-1</sup>, respectively, received chemotherapy on day 1. If these conditions were not met, administration of chemotherapy was delayed 1 week or until recovery. A dose delay for more than 3 weeks resulted in withdrawal from the study. The dose of chemotherapy (gemcitabine and docetaxel) was modified on day 8 according to haematological and non-haematological toxicities as follows: if neutrophil count was >1.5 × 10<sup>9</sup> l<sup>-1</sup> and platelet count was >100 × 10<sup>9</sup> l<sup>-1</sup>, chemotherapy was administered at full dose; for neutrophil count 1.0–1.49 × 10<sup>9</sup> l<sup>-1</sup> or platelet count 75–99 × 10<sup>9</sup> l<sup>-1</sup>, the dose was reduced by 25%; for neutrophil count 0.5–0.99 × 10<sup>9</sup> l<sup>-1</sup> or platelet count 50–74 × 10<sup>9</sup> l<sup>-1</sup> the dose was reduced by 50%; for neutrophil count <0.5 × 10<sup>9</sup> l<sup>-1</sup> or platelet count <50 × 10<sup>9</sup> l<sup>-1</sup>, chemotherapy was omitted. If grade 2 non-haematological toxicity (except for alopecia) was observed, chemotherapy was omitted until resolution and then readministered at the next cycle at doses reduced by 25–50%. In case of grade 2 neurological toxicity or grade 3–4 non-haematological toxicity the patient was withdrawn from the study.

The routine use of granulocyte colony-stimulating factor was not allowed.

### Evaluation criteria

Pretreatment evaluation included medical history, physical examination, assessment of PS, complete blood cell count with differential, routine chemistry, chest radiograph, computed tomography (CT) scan of the chest and abdomen. A comorbidities assessment using the Charlson comorbidity index was carried out (Charlson *et al*, 1987). In particular, the presence or absence of the following disease states was recorded: myocardial infarction, peripheral vascular disease, cerebrovascular disease, chronic pulmonary disease, connective tissue disease, ulcer disease, mild liver disease, diabetes, hemiplegia; each disease state had a corresponding numeric value (Charlson *et al*, 1987), and the value for all present disease states were summed for each patient. The other pathologic conditions (congestive heart failure, dementia, moderate or severe kidney disease, diabetes with organ damage, any other cancer, moderate or severe liver disease, AIDS), used to calculate the Charlson score, were not taken into account because they were precluded by the exclusion criteria for the study.

During treatment, a complete blood cell count was performed before each chemotherapy administration. Eastern Cooperative Oncology Group PS was evaluated at each cycle. Evaluation of tumour response was carried out with CT scan every three cycles. Responses were assessed using standard RECIST criteria (Therasse *et al*, 2000). The best overall response for each patient was reported; all results were reviewed by an independent radiologist

and had to be confirmed 28 days or more after initial documentation of the response. Patients with disease progression after 3 courses of gemcitabine were considered as such in the final analysis. The overall response rate was calculated according to the intention-to-treat analysis.

Haematological toxicity and non-haematological toxicities were recorded at days 1 and 8 of every course of treatment. The worst toxicity grade for each patient in all cycles was reported. Toxicities were assessed using National Cancer Institute common toxicity criteria version 2.0 (NCI-CTC) (National Cancer Institute, 1999).

Quality of life was assessed using the European Organisation for Research and Treatment (EORTC) QLQ-C30 questionnaire (version 3.0) and the lung cancer-specific module (QLQ-LC13). Patients were asked to complete the questionnaires at baseline and at the end of every cycle.

### Statistical analysis

The main objective of the study was to test whether the sequential combination of gemcitabine and docetaxel would improve response rate. Secondary end points were to evaluate toxicity, time to progression and survival. Simon's two-stage minimax design was applied to calculate the sample size. Assuming  $p_0$  (low response rate) 20% and  $p_1$  (target response rate of interest for further investigation) 35% and with an  $\alpha$  error of 0.05 and a  $\beta$  error of 0.20 a total of 31 evaluable patients had to be accrued during stage 1. If at least six objective responses were observed, 22 additional patients were to be enrolled into the study during stage 2. The regimen would be considered for further investigation if  $\geq 15$  objective responses out of 53 evaluable patients were observed.

Time to progression (TTP) was calculated from the date of registration to the date of clinical and/or radiological evidence of progression or death, whichever occurred first. Survival was calculated from registration to death or last follow-up. Survival and TTP were estimated using the Kaplan–Meier method.

Since most patients had more than one comorbid medical condition, the comorbidities count was dichotomised as 0–1 or  $>1$  whereas Charlson score was dichotomised as 0 or  $\geq 1$ , as reported previously (Hesketh *et al*, 2006), for survival comparisons. Cox regression analyses were used to examine any possible relationships between overall survival and comorbidities count or Charlson score.

Data were analysed using SPSS/PC + 11.5 statistical software (SPSS Inc., Chicago, IL, USA).

## RESULTS

### Patient characteristics

From March 2005 to October 2006, 56 patients entered the study (Table 1). Forty-six patients (82%) were males, 10 (18%) were females; median age was 76 years (range 70–84). Seven patients (12.5%) had a PS = 0, 38 patients had a PS = 1 (67.8%), 11 patients had a PS = 2 (19.6%). Squamous carcinoma was the most frequent histology (20 patients); 17 patients had adenocarcinoma, one patient had bronchioloalveolar carcinoma, four patients presented large cell carcinoma and 14 patients had undifferentiated NSCLC. All patients had stage IV disease. Table 2 shows the distribution of comorbid conditions.

### Dose administration

Fifty-six patients received a total of 243 cycles, 159 cycles of gemcitabine and 84 cycles of docetaxel; the median number of total courses (gemcitabine plus docetaxel) was 5 (range, 1–6). Thirty-two patients received docetaxel and the median number of docetaxel courses was 3 (range 1–3). The delivered dose-intensity

**Table 1** Clinical characteristics

	No of patients (%)
No of Patients:	56
Age, years (range)	76 (70–84)
Sex	
Male	46 (82)
Female	10 (18)
Smoking history	
Smokers	48 (85.7)
Never smokers	8 (14.3)
ECOG PS	
0	7 (12.5)
1	38 (67.8)
2	11 (19.6)
Histology	
Adenocarcinoma	17 (30.4)
Squamous	20 (35.7)
Large cell	4 (7.1)
Bronchioloalveolar	1 (1.8)
Unspecified NSCLC	14 (25.0)
Metastatic sites	
Bone	14 (25.0)
Brain	6 (10.7)
Liver	4 (7.1)
Adrenal gland	7 (12.5)
Lymphonodes	29 (51.8)
Pleura	11 (19.6)
Lung	14 (25.0)
Number of comorbidities	
None	3 (5.3)
1	21 (37.6)
2	15 (26.7)
3	12 (21.4)
4	5 (9)
Charlson score	
0	18 (32.1)
1	28 (50.0)
2	10 (17.8)

**Table 2** Comorbid conditions

	No of patients (%)
Myocardial infarction	5 (8.9)
Ischaemic heart disease	6 (10.7)
Valvular heart disease	1 (1.8)
Atrial fibrillation	5 (8.9)
Peripheral vascular disease	3 (5.4)
Cerebrovascular disease	6 (10.7)
Hypertension	26 (46.4)
Chronic pulmonary disease	11 (19.6)
Mild liver disease	3 (5.4)
Peptic ulcer	7 (12.5)
Diabetes	12 (21.4)
Connective tissue disease	1 (1.8)
Osteoporosis	5 (8.9)
Mild depression	2 (3.6)
Mild kidney disease	4 (7.1)
Genitourinary diseases	10 (17.9)

for gemcitabine and docetaxel was 690.4 and 20.3 mg m<sup>-2</sup> per week, respectively, whereas the relative dose intensity was 86.2 and 81.2%, respectively. Twelve delays (7.5% of the courses) were reported during gemcitabine treatment. Toxicity was the reason


for the delays in only three cases (1 episode of grade 2 neutropenia, 1 episode of grade 2 skin toxicity and 1 episode of AST and ALT elevation); the reasons for the other nine delays were fever in three cases and scheduling conflict in six cases. Seventeen delays (20.2% of the courses) were reported during docetaxel treatment. The reasons for the delays were one episode of grade 3 mucositis, two episodes of grade 3 diarrhoea, one episode of grade 3 neutropenia, two episodes of grade 2 skin toxicity, one episode of creatinine increase and one episode of bilirubin increase; in nine cases the reasons were not related to toxicity (fever in five cases and scheduling conflict in four cases). Six patients discontinued treatment due to docetaxel toxicity: one patient experienced grade 4 mucositis, one patient had grade 3 mucositis, one patient experienced grade 4 skin toxicity, two patients reported grade 3 asthenia and one patient had an allergic reaction; three patients didn't discontinue treatment although they reported grade 3 asthenia. The gemcitabine dose was reduced by 25% in six administrations whereas the docetaxel dose was reduced by 25% in 16 administrations, and by 50% in four administrations.

### Response and survival

Fifty-three out of 56 patients were evaluable for response. Three patients could not be evaluated for the following reasons: one patient refused to continue chemotherapy after the first cycle, one patient died before the first evaluation probably due to rapid tumour progression, one patient was lost to the follow-up; these three patients were considered as progression disease in the final analysis.

According to the intention-to-treat analysis, the overall response rate was 16.0% (9 out of 56 patients) (95% CI 7.6–28.3%); twenty-three patients (41.0%) had stable disease and twenty-four patients (43%) had progressive disease. Five patients who had a stable disease after three courses of gemcitabine obtained a conversion to partial response by docetaxel.

Median time to progression was 4.8 months (95% CI 3.6–6.0 months) and median duration of survival was 8.0 months (95% CI 5.6–10.5 months). The 1-year survival rate was 34% (Figure 1). At an exploratory planned analysis, the subgroup of patients with PS 0–1 had a median time to progression of 4.8 months (95% CI 2.6–7.0 months), a median duration of survival of 8.7 months (95% CI 7.4–9.9 months) and 1-year survival rate of 40%; patients with PS 2 reported a median time to progression and a median overall



**Figure 1** Kaplan–Meier plot of overall survival (OS) and time to progression (TTP).

survival of 4.0 months (95% CI 0.6–7.3 months) and 5.4 months (1.3–9.4 months), respectively, with a 1-year survival rate of 15%.

Patients with a Charlson score of 0 (18 patients) or  $\geq 1$  (38 patients) had a median survival of 9.7 months and 7.9 months, respectively ( $P=0.82$ ).

Since several comorbidities were not taken into account by Charlson score we analysed the median survival according to the number of comorbidities: the median survival was 8.6 months for patients with 0–1 comorbidities and 5.4 months for patients with  $>1$  comorbidities ( $P=0.64$ ).

Although planned, quality of life assessment isn't reported because of the lacking of sufficient data due to the high rate of missing questionnaires.

### Toxicity

All 56 patients were evaluable for toxicity. Observed toxicities were mild and the compliance to treatment was good. No grade 4 haematological toxicity was observed and grade 3 neutropenia and thrombocytopenia were reported in 5.4 and 3.6% of the patients, respectively. No patients developed febrile neutropenia or haemorrhages. Non-haematological toxicity consisted mainly in grade 3/4 mucositis and grade 3 diarrhoea both occurring in two (3.6%) patients, grade 3 asthenia observed in five (9%) patients. One patient reported a grade 4 skin toxicity and two patients reported a grade 3 nail toxicity. No treatment-related deaths occurred. Haematological and non-haematological toxicities are summarised in Table 3.

### DISCUSSION

The present trial, targeting elderly (age 70 years and older) advanced-stage NSCLC patients, was designed to test the hypothesis that planned sequential administration of gemcitabine followed by docetaxel may improve response rate with good tolerability. We hypothesised that the introduction of docetaxel in our first line sequential regimen could enhance the response rate with respect to our previous trial carried out in 110 elderly advanced NSCLC patients treated with gemcitabine alone (Tibaldi *et al*, 2005). In the present trial, the overall response rate according to an intention to treat analysis was 16.0% that was similar to our previous trial (13.9%). Nevertheless, it is remarkable that five out of nine responses that we observed were obtained from a conversion of stable disease to partial response by docetaxel.

We observed a median time to progression (TTP) of 4.8 months, a median overall survival (OS) of 8.0 months, and 1-year survival rate of 34%, that can be considered encouraging. In our previous trial, in fact, we reported a median TTP of 3.2 months, a median OS of 5.4 months and 1-year survival rate of 27% (Tibaldi *et al*, 2005). In addition, the percentage of PS 2 patients enrolled in both trials was similar at 20%.

**Table 3** Haematological and non-haematological toxicity per patient

	NCI-CTC grade % (56 patients)			
	1	2	3	4
Neutropenia	12.5	10.7	5.4	0
Thrombocytopenia	8.9	1.8	3.6	0
Anaemia	46.4	18	0	0
Nausea/vomiting	23.2	3.6	0	0
Diarrhoea	14.3	3.6	3.6	0
Mucositis	16.1	3.6	1.8	1.8
Asthenia	20	32	9	0
Nail toxicity	15	3.6	–	–
Skin toxicity	5.4	3.6	0	1.8

The introduction of docetaxel in first-line treatment of advanced NSCLC resulted advantageous in progression-free survival in a recent randomised phase III trial that compared immediate docetaxel with docetaxel upon evidence of progression in non-progressing patients after four cycles of induction therapy with gemcitabine plus carboplatin. The progression-free survival in the immediate docetaxel arm (6.5 months) was significantly greater ( $P=0.0001$ ) than in the delayed docetaxel arm (2.8 months) (Fidias *et al*, 2007).

Haematological and non-haematological toxicities, in our study, were mild and acceptable. The compliance to treatment was in general good. In particular, the present trial confirms our previous observation (Tibaldi *et al*, 2006) that a modified schedule of docetaxel ( $37.5 \text{ mg m}^{-2}$  on days 1 and 8 every 3 weeks) is feasible and well tolerated in elderly advanced NSCLC patients. Our tolerability data confirm the previous observations that weekly docetaxel is advantageous in terms of haematological toxicity with respect to 3-weekly schedule (Gridelli *et al*, 2004; Schuette *et al*, 2005; Camps *et al*, 2006).

Recently, the results of the SWOG (S0027) phase II trial using a sequential regimen of vinorelbine  $25 \text{ mg m}^{-2}$  on days 1 and 8 every 3 weeks for three cycles followed by three courses of docetaxel  $35 \text{ mg m}^{-2}$  on days 1, 8, 15 every 4 weeks in patients 70 years of age and older and in patients with PS two of any age were published (Hesketh *et al*, 2006). Patients aged  $\geq 70$  years with a PS of 0–1 had a median TTP and a median OS, of 4.7 and 9.1 months respectively, comparable to our results. However, some toxicities were more common such as grade 3/4 neutropenia and grade 3/4 fatigue, seen in 32 and 22% of patients, respectively. Another recent randomised phase II trial evaluated chemotherapy with pemetrexed alone vs a sequential pemetrexed/gemcitabine regimen in patients who were elderly ( $\geq 70$  years) or younger than 70 years and ineligible for platinum-based chemotherapy. The median TTP and OS reported were lower than expected and equal, in both arms, to 4.1 and 5.4 months, respectively (Gridelli *et al*, 2007a). A possible explanation for these results might be the high proportion of PS 2 patients (35.6%), who have poorer prognosis, enrolled into this study. However, although some trials grouped elderly patients with PS 2 patients of any age, these cohorts of patients have probably different characteristics and prognosis and should be

included in different and specific dedicated clinical trials in the future.

Two multicenter phase II trials evaluated a weekly combination regimen of docetaxel and gemcitabine in advanced NSCLC; the first trial (Neubauer *et al*, 2005) reported a response rate of 20%, a median TTP and OS of 5.1 and 6.9 months, respectively, that favourably compare with our results; the second trial, that used a patient selection criteria based on age, PS and Charlson score, reported a higher response rate of 34%, but a similar median TTP and OS of 5 and 7 months, respectively (LeCaer *et al*, 2007). However, the frequency of grade III–IV fatigue was high and reported in 30% of patients.

In our study, we did not observe significant differences in overall survival according to Charlson score (0 or  $> 1$ ); although, the Charlson score did not correlate with PS (Dujon *et al*, 2006), it seems to be insufficient to screen elderly advanced NSCLC patients (Maione *et al*, 2005). The use of a comprehensive geriatric assessment, according to the recommendations of the SIOG (Extermann *et al*, 2005), appears crucial to improve the selection and stratification of elderly patients and thereby to allow valid comparisons among different studies.

In this setting, quality of life evaluation has prognostic value for survival (Maione *et al*, 2005); nevertheless, in our study we did not report this analysis because of a lack of sufficient questionnaires and quality of life assessment should be recommended in future trials.

At present, the role of platinum-based combinations in elderly advanced NSCLC is not clear; retrospective subset analyses of a number of phase III trials suggested that combination platinum-based therapy is superior to single agents in both younger and older patients (Langer *et al*, 2002; Lilenbaum *et al*, 2005); however, prospective randomised trials are lacking to confirm this hypothesis and the results of the MILES-2 study, comparing gemcitabine vs cisplatin/vinorelbine and vs cisplatin/gemcitabine, are eagerly awaited.

In conclusion, our results suggest that the sequential use of gemcitabine and docetaxel is another viable, option for elderly advanced NSCLC patients. The use of standardised specific geriatric evaluations appears to be crucial for future trials in this setting.

## REFERENCES

- Camps C, Massutti B, Jimenez AM, Maestu I, Garcia Gomez R, Isla D, Gonzales JL, Almenar D, Blasco A, Rosell R, Carrato A, Vinolas N, Batista N, Garcia Giron C, Galan A, Lopez M, Blanco R, Provencio M, Diz P, Felip E (2006) Randomized phase III study of 3-weekly vs weekly docetaxel in pretreated advanced non-small-cell lung cancer: a Spanish Lung Cancer Group Trial. *Ann Oncol* **47**: 467–472
- Charlson ME, Pompei P, Ales KL, MacKenzie CR (1987) A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* **40**: 373–383
- Day RS (1986) Treatment sequencing, asymmetry, and uncertainty: protocol strategies for combination chemotherapy. *Cancer Res* **46**: 3876–3885
- Dujon C, Azarian R, Azarian V, Petitpretz P (2006) Lung Cancer in the elderly: performance status and/or geriatric indices? *Rev Mal Respir* **23**: 307–318
- Extermann M, Aapro M, Bernabei R, Cohen HJ, Droz JP, Lichtman S, Mor V, Monfardini S, Repetto L, Sorbye L, Topinkova E (2005) Use of comprehensive geriatric assessment in older cancer patients: recommendations from the task force on CGA of the International Society of Geriatric Oncology (SIOG). *Crit Rev Oncol Hematol* **55**: 241–252
- Fidias P, Dakhil S, Lyss A, Loesch D, Waterhouse D, Cunneen J, Ye Z, Tai F, Obasaju C, Schiller J (2007) Phase III study of immediate vs delayed docetaxel after induction therapy with gemcitabine plus carboplatin in advanced non-small cell lung cancer: update report with survival. *Proc Am Soc Clin Oncol* **25**: LBA7516
- Fossella F, Pereira JR, von Pawel J, Pluzanska A, Gorbounova V, Kaukel E, Mattson KV, Ramlau R, Szczesna A, Fidias P, Millward M, Belani CP (2003) Randomized, multinational, phase III study of docetaxel plus platinum combinations vs vinorelbine plus cisplatin for advanced non-small-cell lung cancer: The TAX 326 Study Group. *J Clin Oncol* **21**: 3016–3024
- Fossella FV, DeVore R, Kerr RN, Crawford J, Natale RR, Dunphy F, Kalman L, Miller V, Lee JS, Moore M, Gandara D, Karp D, Vokes E, Kris M, Kim Y, Gamza F, Hammershaimb L (2000) Randomized phase III trial of docetaxel vs vinorelbine or ifosfamide in patients with advanced non-small-cell lung cancer previously treated with platinum-containing chemotherapy regimens. The TAX 320 Non Small Cell Lung Cancer Study Group. *J Clin Oncol* **18**: 2354–2362
- Georgoulas V, Ardavanis A, Tsiafaki X, Agelidou A, Mixalopoulou P, Anagnostopoulou O, Ziotopoulos P, Toubis M, Syrigos K, Samaras N, Polyzos A, Christou A, Kakolyris S, Kouroussis C, Androulakis N, Samonis G, Chatzidakis D (2005) Vinorelbine plus cisplatin vs docetaxel plus gemcitabine in advanced non-small-cell lung cancer: a phase III randomized trial. *J Clin Oncol* **23**: 2937–2945
- Georgoulas V, Papadakis E, Alexopoulos A, Tsiafaki X, Rapti A, Veslemes M, Palamidis P, Vlachonikolis I, Greek Oncology Cooperative Group (GOCC) for Lung Cancer (2001) Platinum-based and non-platinum-based chemotherapy in advanced non-small-cell lung cancer: a randomised multicentre trial. *Lancet* **357**: 1478–1484
- Govindan R, Page N, Morgensztern D, Read W, Tierney R, Vlahiotis A, Spitznagel E, Piccirillo J (2006) Changing epidemiology of small-cell lung

- cancer in the United States over the last 30 years: analysis of the surveillance, epidemiologic, and end results database. *J Clin Oncol* **24**: 4539–4544
- Gridelli C, Gallo C, Di Maio M, Barletta E, Illiano A, Maione P, Salvagni S, Piantedosi FV, Palazzolo G, Caffo O, Ceribelli A, Falcone A, MAzzanti P, Brancaccio L, Capuano MA, Isa L, Barbera S, Perrone F (2004) A randomized clinical trial of two docetaxel regimens (weekly vs 3 week) in the second-line treatment of non small cell lung cancer. The DISTAL 01 study. *Br J Cancer* **91**: 1996–2004
- Gridelli C, Kaukel E, Gregorc V, Migliorino MR, Müller TR, Manegold C, Favaretto A, Martoni A, Caffo O, Schmittel A, Rossi A, Russo F, Peterson P, Muñoz M, Reck M (2007a) Single-agent pemetrexed or sequential pemetrexed/gemcitabine as front-line treatment of advanced non-small cell lung cancer in elderly patients or patients ineligible for platinum-based chemotherapy: a multicenter, randomized, phase II trial. *J Thoracic Oncol* **2**: 221–229
- Gridelli C, Langer C, Maione P, Rossi A, Schild SE (2007b) Lung cancer in the elderly. *J Clin Oncol* **14**: 1898–1907
- Gridelli C, Perrone F, Gallo C, Cigolari S, Rossi A, Piantedosi F, Barbera S, Ferrau F, Piazza E, Rosetti F, Clerici M, Bertetto O, Robbiati SF, Frontini L, Sacco C, Castiglione F, Favaretto A, Novello S, Migliorino MR, Gasparini G, Galetta D, Iaffaioli RV, Gebbia V, MILES Investigators (2003) Chemotherapy for elderly patients with advanced non-small-cell lung cancer: The Multicenter Italian Lung Cancer in the Elderly Study (MILES) Phase III Randomized Trial. *J Natl Cancer Inst* **95**: 362–372
- Havlik RJ, Yancik R, Long S, Ries L, Edwards B (1994) The National Cancer Institute on aging and the National Cancer Institute SEER collaborative study on comorbidity and early diagnosis of cancer in the elderly. *Cancer* **74**(suppl 7): 2101–2106
- Hesketh PJ, Chansky K, Lau DH, Doroshow JH, Moinpour CM, Chapman RA, Goodwin JW, Gross HM, Crowley JJ, Gandara DR (2006) Sequential vinorelbine and docetaxel in advanced non-small cell lung cancer patients age 70 and older and/or with a performance status of 2: A Phase II Trial of the Southwest Oncology Group (S0027). *J Thoracic Oncol* **1**: 537–544
- Kudoh S, Takeda K, Nakagawa K, Takada M, Katakami N, Matsui K, Shinkai T, Sawa T, Goto I, Semba H, Seto T, Ando M, Satoh T, Yoshimura N, Negoro S, Fukuoka M (2006) Phase III study of docetaxel compared with vinorelbine in elderly patients with advanced non-small-cell lung cancer: results of the West Japan Thoracic Oncology Group Trial (WJTOG 9904). *J Clin Oncol* **22**: 3657–3663
- Langer CJ, Manola J, Bernardo P, Kugler JW, Bonomi P, Cella D, Johnson DH (2002) Cisplatin-based therapy for elderly patients with advanced non-small-cell lung cancer: implications of Eastern Cooperative Oncology Group 5592, a randomized trial. *J Natl Cancer Inst* **94**: 173–181
- LeCaer H, Fournel P, Jullian H, Chouaid C, LeTret J, Thomas P, Paillotin D, Perol M, Gimenez C, Vernenege A (2007) An open multicenter phase II trial of docetaxel-gemcitabine in Charlson score and performance status (PS) selected elderly patients with stage IIIB pleura/IV non-small-cell lung cancer (NSCLC): The GFPC 02-02a Study. *Crit Rev Oncol Hematol* **64**: 73–81
- Lilenbaum RC, Herndon JE, List MA, Desch C, Watson DM, Miller AA, Graziano SL, Perry MC, Saville W, Chahinian P, Weeks JC, Holland JC, Green MR (2005) Single-agent vs combination chemotherapy in advanced non-small-cell lung cancer: the Cancer and Leukemia Group B (Study 9730). *J Clin Oncol* **23**: 190–196
- Maione P, Perrone F, Gallo C, Manzione L, Piantedosi FV, Barbera S, Cigolari S, Rosetti F, Piazza E, Robbiati SF, Bertetto O, Novello S, Migliorino MR, Favaretto F, Spatafora M, Ferrau F, Frontini L, Bearz A, Repetto LC (2005) Pretreatment quality of life and functional status assessment significantly predict survival of elderly patients with advanced non-small-cell lung cancer receiving chemotherapy: a prognostic analysis of the Multicenter Italian Lung Cancer in the Elderly Study. *J Clin Oncol* **23**: 6865–6872
- Manegold C, Thatcher N, Kortsik C, Koschel G, Mezger J, Schott Von Römer K, Pilz LR (2006) Sequencing of single-agent (SA) docetaxel (T) and gemcitabine (G) therapy for patients (pts) with advanced non-small cell lung cancer (NSCLC): results of three consecutive randomized trial. *Proc Am Soc Clin Oncol* **24**: 7035
- National Cancer Institute (1999) Cancer Therapy Evaluation Programs (CTEP) Common toxicity criteria-Version 2.0. Bethesda (MD). Division of Cancer Treatment and Diagnosis, National Cancer Institute. Available at: <http://ctep.cancer.gov/reporting/ctc.html>
- Neubauer MA, Garfield DH, Kuerfler PR, Raju RN, Lindquist DL, Soo EW, Khan M, Boehm KA, Asmar L (2005) Results of a phase II multicenter trial of weekly docetaxel and gemcitabine as first-line therapy for patients with advanced non-small cell lung cancer. *Lung Cancer* **47**: 121–127
- Norton L, Simon R (1977) Tumor size, sensitivity to therapy, and design of treatment schedules. *Cancer Treat Rep* **61**: 1307–1317
- Pfister DG, Johnson DH, Azzoli CG, Sause W, Smith TJ, Baker S, Olak J, Stover D, Strawn JR, Turrisi AT, Somerfield MR, American Society of Clinical Oncology (2004) American Society of Clinical Oncology treatment of unresectable non-small-cell lung cancer guideline: update 2003. *J Clin Oncol* **22**: 330–353
- Pujol JL, Breton JL, Gervais R, Rebattu P, Depierre A, Morère JF, Milleron B, Debieuvre D, Castéra D, Souquet PJ, Moro-Sibilot D, Lemarié E, Kessler R, Janicot H, Braun D, Spaeth D, Quantin X, Clary C (2005) Gemcitabine-docetaxel vs cisplatin-vinorelbine in advanced or metastatic non-small-cell lung cancer: a phase III study addressing the case for cisplatin. *Ann Oncol* **16**: 602–610
- Schuette W, Nagel S, Blankenburg T, Lautenschlaeger C, Hans K, Schmidt EW, Dittrich I, Schweisfurth H, Fischer von Weikersthal L, Raghavachar A, Reißig A, Serke M (2005) Phase III study of second-line chemotherapy for advanced non-small-cell lung cancer with weekly compared with 3-weekly docetaxel. *J Clin Oncol* **23**: 8389–8395
- Shepherd FA, Dancey J, Ramlau R, Mattson K, Gralla R, O'Rourke M, Levitan N, Gressot L, Vincent M, Burkes R, Coughlin S, Kim Y, Berille J (2000) Prospective randomized trial of docetaxel vs best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy. *J Clin Oncol* **18**: 2095–2103
- Therasse P, Arbuck SG, Eisenhauer EA, Wanders J, Kaplan RS, Rubinstein L, Verweij J, Van Glabbeke M, van Oosterom AT, Christian MC, Gwyther SG (2000) New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* **92**: 205–216
- Tibaldi C, Bernardini I, Chella A, Russo F, Vasile E, Malventi M, Falcone A (2006) Second-line chemotherapy with a modified schedule of docetaxel in elderly patients with advanced-stage non-small-cell lung cancer. *Clin Lung Cancer* **7**: 401–405
- Tibaldi C, Ricci S, Russo F, Bernardini I, Galli L, Chioni A, Orlandini C, Grosso AM, Pegna AL, Fabbri A, Innocenti F, Ferrari K, Tognarini L, Conte PF, Falcone A, on behalf of the POLTO group (2005) Increased dose-intensity of gemcitabine in advanced non small cell lung cancer (NSCLC): a multicenter phase II study in elderly patients from the 'polmone toscano group' (POLTO). *Lung Cancer* **48**: 121–127