



Published in final edited form as:

Eur J Cancer. 2008 January ; 44(1): 100–109.

Morbidity and mortality in relation to smoking among women and men of Chinese ethnicity: The Singapore Chinese Health Study

Anoop Shankar^{a,*}, Jian-Min Yuan^b, Woon-Puay Koh^a, Hin-Peng Lee^a, and Mimi C. Yu^b

^aDepartment of Community, Occupational, and Family Medicine, Yong Loo Lin School of Medicine, National University of Singapore, Block MD3, 16 Medical Drive, 117597 Singapore, Singapore

^bThe Cancer Center, University of Minnesota, Minneapolis, MN 55455, USA

Abstract

Objectives—We examined the association among cigarette smoking, smoking cessation and a broad range of cancer incidence and all cause and cause-specific mortality in a population-based cohort of adults of Chinese ethnicity in Singapore.

Methods—Subjects were 61,320 participants of the Singapore Chinese Health Study (44.5% men, aged 45–74 years, recruitment from 1993–1998) who were free of cancer at the baseline examination. Main outcomes-of-interest included cancer incidence, all cause and cause-specific mortality as of December 31, 2005.

Results—Cigarette smoking was positively associated with overall cancer incidence, including cancers at the following specific sites: head and neck region, upper gastrointestinal tract, hepatobiliary and pancreas cancer, lung, and bladder/renal pelvis cancer. Compared to never smokers, the relative risk (RR) (95% confidence interval [CI]) of cancer incidence (all cancer sites) among current smokers smoking >22 cigarettes/day was 1.9 (1.7–2.1), p -trend<0.0001. Similarly, cigarette smoking was associated with all cause and cause-specific mortality, including deaths due to cancer, ischemic heart disease, other heart diseases, and chronic obstructive pulmonary disease. Compared to never smokers, RR (95%CI) of all cause mortality among current smokers smoking >22 cigarettes/day was 1.8 (1.6–2.0), p -trend<0.0001. Also, relative to current smokers, ex-smokers experienced reduced cancer incidence and total mortality. The population attributable risk of smoking in men for cancer incidence as well as all-cause mortality was 23%, whereas in women it ranged from 4–5%.

Conclusions—Cigarette smoking is an important risk factor for cancer incidence and major causes of mortality in Chinese men and women of Singapore.

Keywords

Smoking; Smoking cessation; Cancer; Mortality; Population attributable risk; Chinese; Singapore

* Corresponding author: Tel.: +65 6516 4968; fax: +65 6779 1489. Email: ashankar@nus.edu.sg (A. Shankar).

Conflict of interest statement: None declared.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Cigarette smoking is an important public health problem and a major cause of morbidity and mortality(1). Numerous epidemiologic studies in Western populations have reported a positive association between cigarette smoking and cancer, cardiovascular disease, chronic obstructive pulmonary disease (COPD), total and cause-specific mortality(2-26).

In contrast, relatively few prospective studies from Asia, including studies from China(27, 28) and East Asian and other countries(29-37) have examined morbidity and mortality in relation to smoking. Most of these reports exclusively studied men(27,30-32,37); among the few studies that included women(28,29,33-36), only two included women of Chinese ethnicity (28,36). While previous Asian studies examined smoking in relation to coronary heart disease (CHD)(29-31), stroke incidence(29,31,33,35) and cause-specific mortality(27,28,30,32,36), only two prospective studies examined cancer incidence(27,34). Studies in Western populations have examined the effect of smoking cessation on morbidity and mortality (10-18,26); in contrast, only one prospective Asian study examined the relationship between smoking cessation and cardiovascular mortality(33), but not other mortality causes or cancer incidence. In this context, we examined the association among cigarette smoking, smoking cessation and a broad range of cancer incidence and all cause and cause-specific mortality in a population-based cohort of adult Singaporeans of Chinese ethnicity, the Singapore Chinese Health Study.

2. Patients and methods

2.1. Study population

The design of the Singapore Chinese Health Study has been described previously(38). Briefly, the cohort was recruited between 1993 and 1998, drawn from permanent residents or citizens of Singapore who lived in government-built housing (86% of the Singapore population resided in such facilities during enrollment period). Men and women of Chinese ethnicity (restricting to the two major dialect groups, the Cantonese and the Hokkiens), aged 45–74 years, were eligible. A total of 63,257 persons (~85% of eligible subjects) were enrolled. At recruitment, a face-to-face interview was conducted in the subject's home by a trained interviewer using a structured, scanner-readable questionnaire, which requested information on demographics, educational attainment, lifetime use of tobacco, current use of alcohol, current level of physical activity, medical history and family history of cancer. The questionnaire included a validated, semi-quantitative food frequency section listing 165 food items commonly consumed in the study population. The Singapore Food Composition Table(38), which we developed in conjunction with this cohort study, allowed for the estimation of intake levels of approximately 100 nutritive/non-nutritive food components per study subject.

2.2. Exposure assessment

The study population was divided into never, former, and current smokers based on their choice of three possible responses to the following question, 'Have you ever smoked at least 1 cigarette a day for 1 y or longer?' Subjects who answered 'no' were classified as 'non smokers,' those who answered 'yes, but I quit smoking' were classified as 'former smokers,' and those who answered 'yes, and I currently smoke' were classified as 'current smokers.' There were six predefined categories of average number of cigarettes smoked/ day: ≤ 6 cigarettes, 7–12 cigarettes, 13–22 cigarettes, 23–32 cigarettes, 33–42 cigarettes, and ≥ 43 cigarettes. The number of years since quitting smoking for former smokers was categorised as: <1 year, 1–2 years, 3–4 years, 5–9 years, 10–14 years, 15–19 years, and ≥ 20 years. For the current analysis, based on sample size considerations, we categorised cigarette smoking status into four categories: non smoker, former smoker, current smoker with ≤ 22 cigarettes/ day, current smoker with > 22

cigarettes/ day. Also, for former smokers, the number of years since quitting smoking was categorised as follows: <10 years, and 10 or more years.

Age was defined as the age in years at the time of baseline examination; education was categorised into no formal education, primary school, secondary school or above; alcohol intake was assessed as grams of daily ethanol consumption; body mass index (BMI) was calculated as weight (kilograms) divided by the square of height (metres); moderate physical activities included brisk walking, bowling, cycling on level ground, tai chi, or chi kung, and hours/week spent on such activities were categorised into none, ½ to 3 h/week, ≥4 h/week.

2.3. Cancer incidence and cause-specific mortality

We identified cancer cases through the population-based Singapore Cancer Registry which has been continuously included in the International Agency for Research on Cancer (IARC) serial publications on world-wide cancer incidence since 1968(39). Deaths were identified through record linkage with the Singapore Registry of Births and Deaths. In our recent follow-up telephone/in-person interview conducted between 1999 and 2004, among the 61,685 subjects (97.5%) that we had contact or follow-up information, either from themselves, their next-of-kin or death records, only 17 subjects (0.03%) have migrated out of Singapore. This suggests that emigration is negligible among the subjects in the cohort.

Of the 63,257 cohort participants, 1937 had a history of invasive cancer (except for nonmelanoma skin cancer) or *in situ* bladder cancer at recruitment, thus were excluded from the current analysis. As of December 31, 2005, among the 61,320 participants who were free of cancer at baseline, a total of 7982 had died and 5253 had developed either incident invasive cancer of any site or *in-situ* cancer of the bladder and cervix (for females only). Among the cancer cases, 4750 cases (90.4%) had histopathological diagnoses while the remaining 503 cases (9.6%) were diagnosed using clinical or radiological evidence. In addition, histological information on 4407 cancer diagnoses (83.9%) were verified via manual review of all pathology reports and 218 cases (4.2%) were verified via review of medical records by medically trained research staff.

2.4. Statistical analysis

Person-years for each cohort participant was counted from the date of interview to date of cancer diagnosis (for cancer morbidity analysis), or date of death, or December 31 2005, whichever occurred first. We examined two smoking-related exposures, assessed at the baseline examination: (1) categories of smoking status (non smoker [referent category], former smoker, current smoker with ≤22 cigarettes/ day, current smoker with >22 cigarettes/ day), and (2) categories related to smoking cessation (current smoker [referent category], <10 years since quitting, ≥10 years since quitting, never smoker). We were interested in two groups of outcomes: (1) cancer at all sites and site-specific cancer incidence (based on first diagnosed primary site of cancer categorised according to International Classification of Diseases, Tenth Revision [ICD-10] code), and (2) all cause and cause-specific mortality (based on underlying cause of death classification according to International Classification of Diseases, Ninth Revision [ICD-9] code from death certificates). The following cancer sites (ICD-10 code) were studied: all cancer sites (C00-C97), head and neck (C00.0-C14.8, C32.0-C32.9), upper gastrointestinal tract (oesophagus [C15.2-C15.9] and stomach [C16.0-C16.9]), colorectal cancer (colon [C18.0-C18.9] and rectum [C19]), hepatobiliary and pancreas cancer (liver and intrahepatic bile ducts[C22], gallbladder and other biliary tract [C23, 24] and pancreas [C25.0-C25.9]), lung (C34.0-C34.9), bladder and renal pelvis cancer (C65.0-C68.9), and breast cancer (women only) (C50). We counted only invasive cancers in defining incident cancer, except for bladder cancer (C67) and cervical cancer (C53), where *in situ* cancers were also included in the analysis. In the current analysis, we were interested in the global association between

smoking and a broadly defined range of site-specific incident cancers. We initially performed separate analyses for liver (C22), pancreas (C25) and gallbladder/ other biliary tract (C23, 24) cancers. Since these sites showed similar positive association with smoking, we combined these categories into a single group, 'hepatobiliary and pancreas cancer', to obtain adequate sample size. Similarly, after separate analyses that showed similar associations, we chose to combine bladder (C67), and renal pelvis and other urinary organs (C65, C66, C68) into a single group, 'bladder/renal pelvis cancer'. Breast cancer analyses are confined to women. There were very few bladder/renal pelvis cancer among women smokers (35 cases). Therefore, we presented results of both men and women combined. Mortality due to the following specific causes (ICD-9 code) were studied: cancer (140–208), ischemic heart disease (410–414), stroke (430–438), other heart diseases (391–398, 401, 402, 404, 415–417, 420–429), chronic obstructive pulmonary disease (COPD) (490–496), and liver cirrhosis (571). We used multivariable-adjusted proportional hazards models to estimate the relative risk (RR) and 95% confidence intervals (CI), controlling simultaneously for potential confounders, including age (years), gender (males, females), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (grams), and h/week of moderate physical activity (none, ½ to 3 h/week, 4 h/week or more). Additional variables considered as potential confounders but not included in the final multivariable model were as follows: total calorie intake (kcal/day), total fruits (g/day), total vegetables (g/day), total red meats (g/day), dietary fibre (g/day), dietary cholesterol (mg/day), vitamin E (mg/day), weekly vitamin supplement use (yes, no), and among women, menopausal status (premenopausal, postmenopausal), and post-menopausal oestrogen use (current, former, never). We analysed cigarette smoking categories as an ordinal variable in multivariable models to examine trend in the association. We used cross-product interaction terms in regression models to test multiplicative interaction. We calculated the population attributable risk in relation to ever smoking using Levin's formula(40). All analyses were conducted using SAS version 9.2 (SAS Inc; Cary, NC). Statistical significance was defined as 2-sided p -value<0.05.

3. Results

Table 1 presents the baseline characteristics of the cohort by smoking status. Approximately 20% of all cohort participants (36% of men and 6% of women) reported smoking cigarettes at recruitment. Current smokers had lower levels of education and BMI, higher mean intake of alcohol, and were physically less active.

Table 2 presents the relation between categories of smoking status at baseline and cancer incidence. We observed a positive association between increasing smoking levels and risks of total cancer as well as cancer of the head and neck, upper gastrointestinal tract, hepatobiliary and pancreas, lung, and bladder/renal pelvis. Smoking was unrelated to the overall risk of colorectal cancer. A null association between smoking and breast cancer risk was noted in this study population. In a subsidiary analysis, women who began to smoke before 15 years of age had a relative risk of 1.7 (95% CI = 0.97–3.1) compared with never smokers.

In Table 3, we observed a positive association between increasing smoking categories and all cause mortality and the following cause-specific deaths: cancer, ischemic heart disease, and COPD. We also observed a positive association between smoking and other heart disease mortality, the RR estimates were similar to the results for ischemic heart disease (data not shown). A positive association between smoking and mortality due to liver cirrhosis was of borderline statistical significance (data not shown). Compared to never smokers (referent group), the RR (95% CI) of liver cirrhosis mortality ($n=78$) was 1.2 (0.60–2.4) among former smokers, 1.3 (0.70–2.5) among current smokers smoking ≤ 22 cigarettes/day, and 2.0 (0.85–

4.7) among current smokers smoking >22 cigarettes/day, p -trend=0.06. There was no overall association between smoking and stroke mortality.

Table 4 presents the effect of smoking cessation on cancer incidence. For all cancers, we observed a statistically significant risk reduction for former smokers, and the risk reduction was dependent on the number of years of smoking cessation. Compared to current smokers, the RR (95% CI) for all cancers was 0.84 (0.75–0.94) for former smokers with <10 years and 0.67 (0.59–0.75) for former smokers with at least 10 years since smoking cessation. Similarly, a statistically significant, inverse associations was observed between the number of years since smoking cessation and the risk of developing cancer of the head and neck, upper gastrointestinal tract, hepatobiliary and pancreas, lung, and bladder/renal pelvis (Table 4).

Table 5 presents the effect of smoking cessation on all cause and selected cause-specific mortality. A statistically significant reduction in risk of all-cause mortality and cause-specific mortality was noted among former smokers. Compared with current smokers, the RR (95% CI) for all-cause mortality was 0.91 (0.84–0.99) for former smokers with <10 years since smoking cessation and 0.65 (0.60–0.71) for those who quit smoking at least 10 years.

Table 6 presents the population attributable risk of cancer incidence and cause-specific mortality in relation to ever smoking. Smoking accounted for more than one in five cancer cases as well as all-cause deaths in men whereas about 4–5% of cancer and death burdens were attributable to smoking in women.

4. Discussion

In a large population-based cohort of adult Chinese Singaporeans, cigarette smoking was positively associated with risk of developing any cancer as well as certain specific cancer sites including head and neck, upper gastrointestinal tract, hepatobiliary and pancreas, lung, and bladder/renal pelvis. Similarly, cigarette smoking also was associated with all-cause and cause-specific mortality including deaths due to cancer, ischemic heart disease, other heart diseases, and COPD. Smoking cessation was associated with statistically significant reduction in risk of smoking-related cancers and smoking related causes of death. Overall, smoking accounted for one in four cancer cases and deaths in men and <5% of cancer and death burdens in women of the study population.

Prevalence of smoking was historically extremely low among Asian women(41-43), and most previous Asian studies examining smoking included only men(27,30-32,37). The well documented increase in prevalence of cigarette smoking in women in the newly industrialised Asian countries(44), including Singapore, is postulated to be related, at least in part, to the westernisation(45,46) in lifestyles of those Asian populations. While smoking prevalence has declined in many western developed countries, it remains high among men in newly industrialised Asian countries and its prevalence is increasing among women (44). Smoking related lung cancer is the third most frequent cancer among women in Singapore(47). While the age-standardised incidence rate of lung cancer in men has fallen from 60 per 100,000 person-years in 1980 to 45 per 100,000 person-years in 2002, it has remained stable in women at approximately 16 per 100,000 person-years over the last two decades(47). Our data highlights the potential enormous impact of smoking on population health of this continent.

Our results of an overall positive association between cancer incidence and smoking is consistent with previous prospective studies from Asia(27,34) and agrees with recent reports from the US Surgeon General(1,48) and the International Agency for Research on Cancer (44) on the health impact of smoking.

In the current study, cigarette smoking was unrelated to colorectal cancer overall. In a concurrent, separate report, we described in further details the relationship between cigarette smoking and colorectal cancer by smoking intensity and duration, stratified by sub-site and stage of the disease. We noted that smoking was associated with a statistically significant, 50% increase in risk of rectal cancer, but was unrelated to risk of colon cancer (Tsong et al., submitted). Some(34,49-51) but not all(52-54) prior studies have reported a lack of association between smoking and colorectal cancer. Other studies have examined cofactors such as subject's metabolic genotype in tobacco carcinogen metabolism(55,56) and tumour characteristics(55,57) and found smoking to be associated with subgroups defined by these cofactors.

We noted no overall association between cigarette smoking and breast cancer in Singapore Chinese women. However, there was a hint that women who began smoking at a young age (≤ 14 years) might be at risk of breast cancer. This observation is consistent with a recent report that smoking initiation at younger age (adolescence and early adulthood) is associated with increased risk of breast cancer, but that smoking was not associated with breast cancer after first child birth(58). Most studies(59-68), including a pooled analysis of 53 epidemiologic studies(68), reported no association between smoking and female breast cancer although some prior studies have noted a positive association(69-73). Recent reports from the US Surgeon General(1,48) and the International Agency for Research on Cancer(44) concluded that there is no evidence of an association between smoking and breast cancer among women(1,44,48).

Our findings of an overall positive association between smoking and all cause and cause-specific mortality other than stroke are consistent with previous studies from Asia(27,74) and the US Surgeon General's report(1). In the present study, current smoking was not associated with stroke mortality, a finding consistent with many(29,32,75,76), but not all(27,31,33,37,77,78), previous studies from Asia. Our findings on stroke mortality are contrary to the US Surgeon General's report(1) which concluded that smoking is positively related to stroke risk, particularly ischemic stroke and subarachnoid haemorrhage. The US report was predominantly based on studies from Western populations(2,18-26); only two studies from Asian populations were included(31,78). One explanation for the observed lack of association may be the reported heterogeneous nature of the association between smoking and stroke subtypes, including a potential lack of association with intracerebral haemorrhage(79) and embolic brain infarction (80).

The main strengths of the current study include its large sample size, good representation of women, its population-based and prospective follow-up nature, and the high percentage of cancer cases that were diagnosed histopathologically. The nationwide cancer registry has been in place since 1968 and has been shown to be comprehensive in its recording of cancer cases (81). Thus, cancer case ascertainment can be assumed to be complete. However, although women represent 55% of this cohort, only 8.7% ($n=2965$) were ever smokers and most women were light smokers (i.e. under one pack/day). The relatively small number of cancer death cases in women smokers prevented an in-depth analysis of smoking in this gender subgroup. Despite these limitations, our data suggest that female smokers were equally at risk for cancer and other major causes of death from chronic diseases as men.

In conclusion, in a large population-based cohort of Chinese men and women in Singapore, cigarette smoking was positively associated with cancer incidence in multiple sites, and with all cause and cause-specific mortality. Smoking cessation, as opposed to continuation of smoking, was associated with reduction in cancer incidence and all cause and cause-specific mortality.

Acknowledgements

Guarantor statement: 'The guarantor, AS, accepts full responsibility for the work and/or the conduct of the study, had access to the data, and controlled the decision to publish'.

Contributors: All authors contributed to the intellectual development of this paper. AS and MCY had the original idea for the study. AS analysed the data, wrote the first draft paper, and is the guarantor. MCY, JMY provided statistical expertise and critical corrections to the manuscript. MCY, JMY, WPK, HPL provided critical corrections to the manuscript. MCY is the Principal Investigator of the Singapore Chinese Health Study.

Ethical Approval: This study followed the recommendations of the Declaration of Helsinki and was approved by the institutional review boards at the University of Minnesota, Minneapolis and the National University of Singapore, Singapore. Written, informed consent was obtained from all participants.

Details of funding: Supported by National Institutes of Health grants RO1 CA55069, R35 CA53890, and R01 CA80205, from the National Cancer Institute, Bethesda, MD.

Independence of researchers from funding source: The funding agencies had no role in the research presented in the paper and the researchers were fully independent in pursuing this research.

We thank Ms. Siew-Hong Low of the National University of Singapore for supervising the field work of the Singapore Chinese Health Study, and Ms. Kazuko Arakawa of the University of Southern California for the development and management of the cohort study database. We also thank the Singapore Cancer Registry for assistance with the identification of cancer outcomes.

References

1. United States, Public Health Service, Office of the Surgeon General, National Center for Chronic Disease Prevention and Health Promotion. The health consequences of smoking - a report of the Surgeon General. Atlanta, Ga.: U.S. Public Health Service, National Center for Chronic Disease Prevention and Health Promotion; 2004.
2. Doll R, Peto R. Mortality in relation to smoking: 20 years' observations on male British doctors. *Br Med J* 1976;2:1525–1536. [PubMed: 1009386]
3. Rogot E, Murray JL. Smoking and causes of death among U.S. veterans: 16 years of observation. *Public Health Rep* 1980;95:213–222. [PubMed: 7384406]
4. Freund KM, Belanger AJ, D'Agostino RB, Kannel WB. The health risks of smoking. The Framingham Study: 34 years of follow-up. *Ann Epidemiol* 1993;3:417–424. [PubMed: 8275219]
5. Willett WC, Green A, Stampfer MJ, et al. Relative and absolute excess risks of coronary heart disease among women who smoke cigarettes. *N Engl J Med* 1987;317:1303–1309. [PubMed: 3683458]
6. Neaton JD, Wentworth D. Serum cholesterol, blood pressure, cigarette smoking, and death from coronary heart disease. Overall findings and differences by age for 316,099 white men. Multiple Risk Factor Intervention Trial Research Group. *Arch Intern Med* 1992;152:56–64. [PubMed: 1728930]
7. Centers for Disease Control and Prevention. Cigarette smoking-attributable mortality and years of potential life lost—United States, 1990. *MMWR Morb Mortal Wkly Rep* 1993;42:645–649. [PubMed: 8350857]
8. Hammond EC, Seidman H. Smoking and cancer in the United States. *Prev Med* 1980;9:169–174. [PubMed: 7383981]
9. Baron AE, Franceschi S, Barra S, Talamini R, La VC. A comparison of the joint effects of alcohol and smoking on the risk of cancer across sites in the upper aerodigestive tract. *Cancer Epidemiol Biomarkers Prev* 1993;2:519–523. [PubMed: 8268767]
10. Ebbert JO, Yang P, Vachon CM, et al. Lung cancer risk reduction after smoking cessation: observations from a prospective cohort of women. *J Clin Oncol* 2003;21:921–926. [PubMed: 12610194]
11. Enstrom JE. Trends in mortality among California physicians after giving up smoking: 1950-79. *Br Med J (Clin Res Ed)* 1983;286:1101–1105.
12. Enstrom JE. Smoking cessation and mortality trends among two United States populations. *J Clin Epidemiol* 1999;52:813–825. [PubMed: 10529023]

13. Speizer FE, Colditz GA, Hunter DJ, Rosner B, Hennekens C. Prospective study of smoking, antioxidant intake, and lung cancer in middle-aged women (USA). *Cancer Causes Control* 1999;10:475–482. [PubMed: 10530619]
14. Halpern MT, Gillespie BW, Warner KE. Patterns of absolute risk of lung cancer mortality in former smokers. *J Natl Cancer Inst* 1993;85:457–464. [PubMed: 8445673]
15. Anthonisen NR, Connett JE, Kiley JP, et al. Effects of smoking intervention and the use of an inhaled anticholinergic bronchodilator on the rate of decline of FEV1. The Lung Health Study. *JAMA* 1994;272:1497–1505. [PubMed: 7966841]
16. Kawachi I, Colditz GA, Stampfer MJ, et al. Smoking cessation and time course of decreased risks of coronary heart disease in middle-aged women. *Arch Intern Med* 1994;154:169–175. [PubMed: 8285812]
17. Kawachi I, Colditz GA, Stampfer MJ, et al. Smoking cessation and decreased risk of stroke in women. *JAMA* 1993;269:232–236. [PubMed: 8417241]
18. Wannamethee SG, Shaper AG, Whincup PH, Walker M. Smoking cessation and the risk of stroke in middle-aged men. *JAMA* 1995;274:155–160. [PubMed: 7596004]
19. Doll R, Peto R, Wheatley K, Gray R, Sutherland I. Mortality in relation to smoking: 40 years' observations on male British doctors. *BMJ* 1994;309:901–911. [PubMed: 7755693]
20. Kuller LH, Ockene JK, Meilahn E, Wentworth DN, Svendsen KH, Neaton JD. Cigarette smoking and mortality. MRFIT Research Group. *Prev Med* 1991;20:638–654. [PubMed: 1758843]
21. Robbins AS, Manson JE, Lee IM, Satterfield S, Hennekens CH. Cigarette smoking and stroke in a cohort of U.S. male physicians. *Ann Intern Med* 1994;120:458–462. [PubMed: 8311368]
22. Shaper AG, Phillips AN, Pocock SJ, Walker M, Macfarlane PW. Risk factors for stroke in middle aged British men. *BMJ* 1991;302:1111–1115. [PubMed: 1828378]
23. Knekt P, Reunanen A, Aho K, et al. Risk factors for subarachnoid hemorrhage in a longitudinal population study. *J Clin Epidemiol* 1991;44:933–939. [PubMed: 1890435]
24. Petitti DB, Friedman GD. Cardiovascular and other diseases in smokers of low yield cigarettes. *J Chronic Dis* 1985;38:581–588. [PubMed: 4008601]
25. Gillum RF. Risk factors for stroke in blacks: a critical review. *Am J Epidemiol* 1999;150:1266–1274. [PubMed: 10604768]
26. Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ* 2004;328:1519. [PubMed: 15213107]
27. Yuan JM, Ross RK, Wang XL, Gao YT, Henderson BE, Yu MC. Morbidity and mortality in relation to cigarette smoking in Shanghai, China. A prospective male cohort study. *JAMA* 1996;275:1646–1650. [PubMed: 8637137]
28. Lam TH, He Y, Li LS, Li LS, He SF, Liang BQ. Mortality attributable to cigarette smoking in China. *JAMA* 1997;278:1505–1508. [PubMed: 9363970]
29. Kiyohara Y, Ueda K, Fujishima M. Smoking and cardiovascular disease in the general population in Japan. *J Hypertens Suppl* 1990;8:S9–15. [PubMed: 2286856]
30. Yano K, MacLean CJ, Reed DM, et al. A comparison of the 12-year mortality and predictive factors of coronary heart disease among Japanese men in Japan and Hawaii. *Am J Epidemiol* 1988;127:476–487. [PubMed: 3341354]
31. Jee SH, Suh I, Kim IS, Appel LJ. Smoking and atherosclerotic cardiovascular disease in men with low levels of serum cholesterol: the Korea Medical Insurance Corporation Study. *JAMA* 1999;282:2149–2155. [PubMed: 10591337]
32. Kono S, Ikeda M, Tokudome S, Nishizumi M, Kuratsune M. Smoking and mortalities from cancer, coronary heart disease and stroke in male Japanese physicians. *J Cancer Res Clin Oncol* 1985;110:161–164. [PubMed: 4044630]
33. Iso H, Date C, Yamamoto A, et al. Smoking cessation and mortality from cardiovascular disease among Japanese men and women: the JACC Study. *Am J Epidemiol* 2005;161:170–179. [PubMed: 15632267]
34. Wakai K, Hayakawa N, Kojima M, et al. Smoking and colorectal cancer in a non-Western population: a prospective cohort study in Japan. *J Epidemiol* 2003;13:323–332. [PubMed: 14674660]

35. Mannami T, Iso H, Baba S, et al. Cigarette smoking and risk of stroke and its subtypes among middle-aged Japanese men and women: the JPHC Study Cohort I. *Stroke* 2004;35:1248–1253. [PubMed: 15118170]
36. Liaw KM, Chen CJ. Mortality attributable to cigarette smoking in Taiwan: a 12-year follow-up study. *Tob Control* 1998;7:141–148. [PubMed: 9789932]
37. Yamagishi K, Iso H, Kitamura A, et al. Smoking raises the risk of total and ischemic strokes in hypertensive men. *Hypertens Res* 2003;26:209–217. [PubMed: 12675276]
38. Hankin JH, Stram DO, Arakawa K, et al. Singapore Chinese Health Study: development, validation, and calibration of the quantitative food frequency questionnaire. *Nutr Cancer* 2001;39:187–195. [PubMed: 11759279]
39. Parkin, DM.; Whelan, SL.; Ferlay, J.; Teppo, L.; Thomas, DB. *Cancer incidence in five continents*. Lyon: IARC; 2002.
40. Szklo, M.; Nieto, FJ. *Epidemiology beyond the basics*. Gaithersburg, Md: Aspen; 2000.
41. Gu D, Wu X, Reynolds K, et al. Cigarette smoking and exposure to environmental tobacco smoke in China: the international collaborative study of cardiovascular disease in Asia. *Am J Public Health* 2004;94:1972–1976. [PubMed: 15514239]
42. Smith DR, Wei N, Zhang YJ, Wang RS. Tobacco smoking habits among a cross-section of rural physicians in China. *Aust J Rural Health* 2006;14:66–71. [PubMed: 16512792]
43. Shakib S, Zheng H, Johnson CA, et al. Family characteristics and smoking among urban and rural adolescents living in China. *Prev Med* 2005;40:83–91. [PubMed: 15530584]
44. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans, World Health Organization, International Agency for Research on Cancer. *Tobacco smoke and involuntary smoking*. Lyon, France: IARC Press; 2004.
45. Woo KS, Chook P, Raitakari OT, McQuillan B, Feng JZ, Celermajer DS. Westernization of Chinese adults and increased subclinical atherosclerosis. *Arterioscler Thromb Vasc Biol* 1999;19:2487–2493. [PubMed: 10521379]
46. Chen MS Jr. Cancer health disparities among Asian Americans: what we do and what we need to do. *Cancer* 2005;104:2895–2902. [PubMed: 16270313]
47. Seow A, Koh WP, Chia KS, Shi LM, Lee HP, Shanmugaratnam K. Trends in cancer incidence in Singapore, 1968-2002. *The Singapore Cancer Registry Report No 6 Singapore*. 2004
48. United States, Dept. of Health and Human Services, Office on Women's Health. *Surgeon General's report on women and smoking*. Washington, DC: Office on Women's Health; 2001.
49. Sandler RS, Sandler DP, Comstock GW, Helsing KJ, Shore DL. Cigarette smoking and the risk of colorectal cancer in women. *J Natl Cancer Inst* 1988;80:1329–1333. [PubMed: 3172257]
50. Nyren O, Bergstrom R, Nystrom L, et al. Smoking and colorectal cancer: a 20-year follow-up study of Swedish construction workers. *J Natl Cancer Inst* 1996;88:1302–1307. [PubMed: 8797770]
51. Giovannucci E, Colditz GA, Stampfer MJ, et al. A prospective study of cigarette smoking and risk of colorectal adenoma and colorectal cancer in U.S. women. *J Natl Cancer Inst* 1994;86:192–199. [PubMed: 8283491]
52. Mizoue T, Inoue M, Tanaka K, et al. Tobacco smoking and colorectal cancer risk: an evaluation based on a systematic review of epidemiologic evidence among the Japanese population. *Jpn J Clin Oncol* 2006;36:25–39. [PubMed: 16423841]
53. Chyou PH, Nomura AM, Stemmermann GN. A prospective study of colon and rectal cancer among Hawaii Japanese men. *Ann Epidemiol* 1996;6:276–282. [PubMed: 8876837]
54. Giovannucci E, Rimm EB, Stampfer MJ, et al. A prospective study of cigarette smoking and risk of colorectal adenoma and colorectal cancer in U.S. men. *J Natl Cancer Inst* 1994;86:183–191. [PubMed: 8283490]
55. Luchtenborg M, Weijenberg MP, Kampman E, et al. Cigarette smoking and colorectal cancer: APC mutations, hMLH1 expression, and GSTM1 and GSTT1 polymorphisms. *Am J Epidemiol* 2005;161:806–815. [PubMed: 15840612]
56. Welfare MR, Cooper J, Bassendine MF, Daly AK. Relationship between acetylator status, smoking, and diet and colorectal cancer risk in the north-east of England. *Carcinogenesis* 1997;18:1351–1354. [PubMed: 9230278]

57. Miyaki M, Iijima T, Ishii R, et al. Increased frequency of p53 mutation in sporadic colorectal cancer from cigarette smokers. *Jpn J Clin Oncol* 2002;32:196–201. [PubMed: 12110635]
58. Ha M, Mabuchi K, Sigurdson AJ, et al. Smoking cigarettes before first childbirth and risk of breast cancer. *Am J Epidemiol* 2007;166:55–61. [PubMed: 17426039]
59. Wartenberg D, Calle EE, Thun MJ, Heath CW Jr, Lally C, Woodruff T. Passive smoking exposure and female breast cancer mortality. *J Natl Cancer Inst* 2000;92:1666–1673. [PubMed: 11036112]
60. Brunet JS, Ghadirian P, Rebbeck TR, et al. Effect of smoking on breast cancer in carriers of mutant BRCA1 or BRCA2 genes. *J Natl Cancer Inst* 1998;90:761–766. [PubMed: 9605646]
61. Hunter DJ, Hankinson SE, Hough H, et al. A prospective study of NAT2 acetylation genotype, cigarette smoking, and risk of breast cancer. *Carcinogenesis* 1997;18:2127–2132. [PubMed: 9395212]
62. London SJ, Colditz GA, Stampfer MJ, Willett WC, Rosner BA, Speizer FE. Prospective study of smoking and the risk of breast cancer. *J Natl Cancer Inst* 1989;81:1625–1631. [PubMed: 2795691]
63. Schechter MT, Miller AB, Howe GR, Baines CJ, Craib KJ, Wall C. Cigarette smoking and breast cancer: case-control studies of prevalent and incident cancer in the Canadian National Breast Screening Study. *Am J Epidemiol* 1989;130:213–220. [PubMed: 2787599]
64. Rohan TE, Baron JA. Cigarette smoking and breast cancer. *Am J Epidemiol* 1989;129:36–42. [PubMed: 2910070]
65. Chu SY, Stroup NE, Wingo PA, Lee NC, Peterson HB, Gwinn ML. Cigarette smoking and the risk of breast cancer. *Am J Epidemiol* 1990;131:244–253. [PubMed: 2404408]
66. Field NA, Baptiste MS, Nasca PC, Metzger BB. Cigarette smoking and breast cancer. *Int J Epidemiol* 1992;21:842–848. [PubMed: 1468843]
67. Brinton LA, Schairer C, Stanford JL, Hoover RN. Cigarette smoking and breast cancer. *Am J Epidemiol* 1986;123:614–622. [PubMed: 3953540]
68. Hamajima N, Hirose K, Tajima K, et al. Alcohol, tobacco and breast cancer--collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *Br J Cancer* 2002;87:1234–1245. [PubMed: 12439712]
69. Palmer JR, Rosenberg L, Clarke EA, et al. Breast cancer and cigarette smoking: a hypothesis. *Am J Epidemiol* 1991;134:1–13. [PubMed: 1853854]
70. Reynolds P, Hurley S, Goldberg DE, et al. Active smoking, household passive smoking, and breast cancer: evidence from the California Teachers Study. *J Natl Cancer Inst* 2004;96:29–37. [PubMed: 14709736]
71. Kropp S, Chang-Claude J. Active and passive smoking and risk of breast cancer by age 50 years among German women. *Am J Epidemiol* 2002;156:616–626. [PubMed: 12244030]
72. Calle EE, Miracle-McMahill HL, Thun MJ, Heath CW Jr. Cigarette smoking and risk of fatal breast cancer. *Am J Epidemiol* 1994;139:1001–1007. [PubMed: 8178779]
73. Lash TL, Aschengrau A. Active and passive cigarette smoking and the occurrence of breast cancer. *Am J Epidemiol* 1999;149:5–12. [PubMed: 9883788]
74. Woodward M, Lam TH, Barzi F, et al. Smoking, quitting, and the risk of cardiovascular disease among women and men in the Asia-Pacific region. *Int J Epidemiol* 2005;34:1036–1045. [PubMed: 15914503]
75. Okada H, Horibe H, Yoshiyuki O, Hayakawa N, Aoki N. A prospective study of cerebrovascular disease in Japanese rural communities, Akabane and Asahi. Part 1: evaluation of risk factors in the occurrence of cerebral hemorrhage and thrombosis. *Stroke* 1976;7:599–607. [PubMed: 1006736]
76. Tanaka H, Ueda Y, Hayashi M, et al. Risk factors for cerebral hemorrhage and cerebral infarction in a Japanese rural community. *Stroke* 1982;13:62–73. [PubMed: 7064181]
77. Feigin V, Parag V, Lawes CM, et al. Smoking and elevated blood pressure are the most important risk factors for subarachnoid hemorrhage in the Asia-Pacific region: an overview of 26 cohorts involving 306,620 participants. *Stroke* 2005;36:1360–1365. [PubMed: 15933249]
78. Hirayama, T. *Life-style and mortality a large-scale census-based cohort study in Japan*. Basel: Karger; 1990.
79. Shinton R, Beevers G. Meta-analysis of relation between cigarette smoking and stroke. *BMJ* 1989;298:789–794. [PubMed: 2496858]

80. Yip PK, Jeng JS, Lee TK, et al. Subtypes of ischemic stroke. A hospital-based stroke registry in Taiwan (SCAN-IV). *Stroke* 1997;28:2507–2512. [PubMed: 9412641]
81. Seow A, Koh WP, Chia KS, Shi LM, Lee HP, Shanmugaratnam K. Trends in cancer incidence in Singapore, 1968-2002. *The Singapore Cancer Registry Report No 6 Singapore*. 2006

Table 1

Selected baseline characteristics by smoking status^a

	Men (n=27,292)			Women (n=34,028)		
	Never smoker	Former smoker	Current smoker	Never smoker	Former smoker	Current smoker
No. of subjects	11,521 (42.2%)	5838 (21.4%)	9933 (36.4%)	31,062 (91.3%)	842 (2.5%)	2124 (6.2%)
Mean age, years	55.3	59.0	56.7	55.7	63.1	60.8
Highest education						
No formal education	859 (29.0%)	602 (20.3%)	1500 (50.7%)	11881 (86.7%)	471 (3.4%)	1348 (9.8%)
Primary school	4964 (35.6%)	3294 (23.6%)	5682 (40.8%)	12324 (92.8%)	301 (2.3%)	658 (5.0%)
Secondary school or above	5698 (54.8%)	1942 (18.7%)	2751 (26.5%)	6857 (97.3%)	70 (1.0%)	118 (1.7%)
Alcohol intake frequency						
Never	8385 (44.8%)	4151 (22.2%)	6175 (33.0%)	28362 (91.7%)	741 (2.4%)	1835 (5.9%)
Monthly	1293 (44.3%)	584 (20.0%)	1040 (35.7%)	1409 (89.4%)	50 (3.2%)	118 (7.5%)
Weekly	1472 (37.8%)	782 (20.1%)	1643 (42.2%)	991 (88.4%)	34 (3.0%)	96 (8.6%)
Daily	371 (21.0%)	321 (18.2%)	1075 (60.8%)	300 (76.5%)	17 (4.3%)	75 (19.1%)
Mean daily ethanol intake, g	2.0	3.1	5.8	0.3	0.6	0.9
Mean body mass index, kg/m ²	23.3	23.3	22.5	23.3	23.4	22.7
Hours/week of moderate physical activity						
No	8282 (40.2%)	4201 (20.4%)	8098 (39.4%)	24694 (90.8%)	660 (2.4%)	1858 (6.8%)
½ to 3 h/week	2080 (49.6%)	981 (23.4%)	1133 (27.0%)	4029 (93.3%)	121 (2.8%)	168 (3.9%)
4 h/week or more	1159 (46.1%)	656 (26.1%)	702 (27.9%)	2339 (93.6%)	61 (2.4%)	98 (3.9%)

^a Row percentages in parentheses are presented together with number of subjects in each cell, unless otherwise stated.

Table 2

Relation between smoking categories and cancer incidence

Cancer sites	Never smoker	Former smoker	Current smoker		p-trend
			≤22 cigarettes/day	>22 cigarettes/day	
All cancers					
No. of cases, whole cohort (n=5253)	2965	711	1272	305	
No. of cases, men (n=2741)	801	621	1034	285	
No. of cases, women (n=2512)	2164	90	238	20	
Multivariable RR ^a , whole cohort	1 (Referent)	1.2 (1.1-1.3)	1.60 (1.5-1.7)	1.9 (1.7-2.1)	<0.0001
Multivariable RR ^a , men	1 (Referent)	1.2 (1.1-1.4)	1.70 (1.6-1.9)	1.9 (1.7-2.2)	<0.0001
Multivariable RR ^a , women	1 (Referent)	1.3 (1.0-1.6)	1.45 (1.3-1.7)	2.1 (1.4-3.3)	<0.0001
Head and neck cancer					
No. of cases, whole cohort (n=345)	149	48	114	34	
No. of cases, men (n=252)	72	45	104	31	
No. of cases, women (n=93)	77	3	10	3	
Multivariable RR ^a , whole cohort	1 (Referent)	1.2 (0.85-1.7)	2.1 (1.6-2.71)	2.5 (1.6-3.7)	<0.0001
Multivariable RR ^a , men	1 (Referent)	1.1 (0.78-1.7)	2.0 (1.5-2.76)	2.3 (1.5-3.5)	<0.0001
Multivariable RR ^a , women	1 (Referent)	1.7 (0.54-5.6)	2.1 (1.1-4.12)	9.9 (3.1-32.0)	0.0005
Upper gastrointestinal tract cancer					
No. of cases, whole cohort (n=458)	222	75	127	34	
No. of cases, men (n=294)	83	67	112	32	
No. of cases, women (n=164)	139	8	15	2	
Multivariable RR ^a , whole cohort	1 (Referent)	1.2 (0.88-1.6)	1.5 (1.2-1.9)	1.7 (1.2-2.5)	0.0004
Multivariable RR ^a , men	1 (Referent)	1.1 (0.82-1.6)	1.6 (1.2-2.1)	1.8 (1.2-2.7)	0.0003
Multivariable RR ^a , women	1 (Referent)	1.6 (0.78-3.3)	1.1 (0.66-1.9)	2.7 (0.67-11.01)	0.26
Colorectal cancer					
No. of cases, whole cohort (n=931)	550	150	201	30	
No. of cases, men (n=514)	185	138	162	29	
No. of cases, women (n=417)	365	12	39	1	
Multivariable RR ^a , whole cohort	1 (Referent)	1.1 (0.92-1.4)	1.2 (0.97-1.4)	0.83 (0.56-1.2)	0.46
Multivariable RR ^a , men	1 (Referent)	1.2 (0.95-1.5)	1.2 (0.96-1.5)	0.87 (0.58-1.3)	0.47
Multivariable RR ^a , women	1 (Referent)	0.82 (0.46-1.5)	1.2 (0.84-1.7)	0.54 (0.08-3.9)	0.61
Hepatobiliary and pancreas cancer					
No. of cases, whole cohort (n=480)	235	80	136	29	
No. of cases, men (n=322)	107	72	115	28	
No. of cases, women (n=158)	128	8	21	1	
Multivariable RR ^a , whole cohort	1 (Referent)	1.1 (0.85-1.5)	1.5 (1.2-1.9)	1.4 (0.92-2.1)	0.002
Multivariable RR ^a , men	1 (Referent)	1.1 (0.77-1.4)	1.4 (1.1-1.8)	1.3 (0.86-2.0)	0.02
Multivariable RR ^a , women	1 (Referent)	1.6 (0.79-3.4)	1.8 (1.1-2.9)	1.6 (0.22-11.5)	0.009
Lung cancer					
No. of cases, whole cohort (n=905)	254	132	391	128	
No. of cases, men (n=628)	77	110	321	120	
No. of cases, women (n=277)	177	22	70	8	
Multivariable RR ^a , whole cohort	1 (Referent)	2.1 (1.6-2.6)	4.8 (4.0-5.7)	7.9 (6.2-10.0)	<0.0001
Multivariable RR ^a , men	1 (Referent)	2.0 (1.5-2.7)	5.0 (3.9-6.4)	7.9 (5.9-10.6)	<0.0001
Multivariable RR ^a , women	1 (Referent)	2.7 (1.7-4.3)	4.3 (3.2-5.7)	8.3 (4.0-17.1)	<0.0001
Bladder/renal pelvis cancer					
No. of cases, whole cohort (n=146)	61	33	40	12	
Multivariable RR ^a , whole cohort	1 (Referent)	1.5 (0.96-2.5)	1.7 (1.1-2.7)	2.7 (1.4-5.2)	0.002
Breast cancer (women only)					
No. of cases, women (n=534)	491	14	28	1	
Multivariable RR ^a , women	1 (Referent)	1.2 (0.68-2.0)	0.97 (0.66-1.4)	0.58 (0.08-4.1)	0.87

Cancer sites	Current smoker		p-trend
	Never smoker	Former smoker	
	≤22 cigarettes/day	>22 cigarettes/day	

^aRelative risk (95% confidence interval) estimated from a multivariable-adjusted proportional hazards model adjusted for age (years), gender (males, females), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (g), and h/week of moderate physical activity (none, ½ to 3 h/week, 4 h/week or more).

Table 3

Relation between smoking categories and mortality

Cause of death	Never smoker	Former smoker	Current smoker		p-trend
			≤22 cigarettes/day	>22 cigarettes/day	
All causes					
No. of deaths, whole cohort (n=7982)	3945	1340	2228	469	
No. of deaths, men (n=4706)	1325	1146	1802	433	
No. of deaths, women (n=3276)	2620	194	426	36	
Multivariable RR ^a , whole cohort	1 (Referent)	1.3 (1.2-1.4)	1.7 (1.6-1.8)	1.8 (1.6-2.0)	<0.0001
Multivariable RR ^a , men	1 (Referent)	1.3 (1.2-1.4)	1.7 (1.6-1.8)	1.7 (1.5-1.9)	<0.0001
Multivariable RR ^a , women	1 (Referent)	1.7 (1.4-1.9)	1.6 (1.4-1.8)	2.5 (1.8-3.5)	<0.0001
Cancer					
No. of deaths, whole cohort (n=2710)	1271	402	835	202	
No. of deaths, men (n=1618)	403	344	685	186	
No. of deaths, women (n=1092)	868	58	150	16	
Multivariable RR ^a , whole cohort	1 (Referent)	1.3 (1.2-1.5)	2.1 (1.9-2.3)	2.4 (2.0-2.8)	<0.0001
Multivariable RR ^a , men	1 (Referent)	1.3 (1.1-1.4)	2.1 (1.9-2.4)	2.3 (2.0-2.8)	<0.0001
Multivariable RR ^a , women	1 (Referent)	1.7 (1.3-2.3)	1.9 (1.6-2.3)	3.6 (2.2-5.9)	<0.0001
Ischemic heart disease					
No. of deaths, whole cohort (n=1664)	799	300	479	86	
No. of deaths, men (n=1050)	291	258	402	79	
No. of deaths, women (n=634)	508	42	77	7	
Multivariable RR ^a , whole cohort	1 (Referent)	1.3 (1.2-1.5)	1.7 (1.5-1.9)	1.6 (1.3-2.0)	<0.0001
Multivariable RR ^a , men	1 (Referent)	1.3 (1.1-1.6)	1.8 (1.5-2.1)	1.6 (1.2-2.0)	<0.0001
Multivariable RR ^a , women	1 (Referent)	1.7 (1.2-2.3)	1.4 (1.1-1.8)	2.4 (1.1-5.1)	0.0002
Stroke					
No. of deaths, whole cohort (n=832)	476	136	193	27	
No. of deaths, men (n=436)	149	119	145	23	
No. of deaths, women (n=396)	327	17	48	4	
Multivariable RR ^a , whole cohort	1 (Referent)	1.1 (0.89-1.4)	1.2 (1.0-1.4)	0.86 (0.57-1.3)	0.27
Multivariable RR ^a , men	1 (Referent)	1.1 (0.85-1.4)	1.1 (0.89-1.4)	0.74 (0.47-1.2)	0.93
Multivariable RR ^a , women	1 (Referent)	1.0 (0.62-1.7)	1.3 (0.98-1.8)	2.1 (0.80-5.8)	0.03
Chronic obstructive pulmonary disease					
No. of deaths, whole cohort (n=378)	85	97	162	34	
No. of deaths, men (n=282)	36	87	126	33	
No. of deaths, women (n=96)	49	10	36	1	
Multivariable RR ^a , whole cohort	1 (Referent)	3.6 (2.6-5.0)	4.9 (3.7-6.5)	5.6 (3.6-8.6)	<0.0001
Multivariable RR ^a , men	1 (Referent)	3.1 (2.1-4.6)	4.1 (2.8-5.9)	5.0 (3.1-8.0)	<0.0001
Multivariable RR ^a , women	1 (Referent)	4.0 (2.0-8.1)	6.6 (4.3-10.2)	3.5 (0.49-25.6)	<0.0001

^aRelative risk (95% confidence interval) estimated from a multivariable-adjusted proportional hazards model adjusted for age (years), gender (males, females), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (g), and h/week of moderate physical activity (none, 1/2 to 3 h/week, 4 h/week or more).

Table 4

Relation between smoking cessation and cancer incidence

Cancer site	Current smoker	Years since quitting among former smokers		Never smoker
		<10 years	≥10 years	
All cancers	1577	356	355	2965
No. of cases, whole cohort (n=5253)	1 (Referent)	0.84 (0.75–0.94)	0.67 (0.59–0.75)	0.61 (0.57–0.66)
Multivariable RR ^a , whole cohort				
Head and neck cancer	148	31	17	149
No. of cases, whole cohort (n=345)	1 (Referent)	0.81 (0.55–1.2)	0.37 (0.22–0.61)	0.47 (0.36–0.61)
Multivariable RR ^a , whole cohort				
Upper gastrointestinal tract cancer	161	38	37	222
No. of cases, whole cohort (n=458)	1 (Referent)	0.87 (0.61–1.3)	0.69 (0.48–1.0)	0.66 (0.52–0.84)
Multivariable RR ^a , whole cohort				
Hepatobiliary and pancreas cancer	165	31	49	235
No. of cases, whole cohort (n=480)	1 (Referent)	0.67 (0.46–0.99)	0.85 (0.62–1.2)	0.69 (0.54–0.86)
Multivariable RR ^a , whole cohort				
Lung cancer	519	88	44	254
No. of cases, whole cohort (n=905)	1 (Referent)	0.59 (0.47–0.74)	0.23 (0.17–0.32)	0.20 (0.16–0.23)
Multivariable RR ^a , whole cohort				
Bladder/renal pelvis cancer	52	21	12	61
No. of cases, whole cohort (n=146)	1 (Referent)	1.2 (0.75–2.1)	0.52 (0.27–0.98)	0.54 (0.36–0.82)
Multivariable RR ^a , whole cohort				

^aRelative risk (95% confidence interval) estimated from a multivariable-adjusted proportional hazards model adjusted for age (years), gender (males, females), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (g), and h/week of moderate physical activity (none, ½ to 3 h/week, 4 h/week or more).

Table 5

Relation between smoking cessation and mortality

Cause of death	Years since quitting among former smokers		
	Current smoker	<10 years	≥10 years
All causes			
No. of deaths, whole cohort (n=7982)	2697	702	638
Multivariable RR ^a , whole cohort	1 (Referent)	0.91 (0.84-0.99)	0.65 (0.60-0.71)
Cancer			
No. of deaths, whole cohort (n=2710)	1037	222	180
Multivariable RR ^a , whole cohort	1 (Referent)	0.77 (0.67-0.89)	0.50 (0.42-0.58)
Ischemic heart disease			
No. of deaths, whole cohort (n=1664)	565	159	141
Multivariable RR ^a , whole cohort	1 (Referent)	0.96 (0.80-1.1)	0.66 (0.55-0.80)
Chronic obstructive pulmonary disease			
No. of deaths, whole cohort (n=378)	196	59	38
Multivariable RR ^a , whole cohort	1 (Referent)	1.0 (0.75-1.3)	0.50 (0.35-0.71)

^aRelative risk (95% confidence interval) estimated from a multivariable-adjusted proportional hazards model adjusted for age (years), gender (males, females), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (g), and h/week of moderate physical activity (none, 1/2 to 3 h/week, 4 h/week or more).

Population attributable risk (PAR) of cancer morbidity and all cause and cause-specific mortality to cigarette smoking

Table 6

	Relative risk (95% confidence interval) associated with ever smoking ^a		Population attributable risk associated with ever smoking	
	Men	Women	Men	Women
Cancer morbidity				
All cancers	1.5 (1.4–1.7)	1.4 (1.3–1.6)	23.0%	3.6%
Head and neck cancer	1.7 (1.3–2.3)	2.4 (1.3–4.1)	29.0%	10.5%
Upper gastrointestinal tract cancer	1.4 (1.1–1.9)	1.3 (0.85–2.0)	19.8%	2.7%
Hepatobiliary and pancreatic cancer	1.2 (0.97–1.6)	1.7 (1.2–2.6)	11.9%	6.1%
Lung cancer	4.1 (3.2–5.2)	4.0 (3.1–5.1)	63.8%	20.5%
Bladder/renal pelvis cancer	1.8 (1.2–2.8)	1.3 (0.48–3.3)	32.5%	2.2%
Mortality				
All causes	1.5 (1.4–1.6)	1.7 (1.5–1.8)	22.6%	5.3%
Cancer	1.8 (1.6–2.0)	1.9 (1.7–2.2)	30.8%	7.4%
Ischemic heart disease	1.6 (1.4–1.8)	1.5 (1.2–1.8)	24.3%	4.3%
Chronic obstructive pulmonary disease	3.8 (2.6–5.3)	5.7 (3.8–8.6)	61.3%	29.2%

^aRelative risk (95% confidence interval) estimated from a multivariable-adjusted proportional hazards model adjusted for age (years), dialect group (Cantonese, Hokkien), year of recruitment, level of education (no formal education, primary school, secondary school or above), daily ethanol intake (g), and h/week of moderate physical activity (none, 1/2 to 3 h/week, 4 h/week or more).