

The effect of $\alpha 2\text{-}\delta$ and other accessory subunits on expression and properties of the calcium channel $\alpha 1\text{G}$

A. C. Dolphin, C. N. Wyatt, J. Richards, R. E. Beattie*, P. Craig*, J.-H. Lee †, L. L. Cribbs †, S. G. Volsen* and E. Perez-Reyes †

*Department of Pharmacology, University College London, Gower Street, London WC1E 6BT, *Lilly Research Centre, Erl Wood Manor, Windlesham, Surrey GU20 6PH, UK and †Department of Physiology, Loyola University Medical Center, 2160 South 1st Avenue, Maywood, IL 60153, USA*

(Received 1 March 1999; accepted after revision 10 May 1999)

1. The effect has been examined of the accessory $\alpha 2\text{-}\delta$ and β subunits on the properties of $\alpha 1\text{G}$ currents expressed in monkey COS-7 cells and *Xenopus* oocytes.
2. In immunocytochemical experiments, the co-expression of $\alpha 2\text{-}\delta$ increased plasma membrane localization of expressed $\alpha 1\text{G}$ and conversely, the heterologous expression of $\alpha 1\text{G}$ increased immunostaining for endogenous $\alpha 2\text{-}\delta$, suggesting an interaction between the two subunits.
3. Heterologous expression of $\alpha 2\text{-}\delta$ together with $\alpha 1\text{G}$ in COS-7 cells increased the amplitude of expressed $\alpha 1\text{G}$ currents by about 2-fold. This finding was confirmed in the *Xenopus* oocyte expression system. The truncated δ construct did not increase $\alpha 1\text{G}$ current amplitude, or increase its plasma membrane expression. This indicates that it is the exofacial $\alpha 2$ domain that is involved in the enhancement by $\alpha 2\text{-}\delta$.
4. $\beta 1\text{b}$ also produced an increase of functional expression of $\alpha 1\text{G}$, either in the absence or the presence of heterologously expressed $\alpha 2\text{-}\delta$, whereas the other β subunits had much smaller effects.
5. None of the accessory subunits had any marked influence on the voltage dependence or kinetics of the expressed $\alpha 1\text{G}$ currents. These results therefore suggest that $\alpha 2\text{-}\delta$ and $\beta 1\text{b}$ interact with $\alpha 1\text{G}$ to increase trafficking of, or stabilize, functional $\alpha 1\text{G}$ channels expressed at the plasma membrane.

Voltage-dependent calcium channels and currents in native neurons and other cells have been divided into high voltage activated (HVA) and low voltage activated (LVA) (Carbone & Lux, 1984; Nowycky *et al.* 1985). LVA currents can be distinguished by their activation at smaller depolarizations, near to the resting potential, and by their rapid inactivation (Huguenard, 1996). At the single channel level, channels with a small unitary conductance activate over the same voltage range (Carbone & Lux, 1984). Native T-type channels are heterogeneous (Kobrinisky *et al.* 1994; Huguenard, 1996), suggesting that they comprise more than one subtype of channel. A new subfamily of voltage-dependent calcium channel $\alpha 1$ subunit genes (comprising $\alpha 1\text{G}$, $\alpha 1\text{H}$ and $\alpha 1\text{I}$) has recently been cloned, whose structure is superficially similar to the previously cloned HVA $\alpha 1$ subunits A, B, C, D, E and S (Perez-Reyes *et al.* 1998; Cribbs *et al.* 1998; Lee *et al.* 1999), having four domains, each with a voltage sensor and a pore-forming P loop. However, there are a number of regions where the homology is very low, particularly in the

intracellular linkers and the N and C termini. These novel channels, when expressed, form rapidly inactivating LVA currents that also have a small single channel conductance and slowly deactivating tail currents like native T-type currents (Carbone & Lux, 1984; Armstrong & Matteson, 1985). Recently, using an antisense approach, evidence has been obtained that T-type currents in primary sensory neurons are generated by the $\alpha 1\text{G}$, H and I family (Lambert *et al.* 1998).

The HVA channels are all thought to form heteromeric channels with the accessory subunits $\alpha 2\text{-}\delta$, β and possibly γ . The accessory subunits, particularly β subunits, have marked effects on the assembly of functional channels at the plasma membrane. In expression systems, the β subunits increase the number of plasma membrane channels (Chien *et al.* 1995; Shistik *et al.* 1995; Brice *et al.* 1997) and also affect the voltage dependence and kinetics of activation and inactivation (Jones *et al.* 1998). Inactivation kinetics are

particularly affected by different β subunits in a β -dependent manner, with $\beta 2a$ producing a marked slowing of inactivation (De Waard & Campbell, 1995).

It has not yet been examined whether the expression of LVA channel $\alpha 1$ subunits requires any known accessory subunits for trafficking to the plasma membrane, or for functional expression, or whether these accessory subunits influence the biophysical properties of the expressed channels. There are notable structural differences when comparing the $\alpha 1G$, H and I sequence with those of HVA channels, one contrast being that the I–II loop of the LVA channels is very large. Furthermore, they do not contain the full consensus sequence identified to be the binding site for β subunits on the I–II loop, the α interaction domain (AID). This consensus sequence, QQ-E-D/EL-GY--WI--E, is present in all the HVA channels (Walker & De Waard, 1998). The homologous sequence in $\alpha 1G$ is GSCYE**ELLK**YLVYILRKA, with identical residues underlined in bold, and a conserved charge in italics. However, in the C terminal part of the consensus sequence, the W is Y in $\alpha 1G$, a conservative substitution, compared with the W→A mutation identifying this residue as essential for β subunit binding (De Waard *et al.* 1996). Thus the degree of conservation of the consensus sequence may be sufficient for β interaction.

The consensus site(s) on the exofacial loops of HVA $\alpha 1$ subunits to which the $\alpha 2$ - δ subunit binds have not been identified, although repeat III of $\alpha 1S$ has been found to associate with $\alpha 2$ (Gurnett *et al.* 1997). Furthermore, the extracellular region of δ is also involved in the production of certain functional effects, although the region of $\alpha 1$ with which it interacts is not known (Felix *et al.* 1997). Thus it is not possible to determine *a priori* whether $\alpha 1G$ has the capacity to bind to $\alpha 2$ - δ . This study was therefore designed to examine whether $\alpha 1G$ is influenced by any accessory subunits.

METHODS

Materials

The rat $\alpha 2$ - δ_b cDNA (accession number M86621) was obtained from either Dr H. Chin (NIH, Bethesda, MD, USA) or Dr T. Snutch (UBC, Vancouver, Canada), rat $\beta 1b$ (X11394) from Dr T. Snutch and mut-3 GFP (green fluorescent protein) from Dr T. Hughes (Yale, New Haven, CT, USA). Rat $\beta 2a$ (M80545), $\beta 3$ (M88751) and $\beta 4$ (M80545) cDNAs were also used.

Molecular biology

The $\alpha 1G$ cDNA (Perez-Reyes *et al.* 1998) was subcloned into the vertebrate expression vector pMT2 (Genetics Institute, Cambridge, MA, USA) (Swick *et al.* 1992), using standard molecular biological techniques, and the correct orientation of the insert was verified by multiple restriction digests. All other cDNAs were expressed from the same vector for expression in monkey COS-7 cells. The δ subunit (Gurnett *et al.* 1996) was constructed by deletion of the $\alpha 2$ cDNA, using the plasmid pMT2- $\alpha 2$ - δ as a template. The forward primer was ATGGAAGAGGATGACTTCACAGCT, which binds at the beginning of the δ sequence, position 2995. The reverse primer was AGGGAAGGGCTCCTCGCTCGA, which binds at position 238, including only the signal sequence of $\alpha 2$. The PCR product of

6.1 kb was amplified using *Pfu* polymerase for 14 cycles, with the annealing temperature of 50 °C for 30 s, and an extension temperature of 75 °C for 13 min. This product was then treated with polynucleotide kinase and ligated to form the pMT2- δ construct, attached to the signal sequence of $\alpha 2$ - δ , as described previously (Felix *et al.* 1997). The construct was sequenced for confirmation of its identity.

COS-7 cell expression

Cells were cultured and transfected by electroporation, essentially as described previously (Campbell *et al.* 1995a). The $\alpha 1G$, $\alpha 2$ - δ or δ , β and GFP cDNAs in the vector pMT2 were used at 15, 5, 5 and 1 μ g per transfection, respectively. Blank pMT2 vector was included where necessary to maintain the total cDNA at 31 μ g per transfection. In some experiments in which $\alpha 1G$ was expressed alone, the additional pMT2 vector was omitted, with no effect on the current properties. Following transfection, cells were maintained at 37 °C for about 60 h, and then replated using non-enzymatic cell dissociation medium (Sigma), and maintained at 25 °C for between 2 and 8 h prior to electrophysiological recording, or for 1–2 h prior to fixation for immunocytochemistry.

Ba²⁺ currents were recorded using the whole cell patch-clamp technique. The internal and external solutions were similar to those described previously (Campbell *et al.* 1995b). The patch pipette solution contained (mM): caesium aspartate, 140; EGTA, 5; MgCl₂, 2; CaCl₂, 0.1; K₂ATP, 2; Hepes, 10; pH 7.2, 310 mosmol kg⁻¹ with sucrose. The external solution contained (mM): tetraethylammonium (TEA) bromide, 160; KCl, 3; NaHCO₃, 1.0; MgCl₂, 1.0; Hepes, 10; glucose, 4; BaCl₂, 1; pH 7.4, 320 mosmol kg⁻¹ with sucrose. Whole cell currents were elicited from a holding potential (V_h) of -100 mV and recorded using an Axopatch-1D amplifier. Data were filtered at 2–5 kHz and digitized at 10–20 kHz. The junction potential between external and internal solutions was 6 mV; the values given in the figures and text have not been corrected for this. Current records are shown following leak and residual capacitance current subtraction (*P/8* protocol) and series resistance compensation up to 80%. All experiments were performed at room temperature (20–24 °C). Analysis was performed using pCLAMP6 (Axon Instruments) and Origin 5 software (Microcal Software Inc., Northampton, MA, USA).

Xenopus oocyte expression

Oocytes were obtained from *X. laevis* (Xenopus One) using standard techniques (Leonard & Snutch, 1991), which have been approved by the Loyola University Animal Care Committee. Frogs were anaesthetized using 1 g l⁻¹ of tricaine, then oocytes were collected by making a 1 cm incision in the ventral abdomen and removing a portion of the ovary. The frog was sutured both on the rectus abdominus muscle sheath, and then on the skin. It was allowed to recover for 4 h before returning it to the home tank. Oocytes were prepared for microinjection as previously described (Leonard & Snutch, 1991).

Capped cRNA was synthesized using T7 RNA polymerase (Ambion). The rat $\alpha 1G$ (Perez-Reyes *et al.* 1998) was contained in the vector pGEM-HEA. The rat brain $\alpha 2$ - δ_b was contained in the vector pAGA. Each oocyte was injected with 5 ng of either $\alpha 1G$ alone or plus 5 ng $\alpha 2$ - δ cRNA in a volume of 50 nl. Oocytes were voltage clamped using a two-microelectrode voltage clamp amplifier (OC-725B, Warner Instrument Corp.). Voltage and current electrodes (1.5–1.8 M Ω tip resistance) were filled with a cushion of 1% agarose and 3 M KCl (Schreibmayer *et al.* 1994). The bath solution contained the following (mM): 10 BaCl₂, 36 tetraethyl ammonium, 2 CsCl₂ and 5 Hepes, adjusted to pH 7.4 with methanesulfonic acid. Data were filtered at 1 kHz and acquired at

4 kHz using the pCLAMP system (Digidata 1200 and pCLAMP 6.0). All data are expressed as the means \pm s.e.m. and statistical analysis was performed using Student's *t* test.

Immunocytochemistry

COS-7 cells were washed twice in Tris-buffered saline (TBS; 154 mM NaCl, 20 mM Tris, pH 7.4), then fixed in 4% paraformaldehyde in TBS as described (Brice *et al.* 1997). The cells were permeabilized in 0.02% Triton X-100 in TBS, and incubated with blocking solution (20% (v/v) goat serum, 4% (w/v) bovine serum albumin (BSA), 0.1% (w/v) DL-lysine in TBS). The cells were incubated for 14 h at 4 °C with the appropriate primary antibody diluted 1:500 in 10% goat serum, 2% BSA, 0.05% DL-lysine. The VDCC $\alpha 2$ antibody used in this study was raised in rabbits against a specific peptide derived from the sequence of $\alpha 2$. Its specificity has been described previously (Brickley *et al.* 1995; Brice *et al.* 1997). The $\alpha 1\text{G}$ antibody was raised in rabbits against a glutathione S transferase (GST) fusion protein derived from the cytoplasmic loop between IS6 and IIS1 of $\alpha 1\text{G}$, as described in Craig *et al.* (1999), in which the antibody specificity is also described. The anti-GST component of the serum was removed by adsorption on GST Sepharose 4B, and it was then affinity purified on $\alpha 1\text{G}$ -GST Sepharose 4B. The stock concentration of affinity-purified $\alpha 1\text{G}$ antibody was 98 $\mu\text{g ml}^{-1}$. The primary antibodies were detected using biotin-conjugated goat

anti-rabbit IgG (1:200) (Sigma), then streptavidin FITC (1:100) (Molecular Probes, Eugene, OR, USA). Cells were examined on an MRC 1024 laser scanning confocal microscope (Bio-Rad, Hemel Hempstead, UK), with all parameters (gain, aperture) identical between experiments. Images represent 2 μm optical sections midway through the cell. Quantification was performed using Imagequant software (Molecular Dynamics, Sunnyvale, CA, USA), and results are given, following background subtraction, as arbitrary units (a.u.) of pixel density.

RESULTS

The effect of co-expression of $\alpha 2\text{-}\delta$ on the immunolocalization of $\alpha 1\text{G}$

COS-7 cells represent an expression system with a very low background in terms of expressed currents. No Ba^{2+} currents were detected in the absence of expressed $\alpha 1$ subunits (Berrow *et al.* 1997; Brice *et al.* 1997; Meir & Dolphin, 1998). However, by RT-PCR using primers conserved across all species whose sequences have been reported, we found evidence for a low level of mRNA for $\alpha 1\text{G}/\text{H}/\text{I}$, $\alpha 2\text{-}\delta$ and $\beta 3$ (J. Richards and A. C. Dolphin, unpublished results), but





Figure 1. Effect of co-expression of $\alpha 2\text{-}\delta$ on immunolocalization of expressed $\alpha 1\text{G}$ in COS-7 cells. *A*, $\alpha 1\text{G}$ immunolocalization in untransfected COS-7 cells; *B*, $\alpha 1\text{G}$ immunolocalization in cells transfected with $\alpha 1\text{G}$; *C*, $\alpha 1\text{G}$ immunolocalization in cells transfected with $\alpha 1\text{G} + \alpha 2\text{-}\delta$. *D*, lack of immunofluorescence in the absence of the primary antibody. Experiments were performed under identical conditions in parallel cultures, and repeated 3 times with similar results. Scale bar, 20 μm .

Table 1. Biophysical properties of $\alpha 1G$, expressed with different accessory subunit combinations in COS-7 cells

| Transfected subunits | I_{Ba} at -0 mV (pA pF $^{-1}$) | G_{max} (nS pF $^{-1}$) | τ_{inact} at -20 mV (ms) | V_{50} for activation (mV) | V_{50} for steady-state inactivation (mV) |
|--------------------------------------|--------------------------------------|----------------------------|---------------------------------|------------------------------|---|
| $\alpha 1G$ | -19.2 ± 3.1 (15) | 0.55 ± 0.07 (15) | 13.8 ± 1.3 (7) | -27.1 ± 2.7 (10) | -72.9 ± 1.3 (8) |
| $\alpha 1G/\alpha 2-\delta$ | $-40.3 \pm 6.2^{**}$ (17) | $0.75 \pm 0.08^*$ (17) | 13.1 ± 0.3 (6) | -27.3 ± 1.8 (8) | -69.7 ± 1.1 (6) |
| $\alpha 1G/\beta 1b$ | $-39.8 \pm 3.8^{**}$ (5) | $0.87 \pm 0.07^{**}$ (5) | 14.2 ± 1.0 (5) | -22.3 ± 4.5 (5) | -68.7 ± 5.6 (4) |
| $\alpha 1G/\alpha 2-\delta/\beta 1b$ | $-39.5 \pm 4.9^{**}$ (9) | $0.79 \pm 0.09^{**}$ (9) | 13.6 ± 0.4 (4) | -27.4 ± 3.8 (3) | -72.0 ± 2.6 (5) |
| $\alpha 1G/\beta 2a$ | -27.3 ± 5.6 (9) | $0.67 \pm 0.09^*$ (9) | 11.7 ± 0.6 (3) | -22.3 ± 1.0 (3) | -69.4 ± 2.0 (5) |
| $\alpha 1G/\beta 3$ | $-30.7 \pm 3.2^*$ (11) | 0.65 ± 0.06 (11) | 20.3 ± 2.1 (11) | $-17.0 \pm 2.2^*$ (8) | $-65.5 \pm 3.0^*$ (8) |
| $\alpha 1G/\beta 4$ | -23.8 ± 2.2 (6) | 0.62 ± 0.04 (6) | 12.9 ± 1.8 (5) | $-20.7 \pm 1.2^*$ (5) | $-67.3 \pm 2.5^{**}$ (4) |

The parameters determined for the different $\alpha 1G$ and accessory subunit combinations were measured as described in Methods. The data for each subunit combination were determined from at least 3 separate transfections. Individual current density–voltage relationships were fitted with a Boltzmann equation, $I = G_{max}(V - V_{rev}) / (1 + \exp(-(V - V_{1/2})/k))$, where G_{max} is the maximum conductance, V_{rev} is the reversal potential, k is the slope factor and $V_{1/2}$ is the voltage for 50% current activation. No significant differences were observed in the V_{rev} or k values (results not shown). The activation V_{50} values given were determined from tail current analysis (Fig. 4), and steady-state inactivation V_{50} data were obtained using the protocol described in the legend to Fig. 4, and fitted with a single Boltzmann equation of the form $I/I_{max} = 1 / (1 + \exp((V - V_{50})/k))$. The statistical significance of the differences compared with $\alpha 1G$ alone are indicated by * $P < 0.05$; ** $P < 0.01$ (Student's two-tailed t test).

Figure 2. I – V relationships for $\alpha 1G$ and the effect of accessory subunits in COS-7 cells

A shows $\alpha 1G \pm \alpha 2-\delta$ and B shows $\alpha 1G/\beta 1b \pm \alpha 2-\delta$. The left and centre panels show representative traces of $\alpha 1G$ currents without (left) or with (centre) co-expression of $\alpha 2-\delta$. Currents are shown in response to voltage steps (V) from -100 mV to between -70 and -20 mV, in 5 mV steps. The right panels show the corresponding I – V relationships (means \pm S.E.M.) for the numbers of experiments given in parentheses.

immunocytochemistry and Western blotting showed no β subunit protein in untransfected cells (Campbell *et al.* 1995a; Berrow *et al.* 1997; Brice *et al.* 1997; Stephens *et al.* 1997). A very low level of $\alpha 1\text{G}$ immunostaining was observed in untransfected control cells, but this was not localized to the plasma membrane (Fig. 1A; pixel density throughout cell, 5.5 ± 0.8 a.u., of which none was membrane associated, $n = 10$). Expression of $\alpha 1\text{G}$ resulted in much greater immunostaining for $\alpha 1\text{G}$, largely at the plasma membrane (Fig. 1B; pixel density at plasma membrane, 30.8 ± 4.4 a.u., $n = 10$). Co-expression of $\alpha 2\text{-}\delta$ with $\alpha 1\text{G}$ resulted in a further increase in staining intensity at the plasma membrane (Fig. 1C; pixel density at plasma membrane, 42.0 ± 5.1 a.u., $n = 10$, a 36.4% increase). No staining was observed in the absence of the primary antibody (Fig. 1D). These results suggest that heterologously expressed $\alpha 2\text{-}\delta$ is either increasing translation of $\alpha 1\text{G}$, increasing its trafficking to the plasma membrane, or increasing its stability. This raises the possibility that since endogenous $\alpha 2\text{-}\delta$ is present in cells, it may be involved in trafficking heterologously expressed $\alpha 1\text{G}$ when $\alpha 1\text{G}$ is expressed alone.

Properties of $\alpha 1\text{G}$ expressed in COS-7 cells

The heterologous expression of $\alpha 1\text{G}$ alone in COS-7 cells resulted in the observation of T-type currents, using 1 mM

Ba^{2+} as the charge carrier, in a large proportion (usually $>70\%$) of GFP-positive cells (Fig. 2A and Table 1). The peak current was observed between -30 and -20 mV. The inactivation kinetics were voltage dependent at small depolarizations, reaching a voltage-independent minimum at potentials above -30 mV (Fig. 3A and Table 1). Recovery from inactivation occurred with a time constant of 107 ms (Fig. 3B). The voltage for 50% activation (V_{50}) was -27 mV, and the V_{50} for inactivation was -73 mV (Fig. 4 and Table 1). Superimposition of the activation and steady-state inactivation relationships showed no clear window current (inset in Fig. 4B).

The effect of co-expression of $\alpha 2\text{-}\delta$ on the properties of $\alpha 1\text{G}$ currents in COS-7 cells

Co-expression with $\alpha 2\text{-}\delta$ significantly increased the $\alpha 1\text{G}$ current density at -20 mV by 2.1-fold, and increased the maximum conductance, compared with that for $\alpha 1\text{G}$ alone (Fig. 2A and Table 1), while not significantly affecting any of the kinetic parameters measured (Figs 2A and 3A and Table 1). The kinetics of inactivation were very similar for $\alpha 1\text{G}$ and $\alpha 1\text{G}/\alpha 2\text{-}\delta$, reaching identical voltage-independent minima above -30 mV (Fig. 3A and Table 1). The recovery from inactivation also followed an identical time course (Fig. 3B). Furthermore, the reversal potential (Fig. 2A) was




Figure 3. Inactivation kinetics of $\alpha 1\text{G}$ currents: effect of accessory $\alpha 2\text{-}\delta$ subunits in COS-7 cells

A, the inactivation phase of individual currents was fitted to a single exponential, and the mean values were plotted for $\alpha 1\text{G}$ (■) and $\alpha 1\text{G}/\alpha 2\text{-}\delta$ (□), for the numbers of experiments given in parentheses. B, recovery from inactivation was determined using the protocol shown in the inset, which depicts 13 overlaid traces obtained at 30 s intervals. The V_h was -100 mV. A 50 ms test step was given to -20 mV (I_{max}), followed by a 50 ms step to $+100$ mV to produce complete inactivation, followed by a variable interpulse interval (Δt), and a subsequent identical test step ($I_{\Delta t}$). The recovery from inactivation was plotted for $\alpha 1\text{G}$ (■) and $\alpha 1\text{G}/\alpha 2\text{-}\delta$ (□), for the numbers of experiments given in parentheses. The data points were fitted to single exponentials (dotted lines), with τ values of 122.6 ms for $\alpha 1\text{G}$ and 125.1 ms for $\alpha 1\text{G}/\alpha 2\text{-}\delta$.

not significantly increased by $\alpha 2\text{-}\delta$ ($+29.0 \pm 2.3$ mV, $n = 15$, for $\alpha 1\text{G}$ compared with $+34.3 \pm 2.7$ mV, $n = 19$, for $\alpha 1\text{G}/\alpha 2\text{-}\delta$). The voltage dependence of activation (Fig. 4A) and the steady-state inactivation (Fig. 4B) of $\alpha 1\text{G}$ were also not significantly affected by co-expression of $\alpha 2\text{-}\delta$ (Table 1). However, there was a small change in the value of the slope factor, k , of the Boltzmann fits to the activation data (evident from Fig. 4A), which was 9.1 ± 0.5 mV ($n = 10$) for $\alpha 1\text{G}$ and 7.5 ± 0.5 mV ($n = 8$) for $\alpha 1\text{G}/\alpha 2\text{-}\delta$.

In control experiments no calcium channel currents were observed in cells transfected with $\alpha 2\text{-}\delta$ alone, or with $\alpha 2\text{-}\delta$ plus a β subunit (results not shown), indicating that these accessory subunits are not recruiting an endogenous $\alpha 1$ subunit, as we also concluded in a previous work (Meir & Dolphin, 1998).

The effect of co-expression of $\alpha 2\text{-}\delta$ on the properties of $\alpha 1\text{G}$ currents expressed in *Xenopus* oocytes

Co-injection of $\alpha 2\text{-}\delta$ with $\alpha 1\text{G}$ cRNA in *Xenopus* oocytes increased the peak currents at -20 mV from -493 ± 60 nA for $\alpha 1\text{G}$ alone to -784 ± 79 nA for $\alpha 1\text{G}/\alpha 2\text{-}\delta$ ($P < 0.05$, $n = 40$ from 4 separate experiments; Fig. 5). The average

enhancement of the $\alpha 1\text{G}$ current by $\alpha 2\text{-}\delta$ from these four experiments was 1.7 ± 0.2 -fold. The shape of the current–voltage (I – V) relationship was not affected by co-injection with $\alpha 2\text{-}\delta$, as determined following normalization of the peak currents. There was also no effect on the voltage dependence of current activation, determined from I – V relationships (for $\alpha 1\text{G}$ the V_{50} for activation was -31.6 ± 0.6 mV, $n = 40$, and for $\alpha 1\text{G}/\alpha 2\text{-}\delta$ it was -32.2 ± 0.3 mV, $n = 40$). Activation kinetics were not affected (τ at -20 mV was 1.9 ms for both), and neither were inactivation kinetics (τ_{inact} at -20 mV was 11.1 ± 0.3 ms, $n = 8$, for $\alpha 1\text{G}$ and 10.0 ± 0.4 ms, $n = 9$, for $\alpha 1\text{G}/\alpha 2\text{-}\delta$). Furthermore, injection of $\alpha 2\text{-}\delta$ had no significant effect on the voltage dependence of steady-state inactivation as measured with 10 s conditioning prepulses (the V_{50} for steady-state inactivation was -63.8 ± 0.3 mV, $n = 10$, for $\alpha 1\text{G}$ and -62.6 ± 0.5 mV, $n = 14$, for $\alpha 1\text{G}/\alpha 2\text{-}\delta$). All the data in oocytes were therefore in full agreement with the results from COS-7 cells.

Does heterologously expressed $\alpha 1\text{G}$ associate with endogenous $\alpha 2\text{-}\delta$?

Since $\alpha 2\text{-}\delta$ co-expression with $\alpha 1\text{G}$ results in an increase in $\alpha 1\text{G}$ immunostaining at the plasma membrane and an




Figure 4. Voltage dependence of activation and inactivation of $\alpha 1\text{G}$ currents in COS-7 cells: effect of accessory subunits

A, the current activation plots were determined from tail current amplitudes using the protocol shown in the inset, and normalized to the maximum current. Although this protocol, with a step length of 7.5 ms, gives a small error at low depolarizations because the current is not completely activated, this did not affect the V_{50} values. Mean \pm S.E.M. values are shown for $\alpha 1\text{G}$ (■), $\alpha 1\text{G}/\alpha 2\text{-}\delta$ (□), $\alpha 1\text{G}/\beta 4$ (○), with the numbers of experiments shown in the key. For both A and B, the curves are fits to the Boltzmann equation described in the legend to Table 1. B, steady-state inactivation curves were determined by measurement of peak current amplitude at -20 mV, following a 10 s conditioning prepulse to the potentials shown. Data were normalized before averaging the number of experiments given in the key. The symbols are the same as for A. The inset graph shows the region of overlap between the $\alpha 1\text{G}$ and $\alpha 1\text{G}/\alpha 2\text{-}\delta$ activation and steady-state inactivation curves.




Figure 5. Effect of $\alpha 2$ - δ on $\alpha 1G$ expression in *Xenopus* oocytes

The left and centre panels show representative families of $\alpha 1G$ currents recorded at between -60 and -10 mV in *Xenopus* oocytes in the absence (left) and presence (centre) of heterologously expressed $\alpha 2$ - δ . The I - V relationships on the right represent the mean \pm s.e.m. values from 40 experiments (10 oocytes from 4 different experiments) for $\alpha 1G$ (■), and $\alpha 1G + \alpha 2\delta$ (○).

increase in functional expression, it is possible that $\alpha 1G$, when expressed alone, associates with endogenous $\alpha 2$ - δ . We have observed a low level of endogenous $\alpha 2$ immunostaining in untransfected control cells (Fig. 6A; total pixel density of 38.2 ± 6.3 a.u., $n = 14$, from 3 separate transfection experiments), and this was not associated with the plasma membrane in most cells. The total staining for endogenous $\alpha 2$ was significantly increased when $\alpha 1G$ cDNA was transfected into the cells (Fig. 6B; pixel density 82.7 ± 11.9 a.u., $n = 19$, $P < 0.01$ compared with untransfected cells), and the staining always showed some membrane association, indicating an association between the two proteins. These results suggest that expression of $\alpha 1G$ is able to increase expression or stabilize endogenous $\alpha 2$ - δ , and also increase membrane association of endogenous $\alpha 2$ - δ .

Since these immunocytochemical results suggested that transfected $\alpha 1G$ was associating with endogenous $\alpha 2$ - δ , co-expression experiments were also performed with the δ construct, which has been found to influence some properties of $\alpha 1C$ and $\alpha 1A$, but unlike $\alpha 2$ - δ does not increase the current amplitude (Gurnett *et al.* 1996; Felix *et al.* 1997). When the δ -construct was expressed in COS-7 cells, clear immunostaining was observed at the plasma membrane (results not shown), using an antibody directed against an extracellular epitope in δ (Brickley *et al.* 1995). Co-expression of $\alpha 1G$ with δ reduced the plasma membrane expression of $\alpha 1G$ by 25%, from immunocytochemical experiments (pixel density 22.8 ± 1.8 a.u., $n = 10$, for $\alpha 1G/\delta$ compared with 30.8 ± 4.4 a.u. ($n = 10$) for $\alpha 1G$ expressed alone in parallel experiments). The expression level for $\alpha 1G$ when co-expressed




Figure 6. Effect of $\alpha 1G$ on immunolocalization of endogenous $\alpha 2$ - δ in COS-7 cells

A, immunolocalization of endogenous $\alpha 2$ in untransfected COS-7 cells; B, immunolocalization of endogenous $\alpha 2$ in cells transfected with $\alpha 1G$. Experiments were performed under identical conditions in parallel cultures, and transfections were repeated 3 times with similar results.

with δ represents a 45.5% reduction compared with co-expression of $\alpha 1G$ with full-length $\alpha 2-\delta$. In agreement with this, in electrophysiological experiments, co-expression of δ with $\alpha 1G$ did not produce any increase in the expression of $\alpha 1G$ currents, unlike $\alpha 2-\delta$. The current density for $\alpha 1G/\delta$ was 22.1 ± 3.2 pA pF⁻¹ at -20 mV, and the conductance was 0.49 ± 0.04 nS pF⁻¹ ($n = 8$). This result indicates that it is the $\alpha 2$ moiety of $\alpha 2-\delta$ that is responsible for the $\alpha 1G$ current enhancement, in agreement with the results for $\alpha 1C$ (Gurnett *et al.* 1996).

Effect of co-expression of β subunits on the properties of $\alpha 1G$ currents in COS-7 cells

Co-expression of $\alpha 1G$ with the β subunit $\beta 1b$ (Fig. 2B and Table 1) increased the $\alpha 1G$ current density without significant effect on the reversal potential, which was $+31.1 \pm 1.9$ mV ($n = 5$). In contrast, for co-expression of $\beta 2a$, $\beta 3$ or $\beta 4$ with $\alpha 1G$, only a small current enhancement was observed (Table 1). Additional co-expression of $\alpha 2-\delta$ produced no further increment of current density for $\beta 1b$ (Table 1). There was no systematic effect of β subunits on the inactivation kinetics of $\alpha 1G$ currents, the limiting τ_{inact} at -20 mV being similar for all subunit combinations (Table 1). There was also no effect of any β subunit on recovery from inactivation (results not shown). Co-expression with any one of the β subunits produced a small depolarization of the voltage dependence of activation of $\alpha 1G$, determined from tail current amplitudes (see Fig. 4A for $\beta 4$). The voltage for 50% activation (V_{50}) was only depolarized significantly by $\beta 3$ and $\beta 4$ (Table 1). Similarly, the steady-state inactivation of $\alpha 1G$, measured after a 10 s conditioning potential step, was shifted to more depolarized potentials by co-expression of the β subunits $\beta 3$ and $\beta 4$ (Table 1 and, for $\beta 4$, Fig. 4B). In control experiments, $\beta 1b$ did not induce calcium channel currents when expressed alone in COS-7 cells (results not shown).

DISCUSSION

Effect of co-expression of accessory $\alpha 2-\delta$ subunits on $\alpha 1G$ expression

From the literature, diverse effects of $\alpha 2-\delta$ have been reported on the properties of cloned HVA channels. Many studies of the effects of accessory subunits have been performed using oocytes, before it was realized that they contain endogenous calcium channel subunits, including a *Xenopus* $\beta 3$ subunit (Tareilus *et al.* 1997). Therefore, although the effect of $\alpha 2-\delta$ has been reported to increase the membrane expression of HVA $\alpha 1$ subunits, and to increase their open probability (Shistik *et al.* 1995; Gurnett *et al.* 1996) as well as shifting the voltage dependence of activation to more positive potentials (Qin *et al.* 1998), these effects may have been dependent on the presence of endogenous β subunits (Tareilus *et al.* 1997), and for $\alpha 1A$ the effect of $\alpha 2-\delta$ is small in the absence of co-expressed β subunits. In HEK293 cells $\alpha 2-\delta$ produced a small increase in the density of $\alpha 1E$ currents, but had no effect on the

biophysical properties (Jones *et al.* 1998), whereas in COS-7 cells we observed $\alpha 2-\delta$, in the absence of β , to shift activation of $\alpha 1E$ to more positive potentials, while having very little effect on current density (Stephens *et al.* 1997). In both cell types, $\alpha 2-\delta$ was ineffective alone in targeting $\alpha 1C$ or $\alpha 1A$ to the plasma membrane (Brice *et al.* 1997; Gao *et al.* 1999). A recent report has shown that a novel $\alpha 2-\delta$ ($\alpha 2\delta-3$) produces a similar enhancement of $\alpha 1C$ and $\alpha 1E$ current amplitudes to $\alpha 2\delta-1$ (the $\alpha 2-\delta$ used in the current study), but these effects are only evident in the additional presence of a β subunit (Klugbauer *et al.* 1999).

We have previously reported that we could not detect mRNA for $\alpha 2-\delta$ in COS-7 cells (Berrow *et al.* 1997), but since these experiments were performed, additional $\alpha 2-\delta$ sequences have been reported, and using PCR primers conserved across species, we have now observed a faint PCR product in COS-7 cells of the size expected for $\alpha 2-\delta$ (results not shown). Consistent with this, we have also observed a low level of immunostaining for endogenous $\alpha 2-\delta$, which, however, was not specifically plasma membrane associated (Fig. 6A). We have examined the effect of co-expression of $\alpha 2-\delta$ on immunostaining for heterologously expressed $\alpha 1G$, and observed a clear increase in $\alpha 1G$ immunostaining at the plasma membrane (Fig. 1). Furthermore, the effect of heterologous expression of $\alpha 1G$ on immunostaining for endogenous $\alpha 2-\delta$ was also studied, since we found previously that $\alpha 2-\delta$ was trafficked to the plasma membrane by HVA calcium channel $\alpha 1/\beta$ combinations (Brice *et al.* 1997). We consistently found that immunostaining for endogenous $\alpha 2-\delta$ was greater in cells transfected with $\alpha 1G$ than in untransfected cells, and that the staining was more plasma membrane associated, suggesting that $\alpha 1G$ may be associating with, and stabilizing, endogenous $\alpha 2-\delta$ and also trafficking it to the plasma membrane. We still observed $\alpha 1G$ currents when $\alpha 1G$ was co-expressed with a truncated δ construct, lacking the $\alpha 2$ domain, but the current amplitude was not enhanced as it was with $\alpha 2-\delta$, indicating that the extracellular $\alpha 2$ domain is required for current enhancement, in accord with previous results for $\alpha 1C$ (Gurnett *et al.* 1996). In further agreement with this, $\alpha 1G$ immunostaining was slightly reduced by δ , in contrast to the enhancement produced by $\alpha 2-\delta$, again indicating the importance of the $\alpha 2$ moiety in this effect. The δ subunit acted as a 'dominant negative' in previous studies, reducing the effect of $\alpha 2-\delta$ on $\alpha 1A$ expression (Gurnett *et al.* 1996). However, in the present study co-expression of δ with $\alpha 1G$ did not prevent the expression of $\alpha 1G$ currents, indicating either that it was not able to disrupt $\alpha 1G$ interaction with endogenous $\alpha 2-\delta$, or that such an interaction was not essential for functional expression of $\alpha 1G$ currents. It therefore remains unclear whether the presence of endogenous $\alpha 2-\delta$ is essential for the expression of functional $\alpha 1G$ channels, and in the future an antisense approach might be taken to deplete endogenous $\alpha 2-\delta$, in order to answer this point.

In the present study, overexpression of exogenous $\alpha 2-\delta$ produced an approximately 2-fold increase in the amplitude

of $\alpha 1\text{G}$ currents in COS-7 cells, and a 1.7-fold increase in *Xenopus* oocytes, but had no effect on their kinetics or voltage dependence of activation or inactivation, or on recovery from inactivation in either system. This argues either for an effect of $\alpha 2\text{-}\delta$ on trafficking of the nascent $\alpha 1\text{G}$ channels from the endoplasmic reticulum to the plasma membrane, or an effect to stabilize the plasma membrane channels in a functional conformation. Further studies at the single channel level will be necessary to determine whether there are effects on the open probability of the channels.

In a previous study, we have shown that overexpression of $\alpha 2\text{-}\delta$, in undifferentiated NG108-15 cells, induced the appearance of an HVA sustained current. It is likely that this current component represented L-type current that had a very low open probability in the absence of $\alpha 2\text{-}\delta$, as a similar component could also be induced by the L-type channel agonist Bay K8644 (Wyatt *et al.* 1998), and the induced HVA current was blocked by the L-type channel antagonist, nicardipine (C. N. Wyatt and A. C. Dolphin, unpublished results). There were no marked effects of $\alpha 2\text{-}\delta$ on the biophysical properties of the T-type current component itself, in terms of kinetics or voltage dependence (Wyatt *et al.* 1998), consistent with the observation made here. However, it should be noted that NG108-15 cells express substantial endogenous $\alpha 2\text{-}\delta$ at the plasma membrane.

Effect of co-expression of accessory β subunits on $\alpha 1\text{G}$ expression

In the present study we have shown that co-expression of the β subunit, $\beta 1\text{b}$, also has clear effects on expression of functional $\alpha 1\text{G}$ current (Brice *et al.* 1997). In contrast, a recent antisense study in nodose ganglia concluded that native T-type channels were not associated with β subunits (Lambert *et al.* 1997), although in these cells the main β subunits observed were $\beta 2$ and $\beta 3$, with no $\beta 1$ apparently present. However an antisense study in cardiac atrial cells has suggested that loss of β subunits may affect expression of T-type currents (Chen & Best, 1998). The consensus binding site for β subunits, which has been identified on the I–II loop of HVA $\alpha 1$ subunits, is not completely conserved in $\alpha 1\text{G}$ (Perez-Reyes *et al.* 1998). While it has been suggested that there is another region on the C terminus, at least of $\alpha 1\text{E}$ and $\alpha 1\text{A}$ (Tareilus *et al.* 1997; Walker *et al.* 1998), to which certain β subunits may bind, it has been disputed whether this has any functional consequences for the biophysical properties (Jones *et al.* 1998). It is therefore possible that $\alpha 1\text{G}$ may interact transiently with the $\beta 1\text{b}$ subunit, which may serve a chaperone function to traffic the $\alpha 1\text{G}$ channel protein to the plasma membrane, or to stabilize the $\alpha 1\text{G}$ once in the membrane, thus increasing current density, while having only minor effects on the biophysical characteristics of the channels. The finding that the effects of $\beta 1\text{b}$ and $\alpha 2\text{-}\delta$ were not additive in COS-7 cells suggests that they may be acting by a similar chaperone or stabilizing mechanism. We have observed that all β subunits increase the membrane expression of $\alpha 1\text{A}$ (Brice *et al.* 1997),

whereas it has been found that $\beta 1\text{b}$ was the only β subunit that increased the heterologous expression of $\alpha 1\text{S}$ in *Xenopus* oocytes (Ren & Hall, 1997). We were unable to perform experiments on $\alpha 1\text{G}/\beta$ interactions in *Xenopus* oocytes because β subunits stimulate the expression of endogenous oocyte calcium channels (Lacerda *et al.* 1994), making the results uninterpretable. This problem does not arise in COS-7 cells, as endogenous currents were not detected in these cells. It is of interest that a jellyfish $\alpha 1$ subunit has recently been cloned which also has only a rudimentary β binding motif (HMLDDAVKGYLDWINQAS, again with conserved residues in bold, and conserved charges in italics). This channel is also able to express in the absence of β subunits, and co-expression of a β subunit has also been reported to increase expression (Jeziorski *et al.* 1998).

For the HVA channels, all β subunits shift the voltage dependence of activation of the currents to more hyperpolarized potentials, with $\beta 4$ producing the greatest effect for $\alpha 1\text{A}$, and all β subunits producing a similar effect for $\alpha 1\text{E}$ (De Waard & Campbell, 1995; Jones *et al.* 1998). In contrast, the trend in the present study was for β subunits to shift the voltage dependence of activation of $\alpha 1\text{G}$ to more depolarized potentials, although the depolarization of the V_{50} for activation was only statistically significant for $\beta 3$ and $\beta 4$. It remains to be determined whether these subtle effects on the biophysical properties of $\alpha 1\text{G}$ are due to direct interaction with the $\alpha 1\text{G}$ protein.

The effect of all β subunits, except $\beta 2\text{a}$, is to cause an increase in the inactivation rate of HVA channels, and to shift the steady-state inactivation to more hyperpolarized potentials (Stephens *et al.* 1997; Walker & De Waard, 1998), the effect being particularly marked for $\beta 3$ (Jones *et al.* 1998). In contrast, rat $\beta 2\text{a}$, which is palmitoylated, produces an attenuation of inactivation of HVA currents and a shift of steady-state inactivation to more depolarized potentials (De Waard & Campbell, 1995; Costantin *et al.* 1998; Jones *et al.* 1998). No such effects were observed for $\beta 2\text{a}$ on $\alpha 1\text{G}$ currents. This is in agreement with our previous study in which we investigated the effect of overexpression of β subunits on the properties of native T-type currents, expressed in isolation in undifferentiated NG108-15 cells (Wyatt *et al.* 1998). $\beta 2\text{a}$, but not $\beta 1\text{b}$, induced the appearance of a slowly inactivating HVA component of current, but neither had any effect on the biophysical properties of the T-type component. Similarly, it has recently been shown that antisense depletion of β subunits in nodose ganglia had no effect on the biophysical properties of the native T-type currents in these cells (Lambert *et al.* 1997). However, overexpression in *Xenopus* oocytes of either $\beta 2$ or $\beta 4$ slowed inactivation of an endogenous T-type current which had a single channel conductance of 9 pS (Lacerda *et al.* 1994). Part of the effect of $\beta 2$ was attributed in this paper to an effect on channel assembly, stability or trafficking, in agreement with the role suggested here. It is also possible that the single channels observed by Lacerda *et al.* (1994) represent a small conductance mode of HVA channels, with

properties very similar to T-type channels (Meir & Dolphin, 1998).

Conclusion

We have observed a clear effect of $\alpha 2\text{-}\delta$ on the amount of $\alpha 1\text{G}$ channel protein expressed at the plasma membrane, and on the amplitude of the $\alpha 1\text{G}$ current expressed in COS-7 cells and *Xenopus* oocytes. In reciprocal experiments we have also observed that $\alpha 1\text{G}$ increased endogenous $\alpha 2\text{-}\delta$ plasma membrane expression in COS-7 cells. This provides several lines of evidence that $\alpha 1\text{G}$ can interact with $\alpha 2\text{-}\delta$. Nevertheless, from our results and those of others, it appears that $\alpha 1\text{G}$ is able to form functional channels in the absence of co-expressed accessory subunits. However, as *Xenopus* oocytes (Singer-Lahat *et al.* 1992) and most cell lines used for expression studies, including COS-7 cells (present work) and HEK-293 cells (J. Richards and A. C. Dolphin, unpublished results), contain endogenous $\alpha 2\text{-}\delta$ subunits, this requires further study before it can be considered definitive.

- ARMSTRONG, C. M. & MATTESON, D. R. (1985). Two distinct populations of calcium channels in a clonal line of pituitary cells. *Science* **227**, 65–67.
- BERROW, N. S., BRICE, N. L., TEDDER, I., PAGE, K. & DOLPHIN, A. C. (1997). Properties of cloned rat $\alpha 1\text{A}$ calcium channels transiently expressed in the COS-7 cell line. *European Journal of Neuroscience* **9**, 739–748.
- BERROW, N. L., BERROW, N. S., CAMPBELL, V., PAGE, K. M., BRICKLEY, K., TEDDER, I. & DOLPHIN, A. C. (1997). Importance of the different β subunits in the membrane expression of the $\alpha 1\text{A}$ and $\alpha 2$ calcium channel subunits: studies using a depolarisation-sensitive $\alpha 1\text{A}$ antibody. *European Journal of Neuroscience* **9**, 749–759.
- BRICKLEY, K., CAMPBELL, V., BERROW, N., LEACH, R., NORMAN, R. L., WRAY, D., DOLPHIN, A. C. & BALDWIN, S. (1995). Use of site-directed antibodies to probe the topography of the α_2 subunit of voltage-gated Ca^{2+} channels. *FEBS Letters* **364**, 129–133.
- CAMPBELL, V., BERROW, N., BRICKLEY, K., PAGE, K., WADE, R. & DOLPHIN, A. C. (1995a). Voltage-dependent calcium channel β -subunits in combination with $\alpha 1$ subunits have a GTPase activating effect to promote hydrolysis of GTP by $\text{G}\alpha_o$ in rat frontal cortex. *FEBS Letters* **370**, 135–140.
- CAMPBELL, V., BERROW, N. S., FITZGERALD, E. M., BRICKLEY, K. & DOLPHIN, A. C. (1995b). Inhibition of the interaction of G protein G_o with calcium channels by the calcium channel β -subunit in rat neurones. *Journal of Physiology* **485**, 365–372.
- CARBONE, E. & LUX, H. D. (1984). A low voltage-activated fully inactivating Ca channel in vertebrate sensory neurones. *Nature* **310**, 501–502.
- CHEN, C. C. & BEST, P. M. (1998). An antisense oligonucleotide targeting four β subunit genes inhibits the expression of both T and L type calcium currents in cultured atrial myocytes. *Biophysical Journal* **74**, M-Pos350 (abstract).
- CHIEN, A. J., ZHAO, X. L., SHIROKOV, R. E., PURI, T. S., CHANG, C. F., SUN, D., RIOS, E. & HOSEY, M. M. (1995). Roles of a membrane-localized β subunit in the formation and targeting of functional L-type Ca^{2+} channels. *Journal of Biological Chemistry* **270**, 30036–30044.
- COSTANTIN, J., NOCETI, F., QIN, N., WEI, X. Y., BIRNBAUMER, L. & STEFANI, E. (1998). Facilitation by the β_{2a} subunit of pore openings in cardiac Ca^{2+} channels. *Journal of Physiology* **507**, 93–103.
- CRAIG, P. J., BEATTIE, R. E., FOLLY, E. A., BANERJEE, M. D., REEVES, M. B., PRIESTLEY, J. V., CARNEY, S. L., SHER, E., PEREZ-REYES, E. & VOLSSEN, S. G. (1999). Distribution of the voltage-dependent calcium channel $\alpha 1\text{G}$ subunit mRNA and protein throughout the mature rat brain. *European Journal of Neuroscience* (in the Press).
- CRIBBS, L. L., LEE, J.-H., YANG, J., SATIN, J., ZHANG, Y., DAUD, A., BARCLAY, J., WILLIAMSON, M. P., FOX, M., REES, M. & PEREZ-REYES, E. (1998). Cloning and characterization of $\alpha 1\text{H}$ from human heart, a member of the T type Ca^{2+} channel gene family. *Circulation Research* **83**, 103–109.
- DE WAARD, M. & CAMPBELL, K. P. (1995). Subunit regulation of the neuronal α_{1A} Ca^{2+} channel expressed in *Xenopus* oocytes. *Journal of Physiology* **485**, 619–634.
- DE WAARD, M., SCOTT, V. E. S., PRAGNELL, M. & CAMPBELL, K. P. (1996). Identification of critical amino acids involved in $\alpha_1\text{-}\beta$ interaction in voltage-dependent Ca^{2+} channels. *FEBS Letters* **380**, 272–276.
- FELIX, R., GURNETT, C. A., DE WAARD, M. & CAMPBELL, K. P. (1997). Dissection of functional domains of the voltage-dependent Ca^{2+} channel $\alpha 2\delta$ subunit. *Journal of Neuroscience* **17**, 6884–6891.
- GAO, T. Y., CHIEN, A. J. & HOSEY, M. M. (1999). Complexes of the $\alpha 1\text{C}$ and β subunits generate the necessary signal for membrane targeting of class C L-type calcium channels. *Journal of Biological Chemistry* **274**, 2137–2144.
- GURNETT, C. A., DE WAARD, M. & CAMPBELL, K. P. (1996). Dual function of the voltage-dependent Ca^{2+} channel $\alpha_2\delta$ subunit in current stimulation and subunit interaction. *Neuron* **16**, 431–440.
- GURNETT, C. A., FELIX, R. & CAMPBELL, K. P. (1997). Extracellular interaction of the voltage-dependent Ca^{2+} channel $\alpha_2\delta$ and α_1 subunits. *Journal of Biological Chemistry* **272**, 18508–18512.
- HUGUENARD, J. R. (1996). Low-threshold calcium currents in central nervous system neurons. *Annual Review of Physiology* **58**, 329–348.
- JEZIORSKI, M. C., GREENBERG, R. M., CLARK, K. S. & ANDERSON, P. A. V. (1998). Cloning and functional expression of a voltage-gated calcium channel α_1 subunit from jellyfish. *Journal of Biological Chemistry* **273**, 22792–22799.
- JONES, L. P., WEI, S. K. & YUE, D. T. (1998). Mechanism of auxiliary subunit modulation of neuronal α_{1E} calcium channels. *Journal of General Physiology* **112**, 125–143.
- KLUGBAUER, N., LACINOVA, L., MARAIS, E., HOBOM, M. & HOFMANN, F. (1999). Molecular diversity of the calcium channel $\alpha 2\text{-}\delta$ subunit. *Journal of Neuroscience* **19**, 684–691.
- KOBRINSKY, E. M., PEARSON, H. A. & DOLPHIN, A. C. (1994). Low- and high-voltage-activated calcium channel currents and their modulation in the dorsal root ganglion cell line ND7-23. *Neuroscience* **58**, 539–552.
- LACERDA, A. E., PEREZ-REYES, E., WEI, X., CASTELLANO, A. & BROWN, A. M. (1994). T-type and N-type calcium channels of *Xenopus* oocytes: Evidence for specific interactions with β subunits. *Biophysical Journal* **66**, 1833–1843.
- LAMBERT, R. C., MCKENNA, F., MAULET, Y., TALLEY, E. M., BAYLISS, D. A., CRIBBS, L. L., LEE, J.-H., PEREZ-REYES, E. & FELTZ, A. (1998). Low-voltage-activated Ca^{2+} currents are generated by members of the Ca_vT subunit family ($\alpha 1\text{G}/\text{H}$) in rat primary sensory neurons. *Journal of Neuroscience* **18**, 8605–8613.

- LAMBERT, R. C., MAULET, Y., MOUTON, J., BEATTIE, R., VOLSEN, S., DE WAARD, M. & FELTZ, A. (1997). T-type Ca^{2+} current properties are not modified by Ca^{2+} channel β subunit depletion in nodose ganglion neurons. *Journal of Neuroscience* **17**, 6621–6628.
- LEE, J.-H., DAUD, A. N., CRIBBS, L. L., LACERDA, A. E., PEREVERZEV, A., KLOCKNER, U., SCHNEIDER, T. & PEREZ-REYES, E. (1999). Cloning and expression of a novel member of the low voltage activated T type calcium channel family. *Journal of Neuroscience* **19**, 1912–1921.
- LEONARD, J. P. & SNUTCH, T. P. (1991). The expression of neurotransmitter receptors and ion channels in *Xenopus* oocytes. IRL Press, Oxford.
- MEIR, A. & DOLPHIN, A. C. (1998). Known calcium channel $\alpha 1$ subunits can form low threshold, small conductance channels, with similarities to native T type channels. *Neuron* **20**, 341–351.
- NOWYCKY, M. C., FOX, A. P. & TSIEN, R. W. (1985). Three types of neuronal calcium channel with different calcium agonist sensitivity. *Nature* **316**, 440–446.
- PEREZ-REYES, E., CRIBBS, L. L., DAUD, A., LACERDA, A. E., BARCLAY, J., WILLIAMSON, M. P., FOX, M., REES, M. & LEE, J.-H. (1998). Molecular characterization of a neuronal low-voltage-activated T type calcium channel. *Nature* **391**, 896–900.
- QIN, N., OLCESE, R., STEFANI, E. & BIRNBAUMER, L. (1998). Modulation of human neuronal α_{1E} -type calcium channel by $\alpha_2\delta$ -subunit. *American Journal of Physiology* **274**, C1324–1331.
- REN, D. & HALL, L. M. (1997). Functional expression and characterization of skeletal muscle dihydropyridine receptors in *Xenopus* oocytes. *Journal of Biological Chemistry* **272**, 22393–22396.
- SCHREIBMAYER, W., LESTER, H. A. & DASCAL, N. (1994). Voltage clamping of *Xenopus laevis* oocytes utilizing agarose cushion electrodes. *Pflügers Archiv* **426**, 453–458.
- SHISTIK, E., IVANINA, T., PURI, T., HOSEY, M. & DASCAL, N. (1995). Ca^{2+} current enhancement by $\alpha 2/\delta$ and β subunits in *Xenopus* oocytes: Contribution of changes in channel gating and $\alpha 1$ protein level. *Journal of Physiology* **489**, 55–62.
- SINGER-LAHAT, D., LOTAN, I., ITAGAKI, K., SCHWARTZ, A. & DASCAL, N. (1992). Evidence for the existence of RNA of Ca^{2+} -channel α_2/δ subunit in *Xenopus* oocytes. *Biochimica et Biophysica Acta* **1137**, 39–44.
- STEPHENS, G. J., PAGE, K. M., BURLEY, J. R., BERROW, N. S. & DOLPHIN, A. C. (1997). Functional expression of rat brain cloned $\alpha 1E$ calcium channels in COS-7 cells. *Pflügers Archiv* **433**, 523–532.
- SWICK, A. G., JANICOT, M., CHENEVAL-KASTELIC, T., MCLLENITHAN, J. C. & LANE, D. M. (1992). Promoter-cDNA-directed heterologous protein expression in *Xenopus laevis* oocytes. *Proceedings of the National Academy of Sciences of the USA* **89**, 1812–1816.
- TAREILUS, E., ROUX, M., QIN, N., OLCESE, R., ZHOU, J. M., STEFANI, E. & BIRNBAUMER, L. (1997). A *Xenopus* oocyte β subunit: Evidence for a role in the assembly/expression of voltage-gated calcium channels that is separate from its role as a regulatory subunit. *Proceedings of the National Academy of Sciences of the USA* **94**, 1703–1708.
- WALKER, D., BICHET, D., CAMPBELL, K. P. & DE WAARD, M. (1998). A β_4 isoform-specific interaction site in the carboxyl-terminal region of the voltage-dependent Ca^{2+} channel α_{1A} subunit. *Journal of Biological Chemistry* **273**, 2361–2367.
- WALKER, D. & DE WAARD, M. (1998). Subunit interaction sites in voltage-dependent Ca^{2+} channels. *Trends in Neurosciences* **21**, 148–154.
- WYATT, C. N., PAGE, K. M., BERROW, N. S., BRICE, N. L. & DOLPHIN, A. C. (1998). The effect of overexpression of auxiliary calcium channel subunits on native Ca^{2+} channel currents in undifferentiated NG108-15 cells. *Journal of Physiology* **510**, 347–360.

Acknowledgements

We thank Professor C. Hopkins of the Laboratory of Molecular Cell Biology, University College London for access to the confocal microscope. This work was supported by a Wellcome Trust grant to A.C.D. and an NIH grant (HL57828) to E.P.R. We thank N. Balaguer for technical support.

Corresponding author

A. C. Dolphin: Department of Pharmacology, University College London, Gower Street, WC1E 6BT, UK.

Email: a.dolphin@ucl.ac.uk

Authors' present addresses

J. Richards: Department of Biochemistry, Imperial College, London, UK.

C. N. Wyatt: Department of Physiology, Oxford University, UK.