

## Subcellular mechanisms of the positive inotropic effect of angiotensin II in cat myocardium

Martín G. Vila Petroff, Ernesto A. Aiello, Julieta Palomeque, Margarita A. Salas and Alicia Mattiazzi

*Centro de Investigaciones Cardiovasculares, Facultad de Ciencias Médicas, Universidad Nacional de La Plata, La Plata 1900, Argentina*

(Received 21 February 2000; accepted after revision 2 August 2000)

1. Cat ventricular myocytes loaded with  $[Ca^{2+}]_i$ - and  $pH_i$ -sensitive probes were used to examine the subcellular mechanism(s) of the Ang II-induced positive inotropic effect. Ang II ( $1 \mu M$ ) produced parallel increases in contraction and  $Ca^{2+}$  transient amplitudes and a slowly developing intracellular alkalinisation. Maximal increases in contraction amplitude and  $Ca^{2+}$  transient amplitude were  $163 \pm 22$  and  $43 \pm 8\%$ , respectively, and occurred between 5 and 7 min after Ang II administration, whereas  $pH_i$  increase ( $0.06 \pm 0.03$  pH units) became significant only 15 min after the addition of Ang II. Furthermore, the inotropic effect of Ang II was preserved in the presence of  $Na^+-H^+$  exchanger blockade. These results indicate that the positive inotropic effect of Ang II is independent of changes in  $pH_i$ .
2. Similar increases in contractility produced by either elevating extracellular  $[Ca^{2+}]_o$  or by Ang II application produced similar increases in peak systolic  $Ca^{2+}$  indicating that an increase in myofilament responsiveness to  $Ca^{2+}$  does not participate in the Ang II-induced positive inotropic effect.
3. Ang II significantly increased the L-type  $Ca^{2+}$  current, as assessed by using the perforated patch-clamp technique (peak current recorded at 0 mV:  $-1.88 \pm 0.16$  pA  $pF^{-1}$  in control vs.  $-3.03 \pm 0.20$  pA  $pF^{-1}$  after 6–8 min of administration of Ang II to the bath solution).
4. The positive inotropic effect of Ang II was not modified in the presence of either KB-R7943, a specific blocker of the  $Na^+-Ca^{2+}$  exchanger, or ryanodine plus thapsigargin, used to block the sarcoplasmic reticulum function.
5. The above results allow us to conclude that in the cat ventricle the Ang II-induced positive inotropic effect is due to an increase in the intracellular  $Ca^{2+}$  transient, an enhancement of the L-type  $Ca^{2+}$  current being the dominant mechanism underlying this increase.

In the last few years several laboratories have explored the subcellular mechanisms of the positive inotropic effect of angiotensin II (Ang II). The outcome of this work turned out to be largely controversial. Although part of the controversy might arise from species differences (Ishihata & Endoh, 1995), opposite results were also reported in the same species. Experiments by Ikenouchi *et al.* (1994) in isolated rabbit myocytes indicated that the positive inotropic effect of Ang II was exclusively mediated by an increase in the myofilament responsiveness to  $Ca^{2+}$ . In this study, no increase in the intracellular  $Ca^{2+}$  transients was detected. These results are in sharp contrast to those obtained later, showing that Ang II did increase the intracellular  $Ca^{2+}$  transient in rabbit ventricular myocytes (Skolnick *et al.* 1998). Results from Watanabe & Endoh (1998), also in rabbit heart, suggested that the positive inotropic effect of Ang II was due to an increase in both  $[Ca^{2+}]_i$  and myofilament responsiveness to  $Ca^{2+}$ .

The subcellular mechanisms responsible for the described increase in either  $[Ca^{2+}]_i$  or the myofilament responsiveness to  $Ca^{2+}$  are also elusive. The L-type  $Ca^{2+}$  current ( $I_{Ca}$ ) has been shown to be either increased (Kass & Blair, 1981; Allen *et al.* 1988; Kaibara *et al.* 1994), decreased (Habuchi *et al.* 1995) or unchanged (Ikenouchi *et al.* 1994; Ai *et al.* 1998). Moreover, whereas Kaibara *et al.* (1994) concluded that the enhancement in  $I_{Ca}$  was due to the increase in  $pH_i$ , Ikenouchi *et al.* (1994) described in the same species an increase in  $pH_i$  without significant changes in  $I_{Ca}$ . Using pharmacological tools, Talukder & Endoh (1997) concluded that the influx of  $Ca^{2+}$  through the L-type  $Ca^{2+}$  channels was one of the main mechanisms by which Ang II increases  $[Ca^{2+}]_i$ . However more recent experiments of the same group, working with the same species, suggested that the  $Na^+-Ca^{2+}$  exchanger was the mechanism responsible for the Ang II-induced increase in  $[Ca^{2+}]_i$  (Fujita & Endoh, 1999).

Referring to the Ang II-induced increase in myofilament responsiveness to  $\text{Ca}^{2+}$ , the  $\text{pH}_i$  dependence of this mechanism (Ikenouchi *et al.* 1994) has been challenged by the finding that the positive inotropic effect of Ang II was also present in a physiological buffer ( $\text{HCO}_3^-/\text{CO}_2$ ) in which Ang II does not produce significant changes in  $\text{pH}_i$  (Mattiazzi *et al.* 1997).

Considering the discrepant results summarised above, the present study was directed to examining in the cat ventricle the effect of Ang II on the fundamental events of excitation–contraction coupling, i.e. calcium transients, calcium currents, SR  $\text{Ca}^{2+}$  release and  $\text{Na}^+ - \text{Ca}^{2+}$  exchange, in an attempt to comprehensively answer the following questions: Is the positive inotropic effect of Ang II due to an increase in  $[\text{Ca}^{2+}]_i$  or to an increase in myofilament responsiveness to  $\text{Ca}^{2+}$ ? In any of these cases, what is (or are) the underlying mechanism(s) involved?

## METHODS

### Myocyte isolation

All experiments were performed in accordance with the guidelines for Animal Care of the Scientific Committee of the University of La Plata School of Medicine. Cats were anaesthetised by intraperitoneal injection of sodium pentobarbitone (35 mg (kg body weight)<sup>-1</sup>). Myocytes were isolated according to the technique previously described (Aiello *et al.* 1998; Morgan *et al.* 1999) with some modifications. Briefly, the hearts were attached via the aorta to a cannula, excised and mounted in a Langendorff apparatus. They were then retrogradely perfused at 37 °C at a constant perfusion pressure of 70–80 mmHg with Krebs-Henseleit (K-H) solution of the following composition (mM): 146.2 NaCl, 4.7 KCl, 1.35  $\text{CaCl}_2$ , 10.0 Hepes, 0.35  $\text{NaH}_2\text{PO}_4$ , 1.05  $\text{MgSO}_4$ , 10.0 glucose (pH adjusted to 7.4 with NaOH). The solution was continuously bubbled with 100%  $\text{O}_2$ . After a stabilisation period of 4 min, the perfusion was switched to a nominally  $\text{Ca}^{2+}$ -free K-H solution for 6 min. Hearts were then recirculated with collagenase (118 U ml<sup>-1</sup>), 0.1 mg ml<sup>-1</sup> pronase and 1% bovine serum albumin (BSA), in K-H solution containing 50  $\mu\text{M}$   $\text{CaCl}_2$ . Perfusion continued until the hearts became flaccid (15–25 min). They were then removed from the perfusion apparatus by cutting at the atria-ventricular junction. The desegregated myocytes were separated from the undigested tissue and rinsed several times with a K-H solution containing 1% BSA and 500  $\mu\text{M}$   $\text{CaCl}_2$ . After each wash, myocytes were left for sedimentation for 10 min. Myocytes were kept in K-H solution at room temperature (20–22 °C) until use. Only rod-shaped myocytes with clear and distinct striations and an obvious marked shortening and relaxation on stimulation were used. Experiments were performed at room temperature.

### Indo-1 fluorescence and cell shortening measurements

The isolated myocytes were loaded at room temperature with the cell-permeant acetoxymethyl ester (AM) form of indo-1 (17  $\mu\text{M}$  for 9 min) according to the bulk method described by Spurgeon *et al.* (1990) and left for de-esterification for 45 min. Cells were then placed on the stage of an inverted microscope (Nikon Diaphot 200) adapted for epifluorescence. Myocytes were continuously superfused with K-H solution (pH 7.4) at a constant flow of 1 ml min<sup>-1</sup> and field stimulated via two platinum electrodes on either side of the bath (square waves, 2 ms duration and 20% above threshold) at a constant frequency of 30 beats min<sup>-1</sup>. The excitation light was

centred at 350 nm and the fluorescence emitted by the cell was recorded at 410 and 490 nm. Background fluorescence was subtracted from each signal before obtaining the 410 to 490 nm fluorescence ratio. The diastolic fluorescence ratio was measured as the mean value over a 100 ms period after the twitch was completed. Systolic fluorescence ratio was determined directly from the peak of the recorded ratio.

The stage of the microscope was illuminated with red light (640–750 nm) through its normal bright-field illumination optics to allow simultaneous measurements of fluorescence and shortening. Resting cell length and cell shortening were measured by a video-based motion detector (Crescent Electronics, Salt Lake City, UT, USA) and stored by software for an off-line analysis. Cell shortening was simultaneously monitored on a two-channel chart recorder (Gould RS 3200).

Indo-1-loaded myocytes were used to assess myofilament responsiveness to  $\text{Ca}^{2+}$  according to the previously described method (Shah *et al.* 1994). In brief, this method consists in plotting the instantaneous myocyte length *versus* simultaneously measured indo-1 fluorescence during single twitch contractions (phase-plane plot). In mechanically unloaded cells stimulated to contract from slack length, a dynamic equilibrium is achieved between cell relengthening and the decline in cytosolic  $\text{Ca}^{2+}$  during twitch relaxation. This implies that for interventions that increase contractility without changing myofilament responsiveness to  $\text{Ca}^{2+}$  (e.g. an increase in extracellular calcium,  $[\text{Ca}^{2+}]_o$ ) the phase-plane plots describe a common trajectory during the relaxation phase. In contrast, perturbations that decrease myofilament responsiveness to  $\text{Ca}^{2+}$  (e.g. acidosis) shift the relaxation trajectory downward and rightward whereas interventions that increase myofilament responsiveness to  $\text{Ca}^{2+}$  (e.g.  $\text{Ca}^{2+}$ -sensitizing drugs) have the opposite effect. The position of the trajectory of the relaxation phase reflects therefore the relative myofilament responsiveness to  $\text{Ca}^{2+}$ .

### $\text{pH}_i$ measurements

After enzymatic isolation, myocytes were loaded with the membrane-permeant acetoxymethyl ester form of the fluorescent  $\text{H}^+$ -sensitive indicator SNARF-1 AM. Cell suspensions (2 ml) were exposed to a final concentration of 4  $\mu\text{M}$  SNARF-1 AM and 0.6% v/v DMSO. After 10 min, the myocytes were gently centrifuged for 2 min and resuspended in Hepes buffer and stored at room temperature until use.  $\text{pH}_i$  and cell length were monitored on the stage of a modified inverted microscope, as previously described (Blank *et al.* 1992). After excitation at  $530 \pm 5$  nm, the ratio of SNARF-1 AM emission at  $590 \pm 5$  nm to that of  $640 \pm 5$  nm was used as a measure of  $\text{pH}_i$  according to an *in vivo* calibration, obtained from SNARF-1 AM-loaded myocytes exposed to solutions of varying pH values containing 140 mM KCl, 20  $\mu\text{M}$  nigericin, 1  $\mu\text{M}$  valinomycin and 1  $\mu\text{M}$  carbonyl cyanide *p*-(trifluoromethoxy)-phenylhydrazone at room temperature.

To determine  $\text{Na}^+ - \text{H}^+$  exchanger activity, intracellular acidosis was induced by the wash-out (10 min) of a transient (3 min) exposure of cells to 15 mM  $\text{NH}_4\text{Cl}$  (first acid pulse). This procedure was repeated after either a 5 or a 20 min exposure to Ang II (second acid pulse). The rate of  $\text{pH}_i$  recovery obtained by fitting a linear function to the initial recovery from acidosis was used to indicate changes in  $\text{Na}^+ - \text{H}^+$  exchanger activity (Vandenberg *et al.* 1994). The reproducibility of  $\text{pH}_i$  recovery after two consecutive  $\text{NH}_4\text{Cl}$  pulses was established in control experiments.

### Perforated-patch recordings

Isolated cat ventricular myocytes were placed in a perfusion chamber and superfused with bath solution at a flow rate of

1.5 ml min<sup>-1</sup>. The perforated-patch configuration of the patch-clamp technique (Korn *et al.* 1991; Aiello *et al.* 1998) was used for voltage-clamp recordings with a patch-clamp amplifier (Axopatch 200A, Axon Instruments, Foster City, CA, USA). Patch pipettes were pulled with a PP-83 puller (Narishige, Tokyo, Japan) and fire-polished with a MF-83 Microforge (Narishige) to a final resistance of 1–3 M $\Omega$  when filled with pipette solution. The tip of the pipette was positioned above the cell, and its potential and capacitance were nullified. Whole-cell currents (filtered at 1 kHz) were digitally recorded directly to hard disk via an analog-to-digital converter (Digidata 1200, Axon Instruments) interfaced with an IBM clone computer running pCLAMP and Axotape software (Axon Instruments). Data analysis was performed with pCLAMP (Clampfit, Axon Instruments).

Voltage-clamp depolarising pulses (250 ms) were delivered at 0.2 Hz. A holding potential of -80 mV was used in all protocols to prevent slow inactivation and to minimise current rundown (McDonald *et al.* 1994). A 500 ms prepulse to -40 mV, used to inactivate sodium channels and potential T-type calcium channels, preceded the depolarizing test pulses to different potentials. Under the present recording conditions, no whole-cell currents were detected in the absence of Ca<sub>o</sub><sup>2+</sup> (not shown). Nystatin produced good intracellular access after 15–20 min of seal formation. The calcium current amplitude was measured as peak inward current with reference to the current measured at the end of the test pulse. For each cell, capacitive current was recorded to determine the membrane capacitance and the currents were normalised for cell capacitance. The average cell capacitance was 148.7  $\pm$  7.7 pF ( $n$  = 14).

In some experiments the perforated-patch configuration of the patch-clamp technique was used in combination with simultaneous measurements of cell length performed as described above. In these experimental groups the Ang II-induced increase in cell shortening was lower than in the other groups presented in this paper. The reason for this difference is not apparent to us. One possible cause might be the different experimental conditions used, i.e. perforated patch *vs.* intact myocytes.

The superfusion medium used to measure  $I_{Ca}$  had the following composition (mM): CsCl 5, NaCl 133, MgCl<sub>2</sub> 1, MgSO<sub>4</sub> 1.2, Hepes 10, tetraethylammonium chloride (TEA) 10, CaCl<sub>2</sub> 1.35 and glucose 10; pH was adjusted to 7.4 with NaOH. The internal (pipette) solution contained (mM): CsCl 140, MgCl<sub>2</sub> 1, NaCl 10, EGTA 1, Hepes 10 and nystatin 0.4 mg ml<sup>-1</sup>; pH was adjusted to 7.2 with NaOH. The superfusion medium used to measure action potentials and cell length had the following composition (mM): NaCl 138, MgCl<sub>2</sub> 1, MgSO<sub>4</sub> 1.2, Hepes 10, CaCl<sub>2</sub> 1.35 and glucose 10; pH was adjusted to 7.4 with NaOH. The pipette solution used in these experiments contained (mM): KCl 140, MgCl<sub>2</sub> 1, NaCl 10, EGTA 1, Hepes 10 and nystatin 0.4 mg ml<sup>-1</sup>; pH was adjusted to 7.2 with NaOH.

### Materials

Collagenase type B was purchased from Worthington Biochemical Corp. (Lakewood, NJ, USA); pronase from Boehringer Mannheim Corp. GmbH (Mannheim, Germany); BSA (essentially fatty acid free), thapsigargin and angiotensin II from Sigma Chemical Co. (St Louis, MO, USA); indo-1 AM and SNARF-1 AM from Molecular Probes Inc. (Eugene, OR, USA); and KB-R7943 (2-[2-[4-(4-nitrobenzyloxy)phenyl]ethyl]isothiourea methanesulphonate) from Kanebo (Osaka, Japan). All other chemicals were of the purest reagent grade available.

### Statistics

All data are presented as means  $\pm$  s.e.m. Comparisons within groups were assessed by Student's paired or unpaired *t* test as

appropriate. ANOVA was used when required as indicated in the text. A value of  $P < 0.05$  was taken to indicate statistical significance.

## RESULTS

### Effect of Ang II on contraction and pH<sub>i</sub>

The effect of 1  $\mu$ M Ang II on contraction and pH<sub>i</sub> was assessed in electrically stimulated cardiac myocytes. This concentration of Ang II was chosen based on the results of pilot experiments which indicated that the maximal positive inotropic effect of Ang II occurred in the concentration range of 0.5–1  $\mu$ M, in agreement with previous reports (Koch-Weser, 1965; Ishihata & Endoh, 1993; Mattiazzi *et al.* 1997).

Figure 1A shows a representative example of the effect of Ang II on the unloaded myocyte contraction. Ang II induced a positive inotropic response, consisting in an initial increase in contraction that peaked between 5 and 7 min, followed by a slow decay that reached steady state after 20 min of incubation with the drug. The middle panel shows the individual twitch contractions in control and at 5 and 20 min after the addition of Ang II. The lower panel of Fig. 1A depicts a typical tracing of the effect of Ang II on pH<sub>i</sub> monitored continuously during 20 min. Note that 5 min after Ang II administration the positive inotropic effect had reached its maximum whereas pH<sub>i</sub> was not affected.

Figure 1B shows the overall results of these experiments indicating the time course of the effect of Ang II on contraction amplitude and pH<sub>i</sub>. Ang II induced a maximal increase in contraction amplitude of 163  $\pm$  22% before an increase in pH<sub>i</sub> became significant. The increase in pH<sub>i</sub> became statistically significant only after 15 min of administration of the drug reaching a maximal value of 0.06  $\pm$  0.03 pH units after 20 min. At this time the positive inotropic effect of Ang II diminished from its peak value to 129  $\pm$  14% of control.

The increase in pH<sub>i</sub> produced by Ang II in Hepes buffer is due to the activation of the Na<sup>+</sup>-H<sup>+</sup> exchanger (Matsui *et al.* 1995; Camili3n de Hurtado *et al.* 1998; Gunasegaram *et al.* 1999). To further explore the possible dependence of the Ang II-induced positive inotropic effect on the intracellular alkalisation two additional protocols were performed. (1) We examined, using the NH<sub>4</sub>Cl pulse procedure (see Methods), the activity of the Na<sup>+</sup>-H<sup>+</sup> exchanger after 5 min of Ang II application, when the positive contractile response had reached its maximum level, and after 20 min, when the contractile response to Ang II had reached steady state. Figure 2A shows representative pH<sub>i</sub> recordings of two consecutive NH<sub>4</sub>Cl pulses that were superimposed for comparison. The profile of the second pulse, obtained after a 5 min incubation period with Ang II, was similar to that of the first (control) pulse, indicating that after 5 min, Ang II had still not produced activation of the Na<sup>+</sup>-H<sup>+</sup> exchanger. Furthermore, there was no significant difference in the rate


of  $\text{pH}_i$  recovery from acidosis between the first (control) pulse ( $\Delta\text{pH}_i/\Delta t$ :  $0.082 \pm 0.02$  pH units  $\text{min}^{-1}$ ) and the second (Ang II 5 min) pulse ( $0.084 \pm 0.03$  pH units  $\text{min}^{-1}$ ). However, when the cells were pretreated with Ang II for 20 min prior to the second pulse,  $\text{pH}_i$  recovery from acidosis was accelerated (Fig. 2B) such that the rate of  $\text{pH}_i$  recovery from acidosis was significantly increased from  $0.075 \pm 0.02$  pH units  $\text{min}^{-1}$  in the first (control) pulse to  $0.11 \pm 0.01$  pH units  $\text{min}^{-1}$  in the second (Ang II 20 min) pulse. (2) The effect of Ang II on contraction amplitude was examined in the presence and in the absence of the  $\text{Na}^+ - \text{H}^+$  exchanger blocker HOE 642 (HOE) (10 and  $50 \mu\text{M}$ ). Preliminary experiments showed that  $10 \mu\text{M}$  HOE produced a significant inhibition of the rate of  $\text{pH}_i$  recovery from an acid load (results not shown). This indicates that the concentrations of HOE used were enough to inhibit the activity of the  $\text{Na}^+ - \text{H}^+$  exchanger. HOE did not produce significant changes in basal contractility values. Administration of HOE up to  $50 \mu\text{M}$  had no effect when it was applied after the Ang II-induced positive inotropic effect had reached steady state ( $n = 9$ )

(Fig. 3A). Similarly, preincubation of the cells with  $10 \mu\text{M}$  of the blocker did not prevent the positive inotropic effect of Ang II ( $n = 5$ ) (Fig. 3B). These findings in isolated myocytes are in complete agreement with a previously published report from our laboratory in multicellular preparations showing that in feline myocardium the positive inotropic and alkalinising effect of Ang II are not linked phenomena (Mattiuzzi *et al.* 1997).

Since the Ang II-induced positive inotropic effect cannot be attributed to an increase in myofilament responsiveness to  $\text{Ca}^{2+}$  due to intracellular alkalinisation, alternative mechanisms that may be responsible for the Ang II-induced positive inotropic effect were examined, namely an increase in the intracellular  $\text{Ca}^{2+}$  transient and/or a  $\text{pH}_i$ -independent increase in the myofilament responsiveness to  $\text{Ca}^{2+}$ .


#### Effect of Ang II on myocyte contraction and intracellular calcium transient

Using indo-1-loaded cardiac myocytes we investigated the effect of  $1 \mu\text{M}$  Ang II on contraction and the intracellular  $\text{Ca}^{2+}$  transient ( $\text{Ca}_i\text{T}$ ). A representative example of the



**Figure 1. Effect of Ang II on contraction and  $\text{pH}_i$**

A, upper panel, a typical continuous recording of cell length showing the time course of the Ang II-induced positive inotropic effect. Middle panel, individual twitch contractions at the times indicated by the letters a-c on the continuous chart recording. Lower panel, effect of Ang II on  $\text{pH}_i$ . Ang II produced a slow increase in  $\text{pH}_i$ . B, the overall results of these experiments ( $n = 9$ ) indicating the temporal dissociation between the positive inotropic and alkalinising effect of Ang II. The Ang II-induced positive inotropic effect peaked before any significant change in  $\text{pH}_i$  ( $\Delta\text{pH}_i$ ) could be detected. The increase in  $\text{pH}_i$  became significant when the positive inotropic effect of Ang II was already fading. Data points are means of five successive  $\text{pH}_i$  and contraction amplitude measurements (sampling rate 0.5 Hz) from each experiment at every given time. Basal  $\text{pH}_i$  value before Ang II administration was  $7.31 \pm 0.05$ .




**Figure 2. Effect of Ang II on the pH<sub>i</sub> recovery from intracellular acidosis**

Representative pH<sub>i</sub> recordings obtained during two consecutive 15 mM NH<sub>4</sub>Cl pulses. ○, first pulse; ●, second pulse performed after either 5 min (A) or 20 min (B) of exposure of cells to Ang II. The 5 min incubation period with Ang II failed to affect pH<sub>i</sub> recovery (*n* = 4) whereas a 20 min exposure to Ang II significantly accelerated recovery from acidosis (*n* = 5).

Ang II-induced positive inotropic effect and the associated Ca<sub>i</sub>T are depicted in Fig. 4A. The initial increase in contraction amplitude elicited by Ang II followed by the slow decay was closely associated with a similar pattern of rise and slow decay in the Ca<sub>i</sub>T amplitude and peak systolic

[Ca<sup>2+</sup>]<sub>i</sub>. Neither diastolic cell length nor diastolic [Ca<sup>2+</sup>]<sub>i</sub> levels were significantly affected by Ang II. In some cells, the initial increase in contraction amplitude was preceded by a transient negative inotropic effect that was associated with a decrease in the Ca<sub>i</sub>T. Figure 4B shows the overall



**Figure 3. Failure of the Na<sup>+</sup>-H<sup>+</sup> inhibitor HOE 642 to block the Ang II-induced positive inotropic effect**


A, representative example of the effects of HOE added in the continued presence of Ang II. The sustained positive inotropic effect of Ang II was not affected upon addition of 50 μM HOE. B, overall results of the effect of Ang II alone and Ang II in the continued presence of HOE (10 μM) on contraction amplitude, expressed as a percentage of the control value. No significant difference in the positive inotropic effect elicited by Ang II was observed between control and HOE-treated cells.

results of these experiments ( $n = 5$ ) indicating the time course of the effect of Ang II on contraction and  $\text{Ca}_i\text{T}$  amplitude. Similar results were obtained in four other cells using a lower Ang II concentration ( $0.5 \mu\text{M}$ ) (data not shown). Considering the similar time course of the effect of Ang II on contraction and  $\text{Ca}_i\text{T}$  amplitude it seems reasonable to suggest that the Ang II-induced increase in  $[\text{Ca}^{2+}]_i$  constitutes the dominant mechanism for the positive inotropic effect observed.

### Ang II and myofilament response to $\text{Ca}^{2+}$

To assess whether a  $\text{pH}_i$ -independent increase in the myofilament responsiveness to calcium also contributes to the Ang II-induced positive inotropic effect, two different

approaches were used. (1) Phase plane diagrams (loops) of the instantaneous cell length *versus* the simultaneous indo-1 fluorescence (see Methods) at control and after 10 and 20 min of Ang II application (peak and steady state of the Ang II positive inotropic effect, respectively) were compared as illustrated in Fig. 4A, right panel. The relaxation phase of the loops obtained from control twitches, and after 10 and 20 min of Ang II application have a common trajectory indicating that the response of the myofilaments to  $\text{Ca}^{2+}$  was not changed after Ang II administration. Similar phase-plane diagrams were observed in five indo-1-loaded cells exposed to Ang II ( $1 \mu\text{M}$ ). (2) The second approach consisted in comparing the increase in peak systolic  $\text{Ca}^{2+}$  induced by elevating  $[\text{Ca}^{2+}]_o$  with the increase in peak systolic  $\text{Ca}^{2+}$



**Figure 4.** Effect of Ang II on indo-1 transient and contraction amplitudes

A, tracings of indo-1 fluorescence and cell length (lower panels) obtained at the times indicated by the letters in the continuous length recording (upper panel). The control condition (a) is followed by a transient decrease in contraction amplitude and in indo-1 fluorescence transient amplitude (b) and subsequently by a positive inotropic response that peaked and then decreased towards a steady state value. Trace c depicts the peak increase in contraction amplitude and trace d, the steady state contraction reached after 20 min of Ang II administration. These effects are associated with similar changes in indo-1 transient amplitude. On the right are the diagrams of indo-1 fluorescence *vs.* cell length (phase-plane plots) for contractions a, c and d, used to assess the myofilament responsiveness to  $\text{Ca}^{2+}$ . The common trajectory during myocyte relaxation between the control and Ang II loops reflects the lack of effect of Ang II on myofilament responsiveness to  $\text{Ca}^{2+}$ . B, the overall results of the time course of the effect of Ang II on contraction and indo-1 transient amplitude expressed as a percentage of the control value. Data are means  $\pm$  s.e.m. of 5 cells. Both the Ang II-induced increase in contraction amplitude and the indo-1 transient increase and decay in parallel.


evoked by Ang II ( $1 \mu\text{M}$ ). Special care was taken to ensure a similar increase in contraction amplitude induced by either high  $[\text{Ca}^{2+}]_o$  or Ang II. Figure 5 shows a representative example obtained following this protocol. For a similar increase in contraction amplitude induced by either high  $[\text{Ca}^{2+}]_o$  or Ang II there was also a similar increase in peak systolic  $\text{Ca}^{2+}$ . The mean effects ( $n = 8$ ) of either elevating  $[\text{Ca}^{2+}]_o$  (ranging from 3 to 5 mM) or Ang II ( $1 \mu\text{M}$ ) administration on contraction and  $\text{Ca}_i\text{T}$  amplitude and kinetics are provided in Fig. 6A and in Table 1. Figure 6B compares the relationship between the increase in peak systolic indo-1 fluorescence and peak shortening obtained after 10 and 20 min of Ang II and after increasing  $[\text{Ca}^{2+}]_o$ . The slopes of the lines fitted to the three different set of points were not significantly different (analysis of covariance). Taken together, these results are consistent with a lack of effect of Ang II on myofilament responsiveness to  $\text{Ca}^{2+}$  all along the time course of the positive inotropic effect of Ang II.

Since Ang II enhances myocardial contractility by an increase in the  $\text{Ca}_i\text{T}$  without a significant modification of myofilament responsiveness to  $\text{Ca}^{2+}$ , we next explored the possible mechanisms by which this increase in the  $\text{Ca}_i\text{T}$  may occur.

#### Does $\text{Ca}^{2+}$ entry increase due to a prolongation of action potential duration?

Cardiac action potential duration (APD) lengthening after Ang II application has been previously reported (Kass &

Blair, 1981; Morita *et al.* 1995). Prolongation of APD could favour  $\text{Ca}^{2+}$  influx through L-type  $\text{Ca}^{2+}$  channels and/or the reverse mode of the  $\text{Na}^+ - \text{Ca}^{2+}$  exchanger. In both cases, the prolongation of the APD would contribute to the increase in  $\text{Ca}^{2+}$  influx and therefore to the positive inotropic effect of Ang II. Conversely, prolongation of the APD could be the consequence of an increase in  $I_{\text{Ca}}$ . If this were the case, the prolongation of APD *per se* might not contribute to the increase in calcium influx. These possibilities were evaluated in the following current-clamp and voltage-clamp experiments performed in combination with simultaneous measurements of cell length. Figure 7 shows representative traces of action potentials (APs) recorded under current-clamp mode with the perforated-patch configuration of the patch-clamp technique. The corresponding contraction twitches are superimposed. Ang II induced a marked prolongation of APD. This Ang II-induced APD lengthening was associated with an increase of cell shortening within the first 10 min of Ang II treatment. At the peak of the Ang II response, average percentage increase of APD after 30% ( $\text{APD}_{30}$ ), 50% ( $\text{APD}_{50}$ ) and 90% ( $\text{APD}_{90}$ ) of repolarisation time was  $36.7 \pm 8.1$ ,  $27.9 \pm 6.4$  and  $21.7 \pm 5.5\%$ , respectively ( $n = 8$ ,  $P < 0.05$  by repeated-measures ANOVA for paired values). The Ang II effect was significantly more important within the voltage range of the AP plateau than at more repolarised potentials, probably indicating that Ang II is predominantly affecting the currents that underlie plateau potentials, i.e. L-type  $\text{Ca}^{2+}$  currents ( $I_{\text{Ca}}$ ).



**Figure 5.** Lack of effect of Ang II on myofilament responsiveness to  $\text{Ca}^{2+}$

Typical continuous recording of cell length showing similar increases in cell shortening induced by either elevating  $[\text{Ca}^{2+}]_o$  or by addition of Ang II. Below are the actual tracings of the individual twitch contractions and the associated indo-1 fluorescence transients at the times indicated by letters *a-d* on the continuous chart recording.


**Table 1. Effect of high extracellular  $\text{Ca}^{2+}$  and Ang II on contraction and  $\text{Ca}^{2+}$  transient parameters of single cat myocytes**

	Control	High $\text{Ca}^{2+}$	Control	Ang ( $1 \mu\text{M}$ )
$L_0$	$138 \pm 6$	$138 \pm 6$	$138 \pm 6$	$138 \pm 6$
TA (% of $L_0$ )	$1.9 \pm 0.4$	$4.7 \pm 0.6^*$	$1.8 \pm 0.3$	$4.2 \pm 0.5^*$
$t_{1/2}$ contraction (ms)	$134 \pm 7$	$102 \pm 11^*$	$135 \pm 11$	$124 \pm 12$
Indo-1 ratio (410/490)				
Diastolic	$1.01 \pm 0.1$	$1.05 \pm 0.1$	$0.97 \pm 0.1$	$1.05 \pm 0.1$
Systolic	$1.21 \pm 0.1$	$1.37 \pm 0.1^*$	$1.16 \pm 0.1$	$1.32 \pm 0.1^*$
Amplitude	$0.20 \pm 0.03$	$0.32 \pm 0.04^*$	$0.19 \pm 0.02$	$0.27 \pm 0.04^*$
$t_{1/2}$ transient (ms)	$257 \pm 12$	$210 \pm 15^*$	$272 \pm 15$	$253 \pm 14$

$L_0$ , resting cell length; TA, twitch amplitude;  $t_{1/2}$  contraction, half-relaxation time of contraction;  $t_{1/2}$  transient, 50% relaxation of indo-1 fluorescence transient. Values are means  $\pm$  s.e.m.;  $n = 8$  for all values. \*Significant vs. control value ( $P < 0.05$ ).

In order to evaluate whether the Ang II-induced positive inotropic effect was due to an enhancement of  $I_{\text{Ca}}$  and/or of the reverse mode of the  $\text{Na}^+$ - $\text{Ca}^{2+}$  exchange secondary to APD lengthening, a voltage protocol where voltage and time were controlled was used. Figure 8A shows

representative traces of the effect of Ang II on cell shortening at two different simulated APs, a short AP (plateau of 200 ms of duration; Fig. 8A, upper traces) and a long AP (plateau of 400 ms of duration; Fig. 8A, lower traces). Under both these conditions, exposure of the myocyte to Ang II induced a similar increase in cell shortening to that measured under current-clamp configuration. Figure 8B depicts overall results of cell shortening provoked by the current-clamp and voltage-clamp protocols of Figs 7 and 8A. No significant differences were observed in the Ang II-induced positive inotropic effect between contractions provoked by current-clamp stimulus and those produced by protocols in which time duration changes were controlled. These results indicate that Ang II-induced APD lengthening *per se* does not contribute to the Ang II-induced positive inotropic effect.




**Figure 6. Failure of Ang II to modulate myofilament responsiveness to  $\text{Ca}^{2+}$**

A, overall data of the effect of high  $[\text{Ca}^{2+}]_o$  and Ang II on contraction amplitude and peak indo-1 fluorescence ( $n = 8$ ). Both interventions produced similar increases in contraction amplitude and peak fluorescence. B, relationship between the increase in peak systolic indo-1 fluorescence ratio and peak shortening after 10 min (peak) and 20 min (steady state) of the application of Ang II. The results are compared with the relationship obtained by elevating  $[\text{Ca}^{2+}]_o$ . All points follow a similar relationship.

### Does $\text{Ca}^{2+}$ entry increase due to a direct effect of Ang II on $I_{\text{Ca}}$ ?

The following step was directed to the study of the effects of Ang II on  $I_{\text{Ca}}$ . Since results reported previously using the whole-cell configuration of patch-clamp technique are highly controversial (Allen *et al.* 1988; Kaibara *et al.* 1994; Ikenouchi *et al.* 1994; Ai *et al.* 1998), we used, as stated above, a more physiological approach, the perforated-patch technique. Figure 9A shows the time course of the effect of Ang II on the peak  $I_{\text{Ca}}$  evoked at 0 mV. The representative traces corresponding to the points indicated in the figure are shown below. Application of Ang II to the bath induced an increase in  $I_{\text{Ca}}$ . This increase in the current started after 2 min, reached a maximum plateau value (75% above control) after 7–11 min, began to decrease after this period of time, and finally reached a new steady-state value (50% above control) after 16–20 min of exposure of the myocyte to Ang II. Six other myocytes exhibited this behaviour, the mean increase above control  $I_{\text{Ca}}$  being  $77.7 \pm 6.3\%$  at the peak effect of Ang II (8–15 min) and  $62.1 \pm 8.7\%$  after 20 min of exposure of the myocytes to the peptide





**Figure 7.** Representative traces of action potentials (current clamp) and cell shortening recorded simultaneously, before and after 5 and 10 min of application of Ang II to the bath solution


The contraction amplitude was related to that of the control peak shortening which was considered as 100%. Ang II induced an increase in cell shortening and a lengthening of APD.

( $P < 0.05$ ;  $n = 7$ ). Figure 9B depicts the mean current density–voltage relations at control and after 7–8 min of Ang II treatment. A significant increase in current density was observed in the range of voltage between  $-15$  and  $+45$  mV after application of Ang II to the bath solution.

The increase in  $I_{Ca}$  evoked by Ang II and the close temporal association between this effect and the increase in the  $Ca_iT$  would indicate that the increase in  $I_{Ca}$  is a main determinant of the Ang II-induced increase in the  $Ca_iT$  and myocardial contractility. However, other mechanisms that could also contribute to this effect should be considered.

**Role of the Na<sup>+</sup>–Ca<sup>2+</sup> exchanger in the Ang II-induced positive inotropic effect**

Enhanced Ca<sup>2+</sup> entry via the reverse mode of the Na<sup>+</sup>–Ca<sup>2+</sup> exchanger may also play a role in the Ang II-induced positive inotropic effect. To test this possibility, we studied the effect of Ang II in the presence of  $2.5 \mu\text{M}$  of KB-R7943 (KB), a specific blocker of the reverse mode of the Na<sup>+</sup>–Ca<sup>2+</sup> exchanger (Iwamoto *et al.* 1996). This concentration of KB blocked the increase in  $[Ca^{2+}]_i$  after activation of the reverse mode of the Na<sup>+</sup>–Ca<sup>2+</sup> exchanger (Fig. 10A, inset) produced by reduction of external Na<sup>+</sup> from 143 to 70 mM (NaCl replaced with choline chloride). A similar method of



**Figure 8.** Effects of Ang II on contractions evoked by simulated voltage-clamped action potentials


A, representative traces of cell shortening evoked by the voltage protocols shown on the left (voltage clamp), before and after 10 min of exposure of the myocyte to Ang II. B, mean percentage increase of peak shortening in relation to control for contractions evoked by current-clamp stimulus (Fig. 7,  $n = 8$ ) and by voltage-clamp protocols simulating short and long action potentials (A,  $n = 6$  and  $5$ , respectively). No significant changes were observed among these groups.

challenging the cell with a  $\text{Na}^+$  gradient sufficient to make the  $\text{Na}^+$ – $\text{Ca}^{2+}$  exchanger work in the reverse mode was previously used by Ladilov *et al.* (1999). Although KB produced a significant decrease of cell shortening ( $45 \pm 10\%$ ), the percentage increase in contraction amplitude produced by Ang II in the presence of KB was not significantly different from that produced in the absence of the  $\text{Na}^+$ – $\text{Ca}^{2+}$  blocker ( $n = 5$ ) (Fig. 10A, chart recording and Fig. 10C). In the presence of KB Ang II increased the  $\text{Ca}_i\text{T}$  amplitude by  $27 \pm 3\%$ . This increment was not significantly different from that produced by Ang II in the absence of the  $\text{Na}^+$ – $\text{Ca}^{2+}$  exchanger blocker. These results are in agreement

with results obtained in additional experiments performed in cat papillary muscles contracting isometrically. In these preparations KB concentrations up to  $5 \mu\text{M}$  produced only a modest decrease in basal contractility ( $7 \pm 1.9\%$ ,  $n = 12$ ). This concentration of KB failed to significantly affect the positive inotropic effect of  $0.5$  and  $1 \mu\text{M}$  Ang II (results not shown).

#### Role of the SR in the Ang II-induced positive inotropic effect

SR  $\text{Ca}^{2+}$  release plays a key role in the excitation–contraction coupling mechanism of mammalian heart (Fabiato, 1983,




**Figure 9.** Ang II increases  $I_{\text{Ca}}$

*A*, time course of the effects of Ang II on peak  $I_{\text{Ca}}$  current evoked by a step to  $0$  mV in a single myocyte. Representative traces of  $I_{\text{Ca}}$  corresponding to the points indicated in the upper panel are shown below. *B*, current density–voltage relations for mean data of peak current density collected from six myocytes in the absence and the presence of Ang II. A significant increase in  $I_{\text{Ca}}$  was observed after application of Ang II to the bath solution.

1985). We therefore investigated the role of the SR as a possible target for the Ang II-induced positive inotropic effect. For this purpose, the SR was functionally inhibited by pretreatment of the cells with ryanodine (Ry; 500 nM). Caffeine-induced contractions provoked in the presence and in the absence of Ry were used to assess the ability of Ry (500 nM) to inhibit SR function. The tracings in Fig. 10B (inset) show that application of 15 mM caffeine induced a phasic contraction that was not reproducible when caffeine was applied after the incubation with Ry, indicating the failure of the SR to accumulate and release  $Ca^{2+}$  in the presence of Ry. The chart recording in Fig. 10B depicts a

representative example of the effect of Ang II in the continued presence of Ry. Although Ry produced a significant decrease in cell shortening ( $53 \pm 7\%$ ) associated with a prolongation of time to peak shortening ( $37 \pm 7\%$ ) and in half-relaxation time ( $61 \pm 12\%$ ), the percentage increase in contraction amplitude produced by Ang II in the presence of Ry was not significantly different from that produced by Ang II alone. The Ang II-induced positive inotropic effect observed in the presence of Ry was associated with a  $35 \pm 9\%$  increase in the  $Ca_iT$  amplitude that was not significantly different from that produced by Ang II alone. The unexpected finding that the SR seems



**Figure 10. Failure of KB and Ry to prevent the Ang II-induced positive inotropic effect**

*A*, representative example of the effect of Ang II in the continued presence of KB ( $2.5 \mu\text{M}$ ) on myocyte contraction. The continuous chart recording shows a typical biphasic contractile response to Ang II in a myocyte pretreated and continuously perfused with the  $Na^+$ - $Ca^{2+}$  blocker, KB. The inset to the left of the figure shows that KB was able to prevent the increase in  $Ca^{2+}$  entry by activation of the reverse mode of the  $Na^+$ - $Ca^{2+}$  exchanger produced by external  $Na^+$  reduction. *B*, typical continuous chart recording of contraction amplitude in response to Ang II in the continued presence of Ry (500 nM). Ang II exerts a typical positive inotropic response even in the presence of Ry. The inset to the left shows a representative example of the effect of caffeine (15 mM) applied in the absence and presence of Ry (500 nM). Rapid application of caffeine induced a phasic contraction that was not reproducible after cells were treated with Ry. Similar results were obtained in 5 other cells. *C*, overall data of the time course of the contractile effects of Ang II alone, Ang II in the continued presence of KB or Ang II in the continued presence of Ry. Data are means  $\pm$  s.e.m.;  $n = 5$  cells per group.

not to be necessary for the Ang II-induced positive inotropism led us to examine the role of the SR in the presence of a more typical inotropic intervention, like elevating  $[Ca^{2+}]_o$ . These experiments demonstrated that an increase in cell shortening similar to that evoked by Ang II, produced by increasing  $[Ca^{2+}]_o$ , was not blunted in the presence of 500 nM Ry (4 mM  $[Ca^{2+}]_o$ :  $191 \pm 13\%$  vs. 4 mM  $[Ca^{2+}]_o + 500$  nM Ry:  $187 \pm 19\%$ ;  $n = 6$ ). Figure 10C shows the overall results of the effect of 1  $\mu$ M Ang II on contractile amplitude in the absence and in the presence of Ry. Similar results were obtained in three other cells using 500 nM thapsigargin (Thaps) plus 1  $\mu$ M Ry. These results indicate that the presence of a functional SR is not necessary for Ang II to be able to develop a typical positive inotropic response.

## DISCUSSION

Physiological peptides such as endothelin-1 and Ang II have been recently recognised as potential modulators of cardiac contractility, producing positive inotropic effects in several species including the human. Plasma levels of Ang II overlap the range of the threshold concentrations for the positive inotropic effect of the peptide (Koch-Weser, 1965; Catt *et al.* 1970). The physiological importance of Ang II as a modulator of myocardial contractility is strengthened further, in light of experimental evidence showing that Ang II levels in cardiac tissue may reach fairly high concentrations (over 100 times plasma levels; Dell'Italia *et al.* 1997).

The subcellular mechanisms responsible for the positive inotropic effect of Ang II are still a matter of major controversy. Different results obtained in several mammalian species may be due at least in part to species-dependent variations which appear as a common feature among agents that increase phosphoinositide turnover (Endoh *et al.* 1991).

The cat is a mammalian species in which Ang II exerts a pronounced positive inotropic effect (Koch-Weser, 1965; Mattiazzi *et al.* 1997). The underlying mechanisms of this effect have not been previously investigated and constitute the aim of the present study. For this purpose we selected from pilot experiments the concentrations of Ang II that evoked maximal inotropic effects. The findings are the following: (1) the main mechanism involved in the overall shape of Ang II-induced positive inotropic effect is a  $pH_i$ -independent increase in the  $Ca_iT$ ; (2) no significant changes in myofilament responsiveness to  $Ca^{2+}$  were detected; and (3) the increase in the  $Ca_iT$  was mainly determined by an increase in  $I_{Ca}$ .

### Increases in the $Ca_iT$ vs. increases in myofilament responsiveness to $Ca^{2+}$ ; role of $pH_i$

The present experiments have shown that similar contractility increases produced by either Ang II administration or by increasing  $[Ca^{2+}]_o$  were associated with similar increases in the  $Ca_iT$  (Figs 5 and 6). These results, together with those obtained with the phase-plane analysis

of cell shortening vs.  $[Ca^{2+}]_i$  (Fig. 4, right panel) measured either at the peak of the Ang II-induced positive inotropic response or when the positive inotropic effect of Ang II reached steady state, indicate that in the cat ventricle Ang II does not produce changes in myofilament responsiveness to  $Ca^{2+}$  and that at least in this species the overall shape of the Ang II-induced positive inotropic effect is entirely determined by changes in the  $Ca_iT$ .

The temporal dissociation between the increase in  $pH_i$  and in contractility produced by Ang II, plus the fact that the Ang II-induced positive inotropic effect was preserved in the presence of HOE, indicate that Ang II-induced intracellular alkalosis does not modulate myofilament responsiveness to  $Ca^{2+}$ . Considering the results of Fabiato & Fabiato (1978), this is not an unexpected finding, since using skinned cardiac fibres they demonstrated that an increase in  $pH_i$  of 0.4 units was needed to produce only a modest increase in tension (approximately 30% at  $pCa$  6.0). Furthermore, a  $pH_i$  shift towards alkalosis did not produce a significant change in maximal activated force, as acidosis did (Fabiato & Fabiato, 1978). Therefore, if these results can be extrapolated to feline myocardium and to a more intact preparation like the isolated myocytes, the  $pH_i$  increase of 0.06 pH units produced by Ang II in these preparations may not be sufficient to evoke a detectable increase in myofilament responsiveness to  $Ca^{2+}$ . Previous experiments by Ikenouchi *et al.* (1994) in rabbit heart failed to detect a significant increase in the  $Ca_iT$  after Ang II administration. It was concluded from these results that the increase in contractility produced by Ang II (more than 100%) was entirely due to a  $pH_i$ -dependent increase in myofilament responsiveness to  $Ca^{2+}$ . However, the Ang II-induced increase in  $pH_i$  reported in these experiments (approximately 0.2 pH units), although higher than the one detected in the cat ventricle (Mattiazzi *et al.* 1997 and the present results), would again, according to Fabiato & Fabiato's results, not be sufficient to completely account for the increase in contractility observed. Indeed, more recent experiments demonstrated that Ang II also induced an increase in the  $Ca_iT$  in rabbit heart (Skolnick *et al.* 1998). Other experimental evidence indicates that the Ang II-induced activation of the  $Na^+-H^+$  exchanger cannot be taken as a synonym of intracellular alkalosis (Boron *et al.* 1989; Ganz *et al.* 1989; Mattiazzi *et al.* 1997; Camili3n de Hurtado *et al.* 1998). In this scenario, working in the more physiological bicarbonate buffer, Ang II has been shown to activate both an alkalinising mechanism, the  $Na^+-H^+$  exchanger (Mattiazzi *et al.* 1997; Camili3n de Hurtado *et al.* 1998) and an acidifying mechanism, the  $Na^+$ -independent  $Cl^-/HCO_3^-$  exchanger (Camili3n de Hurtado *et al.* 1998), the net result being a lack of change in  $pH_i$ . However, under these conditions, the positive inotropic effect of Ang II was still present (Mattiazzi *et al.* 1997). Therefore the conclusion that the increase in  $pH_i$  plays a predominant role in determining the increase in contractility induced by Ang II is not straightforward and should be supported by increases in  $pH_i$  of a magnitude sufficient to explain a cause-and-effect

relationship. Our findings indicate that the positive inotropic effect of Ang II is not related to a pH<sub>i</sub> change in isolated myocytes from the cat ventricle. This conclusion is in agreement with previous results in the same species in multicellular preparations (Mattiazzi *et al.* 1997).

### Subcellular mechanisms of the Ang II-induced increase in the Ca<sub>i</sub>T

There are different and not mutually exclusive mechanisms by which Ang II may increase the Ca<sub>i</sub>T: (1) an increase in Ca<sup>2+</sup> entry through L-type calcium channels or the reverse mode of the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger; (2) a decrease in Ca<sup>2+</sup> efflux through the forward mode of the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger; and (3) an IP<sub>3</sub>-mediated Ca<sup>2+</sup> release from intracellular calcium stores.

The results obtained in voltage-clamp experiments demonstrated that Ang II produced a significant increase in *I*<sub>Ca</sub> with a time course similar to that of the positive inotropic effect (Fig. 9). The increase in *I*<sub>Ca</sub> evoked by Ang II in cardiac myocytes has been detected in some but not all previous studies. In fact *I*<sub>Ca</sub> has been found to be increased (Kass & Blair, 1981; Allen *et al.* 1988; Kaibara *et al.* 1994), unchanged (Ikenouchi *et al.* 1994; Ai *et al.* 1998) or diminished by Ang II (Habuchi *et al.* 1995). The reason for these discrepant results is not apparent to us. They may be explained at least in part by the different species used and/or the different techniques employed to measure *I*<sub>Ca</sub>. The technique used in the present experiments allows the preservation of the intracellular milieu, which is essential to detect the effects of hormones mediated by second messengers, as is the case for Ang II. Recent experiments have shown, indeed, that the increase in *I*<sub>Ca</sub> evoked by α-adrenoceptor stimulation could be detected only when either perforated patch-clamp or cell-attached single channel recording was used, two conditions that preserve the intracellular environment (Liu & Kennedy, 1998; Zhang *et al.* 1998).

The subcellular mechanisms that determine the increase in *I*<sub>Ca</sub> were not assessed directly in the present study. However, this increase seems unlikely to be mediated by an Ang II-induced increase in pH<sub>i</sub> as suggested by Kaibara *et al.* (1994), since this increase occurred before a significant increment in pH<sub>i</sub> could be detected in parallel experiments.

The present experiments showed in addition that the Ang II-induced prolongation of the APD does not seem to contribute to the positive inotropic effect of the peptide: When the APD was not allowed to increase by using voltage-clamped simulated APs, the Ang II-induced positive inotropic effect was still detected. Thus, the prolongation of the APD produced by Ang II appears to be a consequence rather than a cause of the increase in *I*<sub>Ca</sub>.

Recent experiments in the rabbit heart have suggested that the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger may represent the main mechanism by which the Ca<sub>i</sub>T increases after Ang II administration (Fujita & Endoh, 1999). A possible contribution of the

Na<sup>+</sup>-Ca<sup>2+</sup> exchanger to the positive inotropic effect of Ang II has also been suggested by Barry *et al.* (1995). The rationale behind this suggestion was that the increase in intracellular Na<sup>+</sup> produced by Ang II through the activation of the Na<sup>+</sup>-H<sup>+</sup> exchanger would lead to a decrease in Na<sup>+</sup> gradient which would favour a Ca<sup>2+</sup> influx via the reverse mode of the exchanger. The increase in intracellular Na<sup>+</sup> load produced by Na<sup>+</sup>-H<sup>+</sup> exchanger activation could also produce a reduction of Ca<sup>2+</sup> efflux via the forward mode of the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger and thus increase [Ca<sup>2+</sup>]<sub>i</sub>. Moreover, it has been shown in *in vitro* experiments that Ang II directly activates the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger mechanism (Ballard & Schaffer, 1996). In light of the present experiments it is hard to conceive that the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger plays a significant role in the Ang II-induced increase in the Ca<sub>i</sub>T for the following reasons: (1) no significant changes in pH<sub>i</sub> were detected at the moment in which the positive inotropic effect of Ang II was maximum; (2) the presence of HOE did not modify the positive inotropic effect of Ang II; and (3) the positive inotropic effect and the increase in the Ca<sub>i</sub>T produced by Ang II were both preserved in the presence of the specific blocker of the reverse mode of the Na<sup>+</sup>-Ca<sup>2+</sup> exchanger, KB.

An increase in Ca<sup>2+</sup> release by the SR evoked by IP<sub>3</sub> also seems unlikely since the amplitude of the Ca<sub>i</sub>T produced by Ang II was not significantly modified after functional blockade of the SR by Ry and Thaps. Thus the possibility that IP<sub>3</sub> plays a significant role in the positive inotropic effect of Ang II would require that IP<sub>3</sub> released Ca<sup>2+</sup> from a Ca<sup>2+</sup> store that was insensitive to caffeine, different from that blocked by Ry, like the one that has been described for rat atrial cells (Vigne *et al.* 1990) and in addition, that was insensitive to the SERCA2 inhibitor Thaps. We are not aware of such a reservoir in ventricular cells. The failure of Ry plus Thaps to affect the positive inotropic effect of Ang II shown in the present results is in line with the results of Kentish *et al.* (1990), which showed that Ca<sup>2+</sup> release by IP<sub>3</sub> does not play a significant role in the excitation-contraction coupling of cardiac muscle.

In summary the present results demonstrate for the first time that in feline myocardium the Ang II-induced positive inotropic effect is exclusively mediated by an increase in the Ca<sub>i</sub>T which is mainly due to the activation of *I*<sub>Ca</sub>. Two additional major findings of this study are that the increase in the Ca<sub>i</sub>T and the activation of *I*<sub>Ca</sub> occurred independently of changes in pH<sub>i</sub>.

AI, T., HORIE, M., OBAYASHI, K. & SASAYAMA, S. (1998). Accentuated antagonism by angiotensin II on guinea-pig cardiac L-type Ca currents enhanced by β-adrenergic stimulation. *Pflügers Archiv* **436**, 168–174.

AIELLO, A., VILA-PETROFF, M. G., MATTIAZZI, A. & CINGOLANI, H. E. (1998). Evidence for an electrogenic Na<sup>+</sup>-HCO<sub>3</sub><sup>-</sup> symport in rat cardiac myocytes. *Journal of Physiology* **512**, 137–148.

- ALLEN, I. S., COHEN, N. M., DHALLAN, R. S., GAA, S. T., LEDERER, W. J. & ROGERS T. B. (1988). Angiotensin II increases spontaneous contractile frequency and stimulates calcium current in cultured neonatal rat heart myocytes: insights into the underlying biochemical mechanisms. *Circulation Research* **62**, 524–534.
- BALLARD, C. & SCHAFFER, S. (1996). Stimulation of the  $\text{Na}^+/\text{Ca}^{2+}$  exchanger by phenylephrine, angiotensin II and endothelin 1. *Journal of Molecular and Cellular Cardiology* **28**, 11–17.
- BARRY, W. H., MATSUI, H., BRIDGE, J. H. & SPITZER, K. W. (1995). Excitation-contraction coupling in ventricular myocytes: effects of angiotensin II. *Advances in Experimental and Medical Biology* **382**, 31–39.
- BLANK, P. S., SILVERMAN, H. S., CHUN, O. Y., HOGUE, B. A., STERN, M. D., HANSFORD, R. G., LAKATTA, E. G. & CAPOGROSSI, M. C. (1992). Cytosolic pH measurements in single cardiac myocytes using carboxy-seminaphthorhodafluor-1. *American Journal of Physiology* **263**, H276–284.
- BORON, W. F., BOYARSKY, G. & GANZ, M. (1989). Regulation of intracellular pH in renal mesangial cells. *Annals of the New York Academy of Sciences* **574**, 321–332.
- CAMILIÓN DE HURTADO, M. C., ALVAREZ, B. V., PÉREZ, N. G., ENNIS, I. L. & CINGOLANI H. E. (1998). Angiotensin II activates  $\text{Na}^+$ -independent  $\text{Cl}^-/\text{HCO}_3^-$  exchange in ventricular myocardium. *Circulation Research* **82**, 473–481.
- CATT, K. J., CAIN, M. D., COGHLAN, J. P., ZIMMET, P. Z., CRAN, E. & BEST, J. B. (1970). Metabolism and blood levels of Angiotensin II in normal subjects, renal disease, and essential hypertension. *Circulation Research* **27**, 177–193.
- DELL'ITALIA, L. J., MENG, Q. C., BALCELLS, E., WEI, C., PALMER, R., HAGEMAN, J. D., HANKENS, G. H. & OPARIL, S. (1997). Compartmentalization of angiotensin II generation in the dog heart. Evidence for independent mechanisms in intravascular and interstitial spaces. *Journal of Clinical Investigation* **100**, 253–258.
- ENDO, M., HIRAMOTO, T., ISHIHATA, A., TAKANASHI, M. & INUI, J. (1991). Myocardial  $\alpha_1$ -adrenoceptors mediate positive inotropic effect and changes in phosphatidylinositol metabolism: species differences in receptor distribution and the intracellular coupling process in mammalian ventricular myocardium. *Circulation Research* **68**, 1179–1190.
- FABIATO, A. (1983). Calcium-induced release of calcium from the cardiac sarcoplasmic reticulum. *American Journal of Physiology* **245**, C1–14.
- FABIATO, A. (1985). Time and calcium dependence of activation and inactivation of calcium-induced release of calcium from the sarcoplasmic reticulum of a skinned canine cardiac Purkinje cell. *Journal of General Physiology* **85**, 247–289.
- FABIATO, A. & FABIATO, F. (1978). Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscles. *Journal of Physiology* **276**, 233–255.
- FUJITA, S. & ENDO, M. (1999). Influence of a  $\text{Na}^+/\text{H}^+$  exchange inhibitor ethylisopropylamiloride,  $\text{Na}^+/\text{Ca}^{2+}$  exchange inhibitor KB-R7943 and their combination on the increases in contractility and  $\text{Ca}^{2+}$  transient induced by angiotensin II in isolated adult rabbit ventricular myocytes. *Naunyn-Schmiedeberg's Archives of Pharmacology* **360**, 575–584.
- GANZ, M. B., BOYARSKY, G., STERZEL, R. B. & BORON, W. F. (1989). Arginine vasopressin enhances  $\text{pH}_i$  regulation in the presence of  $\text{HCO}_3^-$  by stimulating three acid-base transport systems. *Nature* **337**, 648–651.
- GUNASEGARAM, S., HAWORTH, R. S., HEARSE, D. J. & AVKIRAN, M. (1999). Regulation of sarcolemmal  $\text{Na}^+/\text{H}^+$  exchanger activity by angiotensin II in adult rat ventricular myocytes: opposing actions via  $\text{AT}_1$  versus  $\text{AT}_2$  receptors. *Circulation Research* **85**, 919–930.
- HABUCHI, Y., LU, L. L., MORIKAWA, J. & YOSHIMURA, M. (1995). Angiotensin II inhibition of L-type  $\text{Ca}^{2+}$  current in sinoatrial node cell of rabbits. *American Journal of Physiology* **268**, H1053–1060.
- IKENOUCI, I., BARRY, W. H., BRIDGE, J. H. B., WEIBERG, E. O., APSTEIN C. S. & LORELL, H. (1994). Effects of angiotensin II on intracellular  $\text{Ca}^{2+}$  and pH in isolated rabbit hearts and myocytes loaded with the indicator indo-1. *Journal of Physiology* **480**, 203–215.
- ISHIHATA, A. & ENDO, M. (1993). Pharmacological characteristics of the positive inotropic effect of angiotensin II in the rabbit ventricular myocardium. *British Journal of Pharmacology* **108**, 999–1005.
- ISHIHATA, A. & ENDO, M. (1995). Species-related differences in inotropic effects of angiotensin II in mammalian ventricular muscle: receptors, subtypes and phosphoinositide hydrolysis. *British Journal of Pharmacology* **114**, 447–453.
- IWAMOTO, T., WATANO, T. & SHIGEKAWA, M. (1996). A novel isothiourea derivative selectively inhibits the reverse mode of  $\text{Na}^+/\text{Ca}^{2+}$  exchange in cells expressing NCX1. *Journal of Biological Chemistry* **271**, 22391–22397.
- KAIBARA, M., MATIARAI, S., YANO, K. & KAMEYAMA, M. (1994). Involvement of  $\text{Na}^+/\text{H}^+$  antiporter in the regulation of L-type  $\text{Ca}^{2+}$  channel current by angiotensin II in rabbit ventricular myocytes. *Circulation Research* **75**, 1121–1125.
- KASS, R. S. & BLAIR, M. L. (1981). Effects of angiotensin II on membrane current in cardiac Purkinje fibers. *Journal of Molecular and Cellular Cardiology* **13**, 797–809.
- KENTISH, J. C., BARSOTTI, R. J., LEA, T. J., MULLIGAN, I. P., PATEL, J. R. & FERENCZI, M. A. (1990). Calcium release from cardiac sarcoplasmic reticulum induced by photorelease of calcium or  $\text{Ins}(1,4,5)\text{P}_3$ . *American Journal Physiology* **258**, H610–615.
- KOCH-WESER, J. (1965). Nature of the inotropic action of angiotensin II on ventricular myocardium. *Circulation Research* **16**, 230–237.
- KORN, S. J., MARTY, A., CONNOR, J. A. & HORN, R. (1991). Perforated patch recording. *Methods in Neurosciences* **4**, 364–373.
- LADILOV, Y., HAFNER, S., BALSER-SCHAFFER, C., MAXEINER, H. & PIPER, H. M. (1999). Cardioprotective effects of KB-R7943: a novel inhibitor of the reverse mode of  $\text{Na}^+/\text{Ca}^{2+}$  exchanger. *American Journal of Physiology* **276**, H1868–1876.
- LIU, S. J. & KENNEDY, R. H. (1998).  $\alpha_1$ -Adrenergic activation of L-type Ca current in rat ventricular myocytes: perforated patch-clamp recordings. *American Journal of Physiology* **274**, H2203–2207.
- MCDONALD, T. F., PELZER, S., TRAUTWEIN, W. & PELZER, D. J. (1994). Regulation and modulation of calcium channels in cardiac, skeletal, and smooth muscle cells. *Physiological Reviews* **74**, 365–507.
- MATSUI, H., BARRY, W. H., LIVSEY, C. & SPITZER, K. W. (1995). Angiotensin II stimulates sodium-hydrogen exchange in adult rabbit ventricular myocytes. *Cardiovascular Research* **29**, 215–221.
- MATTIAZZI, A., PÉREZ, N. G., VILA-PETROFF, M. G., ALVAREZ, B. V., CAMILIÓN DE HURTADO, M. C. & CINGOLANI, H. E. (1997). Dissociation between the positive and alkalizing effects of angiotensin II in feline myocardium. *American Journal of Physiology* **272**, H1131–1136.
- MORGAN, P. E., AIELLO, E. A., CHIAPPE DE CINGOLANI, G. E., MATTIAZZI, A. R. & CINGOLANI, H. E. (1999). Chronic administration of nifedipine induces upregulation of functional calcium channels in rat myocardium. *Journal of Molecular and Cellular Cardiology* **31**, 1873–1883.
- MORITA, H., KIMURA, J. & ENDO, M. (1995). Angiotensin II activation of a chloride current in rabbit cardiac myocytes. *Journal of Physiology* **483**, 119–130.

- SHAH, A. M., SPURGEON, H. A., SOLLOTT, S. J., TALO, A. & LAKATTA, E. G. (1994). 8-Bromo-cGMP reduces the myofilament response to Ca<sup>2+</sup> in intact cardiac myocytes. *Circulation Research* **74**, 970–978.
- SKOLNICK, R. L., LITWIN, S. E., BARRY, W. H. & SPITZER, K. W. (1998). Effect of angiotensin II on pH<sub>i</sub>, [Ca<sup>2+</sup>]<sub>i</sub>, and contraction in rabbit ventricular myocytes from infarcted hearts. *American Journal of Physiology* **275**, H1788–1797.
- SPURGEON, H. A., STERN, M. D., BAARTZ, G., RAFFAELI, S., HANSFORD, R. G., TALO, A., LAKATTA, E. G. & CAPOGROSSI, M. C. (1990). Simultaneous measurements of Ca<sup>2+</sup>, contraction, and potential in cardiac myocytes. *American Journal of Physiology* **258**, H574–586.
- TALUKDER, M. A. H. & ENDOH, M. (1997). Pharmacological differentiation of synergistic contribution of L-type Ca<sup>2+</sup> channels and Na<sup>+</sup>/H<sup>+</sup> exchange to the positive inotropic effect of phenylephrine, endothelin-3 and angiotensin II in rabbit ventricular myocardium. *Naunyn-Schmiedeberg's Archives of Pharmacology* **355**, 87–96.
- VANDENBERG, J. I., METCALFE, J. C. & GRACE A. A. (1994). Intracellular pH recovery during respiratory acidosis in perfused hearts. *American Journal of Physiology* **266**, C489–497.
- VIGNE, P., BREITMAYER, J. P., MARSAULT, R. & FRELIN, C. (1990). Endothelin mobilizes Ca<sup>2+</sup> from a caffeine- and ryanodine-insensitive intracellular pool in rat atrial cells. *Journal of Biological Chemistry* **265**, 6782–6787.
- WATANABE, A. & ENDOH, M. (1998). Relationship between the increase in Ca<sup>2+</sup> transient and contractile force induced by angiotensin II in aequorin-loaded rabbit ventricular myocardium. *Cardiovascular Research* **37**, 524–531.
- ZHANG, S., HIRAOKA, M. & HIRANO, Y. (1998). Effects of α1-adrenergic stimulation on L-type Ca<sup>2+</sup> current in rat ventricular myocytes. *Journal of Molecular and Cellular Cardiology* **30**, 1955–1965.

### Acknowledgements

This study was supported by a grant (PICT 05-00149-02423) from the Agencia Nacional de Promoción Científica y Tecnológica, Argentina. We are most grateful to Dr H. E. Cingolani for his helpful comments on the manuscript and to Dr Wolfgang Scholz from Hoechst AG for kindly supplying the drug HOE 642. The technical assistance of Mrs Mónica Rando is gratefully acknowledged. M.G.V.P., E.A.A. and A.R.M. are Established Investigators of The Consejo Nacional de Investigaciones y Técnicas (CONICET), Argentina.

### Corresponding author

A. Mattiazzi: Centro de Investigaciones Cardiovasculares, Facultad de Ciencias Médicas, 60 y 120, La Plata 1900, Argentina.

Email: ramattia@atlas.med.unlp.edu.ar