

Analysis of mixed foot-and-mouth disease virus infections in Saudi Arabia: prolonged circulation of an exotic serotype

E. L. WOODBURY¹, A. R. SAMUEL^{2*}, N. J. KNOWLES², S. M. HAFEZ³
AND R. P. KITCHING³

¹*Kings College, University of London, Campden Hill Road, Kensington,
London W8, UK*

²*AFRC Institute for Animal Health, Pirbright Laboratory, Ash Road,
Pirbright, Surrey GU24 0NF, UK*

³*C/O National Agriculture and Water Research Centre, PO Box 17285,
Riyadh 11484, Saudi Arabia*

(Accepted 5 August 1993)

SUMMARY

Plaque purification of foot-and-mouth disease (FMD) type O viruses isolated from cattle in Saudi Arabia showed the presence of mixed serotype infections. Sixteen out of 31 samples collected between 1985 and 1991 also contained Asia 1 virus, a serotype which had previously only been isolated from a single outbreak in that country in 1980. Nucleotide sequences of the Asia 1 component of all these samples revealed little variation and showed that they were closely related to both a Russian lapinized vaccine virus strain (Asia 1/Tadzhikistan/64), and to a field isolate from Turkey (Asia 1/TUR/15/73). Although mixed FMD infections have been observed previously this is a first report of a serotype, considered to be exotic to a country, co-existing undetected for an extended period of time.

INTRODUCTION

Foot-and-mouth disease (FMD) is endemic in most Middle Eastern countries with serotype O being the most prevalent. Outbreaks due to serotypes A and Asia 1 occur sporadically whereas those due to serotypes C, SAT 1 and SAT 2 are rare. Some epizootics have swept through the Middle East to threaten parts of Europe, notably SAT 1 in 1962–5, A₂₂ in 1964–5 [1, 2] and Asia 1 in 1973 [3] and 1983–4 [4]. Recent studies have shown that the Middle East strains of FMDV serotype O are serologically distinct from those in Europe and South America. During the period 1987–8 an antigenic shift appears to have occurred [5].

In Saudi Arabia outbreaks of FMD attributable to types O, A, C, Asia 1 and SAT 1 have occurred. Recent control campaigns have utilized vaccines incorporating serotypes O, A, C and Asia 1. Since 1971, when samples were first submitted to the OIE/FAO World Reference Laboratory (WRL) for FMD from Saudi Arabia, serotype O has been recorded in 1971–3, 1978, 1980–93; serotype A in

* To whom reprint requests should be sent.

1973, 1976, 1984, 1986–7 and 1991–3; serotype Asia 1 in 1980 and 1992; serotype C in 1984; and serotype SAT 1 in 1962 and 1970. During 1988 the number of reported outbreaks increased significantly particularly in the large dairy farms around Riyadh.

Antigenic analysis of serotype O viruses isolated from Saudi Arabia showed that while some isolates reacted strongly with a monoclonal antibody (MAb) to O₁/Lausanne/Switzerland/65 others exhibited little or no reactivity [6]. The reactivity of this MAb has been mapped to the G–H loop on 1D^{cap} (VP1) which lies between amino acid residues 129 and 165 [7]. It was considered that plaque purification and sequencing of these phenotypically different strains might help to identify the cause of this phenomenon. This paper reports results obtained during such a study.

MATERIALS AND METHODS

Cell cultures

Monolayers of primary bovine thyroid (BTy), secondary calf kidney (CK) or continuous pig kidney (IB-RS-2) cells were used in the original isolation procedures. Subsequently baby hamster kidney (BHK-21) cells were also used. All the cells were grown in Eagle's MEM containing 5% normal bovine serum and antibiotics. Serum-free medium was used during virus growth.

Viruses

The origin of the virus isolates used in this study are shown in Table 1. Stock viruses were passaged on 175 cm² flasks of IB-RS-2 monolayers. When CPE was complete the supernatant was clarified at 2000 g for 5 min, an equal volume of sterile glycerol added and the virus stored at –20 °C.

Plaque purification

Six-well tissue culture plates confluent with IB-RS-2 cells were washed twice with calcium and magnesium-free PBS. Virus dilutions were made in the range 10⁻¹–10⁻⁶ in Eagle's MEM. Aliquots of the virus dilution (200 µl) were added and left to adsorb for 30 min at 37 °C. The inoculum was removed and the cell sheet washed twice with PBS and overlaid with 3 ml of Eagle's MEM containing 1.6% agar and 1% normal bovine serum at 43 °C. When the agar had set the plates were covered in 'cling film', inverted and incubated for 18 h at 37 °C. Plaques were picked using a pasteur pipette and resuspended in 1 ml of Eagle's MEM. Aliquots (200 µl) were inoculated onto flasks containing confluent monolayers of IB-RS-2 cells. When CPE was complete the supernatant was treated as described above. These viruses were subjected to a further two plaque purification procedures. In some of the later procedures a 1/50 dilution of either anti-O₁/BFS 1860/UK/67 or anti-Asia 1/PAK/1/54 guinea-pig serum was added to the overlay medium to suppress the growth of the homologous serotype.

Monoclonal antibody analysis

A panel of MAbs raised against the reference strain O₁/Lausanne/Switzerland/65 [8, 9] and characterized by sequencing neutralization escape mutant viruses [7, 10], was used in a trapping (capture) ELISA [11] to obtain antigenic profiles.

Table 1. Foot-and-mouth disease virus isolates analysed in this study

WRL ref. no.	Geographical location	Date collected§	Animal	Serotypes isolated
SAU/2/83	Al Kharj, Saudi Arabia	00/00/83	Cattle	O
SAU/4/83	Riyadh, Saudi Arabia	00/00/83	Cattle	O
SAU/11/85	Al Kharj, Saudi Arabia	26/11/85	Cattle	O, Asia 1
SAU/17/86	Al Kharj, Saudi Arabia	07/09/86	Cattle	O, Asia 1
SAU/1/87	Al Jouf, Saudi Arabia	00/00/87	Sheep	O, Asia 1
SAU/34/87	Haradh, Saudi Arabia	00/00/87	Cattle	O
SAU/1/88	Al Qatif, Saudi Arabia	19/01/88	Cattle	O, Asia 1
SAU/8/88	Al Kharj, Saudi Arabia	28/09/88	Cattle	O, Asia 1
SAU/24/88	Al Kharj, Saudi Arabia	29/10/88	Cattle	O, Asia 1
SAU/28/88	Al Kharj, Saudi Arabia	01/01/88	Cattle	O, Asia 1
SAU/30/88†	Riyadh, Saudi Arabia	100/10/88	Cattle	O
SAU/33/88	Riyadh, Saudi Arabia	15/12/88	Cattle	O
SAU/34/89	Al Kharj, Saudi Arabia	00/00/89	Not known	O, Asia 1
SAU/35/89	Al Kharj, Saudi Arabia	00/00/89	Not known	O, Asia 1
SAU/36/89	Al Kharj, Saudi Arabia	00/00/89	Not known	O, Asia 1
SAU/40/89	Al Kharj, Saudi Arabia	00/00/89	Not known	O
SAU/56/89	Todhia, Saudi Arabia	25/08/89	Cattle	O, Asia 1
SAU/17/90	Al Jouf, Saudi Arabia	00/00/90	Sheep	O
SAU/18/90	Afif, Saudi Arabia	00/00/90	Sheep	O
SAU/19/90	Qatif, Saudi Arabia	00/00/90	Cattle	O
SAU/25/90	Thadiq, Saudi Arabia	29/07/90	Cattle	O
SAU/26/90	Al Hair, Saudi Arabia	26/11/90	Cattle	O
SAU/30/90‡	Al Kharj, Saudi Arabia	11/09/90	Cattle	O
SAU/35/90	Al-Majmaa, Saudi Arabia	05/12/90	Sheep	O
SAU/36/90	Riyadh, Saudi Arabia	10/12/90	Cattle	O, Asia 1
SAU/3/91	Al Medyan, Saudi Arabia	19/02/91	Cattle	O
SAU/7/91‡	Al Kharj, Saudi Arabia	05/01/90	Cattle	O, Asia 1
SAU/15/91	Al Kharj, Saudi Arabia	29/06/91	Cattle	O, Asia 1
SAU/16/91	Al Kharj, Saudi Arabia	30/06/91	Cattle	O, Asia 1
SAU/40/91	Quasim, Saudi Arabia	10/08/91	Cattle	O
SAU/42/91	Gizan, Saudi Arabia	27/08/91	Cattle	O, A, Asia 1
LIB/6/88	Bin Walid, Libya	23/08/88	Not known	O
TUR/8/88	Oguzeli, Gaziantep, Turkey	11/05/88	Cattle	O
EGY/2/89	Ismailia, Egypt	00/00/88	Buffalo	O
TUN/3/89	Tunisia	00/00/89	Cattle	O
ALG/3/90	Algeria	15/05/90	Cattle	O
MOR/1/91	Morocco	00/00/90	Sheep	O

* OIE/FAO World Reference Laboratory for Foot-and-Mouth Disease reference number.

† nomadic animals.

‡ carrier animals.

§ 00, date not known.

Briefly, viruses to be tested were reacted with a panel of MABs and the reactivity compared to that of the homologous virus. Results were expressed as a percentage of the homologous reaction.

Enzyme-linked immunosorbent assay

Typing assays were performed using antisera prepared against all seven serotypes of FMDV as previously described [12].

The viruses were compared by one-way liquid phase blocking sandwich ELISA

using sera from vaccinated cattle according to the method of Kitching and colleagues [13]. The assay measures the residual antigen remaining after overnight reaction between dilutions of a reference antiserum and pre-titrated antigen prepared from the field isolate or the homologous reference strain. The serum titre obtained at 50% of the maximum optical density is used to calculate the relationship (r) value and interpreted according to the criteria described by Samuel and colleagues [5].

Nucleotide sequence analysis

Two oligonucleotide primers designated pNK10 (5'-GAAGGGCCCAGGGT-TGGACTC) complementary to the last 15 nucleotides of the 2A gene and the first six nucleotides of the 2B gene and pCP14 (5'-CCTTCACAAATCTGTCC) complementary to nucleotides 108–124 in the 1D^{cap} gene were used.

The sequence of approximately 170 nucleotides at the 3' end of the 1D^{cap} gene was determined for all the plaque isolates by direct RNA sequencing using the dideoxy chain-termination method [14–16]. Additionally the sequence of the region spanning the 1C/1D junction was determined for one of the isolates.

Sequences were analysed using computer programs written by one of the authors (NJK) and from the PHYLIP phylogeny package [17].

RESULTS

Plaque morphology

Virus isolated on BTy cells from the original epithelium of the sample designated O/SAU/8/88 was subjected to plaque purification and 10 plaques were picked. Five had a large plaque phenotype (4 mm diameter) and were designated P1–P5 and five had a small plaque phenotype (1 mm diameter) and designated P6–P10.

Monoclonal antibody analysis (antigenic profiling)

Plaque isolate SAU/8/88 (P1) exhibited reduced binding to all MAbs and approximately one third of the expected binding to the polyclonal positive control. Plaque isolate SAU/8/88 (P7) showed expected binding levels on the antigen controls (approx. 1.2 o.d.) and to the MAbs.

Enzyme-linked immunosorbent assay

SAU/8/88 (P1) and SAU/8/88 (P7) were chosen as representative plaque isolates for examination by ELISA. They were compared with the O₁/Manisa/Turkey/69 and O₁/BFS 1860/UK/67 reference strains. Sucrose density gradient purified material that had been quantified spectrophotometrically was used in the ELISA so that a known amount of each virus was reacted in the assay. Isolate P7 was found to be most closely related to the O₁/Manisa/Turkey/69 reference strain ($r > 1.0$). The reactivity of SAU/8/88 (P1) in the assay was too low for useful comparisons to be made. It was concluded that this isolate was so different antigenically that the polyclonal trapping antibody failed to bind the virus at sufficient levels.

(a)

Tadzhik. /64	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	
TUR/15/73	TTA	CCC	ACC	TCC	TTC	AAC	TAC	GGG	GCT	GTC	AAG	GCT	GAC	ACC	ATC	ACT	GAG	CTG	TTC	ATC	CGC	ATG	AAA	CGC	CGC	GAG	ACT	TAC	TCC	CCT	
SAU/11/85	C	---	---	---	---	---	---	C	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/17/86	C	---	---	---	---	---	---	C	---	---	C	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/1/87																															
SAU/1/88																															
SAU/8/88A	C	---	---	---	---	---	---															T									
SAU/8/88B																															
SAU/8/88C	C-C	---	---	---	---	---	---	C	---	---	---	---	---				*														
SAU/24/88	C-G	---	T	---	T	---	---	*			*																				*
SAU/28/88																															
SAU/34/89	C	-*	---	---	---	---	---	*																		*	*	*	*	*	*
SAU/35/89	C	---	---	---	---	---	---																			*	*	*	*	*	*
SAU/36/89								***																	*	*	*	*	*	*	*
SAU/56/89	C	---	---	---	---	---	---	C	---	---														A	---						
SAU/36/90	C	*	---	---	---	---	C	*	---	---																*	*	*	*	*	*
SAU/7/91	C	*	---	---	---	---	C	*	---	---																*	*	*	*	*	*
SAU/15/91	C	---	---	---	---	---	C	---	---	---																					-C
SAU/16/91																															
SAU/42/91		*	---	---	---	---	---	***																							
SAU/9/92	C-G	---	---	---	---	---	---	-C	---	-A	---	-B	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/10/92	C-G	---	---	---	---	---	---	-C	---	-A	---	-A	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/2/80	C-T	---	---	---	---	---	---	-T	---	*	---	*	---																		---

Tadzhik. /64	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211
TUR/15/73	AGA	CCT	TTC	TTC	GCC	CTG	GAC	ACC	ACC	CAG	GAC	CGC	CGC	AAA	CAG	GAG	ATC	ATT	GCA	CCT	GAG	AAG	CAG	GCT	CTG
SAU/11/85	---	---	-A	-A	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/17/86	---	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/1/87	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/1/88	---	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/8/88A	*	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/8/88B	---	---	-A	*	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	*	*	---	---
SAU/8/88C	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/24/88	---	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/28/88	---	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/34/89	---	---	---	---	---	---	-A	---	---	*	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/35/89	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/36/89	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/56/89	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/36/90	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/7/91	---	---	**	*	*	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/15/91	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/16/91	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/42/91	---	---	---	---	---	---	---	---	---	*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/9/92	-G	---	C-A	---	T	---	T	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/10/92	-G	---	C-A	---	T	---	T	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
SAU/2/80	---	-C	---	-A	---	-T	-T	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

(b)

Tadzhikistan/64	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210
SAU/8/88	ATC	TAC	CAG	ATC	ACA	CAC	GGC	AAA	GCT	GAA	GGA	GAC	GCA	CTG	GTC	GTT	TCC	GTC	AGC	GCC	AAG	GAC	TTT	GAG	---
Tadzhikistan/64	211	212	213	214	215	216	217	218	219	220/001	002	003	004	005	006	007	008	009	010	011	012	013	014	015	
SAU/8/88	TTT	CGC	TTC	CCA	GTC	GAT	GCG	GCC	CAG	CAG/ACT	ACT	ACC	GCT	GGC	GAA	TCT	GCA	GAT	CCA	GTT	ACA	ACC	ACG	GTT	
Tadzhikistan/64	016	017	018	019	020	021	022	023																	
SAU/8/88	GAG	AAT	TAC	GGA	GCG	GAG	ACT	CAG																	

Fig. 1. (a) Nucleotide sequences of the 3' end of the 1D gene of the FMDV Asia 1 isolates from Saudi Arabia. (b) Nucleotide sequences of the 3' end of the 1C gene of the FMDV Asia 1 isolate SAU/8/88 compared to that of Asia 1/Tadzhikistan/64. *Sequence ambiguity (identity of nucleotide could not be determined).

Nucleotide sequence analysis

The nucleotide sequence of the 3' end of the ID^{cap} coding region of the virion genome was determined and the sequences obtained for SAU/8/88 (P1) and SAU/8/88 (P7) were compared. The nucleotide sequence analysis revealed that whilst the SAU/8/88 (P7) sequence was the same as that previously determined for the original unplaqued O/SAU/8/88 isolate [6], the SAU/8/88 (P1) sequence showed that this isolate belonged to the Asia 1 serotype (Fig. 1).

The nucleotide sequence obtained for SAU/8/88 (P1) was compared to other available Asia 1 sequences and was shown to be most closely related to the sequence of a virus isolated from Tadzhikistan SSR in 1964 [18] and to TUR/15/73 [19]. These viruses were not closely related to either SAU/2/80, the first occurrence of Asia 1 in Saudi Arabia, or to SAU/9/92 and SAU/10/92 which were isolated from samples received during the progress of this investigation (Fig. 2).




Fig. 2. Dendrogram showing the relationships between the Saudi Arabian FMDV Asia 1 viruses, Asia 1/Tadzhikistan/64 and Asia 1/Turkey/15/73.

Serotyping by ELISA

Virus typing was performed on the 10 plaque isolates of SAU/8/88 (P1–P10). Results showed that plaques P1–P5 (the large plaque phenotype) were of serotype Asia 1 and plaques P6–P10 were serotype O.

Passage on different cell cultures

The isolate SAU/8/88 was serially passaged five times on either BTy, CK, IB-RS-2 or BHK-21 cells and the supernatant fluids were then serotyped by ELISA as described by Roeder and Le Blanc Smith [12]. All samples typed as serotype O. The Asia 1 component had not been amplified to an extent where it was detectable by ELISA.

Examination of additional FMDV type O isolates

The study was expanded to include virus isolates from different regions within Saudi Arabia and during 1983–91 (Table 1; Fig. 3). Because serotype O strains circulating in North Africa are related to the Saudi Arabian group of isolates (A. R. Samuel and N. J. Knowles, unpublished data), viruses from Algeria (O/ALG/3/90), Tunisia (O/TUN/3/89), Morocco (O/MOR/1/91), Libya (O/LIB/6/88), Turkey (O/TUR/8/88) and Egypt (O/EGY/2/89) were subjected to the plaque isolation procedure described using type O antiserum. In no case was Asia 1 virus isolated.

Many of the samples from Saudi Arabian farms were negative for dual infection (Table 1). However, one sample (SAU/42/91) from Gizan in the south-west of the country contained three different serotypes – O, A and Asia 1. FMDV type Asia 1 was isolated from samples that had originally been typed as O during successive




Fig. 3. Map showing the origin of the samples from Saudi Arabia which were tested for mixtures of FMDV serotypes. Type O viruses were isolated from all the samples, types O and Asia 1 from those contained in a single box and types O, A and Asia 1 from the isolate in a double box.

years from 1985 until 1991. Most of these samples were from farms around the Riyadh and Al Kharj areas, although SAU/1/87 was from Al Jouf in the north, SAU/1/88 from Quatif in the east and SAU/42/91 in the south west (Fig. 3).

DISCUSSION

The phenomenon of mixed serotype FMDV infections is not new. Hedger [20] showed that it was possible to isolate viruses of serotypes SAT 1, SAT 2 and SAT 3 from oesophageal/pharyngeal fluid collected from African buffalo (*Syncaerus caffer*) in Botswana. Here, however, we show for the first time that a serotype thought not to be present in a region has persisted undetected in cattle alongside another endemic serotype. It has been reported that during the course of an epizootic of FMD the serotype of the virus changed. It had been postulated that high mutation rates could be responsible or that latent infections had been triggered when a secondary infection of FMDV had occurred [see refs in 21]. Also, serological tests which gave reactivities with two or more serotypes were sometimes explained as being due to non-specific cross-reactions. It has been usual to passage a virus isolate showing minor cross-reactions in serological assays (complement fixation or ELISA) until a clear typing was obtained.

The possibility of laboratory contamination was discounted by repeating the plaque isolation from the SAU/8/88 sample after re-isolation from the original epithelial material received by the WRL. Isolation and plaquing from the

epithelial material was also carried out independently. In all cases, the viruses were initially identified as serotype O, yet Asia 1 virus was recovered from the samples.

All the plaque isolates examined were closely related to each other and to a rabbit attenuated Russian vaccine virus strain [18] which originated in Tadzhikistan in 1964 and to an isolate from an outbreak in Turkey in 1973 (TUR/15/73) [19] (Fig. 2). They were not closely related to isolates of Asia 1 received from FMD outbreaks in Saudi Arabia in 1992 (> 9% nucleotide difference). These 1992 isolates are most closely related to another vaccine strain Asia 1/IND/8/79 [19]. Interestingly, some of these samples were typed by ELISA as mixtures of type O and Asia 1. The plaque isolates were not related to the isolate Asia 1/SAU/2/80 which was isolated from the first outbreak of Asia 1 in Saudi Arabia in 1980 [22].

Finding type A in one of the isolates was less surprising since outbreaks due to this serotype have occurred in Saudi Arabia since 1991. Sequence analysis showed that the type A component of the SAU/42/91 isolate was most closely related to a group of viruses which caused outbreaks in Turkey during 1991–2 [23] and Saudi Arabia in 1991 (A. R. Samuel, unpublished data). The Saudi Arabian viruses were not closely related to the previous type A outbreak strains from 1986–7 [24] which were more closely related to viruses from the Indian sub-continent [25]. Recent samples received from Saudi Arabia in 1993 have been typed as mixtures of type O and A and similarly samples from two outbreaks in Turkey in 1991 have also been found to be mixtures of serotypes O and A. During the 1984 outbreak of FMD type C in Saudi Arabia, one isolate, SAU/1/84, was found to be a mixture of types C and O (WRL records). Table 2 lists samples from cattle which have been received by the WRL since 1967 and which were identified as mixtures of two or more serotypes; however, the possibility that some were laboratory contaminants cannot be discounted.

Preliminary animal experiments at this institute using SAU/8/88 have shown that both the O and Asia 1 components of the sample can transmit together (F. Davidson, J. Salt, unpublished data).

Possible origins of the Saudi Arabian Asia 1 virus include: (i) the virus has persisted in a sub-clinical form since its initial introduction into Saudi Arabia in 1980; (ii) the virus has been introduced on one or a number of occasions from an area which has been using the Asia 1/Tadzhikistan/64 strain (or a closely related virus), either as a live attenuated vaccine or as an improperly inactivated vaccine; (iii) the virus has been circulating as a sub-population of the type O virus for some years; (iv) parts or all of the capsid-coding region of an inactivated Asia 1 vaccine strain have been rescued by recombination with a type O field virus.

It is improbable that the Asia 1 virus has persisted in Saudi Arabia since 1980 because the genetic difference between the viruses isolated between 1985 and 1991 and the field virus from 1980 (SAU/2/80) is so large (*c.* 5.5% nucleotide difference) and because of the close relationship between the 1985–91 isolates and Tadzhikistan/64 (< 2% nucleotide difference). Asia 1 viruses similar to Tadzhikistan/64 have, so far, not been recovered from type O infections occurring in countries other than Saudi Arabia, even from very closely related type O viruses. Although evidence of experimental and natural heterotypic recombination has been documented [26, 27] and genome fragments may be recovered from

Table 2. *Previously identified mixed serotype infections of foot-and-mouth disease virus**

Year	Country	WRL ref. no.	Serotypes isolated
1971	India	IND/7/71	A + O
1972	Afghanistan	AFG/1/72	O + Asia 1
1972	Uganda	UGA/81/72	O + SAT 1
1973	Malawi	MAL/43/73	SAT 2 + A
1984	Saudi Arabia	SAU/1/84	C + O
1990	Nepal	NEP/115/90	O + C
1991	Kenya	KEN/22/91	O + A + SAT 2
1991	Kenya	KEN/26/91	A + SAT 2
1991	Kenya	KEN/31/91	O + SAT 2
1991	Saudi Arabia	SAU/46/91	O + Asia 1
		SAU/50/91	
1991	Saudi Arabia	SAU/32/91	O + A
1991	Turkey	TUR/9/91	O + A
		TUR/12/91	
1992	Saudi Arabia	SAU/7/92	O + Asia 1
		SAU/8/92	
		SAU/11/92	
1992	Saudi Arabia	SAU/34/92	O + A
		SAU/35/92	
		SAU/36/92	
		SAU/42/92	
1993	Saudi Arabia	SAU/6/93	O + A
		SAU/7/93	
		SAU/9/93	

* Compiled from the records of the OIE/FAO World Reference Laboratory for Foot-and-Mouth Disease.

inactivated vaccines by polymerase chain reaction amplification [28] the viruses recovered show no relationship to Asia 1 vaccine strains known to have been used in Saudi Arabia between 1985 and 1991. We therefore feel that the second possibility is the most likely explanation.

Because the Asia 1 serotype is co-infecting and is able to survive within these populations of animals in the farms around Riyadh, the potential exists for the Asia 1 to cause disease when the opportunity presents itself. At the moment there is no evidence that any outbreaks of FMD have been caused by this virus even though Asia 1 has been isolated from samples from as early as 1985. However, it could be possible that the clinical signs of this particular strain are so mild that infection occurs but remains undiagnosed or are not reported to the authorities. We plan to experimentally infect animals at Pirbright with one of the Asia 1 plaque isolates to ascertain its pathogenicity.

Animals in Saudi Arabia are immunized with vaccines containing an Asia 1 component and this may have suppressed outbreaks of this serotype. However, the danger of outbreaks due to this Asia 1 strain could increase if it were to spread to areas where an Asia 1 strain is not included in the vaccine used to control disease, as is the case in North Africa where outbreaks of type O have recently been extensive.

Demonstration of the phenomenon of dual infections in FMD samples from

cattle in endemic areas shows the need for further virological investigation when diagnostic laboratories find cross-reactivity in serotyping tests such as the ELISA. This policy has now been adopted by the WRL.

ACKNOWLEDGEMENTS

This work was supported by funding from the Ministry of Agriculture, Fisheries and Food, UK. We would like to thank N. Ferris for performing the typing assays and M. Goodlet and R. Osland for the Saudi Arabian field samples.

REFERENCES

1. Arrowsmith AEM. Variation among strains of type A foot-and-mouth disease virus in the Eastern Mediterranean region 1964-1972. *J Hyg* 1975; **75**: 387-97.
2. Hedger RS, Herniman KA. The Middle East foot-and-mouth disease type A epizootic 1964-65. I. Epizootiology and antigenic studies. *Bull Off Int Epiz* 1966; **65**: 303-12.
3. Brooksby JB. Exotic strain of FMD in Turkey. *Vet Rec* 1973; **93**: 320-1.
4. Anonymous. Greek outbreak in Thrace buffer zone. *Bull Off Int Epiz* 1984; **96**: 20-3.
5. Samuel AR, Ouldrige EJ, Arrowsmith AEM, Kitching RP, Knowles NJ. Antigenic analysis of serotype O foot-and-mouth virus isolates from the Middle East, 1981 to 1988. *Vaccine* 1990; **8**: 390-6.
6. Samuel AR, Knowles NJ, Kitching RP. Preliminary antigenic and molecular analysis of strains of foot-and-mouth disease virus serotype O isolated from Saudi Arabia in 1988 and 1989. Report of the Session of the Research Group of the Standing Technical Committee of the European Commission for the Control of Foot-and-Mouth Disease. Lindholm, Denmark. Rome: FAO, 1990: 139-45.
7. Kitson JDA, McCahon D, Belsham GJ. Sequence analysis of monoclonal antibody resistant mutants of type O foot-and-mouth disease virus: evidence for the involvement of the three surface exposed capsid proteins in four antigenic sites. *Virology* 1990; **179**: 26-34.
8. Brocchi E, Civardi A, De Simone F, Panina GF. Characterisation of foot-and-mouth disease virus antibodies. 20th Congress of the Italian Society of Microbiology, Gardone, Italy. *Atti delle Societa Italiana delle scienze veterinarie*, 1983; **36**: 576-8.
9. Cappuchi L, Brocchi E, De Simone F, Panina GF. Characterisation of monoclonal antibodies produced against foot-and-mouth disease viruses. Report of the Session of the Research Group of the Standing Technical Committee of the European Commission for the Control of Foot-and-Mouth Disease, Brescia, Italy. Rome: FAO, 1984: 17-19.
10. Xie QC, McCahon D, Crowther JR, Belsham GJ, McCullough KC. Neutralisation of foot-and-mouth disease virus can be mediated through any of at least three separate antigenic sites. *J Gen Virol* 1987; **68**: 1637-47.
11. Samuel AR, Knowles NJ, Samuel GD, Crowther JR. The evaluation of a trapping ELISA for the differentiation of foot-and-mouth disease virus strains using monoclonal antibodies. *Biologicals* 1991; **19**: 299-310.
12. Roeder PL, Le Blanc Smith PM. Detection and typing of foot-and-mouth disease virus by enzyme-linked immunosorbent assay: a sensitive, rapid and reliable technique for primary diagnosis. *Res Vet Sci* 1987; **43**: 225-32.
13. Kitching RP, Rendle R, Ferris NP. Rapid correlation between field isolates and vaccine strains of foot-and-mouth disease virus. *Vaccine* 1988; **6**: 403-8.
14. Sanger F, Nicklen S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* 1977; **74**: 5463-7.
15. Zimmern D, Kaesberg P. 3'-Terminal nucleotide sequence of encephalomyocarditis virus RNA determined by reverse transcriptase and chain-terminating inhibitors. *Proc Natl Acad Sci USA* 1978; **75**: 4257-61.
16. Knowles NJ. A method for direct nucleotide sequencing of foot-and-mouth disease virus RNA for epidemiological studies. Report of the Session of the Research Group of the Standing Technical Committee of the European Commission for the Control of Foot-and-Mouth Disease, Lindholm, Denmark. Rome: FAO, 1990: 106-12.

17. Felsenstein J. PHYLIP – Phylogeny inference package (version 3.2). *Cladistics* 1989; **5**: 164–6.
18. Sosnovtsev SV, Onischenko AM, Petrov NA, et al. Primary structure of the gene of the VP1 protein of epidemic stomatitis virus of Asia 1 serotype. *Molekulyarnaya Genetika, Microbiologiya i Virusologiya* 1989; **12**: 44–6.
19. Ansell DM, Samuel AR, Carpenter WC, Knowles NJ. Genetic relationships between foot-and-mouth disease type Asia 1 viruses. *Epidemiol Infect* 1994; **112**: 213–24.
20. Hedger RS. Foot-and-mouth disease and the African buffalo (*Syncerus caffer*). *J Comp Path* 1972; **82**: 19–28.
21. Moosbrugger GA. 1966. Changes in type of foot-and-mouth disease virus during an outbreak. *Bull Off Int Epiz* 1966; **65**: 2023–34.
22. Overby E. An outbreak of FMD of serotype Asian 1 in vaccinated (against virus types O, A and C) cattle in Saudi Arabia (case report). *Vet Med Rev* 1984; **1**: 23–8.
23. Armstrong RM, Samuel AR, Knowles NJ, Uluturk S. Genetic studies on foot-and-mouth disease viruses isolated from samples collected in Turkey. Report of the Session of the Research Group of the Standing Technical Committee of the European Commission for the Control of Foot-and-Mouth Disease, Berne, Switzerland. Rome: FAO, 1992.
24. Samuel AR, Knowles NJ, Kitching RP. Serological and biochemical analysis of some recent type A foot-and-mouth disease virus isolates from the Middle East. *Epidemiol Infect* 1988; **101**: 577–90.
25. Armstrong RM, Samuel AR, Carpenter WC, Rama Kant, Knowles NJ. A comparative study of serological and biochemical methods for strain differentiation of foot-and-mouth disease type A viruses. *Vet Microbiol* 1993; in press.
26. Krebs O, Marquardt O. Identification and characterisation of foot-and-mouth disease virus O Burgwedel/1987 as an intertypic recombinant. *J Gen Virol* 1992; **73**: 613–19.
27. Marquardt O. Methods that identify recombinant FMDV strains. Report of the Session of the Research Group of the Standing Committee of the European Commission for the Control of Foot-and-Mouth Disease, Berne, Switzerland. Rome: FAO, 1992.
28. Amaral-Doel CMF, Owen NE, Ferris NP, Kitching RP, Doel TR. Detection of foot-and-mouth disease viral sequences in clinical specimens and ethyleneimine-inactivated preparations by the polymerase chain reaction. *Vaccine* 1993; **11**: 415–21.