

The *Caenorhabditis elegans* *unc-93* Gene Encodes a Putative Transmembrane Protein That Regulates Muscle Contraction

Joshua Z. Levin and H. Robert Horvitz

Howard Hughes Medical Institute, Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Abstract. *unc-93* is one of a set of five interacting genes involved in the regulation or coordination of muscle contraction in *Caenorhabditis elegans*. Rare altered-function alleles of *unc-93* result in sluggish movement and a characteristic "rubber band" uncoordinated phenotype. By contrast, null alleles cause no visibly abnormal phenotype, presumably as a consequence of the functional redundancy of *unc-93*. To understand better the role of *unc-93* in regulating muscle contraction, we have cloned and molecularly characterized this gene. We isolated transposon-insertion alleles and used them to identify the region of DNA encoding the *unc-93* protein. Two *unc-93* proteins differing at their NH₂ termini are potentially encoded by transcripts that differ at their 5' ends. The putative *unc-93* proteins are 700 and 705 amino acids in length

and have two distinct regions: the NH₂ terminal portion of 240 or 245 amino acids is extremely hydrophilic, whereas the rest of the protein has multiple potential membrane-spanning domains. The *unc-93* transcripts are low in abundance and the *unc-93* gene displays weak codon usage bias, suggesting that the *unc-93* protein is relatively rare. The *unc-93* protein has no sequence similarity to proteins listed in current databases. Thus, *unc-93* is likely to encode a novel membrane-associated muscle protein. We discuss possible roles for the *unc-93* protein either as a component of an ion transport system involved in excitation-contraction coupling in muscle or in coordinating muscle contraction between muscle cells by affecting the functioning of gap junctions.

ANIMAL behavior is produced through a complex series of events that result in specific muscle contractions and relaxations. Neuronal inputs to a muscle cell can cause membrane excitation, which is coupled to the mechanical sliding of the thick and thin filaments of the myofilament lattice (Shepherd, 1988). Although much is known about the structure and function of the myofilaments (Leavis and Gergely, 1984; Kamm and Stull, 1985; Wang, 1985; Warrick and Spudich, 1987) and of some of the proteins involved in excitation-contraction coupling (Catterall et al., 1988; Fill and Coronado, 1988; Jan and Jan, 1989), a complete description of the components of this signal transduction pathway has not yet been achieved.

The nematode *Caenorhabditis elegans* is particularly appropriate for the study of muscle, as genetic, biochemical, and morphological analyses of *C. elegans* muscle are all straightforward. More than 30 genes have been identified by mutations that affect muscle structure and function in *C. elegans* (reviewed by Waterston (1988)). Some of these genes encode structural components of the myofilaments, such as myosin heavy chains (Epstein et al., 1974; MacLeod et al., 1977, 1981; Waterston, 1989), actin (Landel et al. 1984), and paramyosin (Waterston et al., 1977; Kagawa et al., 1989). Mutations in other genes severely decrease motility with comparatively minor structural defects in the myofilament lattice; these mutations presumably disrupt the regula-

tion or coordination of muscle contraction. For example, mutations in the gene *unc-22* result in an uncontrolled, almost incessant, twitching of the body-wall muscles, which suggests a regulatory defect (Brenner, 1974; Waterston et al., 1980). *unc-22* encodes a huge 750-kD protein, named twitchin, with a protein kinase domain and several copies of two motifs found in members of the immunoglobulin superfamily, myosin light chain kinase, and titin (Benian et al., 1989). Both genetic and molecular studies indicate that *unc-22* interacts with the *unc-54* myosin heavy chain (Moerman et al., 1982; Mori et al., 1988), which suggests that *unc-22* acts in conjunction with myosin at the end of the excitation-contraction coupling signal transduction pathway.

unc-93 is one of a set of five interacting genes (*unc-93*, *sup-9*, *sup-10*, *sup-11*, and *sup-18*) involved in muscle structure and function (Greenwald and Horvitz, 1980, 1982, 1986). *unc-93(e1500)*, *unc-93(n200)*, and *sup-10(n983)* are rare altered-function mutations that confer a distinctive abnormality in the regulation of *C. elegans* muscle contraction termed "rubber band" (Greenwald and Horvitz, 1980, 1986). When a rubber band mutant is touched on its head, the animal contracts and then quickly relaxes without moving backwards, whereas a wild-type worm simply moves backwards. Thus, rubber band mutants can contract their body wall muscles, but the regulation or coordination of the contraction is defective. These mutants are sluggish and flaccid, character-

istics typical of *C. elegans* muscle mutants (Waterston et al., 1980), but have only minor structural defects in their body wall muscles (Greenwald and Horvitz, 1980; Waterston, 1988). Rubber band mutants are defective in egg laying, which indicates that there are abnormalities in the vulval and uterine muscles as well as in the body wall muscles (Greenwald and Horvitz, 1980, 1986). Genetic mosaic analysis has shown that *sup-10* functions within muscle cells (Herman, 1984).

The rubber band mutations do not eliminate gene activity, but rather produce abnormal protein products that disrupt muscle function (Greenwald and Horvitz, 1980, 1986). Loss-of-function, or null, mutations in *unc-93* or *sup-10* confer no visibly abnormal phenotype, presumably because these two genes are functionally redundant with another gene (or set of genes) that has sufficient overlap with *unc-93* and *sup-10* in regulating muscle contraction so as to maintain normal muscle function. Null mutations in *unc-93*, *sup-10*, *sup-9*, and *sup-18* were identified as recessive suppressors of the rubber band mutations (Greenwald and Horvitz, 1980, 1986). Further evidence that *unc-93* is likely to function in muscle is derived from this mutual suppression of *unc-93* and *sup-10* (Greenwald and Horvitz, 1980, 1986), which suggests that the two genes function in the same cell. Based upon previous detailed genetic analyses of these four genes, our model for their action is as follows. First, these genes function together to regulate muscle contraction, and their protein products interact with each other, possibly as a single protein complex. Second, the rubber band mutations produce aberrant gene products that cause defects in the regulation of muscle contraction. Third, the absence of the protein products of these genes does not adversely affect muscle contraction because these genes control a process that is functionally redundant with another process that regulates muscle contraction.

To understand how *unc-93* and the other functionally related genes regulate muscle contraction, we cloned *unc-93* using transposon tagging. In addition, we molecularly characterized the *unc-93* gene and identified the DNA sequence alterations in the rubber band mutations.

Materials and Methods

General Methods and Strains

General methods for the handling and culturing of *C. elegans* strains have been described by Brenner (1974). *C. elegans* was grown at 20°C, except in experiments with *daf-2(e1370ts)*, in which strains were maintained at 15°C and grown at 22.5° or 25°C to observe the Daf phenotype (Riddle et al., 1981; Swanson and Riddle, 1981). *C. elegans* variety Bristol strain N2 is the wild-type parent of all strains used in this work, except as described for the strains isolated in the *mut-2* genetic background (see below). The genetic markers used were as follows (unless otherwise noted, mutations are described by Hodgkin et al. [1988]): LGI: *unc-13(e51)*; LGII: *sup-9(n180)* and the new *sup-9* alleles identified in this work (*n1330*, *n1414*, *n1421*, *n1424*, *n1426*, *n1428*, *n1430*, *n1469*); LGIII: *daf-2(e1370)*, *unc-93(e1500, n200, e1500 n234, e1500 n243, e1500 n244, e1500 n246, e1500 n248, e1500 n254, e1500 n255)* and the new *unc-93* alleles identified in this work (*e1500 e2128* [M. Shen, personal communication]), *e1500 n1412*, *e1500 n1415*, *e1500 n1418*, *e1500 n1419*, *e1500 n1420*, *e1500 n1422*, *e1500 n1423*, *e1500 n1425*, *e1500 n1427*, *e1500 n1429*, *e1500 n1431*, *n1470*, *n1474*, *n1500*, *n1623* and *n1624* [D. Parry, personal communication], *e1500 n1907* and *e1500 n1912* [J. Thomas, personal communication]), *dpy-17(e164)*, *sup-18(n1010)* and the new *sup-18* allele identified in this work (*n1539*); LGV: *him-5(e1490)*; LGX: *sup-10(n183, n983)* and the new *sup-10* alleles identi-

fied in this work (*e2127*, *e2130*, and *e2131* [M. Shen, personal communication]) and *n1413* and *n983 n1468*).

Isolation of Gamma Ray-induced Alleles

We mutagenized *sup-10(n983)* L4 hermaphrodites on Petri plates with gamma rays using a dose of 7,500 rads from a ⁶⁰Co source, as previously described by Greenwald and Horvitz (1980), and screened the F2 progeny for wild-type revertants. In the first experiment, *unc-93(n1470)* and *unc-93(n1474)* were isolated from about 12,000 haploid genomes screened. In a similar gamma-ray mutagenesis of *sup-10(n983)* animals, *unc-93(n1623)*, and *unc-93(n1624)* were isolated (D. Parry, personal communication). *unc-93(e1500 n1907)* and *unc-93(e1500 n1912)* were isolated as wild-type revertants among the F2 progeny of gamma ray-irradiated *unc-93(e1500)* hermaphrodites (J. Thomas, personal communication).

Isolation and Characterization of Putative Transposon-insertion Alleles

To isolate transposon-insertion alleles of *unc-93*, we constructed strains containing *unc-93(e1500)* or *sup-10(n983)* in a *mut-2* mutator background. These constructions were done using MT2879, a three-times backcrossed *mut-2(r459)* strain derived from TR674 (Finney, 1987; Collins et al., 1987). Because the *mut-2* activity in this strain is known to map very close to *unc-13* on LGI and because the males are sufficiently healthy to mate, this strain allows easy genetic manipulation of the *mut-2* activity (Finney, 1987). *mut-2* males were crossed with *unc-13*; *unc-93(e1500)* hermaphrodites, F1 cross-progeny of genotype *mut-2 +/+ unc-13*; *+unc-93* were picked, and F2 *Unc-93* non-*Unc-13* progeny that did not segregate *Unc-13* progeny (genotype *mut-2 +*; *unc-93*) were identified and used to establish a strain. This strain was screened for phenotypically wild-type revertants. The same strategy was used to construct a strain of genotype *mut-2(r459)*; *sup-10(n983)*. One attempt to construct these strains yielded the mutation *sup-9(n1330)* as a preexisting mutation in a four times backcrossed *mut-2(r459)* strain. We obtained a total of 26 additional suppressor mutations and showed that these mutations failed to complement alleles of known genes. Of the revertants, 18 were derived from the *mut-2*; *unc-93(e1500)* strain: *sup-9(n1414, n1421, n1424, n1426, n1428, n1430)*, *unc-93(e1500 n1412, e1500 n1415, e1500 n1418, e1500 n1419, e1500 n1420, e1500 n1422, e1500 n1423, e1500 n1425, e1500 n1427, e1500 n1429, e1500 n1431)*, and *sup-10(n1413)*. Four revertants were derived from the *mut-2*; *sup-10(n983)* strain: *sup-9(n1469)*, *unc-93(n1500)*, *sup-18(n1539)*, and *sup-10(n983 n1468)*. Four revertants were isolated from a strain with *unc-93(e1500)* in a mutator background derived from the mutator strains TR403, a wild isolate containing mutator activity, and TR679, which contains the *mut-2(r459)* mutation (M. Shen, personal communication; Collins et al., 1987): *unc-93(e1500 e2128)* and *sup-10(e2127, e2130, e2131)*. Because a transposon insertion event can occur in any generation, but the insertion will be detected only when homozygous, we isolated only one revertant from among the F2 progeny of any given worm in an attempt to insure the independence of the mutations. Among the *unc-93* alleles, only *unc-93(e1500 n1418)* and *unc-93(e1500 n1425)* could possibly be re-isolates of the same mutational event.

To remove the additional unlinked Tc1 transposons present in the original isolate of *unc-93(e1500 n1415)* and thereby allow the detection of additional transposons by genomic Southern blot analysis, we backcrossed the original isolate of the *unc-93(e1500 n1415)* mutation to N2 worms. We crossed N2 males with *unc-93(e1500 n1415)* hermaphrodites, mated the cross-progeny males (genotype *unc-93(e1500 n1415)/+*) with *unc-93(e1500) dpy-17* hermaphrodites, picked *Unc* non-*Dpy* hermaphrodites (genotype *unc-93(e1500)/dpy-17/unc-93(e1500 n1415) +*), and, from the self progeny of these worms, picked phenotypically wild-type hermaphrodites (genotype *unc-93(e1500 n1415)* now backcrossed with Bristol strains two times). We generated males of this twice backcrossed *unc-93(e1500 n1415)* strain by heat shock treatment (Hodgkin, 1983), crossed these males to *daf-2 dpy-17* hermaphrodites, and mated the cross-progeny males (genotype *daf-2 + dpy-17/+ unc-93 +*) to *daf-2 dpy-17* hermaphrodites. (*daf-2* maps 4.2 map units to the left of *unc-93* and *dpy-17* map 3.4 units to the right of *unc-93*) (Edgley and Riddle, 1990). By repeating this cross a total of eight times, we constructed a strain containing *unc-93(e1500 n1415)* that is congenic with the Bristol N2 strain except in the region around *unc-93*. In the last cross, we picked phenotypically wild-type cross-progeny (genotype *daf-2 + dpy-17/+ unc-93 +*) hermaphrodites instead of males, and from their self progeny picked wild-type hermaphrodites that did not segregate any *Daf* or *Dpy* progeny (genotype *+ unc-93 +/+ unc-93 +*). We confirmed the genotype of these worms by crossing them with *unc-93(e1500)*; *sup-10(n183)* males and observing only *Unc* cross-progeny males.

Molecular Biology

Standard techniques for molecular biology were used (Sambrook et al., 1989).

To clone the 6.7-kb *Tcl*-containing *EcoRI* fragment from the *unc-93(e1500 n1415)* strain, we purified *EcoRI* cut genomic DNA of approximately the desired size from an agarose gel, cloned the DNA into λ NM1149, screened the resulting clones by hybridization with a *Tcl* probe (*pTcl*, construction described in Finney et al., 1988; Emmons, 1983) and identified a *Tcl*-positive clone with a 6.7-kb insert. We subcloned the 6.7 *EcoRI* insert in this phage, ϕ 93-1, into *pBS+* (Stratagene, La Jolla, CA) to construct *p93-1*. By cutting *p93-1* with *EcoRV*, which cleaves very close to both ends of *Tcl*, and religating the DNA, we removed the *Tcl* DNA along with 183 bp (#2590-2773) of *unc-93* DNA to construct *p93-2*. We used *p93-2* to probe the "JA N2" lambda library containing *C. elegans* wild-type genomic DNA (kindly provided by A. Coulson and J. Sulston) to isolate and characterize two overlapping phage clones, ϕ 93-2 and ϕ 93-3. These two phage clones were analyzed by A. Coulson and J. Sulston (personal communication) and placed on a contig, a set of overlapping DNA clones, that now spans ~10,000 kb of the physical map of LGIII (Coulson et al., 1986, 1988; and personal communication). To characterize the region of DNA we suspected to contain the *unc-93* gene, we subcloned a 6.3-kb *BamHI* fragment and the adjacent 3.5-kb *EcoRI*-*BamHI* fragment from ϕ 93-2 into *pBS+* to construct *p93-3* and *p93-11*, respectively (restriction map shown in Fig. 7).

We used *p93-2* to probe a mixed stage cDNA library (Kim and Horvitz, 1990) and isolated five positive clones from 165,000 clones screened. Two classes of positively hybridizing clones were isolated. Four clones are ~300 bp in length and hybridize to the 0.6-kb *HindIII* fragment shown on the left in Fig. 7. These clones are not likely to be related to *unc-93*, because they are derived from a region of DNA that is unchanged in *unc-93* mutants, except in complex rearrangements in which bona fide *unc-93* sequences are also rearranged (Fig. 7 and our unpublished data). One 2.6-kb clone is colocalized with the DNA polymorphisms observed in *unc-93* mutants and hybridizes to a transcript altered in animals carrying the *Tcl* insertion mutation *unc-93(e1500 n1415)* (Figs. 2 and 7). Thus, this cDNA is probably derived from the *unc-93* transcript. Based on Southern blot analysis and restriction enzyme mapping, this cDNA clone has an unrelated 400 bp of DNA at its 3' end after the polyA tail of the *unc-93* cDNA, probably as a result of the ligation of two unrelated fragments during the construction of the cDNA library. The cDNA was subcloned as two separate *EcoRI* fragments into *pBS+* to construct *p93-5* and *p93-6*—the former contains the 3' half of the cDNA clone and the latter contains the 5' half (see Fig. 7 for the *unc-93* cDNA position).

RNA was isolated as described by Kim and Horvitz (1990). Staged animals were obtained as described by Meyer and Casson (1986) and Kim and Horvitz (1990). The L1 larvae were not fed before they were harvested, and the effect of starvation on *unc-93* RNA levels is not known. Northern blot analysis was performed as described by Meyer and Casson (1986) and modified by Miller et al. (1988).

The cDNA sequence is derived from clones *p93-5* and *p93-6*, and the genomic sequence is derived from clones *p93-3* and *p93-11*. For DNA sequencing, we constructed a series of nested deletions by the method of Henikoff (1984) and also used synthetic oligonucleotides derived from the *unc-93* sequence as primers. The sequencing reactions were done by the method of Sanger et al. (1977) with double-stranded DNA templates using the USB Sequenase kit (U.S. Biochemicals, Cleveland, OH) according to its instructions. All DNA sequences were determined for both strands, except for bases 1 to 41 and 5018 to 5055. The sequence of the *unc-93* cDNA and the genomic DNA are in full agreement, except at position 2963 (C in the genomic sequence and T in the cDNA sequence); because this cDNA sequence corresponds to a TAA stop codon in frame within the open reading frame, the T in the cDNA sequence is likely to be an artifact of the cloning process.

Primer extension, ribonuclease protection, and RNA polymerase chain reaction (PCR)¹ experiments were performed to determine the 5' ends of the *unc-93* transcripts. For primer extension (Kingston, 1991), a ³²P end-labeled oligonucleotide (positions 542-513) was used as a primer with 50 μ g of total RNA from L1 larvae or eggs. The sizes of the resulting DNA products were determined by comparison with the sizes of the products of DNA sequencing reactions in adjacent lanes on a 6% polyacrylamide/urea (sequencing) gel. For ribonuclease protection experiments, a ³²P-labeled anti-sense RNA probe from positions 549-283 was synthesized using the T3

promoter to express a *SspI*-digested deletion subclone of *p93-11* missing DNA from positions 550-1336, according to the directions included with a RNA transcription kit (Stratagene, La Jolla, CA). The labeled probe (between 0.03 and 3 ng) was hybridized to 25 μ g of total RNA from L1 larvae or eggs and then digested with ribonucleases A and T1 (Gilman, 1991). The sizes of the products were determined by comparison with the sizes of the products of DNA sequencing reactions and ³²P-labeled T3 RNA transcripts of known sizes on a 6% polyacrylamide/urea gel. Ribonuclease protection products that were present when yeast RNA was substituted for worm total RNA were considered to be unrelated to *unc-93* RNA. RNA PCR was carried out with the reagents in the GeneAmp ThermoStable rTth Reverse Transcriptase RNA PCR kit (Perkin Elmer Cetus, Norwalk, CT) according to the instructions included. Reverse transcriptase reactions with 50 ng of polyA⁺ egg RNA and an oligonucleotide primer were incubated at room temperature for 5 min, 60°C for 5 min and 70°C for 10 min. Reagents for PCR including the other oligonucleotide primer were added to these reactions and PCR was performed as follows: annealing at 55°C for 1 min, extension at 72°C for 1 min for the first 20 cycles and 2 min for the last 20 cycles, and denaturation for 1 min at 92°C, all for 40 cycles. With one oligonucleotide from exon 4 (positions 1313-1294) and the other in exon 1 upstream of the cDNA start site (positions 388-408), an ~630-bp fragment was generated from the larger *unc-93* transcript beginning at position 233. This fragment contained a single *PvuII* site as expected at position 700. In addition, a second round of PCR amplification with the same 5' end primer and either of two primers from exon 2 (positions 660-644 and 729-713) yielded products of ~220 and 290 bp, as expected. RNA PCR with a primer from either SL1 (positions 2-22) (Krause and Hirsh, 1987) or SL2 (positions 1-22) (Huang and Hirsh, 1989) and a primer from exon 4 (positions 1313-1294 or 1293-1279) yielded multiple fragments, as is often observed for RNA PCR with SL1 and SL2 (M. Nonet, personal communication). A second round of PCR amplification with the same 5' end primer and either of the two exon 2 primers yielded a single product for each reaction of the size expected for a *trans*-spliced leader added at position 497.

DNA sequence analysis was done using the DNA Inspector program (Textco, West Lebanon, NH), the DNA Strider program (Marck, 1988), and the University of Wisconsin GCG package (Devereux et al., 1984). We searched the Genpept(R) protein database (GenBank database, v67.0 and new sequences through 3/91) with the *fasta* program and the combined GenBank/EMBL DNA database (GenBank v68.0, EMBL v26.0) with the *tfasta* program. In addition, the BLAST program (available through NCBI) was used to search a combined database of PIR (v30.0), SwissProt (v19.0), GenPept (v69.0 and new sequences through 12/3/91) (Altschul et al., 1990). None of these searches yielded any proteins with significant sequence similarity to the *unc-93* protein. No significant sequence similarity was found between DNA sequences in the GenBank/EMBL DNA database and the *unc-93* genomic DNA sequence using the *fasta* program. In addition, the sequences in intron 5 and intron 10, containing repeated sequences 3 and 2, respectively (see Fig. 3), were each used to search the GenBank/EMBL DNA database. For intron 10, the eight nucleotide repeat is similar to repeats of two and four nucleotides found in many other genes, but we do not know of any function for these repeats. Intron 5 did not show any significant sequence similarity to any DNA sequences. The *naq/match* program was used to search for *C. elegans* genes with multiple copies of the three repeated sequences. These sequences are found in several genes, but never tandemly repeated or even repeated within a 200-bp interval. For these short sequences, a single copy seems likely to be a random occurrence.

We used Southern blot analysis and PCR amplification (Saiki et al., 1988) to map the locations of *unc-93* mutations. PCR amplification reactions used the *AmpliTaq* polymerase (Perkin-Elmer Cetus, Norwalk, CT) according to the instructions supplied with the enzyme. Conditions for the PCR reactions were: annealing at 50°C for 1 min, extension at 72°C for 3 min for the first 20 cycles and 4 min for the last 10 cycles, and denaturation for 1 min at 92°C, all for 30 cycles. For the PCR amplification experiments, we used sets of primers that covered the entire sequenced region of *unc-93* except for 30 bp at the 5' end, although all combinations of primers were not tested for each mutation. The *unc-93* primers were derived from the following sequences: positions 30 to 55, 388 to 408, 505 to 525, 514 to 533, 1293 to 1274, 1298 to 1279, 1313 to 1294, 2355 to 2379, 2413 to 2392, and 5055 to 5036. The two *Tcl* primers chosen were derived from sequences not in the inverted terminal repeats—positions 105 to 83 and 1396 to 1417 (Rosenzweig et al., 1983). DNA from all of the *Tcl* insertion alleles was PCR-amplified with primers from either end of *Tcl* and the PCR products were checked by digestion with *EcoRV*.

The DNA sequence alterations in strains containing the *unc-93(e1500 n234)* and *unc-93(n200)* mutations were determined from the products of PCR amplification. Ethyl methanesulfonate was used to generate the *e1500*

1. Abbreviation used in this paper: PCR, polymerase chain reaction.

and *n200* mutations and diethyl sulfate was used to generate *n234* (Greenwald and Horvitz, 1980). Genomic DNA was amplified with two sets of primers that spanned the *unc-93* region. The first set of primers was derived from positions 30 to 55 and 2413 to 2392; the second set of primers was derived from positions 2355 to 2379 and 5055 to 5036. The PCR reactions were done as described above. To avoid possible DNA sequence changes introduced by the Taq polymerase, we pooled the products of 10 separate PCR reactions for each of the PCR products. These pools were subcloned into pBS+, and at least four isolates of each of the PCR products were combined for use as templates in DNA sequencing reactions using oligonucleotide primers covering the *unc-93* region. We determined the DNA sequence of these subclones from position 196 to 5055, except for intron 14 from position 4264 to 4330. We ignored DNA sequence alterations that did not appear in all isolates of a particular subclone.

Results

Isolation of Transposon-insertion Alleles

We used the method of transposon tagging (Greenwald, 1985; Moerman et al., 1986) to clone the *unc-93* gene. *C. elegans* strains containing mutator mutations display elevated levels of transposition of the transposable elements Tc1, Tc3, Tc4, and Tc5 (Collins et al., 1987; Finney et al., 1988; Collins et al., 1989; Yuan et al., 1991; J. Collins and P. Anderson, personal communication). We constructed strains carrying either the *unc-93(e1500)* or the *sup-10(n983)* mutation in a mutator background. Because null alleles of *unc-93*, *sup-9*, *sup-10*, and *sup-18* can suppress the rubber band phenotype caused by these mutations, a worm carrying a transposon insertion in one of these four genes can be identified as a phenotypically wild-type revertant. In this way, 27 suppressor mutations that included alleles of all four genes were isolated: 13 alleles of *unc-93*, eight of *sup-9*, five of *sup-10*, and one of *sup-18* (Table I). Since *sup-18* mutations only partially suppress *e1500* but completely suppress *n983* (Greenwald and Horvitz, 1986), *sup-18* mutations would have been detected in these experiments only as suppressors of *n983*.

Identification of a Tc1 Insertion in the *unc-93* Gene

These putative transposon-insertion mutations were isolated in mutator backgrounds, which have as many as several hundred copies of Tc1 (Emmons et al., 1983; Collins et al.,

1987). To identify an insertion in *unc-93*, these mutations were backcrossed into the wild-type Bristol (N2) background (see Materials and Methods), which contains 30 copies of Tc1. We used Tc1 to probe a Southern blot containing genomic DNA from a 10 times backcrossed strain containing the *mut-2*-derived allele *unc-93(e1500 n1415)* and detected one extra Tc1-hybridizing band (data not shown). This Tc1 insertion mapped within about 0.8 map units of the *unc-93* locus, since 26 of 26 recombination events within an ~7 map unit interval spanning *unc-93* failed to separate the Tc1 from the *unc-93* gene (Table II). Thus, this Tc1 inserted in or very near the *unc-93* gene. We cloned a 6.7-kb EcoRI genomic fragment containing this Tc1 insertion to yield the plasmid p93-1. By removing the Tc1 element from p93-1, we constructed the plasmid p93-2, which contains the genomic DNA flanking this Tc1. When p93-2 was used to probe a Southern blot containing genomic DNA, the wild-type N2 strain contained a 5.1-kb EcoRI fragment, whereas *unc-93(e1500 n1415)* contained the 6.7-kb EcoRI fragment, as expected for a 1.6-kb Tc1 insertion into this fragment in the *unc-93(e1500 n1415)* strain (Fig. 1).

Identification of Polymorphisms Associated with *unc-93* Alleles

To obtain clones with wild-type *unc-93* genomic DNA, we used p93-2 to probe a lambda library containing genomic DNA from the wild-type N2 strain. We isolated two overlapping phage clones containing genomic DNA from the region around *unc-93*, based on common restriction maps and genomic Southern blots (data not shown). To identify the site of the *unc-93* gene, we examined DNA from *unc-93* mutants generated by gamma rays for allele-specific polymorphisms. These types of alleles are likely to have polymorphisms detectable by Southern blot analysis (Graf and Chasin, 1982; Grosovsky et al., 1986; Moerman et al., 1986; Collins et al., 1987). Using Southern blot analysis, we identified five gamma ray-induced *unc-93* mutations that alter the wild-type 5.1-kb EcoRI fragment, in addition to the Tc1 insertion in *unc-93(e1500 n1415)*; three of these changes are shown in

Table I. Genetic Screen Used to Isolate Putative Transposon-insertion Alleles

	Phenotype	Genotype*	Number of alleles†
Initial strain	Unc	<i>unc-93(e1500)</i>	
Revertant strains	WT	<i>unc-93(0)</i>	12
	WT	<i>sup-9(0); unc-93(e1500)</i>	6
	WT	<i>unc-93(e1500); sup-10(0)</i>	4
Initial strain	Unc	<i>sup-10(n983)</i>	
Revertant strains	WT	<i>sup-10(0)</i>	1
	WT	<i>sup-9(0); sup-10(n983)</i>	1
	WT	<i>unc-93(0); sup-10(n983)</i>	1
	WT	<i>sup-18(0); sup-10(n983)</i>	1

We constructed strains with *unc-93(e1500)* or *sup-10(n983)* in a mutator background and isolated spontaneous non-Unc revertants, which proved to be of the genotypes indicated.

* Each strain also contains a mutator background (see Materials and Methods).

† One additional suppressor mutation, *sup-9(n1330)*, was isolated from the background of a *mut-2* parental strain (see Materials and Methods).

Table II. Mapping of the *unc-93(e1500 n1415)* Tc1 Insertion

Genotype of parent	Phenotype of recombinant	Genotype of homozygous recombinant	Tc1 present/total
<i>daf-2 + dpy-17</i>	Daf non-Dpy	<i>daf-2 unc-93</i>	6/6
+ <i>unc-93 +</i>		<i>daf-2</i>	0/8
	Dpy non-Daf	<i>unc-93 dpy-17</i>	3/3
		<i>dpy-17</i>	0/9

To map the Tc1 insertion in the ten-times backcrossed *unc-93(e1500 n1415)* strain, we identified recombination events in the interval between *daf-2* and *dpy-17*. The distances between *daf-2* and *unc-93* and between *unc-93* and *dpy-17* are 4.2 map units and 3.4 map units, respectively (Edgley and Riddle, 1990). We isolated Dpy non-Daf and Daf non-Dpy recombinants from the F1 progeny of *daf-2 + dpy-17/ + unc-93 +* heterozygotes. Animals homozygous for the recombinant chromosomes were picked in the F2 generation and tested for the presence of *unc-93(e1500 n1415)* by mating them with *sup-9(n180)*; *unc-93(e1500)* males or *unc-93(e1500)*; *sup-10(n183)* males and observing either all Unc cross-progeny (F2 genotype *unc-93(e1500 n1415)*) or all non-Unc cross-progeny (F2 genotype *unc-93(+)*). The presence of the Tc1 insertion in the homozygous recombinants was determined by Southern blot analysis using pTc1 as a probe.




Figure 1. *unc-93* Southern blot. EcoRI-digested DNA from the wild-type N2 and *unc-93* mutant strains and from the phage clone, $\phi 93-1$, was electrophoretically separated on an agarose gel, transferred to a nylon membrane, and probed with ^{32}P -labeled p93-2. The digested N2 DNA contains a 5.1-kb fragment and the *unc-93(e1500 n1415)* DNA contains a 6.7-kb fragment, consistent with the insertion of a 1.6-kb TcI element. The phage clone $\phi 93-1$ contains an insert of the 6.7-kb fragment from *e1500 n1415*. Three other *unc-93* strains show polymorphisms: *e1500 n244* has a 6.9-kb fragment, *e1500 n248* has the wild-type 5.1-kb fragment as well as a 5.8-kb band—a pattern consistent with a tandem duplication—and *e1500 n254* has a 4.3-kb fragment.

Fig. 1. (See below for detailed physical mapping of *unc-93* mutations.) These data suggest that part of the *unc-93* gene is contained in the 5.1-kb EcoRI fragment.

Identification and Characterization of the *unc-93* Transcript

Using p93-2 to probe a mixed-stage cDNA library, we isolated one candidate *unc-93* cDNA clone from 165,000 clones screened. We used p93-6, a subclone containing the 5' half of this cDNA, to probe a Southern blot containing wild-type genomic DNA and detected only DNA from the same genomic region detected by p93-2 (data not shown). Thus, the cDNA is derived from single-copy DNA in the *unc-93* region. This cDNA detects a single 2.2-kb transcript when used to probe a Northern blot of polyA⁺ RNA from the wild type (Fig. 2). In animals carrying the TcI insertion mutation *unc-93(e1500 n1415)*, a transcript larger than the 3.5-kb ribosomal RNA band is detected instead (Fig. 2 a), providing further support that the 2.2-kb RNA is the *unc-93* transcript.

In a population of worms of mixed stages, the *unc-93* transcript is present at a level roughly 100- to 250-fold lower than that of the *unc-54* transcript, which encodes the major body wall myosin heavy chain (MacLeod et al., 1981) (Fig. 2 a).

The low abundance of the RNA transcript suggests that the *unc-93* protein is also present at low levels. The *unc-93* transcript accumulates at a higher level in L1 larvae than in eggs or in a mixed stage population consisting primarily of adults by weight (Fig. 2 b). This pattern of expression is consistent with the phenotype of *unc-93(e1500)* and *unc-93(n200)* rubber band mutants, which we observed to be most uncoordinated as L1 larvae and to move progressively better as they grow older.

unc-93 cDNA and Genomic DNA Sequences

We determined the sequence of the *unc-93* cDNA and identified one long open reading frame (Fig. 3). In addition, we determined the sequence of 5055 bp of genomic DNA encompassing the *unc-93* cDNA (Fig. 3 a). A comparison of the two DNA sequences yields the exon/intron boundaries for the *unc-93* gene and defines 15 introns ranging in size from 45 to 605 bp. As the *unc-93* cDNA clone is 2168-bp long and the *unc-93* transcript is 2.2-kb long, this cDNA corresponds to very nearly all of the full-length mRNA transcript. The *unc-93* open reading frame ends in a TGA stop codon that is closely followed by an AATAAA polyadenylation signal (Proudfoot and Brownlee, 1976) and the site for the addition of a polyA tail.




Figure 2. *unc-93* Northern blot analysis. (a) Identification of the *unc-93* transcript. Each lane contains polyA⁺ RNA purified from ~1.7 mg of total RNA isolated from a mixed-stage culture of worms. A transcript of 2.2 kb is detected in the wild type (N2) lane. In the *unc-93(e1500 n1415)* lane, there is a transcript larger than the 3.5-kb ribosomal RNA band instead. This blot was probed first with ³²P-labeled p93-6 to detect the *unc-93* transcript and then with a ³²P-labeled 1.0-kb EcoRI-BamHI fragment from the *unc-54* plasmid pMRF (kindly provided by D. Hsu and A. Fire, personal communication). *unc-54* encodes the major body wall myosin heavy chain (MacLeod et al., 1981). Based on the relative intensities and lengths of exposure of the *unc-93* and *unc-54* bands, we estimate that the


To define the 5' end of the *unc-93* transcript(s), we performed primer extension, ribonuclease protection, and RNA PCR experiments (see Materials and Methods). Primer extension from an oligonucleotide primer in exon 1 (positions 542-513) yielded two products: an approximately 310 nucleotide product that could correspond to a 2432-bp transcript beginning at position 233 and a 69 nucleotide product that could correspond to a 2191-bp transcript including a *trans*-spliced leader of 22 nucleotides added at position 496 (Figs. 3 and 4). Both SL1 and SL2 *trans*-spliced leaders are 22 nucleotides in length (Krause and Hirsh, 1987; Huang and Hirsh, 1989). The 3' acceptor splice site at position 496 (TTTCAGA) matches the consensus for 3' acceptor splice sites (TTTCAGY) in *C. elegans* (Emmons, 1988). We also identified two products in a ribonuclease protection experiment with an RNA probe derived from positions 549-283. An approximately 260 nucleotide fragment that resulted from protection of the entire *unc-93* portion of the probe corresponds to the larger transcript and an approximately 53 nucleotide fragment that would result from a splicing event at position 496 corresponds to the smaller *trans*-spliced transcript (data not shown). Position 233 is likely to be the 5' end of the larger transcript because there are no potential 3' acceptor splice sites (Emmons, 1988) between the end of the ribonuclease protection probe (position 283) and the predicted 5' end of the larger transcript (position 233) (Fig. 3 a). RNA PCR experiments with a primer in exon 4 (positions 1313-1294) and a primer upstream of the *trans*-splice site acceptor (positions 388-410) yielded a product of about

unc-93 transcript is present at a level roughly 100- to 250-fold less than that of the *unc-54* transcript. (b) *unc-93* expression at different developmental stages. Northern blot analysis was performed essentially as in a, except that each lane contained polyA⁺ RNA purified from ~1 mg of total RNA. Because of variable yields in purification of polyA⁺ RNA, the amounts of RNA loaded in the different lanes might not be equal. Relative to the *unc-54* transcript, the *unc-93* transcript appears to be more abundant in RNA from L1 larvae than in RNA from eggs and in RNA from a mixed stage population. The 2.2-kb size of the *unc-93* transcript was determined by comparison with the sizes of ethidium bromide-stained ribosomal RNA bands of 1.75 and 3.5 kb (Files and Hirsh, 1981) in parallel lanes of total worm RNA and with a BRL 0.24-9.5 kb RNA ladder (Bethesda Research Laboratory, Gaithersburg, MD).

Figure 3. *unc-93* sequence and structure. (a) *unc-93* nucleotide sequence and predicted amino acid sequence. The top numbers refer to amino acids, and the bottom numbers refer to nucleotides. The exon and intron boundaries are based on the comparisons between sequences of the *unc-93* cDNA and genomic DNA. The conserved GT and AG dinucleotides of splice junctions are underlined, as is the polyadenylation signal at position 4914 to 4919. The beginning of the cDNA (base 497) and the end of the cDNA (base 4932) are indicated by a © beneath the appropriate nucleotides. The 5' end of the larger transcript (base 233) and the 3' splice acceptor site for the *trans*-splice leaders (base 496) are indicated by a → beneath the appropriate nucleotide. A diamond beneath positions 2773 to 2774 indicates the Tc1 insertion site in *unc-93(e1500 n1415)* animals. A heart beneath positions 2960, 3689, 681, and 3773 indicates the locations of the *e1500*, *n234*, and *n200* (two changes) mutations, respectively. Repeated sequence 2 is indicated by arrows beneath the DNA sequences in intron 10 (positions 3190 to 3311), and repeated sequence 3 is similarly marked in intron 5 (positions 1575 to 2179). The orientations of repeated sequence 3 in intron 5 are indicated: an arrow to the right corresponds to the orientation shown in Fig. 5 a, and an arrow to the left corresponds to the inverse orientation. These sequence data are available from EMBL/GenBank/DBJ under accession number X64415. (b) *unc-93* gene structure. The top line shows the exons as thick lines and the introns between them. The extended hollow line for exon 1 represents the larger transcript. The small thick line above the top line connected to exon 1 at position 496 represents the *trans*-spliced transcripts. The location of repeated sequence 2 in intron 10 and repeated sequence 3 in intron 5 is also shown. The bottom line shows the two regions of the *unc-93* protein. Amino acid 1 is the NH₂ terminus of the protein encoded by the larger transcript and amino acid 6 is the NH₂ terminus of the protein encoded by the *trans*-spliced transcripts. The location in the protein of the mutations *e1500*, *n200*, and *n234* are indicated by arrows. *n200* has two arrows because two sequence changes were detected.

A

```
GATTTTATATGAATTTTGGAGATTTTTGACATTTCGCACATTTTCCGGAAAACTTGCTATTTTCAAATCTTAAAAAGACACAGAAAATTTAAATTTATATTAATATGCATTTTCC 120
TGGATTTTGAACGCTTTTTAGTTATTTTATTCATTAATAAATAATGAGATTTTGCACGATGATTTTGTGAAATTCGAAATTAATTTGTGGCTTTTCTGTTTTTTTTGGAAAAAGTTTFTA 240
CATTAAATATAATAATAAATAAATAAATTCAACATTTTCTGCTAATTTTATGAAAACTTTATGACAAAAAACCCTGTAATAAATTCGAAAAATTAGACTTTTACGATATAAAATTTAAAAAAA 360
CTGGTTTGTTCGGAGAGGAGTACACCGCGGATGCGCTTTGCGACTTTTGCQCGTAATCAAATTTGAATTTCCGCATAGTTGAATTTAGCGTATTCTTAGGGCAAAATTAACAGAAAAAA 480
  M K F Q K M G D S N T W D L V G E Q Q O R K K S R S P S R A S R V D S E          360
TGTATGAATTTTCAAGAAATGGGTGATTCAACACCTTGGGACTTTGGAGAGCAGCAGCAACGGAAGAAGTAGATACCGTCAAGAGCAAGCGTGTGGATTCCGAGTAATTTTA 600
AGCGCATCAATATGTACTAATTAAGGTGCATACACTTTCACTACTTGTGGTGTGTAAGACCGCGCTGAACCTTTGAAAGCTTTGGCCTAGGAAAAAGAGCAGCTGGAAAAAGAAAGCCAG 720
  R H K K Q R S R K L P A L N K L R K T S I H L L Q K F G A I P K K K D D S V L L F          102
ACGGCACAATAAAGAGCGGAGCAAGTCCGCGCTCAGAGAATAATTGAAAAAATTCGATTCATCTTTTGCAAAAATTTGGAGCGATTCCAAAAAGAAAGAGTAGATAGTGTTTTACTTTT 840
  R F H D I P D I P L E S L C I R              P K E F F E E          125
TAGATTTTCAATGATATTCCTGATATTCCTTGGAAATCTTATGATCAAGGTGAGAAAAAGATTGCAAAATTTATTTTGAATAATTCGAAAAATAATTTTCAAGCGGAAAGAGTTTTCGGAGA 960
  P K V F S F R D M G R E Q Q K A E V N E K K          147
GCCCAAATGATTTTCTTCGAGACGAGCAGCAGCAAGCGAGGTAAATGAGAAGTCTTAAAGGGTTTTGAATTTGCGGCTCAAATTCGGTGTTCATTTTCCAAAGCTC 1080
GAGCGCAAATGGCCGAGGATTTTTCTTGGCAGATAGAAAAACCAGTCTTCTTGGCTTCAAGTTGCTAAGCGCCGAAAAAAAACATCGGTGGCCGAGTTTTCCAATTTTCGGCCAC 1200
  S Y L F R G T S F D H D D H F E L P T E T G R V P E Y D H F C          178
GAATCAAAAATTCGAAATTTATTTTCACTTCTCGTGGCACCCTTACGACAGCAGCATCTTCCAAGCTTCCACTGAAACAGCCCTGTTCCAGAAATATGATCATTTTTTG 1320
  P I H G S R R R L V T M O T L M H S V D D E D N E D L A Y I Y G          215
CCCGATTCAATGATTCAGCGAAGACTTCCAGTAAATAATTTGGTCACAAATGCAAACTCAGTATCGGTAGATGATGAAAGATTAAGAGATTTGGCTTATATTTATGGGTTAGTTTT 1440
  H D F L A K L V R K K K R E M M S G T E K E R A N K I T          243
CAGAGAAATTTGAATGAAATTTTAAATCGGTTTTCAGGCACTTTCTGGCAATTTGTAGGAAAGAAAGCGAAGGAGATGATGAGTGAACGAAAAAGAAAGAAATTAAGATT 1560
  R K I M              247
AAAGGAAATTAATGGTAAATTTCAAACGTCGCTCGCAATGGGAAAATAGGCCACCACAGAAAAATAGGCCACCAGAGAAAAATAGGCCCGTAGAAAGATAAACTAGGCCACTAGAAC 1680
GATAAACAGTAGACCGCTTGAAAAACTAGGCCAGTCAAAAAAGCTACTACTATTGAAAAAAAATAGCACCTAGCAGAAAAAGCTAGGCCGTTGAAAAAAGACACTAGAAAAATACT 1800
AGGCAGTTTTTAAATTAATTTCAATTAATTAATGCAATTAATCAAGTTCAAAAATGCAAACTCGGCCACCGACTTTTTCGCGCCGCGGTGTTCAAAAACTAGAAAAAATAACAG 1920
CAAAGTCGGTGGCTAGTTTATTTGCTCTGCAAAATGACACATACGATTTGAAAAATGGCTGGGAAAAATCAGTGGCCTAGTTTTCCTTAGTGCCCTATAATTTCTCTAGCGCCCTAGT 2040
CTTGATTTGCTAGTGCCCTAGTTTTTTTTTCTTTCGGTGGCGGAGTTTCTCTCAATTAGGATATAGTAGGATATAAGTAGGCGGAGTTCCTGTTTCGCCACTTCAACTAGACCC 2160
  S N L W I L S V A L F L F T A F N G L Q N L Q T S V N G D L R          281
CTAATCGAATCAATTTTCAGTCAAACTCTCGGATCTGCGCTGGCTTTCTCTTCCGTCACAGCTATTAAGGACTTCCAAAATCTTCAAACTCCGCTCAATGAGAGTCTCGGATCTGA 2280
  S L V              306
TAGTTTTGTGTAAGAAGCTTTGAATTTCAATTTGAAAAATAATAATTTTTCAGGACCCTATATTTATCACATTGCCATATCATCATATTTTGTTCATCATTTTATGATNAATTCGCCCTGG 2400
  C K L T F L I A I F V Y F L Y I V I N L R P T          329
GTGCAATTAACATTTCTGCAATTTCTGATTTCTATACATTTGTCATTAATTTGAGCCGACCTAAGTTTTCGGTTTTCGAGCGCTTGTAAGCTAGAACTAAATAGTTT 2520
GCAAAAACTGCTCAACAAAATTAATTAATCGATTTTGATCTACATTTTCGATGTAGCAAACTTTTGTATATCTTTCACCTATAAGAAGTTATGTCATTTGAAGAGCTAGAGATATA 2640
  Y S S M I P A S I F C G I          342
AAAAATTTATGGCCATTTTTCCTGTTAGAACTTTTTCGTTTAGCATTGTGATGTCCTAGCTAGCGCTTTTCCAGATACCTTCTATGATCCCTGCTTCAATATCTCTGGGAA 2760
  A A S C I W G A K C A Y I T E M G I R Y A S L N F E S O T T V I V R          376
CGCGGCTCATGTTACTCGGGAGCAAAATGCGCTGACATTTACAGAAATGGAAATTCGATATGCTAGCCCTGAACCTTTGAAAGCTCAGACTACTGTTATTTGTTAGGTAATGGGGCTGATTTTCT 2880
  F F G V F F M I V H C G Q V V G N M V S S Y I F T          401
AAACTTTTTCAAGCTTTTCAGCTTTTAATCAAGCCAAATTCAGTATTTTCGGATATTTTTCATGATTTGATCACTGCGGACCAAGCTGTCGGAAAAATGATGATCTCTCTTTATTTAC 3000
  L S Y S Q A L R G P E D S I Y D S              C G Y Q F P K K N          426
ACTGCTTATTCGAACGCTCGGGGTCCTGAAGATTCTATATACGATAGCTTTTTCGCGAAATTTTTCGAAAGTTTAAATTTTCATTTTTCAGCTGTGGCTAGCCATTTTCAAAA 3120
  L S D L T E L A E S N L A R P P Q K V Y V A V          449
ATTTATCAGATTTTACAGGTTGGCGGAGAGCAATTTGCTGCGCCACCAAGAAAGTTTATGATAGAGTGGAGCCTACATACCTACATACCTACACTACACTACACTACA 3240
  C L A Y L A C V I S G M I M S          465
AACCTACATACCTACATACCTACATACCTACATACCTACCTACCTATACCCAAATTTTTTTTAAATTTGCTAGTGTTCGGCATACTCCGTGTTGTAATATCAGTGGAAATGATAATGAG 3360
  M F L N A L A K D A R N R              K M A O K F N S E I F          489
CATGTTCTGTAATGCTTTGGCAAAAGATGCTCGGAACAGGTTTTCCTACTTCCAGTCTTCTCTTAAATTTGGCTCTGAGGCTTTCAGAAAAATGGCTCAAAAGTTCAAACTCGGAAAAATTTTC 3480
  Y L M L K K H L I N I K F M L L V L P L T I F N G L E Q A F L V G V Y T K          524
TACCCTACGCTCAAACTAAATAACAATATAAAAGTTCATGTTTGGGTCGCTGACGATTTTCAATGTTTGGAGCAAGCTTTTCTAGTTGGAGTTTATACAAAAGCTTAGTGTTCAGCTGG 3600
CCGTTAAGAGGGTAAACTCGGCCAGTCACTAAATCGCAAAACTGAAAAATTTTCTTCCAGGACTTCGGATTCGGAGTGGCCTTTGGAAATTTGGCAAAATAGGCTTTGTGATGGCCTGCTTCGGA 544
I S D A V C S L V F G P L I K L F G R M P L F V F G A V V N L L M I V T L M          582
ATCTCCGACGAGTCTGTTTCCCTAGTCTCTGGGCACTGATCAAGCTTTTCGCTCGGATGCTCTTTCGTTTCGGAGCAGTAGTGAATCTATTAAATGATTTGATACACTTATGATGA 3840
  V W P L N A A D T Q I A F Y V V A A M W G M A D G V W N          609
TGCACCATATTTGACCGCAATTTTCAATCTCTCTCTCAGGTTTGGCCATAAATCGACCGACTCAAAATTTCTACGTTAGTCGCTGCAATGTGGGGTATGGCAGATGTGCTGGAAT 3960
T Q I N G F W V A L V G R Q S L O F A F T K Y R F W E S L G I A I G F A L I R          648
ACTGCAATTAATGAGTTTTTGGTTGCATTTGTTGAAGACGACTCTACAGTTTTCAGTTTCACTAATAATTCGATTTTGGAAATCTTTGGAAATCTGGAATTTGGATTTGGATTTGCTCTCGTGA 4080
ATTTTTTTTTTGGCTTTTGGCGCATAAAAATTTTATGGGGAGCAGTCCATTTTTGTAATCGATGTAAGAAAGCGCTCCACGGGCAATTTGAAAACGCTCCCGCCCTACAGTAGGGTCTCGT 4200
TAGGTAGTGTGGTGGGACCTTTGTAATTTCAACTTTTCAATTAGTTCGCGGATTTCCATGCAATTTTTCGTTTTTTTTTTGTTATATTTTTCCGTTCTTTGTAAGGATTTTTTCCGCGAAA 4320
TTGATGAAATTAAGTGGAAATTTAAAAATTTGAACTATTTTAACTAAAAAGTATTTTAAACATAAATAGTGGAAAAATGAGCAAAAAACATTTTATGAAAACAAAAAATCAATTTAAA 4440
ACGTTAAAATTCGCGTTTTTGGTCCCAATACCTAACGAGACCCAGCTTTTGGGGGAGGACATTGCATTTTACCGGTGGAGCACGCTTGCACTCAATCTGACTATTCCGATTTTTTTCG 4560
TCTCCATAAATGATGTCGGGTAAGAAAAAGAGGTTGCTTTTAAATTTCCGGCCAGCTAGATTTAACCTGAAAAAATAATGAATTTTTCAGCACAGCTCAGTGTGAAATA          654
Y L L I T F F M L L L G M C G F L A I E N F D H I I              H V T V E I          4680
TATTTGCTGATAACATCTCTCATGCTACTTTTAGAATGTGCGGATTTTTCAGCAATGAAAAATTTGATTCATTTTAGGCTGATTCGAAAGAAATTAATTTTTCATTTTAAAT 689
  K F W H H L I H T S C P E K E P L D D R N S D F E          705
TTTCCAGAAAATTTGGCATCACTTAAATCAACAGTCAATGTCAGAGAAGGAACCCTGAGTATGAGAAATTCAGATTTTGAAATGAAATAATTAATAAAATCAAAAAATCAAGTTAATAAAT 4920
TCGTACTTTATGTTTTTATTAATAAAGTCCAAAAAGTCAGAAAAATTAACAATAATGCAAAAAATAGAAAAATAGAAAAATTAATTAATAAAGTGGCGGGTGGTAAATGAGTTTCAAAAT 5040
GCTAGACTTTCGCTT 5055
```






Figure 7. Co-localization of *unc-93* mutations and the *unc-93* cDNA. The location of the cDNA is based on a comparison of the sequences of the cDNA and the genomic DNA. For the cDNA, the boxes correspond to exons and the arrowhead indicates the direction of transcription. The location of each mutation has been determined by Southern blot analysis and PCR amplification with sets of primers from the *unc-93* region (see Materials and Methods). For the TcI insertions, an arrow pointing to the right indicates that the TcI DNA has inserted with its reading frame in the same direction as that of *unc-93*, and an arrow pointing to the left indicates the opposite orientation. The positions of the TcI insertions are accurate to within ~50 bp. For the other mutations, the double-arrowed region indicates the interval in which the mutation lies. Any DNA that is not indicated as changed in a given mutation was shown to be wild-type at the gross level of agarose gel

electrophoresis. Complex rearrangements that were not localized precisely were found in three *unc-93* mutants: *e1500 n244*, *e1500 n254*, and *e1500 n255*. The p93-3 plasmid contains the only BamHI fragment shown, which is 6.3 kb, and the p93-11 plasmid contains the 3.5-kb EcoRI-BamHI fragment to the left of the 6.3-kb BamHI fragment. Additional EcoRI sites might be present to the right of the double break shown; the order of the rightmost HindIII and EcoRI sites has not been determined. (A) Aval; (B) BamHI; (H) HindIII; (R) EcoRI.

pairs of primers, one from TcI and the second from the flanking *unc-93* DNA (see Materials and Methods). Eleven of 13 mutator-derived *unc-93* alleles show polymorphisms, and nine of these are TcI insertions (Fig. 7). There are five different insertion sites, within a resolution of about 50 bp. At each insertion site, only one orientation of TcI insertion is found in these mutants. The other two mutator-derived alleles showing polymorphisms, *e1500 n1419* and *e1500 n1429*, are complex rearrangements. Six of twelve gamma ray-induced *unc-93* mutations show polymorphisms. Of the six gamma ray-induced mutations showing polymorphisms, five are complex rearrangements and one is an ~300-bp deletion. The identification of 17 *unc-93* alleles that show alterations in this region confirms that we have cloned the *unc-93* locus. The colocalization of the cDNA and the DNA polymorphisms found in *unc-93* mutants provide strong evidence that this cDNA corresponds to the *unc-93* transcript (Fig. 7).

DNA Sequence Alterations in *unc-93* Mutations

We determined the sequence of part of p93-1 and identified the site of the TcI insertion in *unc-93* (*e1500 n1415*) between bases 2773 and 2774 in exon 8 (Fig. 3). This TcI insertion caused a duplication of the TA dinucleotide on either side of

the insertion, as has been observed for other TcI insertions (Rosenzweig et al., 1983). The sequence of this insertion site (CATGTAATCT) is similar to the consensus sequences for TcI insertion sites—GA(G/T)(A/G)TA(T/C)(G/C)T and GA(T/G)ATAATGT—derived by Eide and Anderson (1988) and Mori et al. (1988), respectively.

To identify the DNA sequence changes in presumptive *unc-93* point mutants, we used PCR to amplify DNA from these mutants, cloned the PCR products into a plasmid vector, and determined the sequence of the *unc-93* region (see Materials and Methods). The *e1500* rubber band mutation has a G→A transition at base pair 2960 that changes amino acid 388 from Gly to Arg (Fig. 3). The *n200* rubber band mutation has a C→T transition at base pair 681 that changes amino acid 49 from Ala to Val and a G→T transversion at base pair 3773 that changes amino acid 562 from Gly to Val (Fig. 3); we do not know whether one or both of these DNA changes is responsible for the phenotype caused by *n200*. The *n234* mutation has a G→A transition at base pair 3689 that changes amino acid 534 from Trp to an amber stop codon (Fig. 3). Because the *n234* mutation has been shown to be suppressed by mutations in the tRNA amber suppressor gene *sup-7* (Greenwald and Horvitz, 1980), this sequence change further confirms the identity between the

unc-93 genetic locus and the DNA that we have cloned. This mutation also suggests that the COOH-terminal 170 amino acids are required for *unc-93* function.

Discussion

We cloned the *C. elegans* muscle gene *unc-93* by transposon tagging. *unc-93* is likely to encode a novel membrane-associated protein involved in muscle contraction. The putative transposon-insertion alleles of *sup-9*, *sup-10*, and *sup-18* generated in this study should facilitate the cloning of these genes. The continued molecular characterization of this set of interacting genes should provide us with additional insight into mechanisms that regulate muscle contraction.

Two aspects of *unc-93* gene structure are striking. First, the *unc-93* gene has 15 introns and produces a 2.2-kb mRNA transcript, which is an unusually high density of introns for a *C. elegans* gene (Emmons, 1988). *C. elegans* genes expressed at high levels, such as those that encode actins (Edwards and Wood, 1983) or myosin heavy chains (MacLeod et al., 1981), have a much lower density of introns. Perhaps *C. elegans* genes expressed at low levels can tolerate a large number of introns and any resulting inefficiencies in RNA splicing. The second intriguing aspect of *unc-93* gene structure is the three repeated sequences found in its introns. Some introns have enhancer elements composed of repeated DNA sequences that regulate gene expression (Atchison, 1988), and it is possible that the *unc-93* repeated sequences regulate *unc-93* expression. However, no role can be assigned to these repeats at present. It is noteworthy that the exons encoding the two distinct regions of the *unc-93* protein are separated by the largest intron of the *unc-93* gene. Intron 5, which contains 14 copies of repeated sequence 3 and could potentially form a stem-loop structure (Figs. 3 *b* and 5), might have joined the two distinct coding regions by a recombination event.

The rubber band phenotype caused by the altered-function *unc-93* alleles suggests a defect in the regulation or coordination of muscle contraction. The adult contains 95 mononucleate body wall muscle cells organized into four quadrants—two dorsal and two ventral (Sulston and Horvitz, 1977). Locomotion is achieved through the propagation of a wave of contraction and relaxation along the length of the worm, such that at a given time some of the dorsal muscle cells are contracted and the ventral muscle cells opposite them are relaxed, while adjacent dorsal cells are relaxed and the ventral cells opposite them are contracted (Chalfie and White, 1988). A defect in the muscle cells that disrupted the propagation of this wave of contraction and relaxation might result in a rubber band phenotype, in which both the anterior and posterior ends of the worm contract at the same time without any backwards movement. Such a defect might occur if the *unc-93* protein were localized within the muscle cell membrane and if *unc-93* rubber band mutations disrupted communication among muscle cells, perhaps by affecting gap junctions, which are known to connect body wall muscle cells within a quadrant (White et al., 1986). The effect of rubber band mutations on the egg-laying muscles could be similarly explained, since these muscles are interconnected by gap junctions and presumably communicate with each other (White et al., 1986).

Alternatively, *unc-93* could regulate muscle contraction by functioning in the response of muscle cells to neuronal inputs in excitation-contraction coupling. Studies of excitation-contraction coupling in mammalian skeletal muscle have defined the steps of excitation-contraction coupling. The binding of a neurotransmitter by its receptor in the muscle cell membrane leads to an influx of sodium ions that triggers the depolarization of the muscle cell membrane (Shepherd, 1988). The dihydropyridine receptor, a calcium ion channel in the transverse tubules, acts as a voltage sensor to signal the ryanodine receptor (the calcium release channel) to release calcium ions from the sarcoplasmic reticulum into the cytoplasm (Catterall et al., 1988; Fill and Coronado, 1988; Jan and Jan, 1989). Calcium ions bind to troponin C in the thin filaments, which causes myosin to slide against actin to generate a contraction (Zot and Potter, 1987). The store of calcium ions in the sarcoplasmic reticulum is replenished by a calcium-dependent ATPase that pumps calcium ions from the cytoplasm back into the sarcoplasmic reticulum (MacLennan, 1970). Some of the details of excitation-contraction coupling differ in *C. elegans*, but the overall mechanism of muscle contraction is likely to be similar (Waterston, 1988). The contractile process in *C. elegans* is likely to be regulated via the release of calcium from the sarcoplasmic reticulum. However, *C. elegans* apparently does not have an equivalent to the transverse tubule system, possibly because the sarcoplasmic reticulum is in close proximity to the plasma membrane (Waterston, 1988). *C. elegans* uses both myosin and thin filament-linked calcium regulation of muscle contraction (Harris et al., 1977). For *C. elegans*, the details of the excitation-contraction coupling pathway between the acetylcholine receptor and the interaction of calcium ions with the thick and thin filaments in muscle cells are not yet known. Excitation-contraction coupling involves ion transport across both the muscle cell membrane and the membrane of the sarcoplasmic reticulum. Thus, *unc-93* might encode an ion transport protein or a protein that interacts with an ion transport protein localized to either of these membranes. If so, the rubber band phenotype could be caused by an ion channel with altered gating properties that disrupt muscle contraction. It is interesting to note that muscimol (Eldefrawi and Eldefrawi, 1987), a GABA agonist that seems likely to open GABA_A chloride channels in body wall muscle (S. McIntire, E. Jorgensen, and H. R. Horvitz, manuscript in preparation), causes wild-type worms to behave like rubber band mutants (our unpublished data; E. Jorgensen, personal communication). This observation suggests that the rubber band mutant phenotype could be caused by a hyperpolarization of body wall muscle cells. Alternatively, *unc-93* could encode some other novel type of muscle membrane protein.

The rubber band mutation *e1500* changes amino acid 388 from Gly to Arg in a possible membrane-spanning domain (roughly amino acids 376 to 400) (Fig. 6 *b*). The altered function of the *unc-93* protein in *e1500* animals could be a result of the introduction of a charged amino acid in a hydrophobic region and/or the substitution of a bulky amino acid for the compact glycine. Because *n200* animals have two DNA changes in the *unc-93* gene, it is not possible to state whether both changes or only one of the two changes is responsible for the rubber band phenotype. In *n200* animals,

the change of amino acid 49 from Ala to Val is in the highly charged NH₂-terminal putative cytoplasmic region and the change of amino acid 562 from Gly to Val probably affects the protein between two membrane-spanning domains. Both of the changes in *n200* animals and the change in *el500* animals substitute a larger amino acid for a smaller one. In addition, because of its conformational flexibility, glycine (which is affected in both *n200* and *el500* animals) is often used as a hinge between protein domains (Chou and Fasman, 1978). Thus, the *el500* and *n200* mutations might directly or indirectly disrupt the configuration of *unc-93* membrane-spanning domains, thereby changing the interaction of these domains with each other or with other proteins.

Rare altered-function *unc-93* and *sup-10* alleles cause the rubber band phenotype and a disruption of muscle contraction (Greenwald and Horvitz, 1980, 1986). Because mutants that lack *unc-93*, *sup-9*, *sup-10*, or *sup-18* gene function display no visibly abnormal phenotype (Greenwald and Horvitz, 1980, 1986), there is likely to be one or more other proteins that can function in parallel to regulate the same aspect of muscle contraction. The functional redundancy for *unc-93* could reflect the ability of a single alternative gene or of a group of genes to replace *unc-93* function in *unc-93* null mutants. The *C. elegans* actin genes *act-1*, *act-2*, and *act-3* and collagen genes *rol-6* and *sqt-1* also have null alleles that result in a wild-type phenotype and altered-function alleles that result in visibly abnormal phenotypes (Waterston et al., 1984; Landel et al., 1984; Park and Horvitz, 1986; Kusch and Edgar, 1986). The function of each of these genes is redundant because it can be provided by other members of a homologous gene family (Landel et al., 1984; Kramer et al., 1988, 1990). Similarly, the functional redundancy of *unc-93* might be explained by an *unc-93* gene family, with an *unc-93* homolog able to function in place of *unc-93*. However, the functionally redundant protein is not likely to be an *unc-93* homolog because it can still properly regulate muscle contraction in the absence of *sup-9* or *sup-10*, whereas an *unc-93* homolog presumably would interact with the products of *sup-9* and *sup-10*. In the absence of *sup-9* or *sup-10* gene function, the *unc-93*(*el500*) mutant protein does not disrupt the regulation of muscle contraction (Greenwald and Horvitz, 1980). This observation suggests that the *unc-93*(+) protein requires *sup-9* and *sup-10* proteins to function. Furthermore, genomic Southern blots probed at low stringency (e.g., 55°C, 0.75 M NaCl) with *unc-93* do not show any additional hybridizing bands (our unpublished data; M. Nadal-Vicens, personal communication). Thus, there is no evidence to support the existence of an *unc-93* gene family. Rather, we suggest that the functional redundancy of *unc-93* is due to a gene or set of genes unrelated by DNA sequence that can perform the same function. Based on the common phenotypes and suppression patterns observed among mutations in *unc-93*, *sup-9*, *sup-10*, and *sup-18*, these four genes are likely to act as a protein complex or in a common process in the membranes of muscle cells. The apparent functional redundancy of each of these genes could be due to proteins unrelated by sequence that act in parallel in a separate protein complex or pathway. In models in which the *unc-93* protein interacts with gap junctions or ion channels, the rubber band mutations could be altering their gating properties to disrupt the regulation of muscle contraction even in the pres-

ence of a functionally redundant alternative protein complex or pathway.

We thank M. Finney for the three times backcrossed *mut-2* strain MT2879; M. Shen, D. Parry, and J. Thomas for providing strains; A. Coulson and J. Sulston for the *C. elegans* genomic DNA library; S. Kim for the cDNA library; and D. Hsu and A. Fire for the pMRF DNA. We thank C. Bargmann, E. Jorgensen, and M. Stern for comments concerning this manuscript.

This work was supported by research grant GM 24663 from the U.S. Public Health Service. J. Z. Levin was supported by National Institutes of Health. Pre-doctoral training grant GM 07287 and the Lucille P. Markey Charitable Trust. H. R. Horvitz is an Investigator of the Howard Hughes Medical Institute.

Received for publication 4 September 1991 and in revised form 13 January 1992.

References

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment sequence tool. *J. Mol. Biol.* 215:403-410.
- Atchison, M. L. 1988. Enhancers: mechanisms of action and cell specificity. *Annu. Rev. Cell Biol.* 4:127-153.
- Benian, G. M., J. E. Kiff, N. Neckelmann, D. G. Moerman, and R. H. Waterston. 1989. Sequence of an unusually large protein implicated in regulation of myosin activity in *C. elegans*. *Nature (Lond.)* 342:45-50.
- Bennetzen, J. L., and B. D. Hall. 1982. Codon selection in yeast. *J. Biol. Chem.* 257:3026-3031.
- Brenner, S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* 77:71-94.
- Catterall, W. A., M. J. Seagar, and M. Takahashi. 1988. Molecular properties of dihydropyridine-sensitive calcium channels in skeletal muscle. *J. Biol. Chem.* 263:3535-3538.
- Chalfie, M., and J. White. 1988. The nervous system. In *The Nematode Caenorhabditis elegans*. W. Wood and the community of *C. elegans* researchers, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 337-392.
- Chou, P. Y., and G. D. Fasman. 1978. Empirical predictions of protein conformation. *Ann. Rev. Biochem.* 47:251-276.
- Collins, J., B. Saari, and P. Anderson. 1987. Activation of a transposable element in the germ line but not the soma of *Caenorhabditis elegans*. *Nature (Lond.)* 328:726-728.
- Collins, J., E. Forbes, and P. Anderson. 1989. The Tc3 family of transposable genetic elements in *Caenorhabditis elegans*. *Genetics* 121:47-55.
- Coulson, A., J. Sulston, S. Brenner, and J. Karn. 1986. Towards a physical map of the genome of the nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 83:7821-7825.
- Coulson, A., R. Waterston, J. Kiff, J. Sulston, and Y. Kohara. 1988. Genome linking with yeast artificial chromosomes. *Nature (Lond.)* 335:184-186.
- Devereux, J., P. Haeblerli, and O. Smithies. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.* 12:387-395.
- Edgley, M. L., and D. L. Riddle. 1990. The nematode *Caenorhabditis elegans*. In *Genetic Maps*. S. J. O'Brien, editor. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 3:111-3.133.
- Edwards, M. K., and W. B. Wood. 1983. Location of specific messenger RNAs in *Caenorhabditis elegans* by cytological hybridization. *Dev. Biol.* 97:375-390.
- Eide, D., and P. Anderson. 1988. Insertion and excision of *Caenorhabditis elegans* transposable element Tc1. *Mol. Cell. Biol.* 8:737-746.
- Eldefrawi, A. T., and M. E. Eldefrawi. 1987. Receptors for gamma-aminobutyric acid and voltage-dependent chloride channels as targets for drugs and toxicants. *FASEB (Fed. Am. Soc. Exp. Biol.) J.* 1:262-271.
- Emmons, S. W. 1988. The genome. In *The Nematode Caenorhabditis elegans*. W. Wood and the community of *C. elegans* researchers, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 47-79.
- Emmons, S. W., L. Yesner, K. S. Ruan, and D. Katzenberg. 1983. Evidence for a transposon in *Caenorhabditis elegans*. *Cell* 32:55-65.
- Epstein, H. F., R. H. Waterston, and S. Brenner. 1974. A mutant affecting the heavy chain of myosin in *Caenorhabditis elegans*. *J. Mol. Biol.* 90:291-300.
- Files, J. G., and D. Hirsh. 1981. Ribosomal DNA of *Caenorhabditis elegans*. *J. Mol. Biol.* 149:223-240.
- Fill, M., and R. Coronado. 1988. Ryanodine receptor channel of sarcoplasmic reticulum. *Trends Neurosci.* 11:453-457.
- Finney, M. 1987. The genetics and molecular biology of *unc-86*, a *C. elegans* cell lineage gene. Ph.D. Thesis, Massachusetts Institute of Technology, Cambridge, MA. 153-174.
- Finney, M., G. Ruvkun, and H. R. Horvitz. 1988. The *C. elegans* cell lineage and differentiation gene *unc-86* encodes a protein with a homeodomain and extended similarity to transcription factors. *Cell* 55:757-769.

- Gilman, M. 1991. Ribonuclease Protection Assay. In *Current Protocols in Molecular Biology*. F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl, editors. John Wiley & Sons, New York. 4.7.1-4.7.8.
- Graf, L. H., and L. A. Chasin. 1982. Direct demonstration of genetic alterations at the dihydrofolate reductase locus after gamma irradiation. *Mol. Cell Biol.* 2:93-96.
- Greenwald, I. 1985. *lin-12*, a nematode homeotic gene, is homologous to a set of mammalian proteins that includes epidermal growth factor. *Cell.* 43:583-590.
- Greenwald, I., and H. R. Horvitz. 1986. A visible allele of the muscle gene *sup-10 X* of *C. elegans*. *Genetics.* 113:63-72.
- Greenwald, I. S., and H. R. Horvitz. 1980. *unc-93(e1500)*: a behavioral mutant of *Caenorhabditis elegans* that defines a gene with a wild-type null phenotype. *Genetics.* 96:147-164.
- Greenwald, I. S., and H. R. Horvitz. 1982. Dominant suppressors of a muscle mutant define an essential gene of *Caenorhabditis elegans*. *Genetics.* 101:211-225.
- Grosovsky, A. J., E. A. Drobetsky, P. J. deJong, and B. W. Glickman. 1986. Southern analysis of genomic alterations in gamma-ray-induced *aprt*-hamster cell mutants. *Genetics.* 113:405-415.
- Harris, H. E., M. Y. Tso, and H. F. Epstein. 1977. Actin and myosin-linked calcium regulation in the nematode *Caenorhabditis elegans*. Biochemical and structural properties of native filaments and purified proteins. *Biochemistry.* 16:859-865.
- Hartmann, E., T. A. Rapoport, and H. F. Lodish. 1989. Predicting the orientation of eukaryotic membrane-spanning proteins. *Proc. Natl. Acad. Sci. USA.* 86:5786-5790.
- Henikoff, S. 1984. Unidirectional digestion with exonuclease III in DNA sequence analysis. *Gene.* 28:351-359.
- Herman, R. K. 1984. Analysis of genetic mosaics of the nematode *Caenorhabditis elegans*. *Genetics.* 108:165-180.
- Hodgkin, J. 1983. Male phenotypes and mating efficiency in *Caenorhabditis elegans*. *Genetics.* 103:43-64.
- Hodgkin, J., M. Edgley, D. L. Riddle, and D. G. Albertson. 1988. Appendix 4: Genetics. In *The Nematode Caenorhabditis elegans*. W. Wood and the community of *C. elegans* researchers, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 491-586.
- Huang, X.-Y., and D. Hirsh. 1989. A second *trans*-spliced RNA leader sequence in the nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA.* 86:8640-8644.
- Jan, L. Y., and Y. N. Jan. 1989. Voltage-sensitive ion channels. *Cell.* 56:13-25.
- Kagawa, H., K. Gengyo, A. D. McLachlan, S. Brenner, and J. Karn. 1989. Paramyosin gene (*unc-15*) of *Caenorhabditis elegans*. Molecular cloning, nucleotide sequence and models for thick filament structure. *J. Mol. Biol.* 207:311-333.
- Kamm, K. E., and J. T. Stull. 1985. The function of myosin and myosin light chain kinase phosphorylation in smooth muscle. *Ann. Rev. Pharmacol. Toxicol.* 25:593-620.
- Kim, S. K., and H. R. Horvitz. 1990. The *Caenorhabditis elegans* gene *lin-10* is broadly expressed while required specifically for the determination of vulval cell fates. *Genes Dev.* 4:357-371.
- Kingston, R. E. 1991. Primer Extension. In *Current Protocols in Molecular Biology*. F. M. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl, editors. John Wiley & Sons, New York. 4.8.1-4.8.3.
- Klein, P., M. Kanehisa, and C. DeLisi. 1985. The detection and classification of membrane-spanning proteins. *Biochim. Biophys. Acta.* 815:468-476.
- Kramer, J. M., J. J. Johnson, R. S. Edgar, C. Basch, and S. Roberts. 1988. The *sqt-1* gene of *C. elegans* encodes a collagen critical for organismal morphogenesis. *Cell.* 55:555-565.
- Kramer, J. M., R. P. French, E. C. Park, and J. J. Johnson. 1990. The *Caenorhabditis elegans rol-6* gene, which interacts with the *sqt-1* collagen gene to determine organismal morphology, encodes a collagen. *Mol. Cell Biol.* 10:2081-2089.
- Krause, M., and D. Hirsh. 1987. A *trans*-spliced leader sequence on actin mRNA in *C. elegans*. *Cell.* 49:753-761.
- Kusch, M., and R. S. Edgar. 1986. Genetic studies of unusual loci that affect body shape of the nematode *Caenorhabditis elegans* and may code for cuticle structural proteins. *Genetics.* 113:621-639.
- Kyte, J., and R. F. Doolittle. 1982. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* 157:105-132.
- Landel, C. P., M. Krause, R. H. Waterston, and D. Hirsh. 1984. DNA rearrangements of the actin gene cluster in *Caenorhabditis elegans* accompany reversion of three muscle mutants. *J. Mol. Biol.* 180:497-513.
- Leavis, P. C., and J. Gergely. 1984. Thin filament proteins and thin filament-linked regulation of vertebrate muscle contraction. *CRC Crit. Rev. Biochem.* 16:235-305.
- MacLennan, D. H. 1970. Purification and properties of an adenosine triphosphatase from sarcoplasmic reticulum. *J. Biol. Chem.* 245:4508-4518.
- MacLeod, A. R., R. H. Waterston, and S. Brenner. 1977. An internal deletion mutant of a myosin heavy chain in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA.* 74:5336-5340.
- MacLeod, A. R., J. Karn, and S. Brenner. 1981. Molecular analysis of the *unc-54* myosin heavy-chain gene of *Caenorhabditis elegans*. *Nature (Lond.)*. 291:386-390.
- Marck, C. 1988. 'DNA Strider': a 'C' program for the fast analysis of DNA and protein sequences on the Apple Macintosh family of computers. *Nucleic Acids Res.* 16:1829-1836.
- Meyer, B. J., and L. P. Casson. 1986. *Caenorhabditis elegans* compensates for the difference in X chromosome dosage between the sexes by regulating transcript levels. *Cell.* 47:871-881.
- Miller, L. M., J. D. Plenefisch, L. P. Casson, and B. J. Meyer. 1988. *xol-1*: a gene that controls the male modes of both sex determination and X chromosome dosage compensation in *C. elegans*. *Cell.* 55:167-183.
- Moerman, D. G., S. Plurad, R. H. Waterston, and D. L. Baillie. 1982. Mutations in the *unc-54* myosin heavy chain gene of *Caenorhabditis elegans* that alter contractility but not muscle structure. *Cell.* 29:773-781.
- Moerman, D. G., G. M. Benian, and R. H. Waterston. 1986. Molecular cloning of the muscle gene *unc-22* in *Caenorhabditis elegans* by Tc1 transposon tagging. *Proc. Natl. Acad. Sci. USA.* 83:2579-2583.
- Mori, I., G. M. Benian, D. G. Moerman, and R. H. Waterston. 1988. Transposable element Tc1 of *Caenorhabditis elegans* recognizes specific target sequences for integration. *Proc. Natl. Acad. Sci. USA.* 85:861-864.
- Park, E. C., and H. R. Horvitz. 1986. Mutations with dominant effects on the behavior and morphology of the nematode *Caenorhabditis elegans*. *Genetics.* 113:821-852.
- Parks, G. D., and R. A. Lamb. 1991. Topology of eukaryotic type II membrane proteins: importance of N-terminal positively charged residues flanking the hydrophobic domain. *Cell.* 64:777-787.
- Proudfoot, N. J., and G. G. Brownlee. 1976. 3' non-coding region sequences in eukaryotic messenger RNA. *Nature (Lond.)*. 263:668-671.
- Rao, J. K., and P. Argos. 1986. A conformational preference parameter to predict helices in integral membrane proteins. *Biochim. Biophys. Acta.* 869:197-214.
- Riddle, D. L., M. M. Swanson, and P. S. Albert. 1981. Interacting genes in nematode dauer larva formation. *Nature (Lond.)*. 290:668-671.
- Rosenzweig, B., L. W. Liao, and D. Hirsh. 1983. Sequence of the *C. elegans* transposable element Tc1. *Nucleic Acids Res.* 11:4201-4209.
- Saiki, R. K., D. H. Gelfand, S. Stoffel, S. J. Scharf, R. Higuchi, G. T. Horn, K. B. Mullis, and H. A. Erlich. 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science (Wash. DC)*. 239:487-494.
- Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F., S. Nicklen, and A. Coulson. 1977. DNA Sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA.* 74:5463-5467.
- Shepherd, G. M. 1988. *Neurobiology*. Oxford Univ. Press, Oxford.
- Sulston, J. E., and H. R. Horvitz. 1977. Post-embryonic cell lineages of the nematode, *Caenorhabditis elegans*. *Dev. Biol.* 56:110-156.
- Swanson, M. M., and D. L. Riddle. 1981. Critical periods in the development of the *Caenorhabditis elegans* dauer larva. *Dev. Biol.* 84:27-40.
- Wang, K. 1985. Sarcomere-associated cytoskeletal lattices in striated muscle. Review and hypothesis. *Cell Muscle Motil.* 6:315-369.
- Warrick, H. M., and J. A. Spudich. 1987. Myosin structure and function in cell motility. *Ann. Rev. Cell Biol.* 3:379-421.
- Waterston, R. H. 1988. Muscle. In *The Nematode Caenorhabditis elegans*. W. Wood and the community of *C. elegans* researchers, editors. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 281-335.
- Waterston, R. H. 1989. The minor myosin heavy chain, mhca, of *Caenorhabditis elegans* is necessary for the initiation of thick filament assembly. *EMBO (Eur. Mol. Biol. Organ.) J.* 8:3429-3436.
- Waterston, R. H., R. M. Fishpool, and S. Brenner. 1977. Mutants affecting paramyosin in *Caenorhabditis elegans*. *J. Mol. Biol.* 117:679-697.
- Waterston, R. H., J. N. Thomson, and S. Brenner. 1980. Mutants with altered muscle structure of *Caenorhabditis elegans*. *Dev. Biol.* 77:271-302.
- Waterston, R. H., D. Hirsh, and T. R. Lane. 1984. Dominant mutations affecting muscle structure in *Caenorhabditis elegans* that map near the actin gene cluster. *J. Mol. Biol.* 180:473-496.
- White, J. G., E. Southgate, J. N. Thomson, and S. Brenner. 1986. The structure of the nervous system of *Caenorhabditis elegans*. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 314:1-340.
- Yuan, J., M. Finney, N. Tsung, and H. R. Horvitz. 1991. Tc4, a *Caenorhabditis elegans* transposable element with an unusual fold-back structure. *Proc. Natl. Acad. Sci. USA.* 88:3334-3338.
- Zot, A. S., and J. D. Potter. 1987. Structural aspects of troponin-tropomyosin regulation of skeletal muscle contraction. *Annu. Rev. Biophys. Biophys. Chem.* 16:535-559.