

1-Aminocyclopropane-1-Carboxylate Deaminase Enhances *Agrobacterium tumefaciens*-Mediated Gene Transfer into Plant Cells[∇]

Satoko Nonaka,¹ Masayuki Sugawara,² Kiwamu Minamisawa,² Ken-ichi Yuhashi,¹ and Hiroshi Ezura^{1*}

Graduate School of Life and Environmental Sciences, University of Tsukuba, Tennodai 1-1-1, Tsukuba 305-8572, Japan,¹ and Graduate School of Life Sciences, Tohoku University, Katahira, Aoba-ku, Sendai, Japan²

Received 3 October 2007/Accepted 20 February 2008

***Agrobacterium*-mediated gene transfer is widely used for plant molecular genetics, and efficient techniques are required. Recent studies show that ethylene inhibits the gene transfer. To suppress ethylene evolution, we introduced 1-aminocyclopropane-1-carboxylate (ACC) deaminase into *Agrobacterium tumefaciens*. The ACC deaminase enhanced *A. tumefaciens*-mediated gene transfer into plants.**

Agrobacterium-mediated gene transfer is widely used for plant molecular genetics and its applications (14). In particular, efficient systems of genetic transformation are required for plant functional genomics and molecular breeding to improve traits (20, 21). Recent studies showed that ethylene is one of the factors that inhibits *Agrobacterium*-mediated gene transfer (1, 3, 5). Therefore, if *Agrobacterium tumefaciens* has the ability to decrease the ethylene level in the host plant, the frequency of gene transfer will increase. To suppress ethylene evolution in plant cells during cocultivation, we introduced the 1-aminocyclopropane-1-carboxylate (ACC) deaminase gene from *Pseudomonas* sp. strain ACP (7, 18) into *A. tumefaciens*. ACC deaminase cleaves ACC (the immediate ethylene precursor) into α -ketobutyrate and ammonia, and as a result, the ethylene level is decreased (4, 12, 16).

The ACC deaminase gene was amplified and cloned into pBBR1MCS-5 (10), a broad-host-range plasmid, to generate a *lacZ::accD* translational fusion (Fig. 1A). The resulting plasmid was designated pBBR*accD* and was introduced into *A. tumefaciens* C58 (17) or C58C1Rif^R (2) by electroporation (19). The binary vector pIG121-Hm, involved in T-DNA transfer (6), was also harbored in *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm). The ACC deaminase activity in *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm) was assayed according to the method of Honma and Shimomura (7). The amount of α -ketobutyrate in the reaction buffer was estimated from a standard curve based on a dilution of 10 to 400 μ M (detected at 340 nm). The controls for this experiment were C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) and samples in reaction buffer without the substrate 200 μ M ACC. The accumulation of α -ketobutyrate was observed only in the lysate from *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm) in the presence of the substrate (Fig. 1B). Therefore, we succeeded in conferring ACC deaminase activity on *A. tumefaciens*.

Surface-sterilized melon (*Cucumis melo* L. var. *cantaloupensis* cv. Vedrantaïs) seeds were sown on a half-strength preparation of

Murashige and Skoog's medium (MS) (13) and incubated at 25°C with 16 h of light per day for 5 days. Cotyledons from the germinated seedlings were transversely sectioned by hand into five pieces, and among these five, three internal pieces were inoculated. The segments were soaked in an *A. tumefaciens* cell suspension of 10⁷ cells ml⁻¹ for 20 min and then placed on cocultivation medium (MS containing 1.0 mg of 6-benzylamino-purine liter⁻¹, 2% glucose, and 0.4% Gelrite [Wako, Tokyo, Japan], pH 5.5) in a gas vial with 16 h of light per day. Thirty melon cotyledon segments were inoculated with *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm) per experiment. The experiments were repeated three times. After 24 h of incubation, ethylene evolution in melon cotyledon segments was measured by gas chromatography (Fig. 2A). Compared to that in the uninoculated controls, ethylene evolution in the melon segments inoculated with *A. tumefaciens* C58C1Rif^R(pIG121-Hm) and C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) was enhanced. The application of 1 μ M aminoethoxyvinylglycine (AVG), an ethylene biosynthesis inhibitor, reduced ethylene evolution in the inoculated segments. Inoculation with *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm) suppressed ethylene evolution in melon cotyledon segments, and the ethylene accumulation rate was the same as that in the control and AVG-treated samples. These results indicated that *A. tumefaciens* with ACC deaminase activity reduced ethylene evolution in plants (Fig. 2A).

Three days after inoculation, the level of gene transfer was estimated (Fig. 2B). The pIG121-Hm plasmid has a reporter gene (35S-*uidA* intron) in the T-DNA region. Because the *uidA* reporter gene possesses an intron sequence, it can produce active protein only in plant cells, thereby making it a marker for gene transfer (15). Gene transfer was evaluated using a fluorometric β -glucuronidase (GUS) assay according to the method of Jefferson et al. (8). Melon segments inoculated with C58C1Rif^R(pIG121-Hm) and C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) showed higher levels of GUS activity than controls. These higher levels of GUS activity indicated that the gene was transferred. The addition of AVG (1 μ M) increased GUS activity two times over that in the absence of AVG. Inoculation with *A. tumefaciens* C58C1Rif^R(pBBR*accD*, pIG121-Hm) yielded approximately six-times-higher levels of GUS activity than inoculation with C58C1Rif^R(pIG121-Hm). Thus, ACC deami-

* Corresponding author. Mailing address: Graduate School of Life and Environmental Sciences, University of Tsukuba, Tennodai 1-1-1, Tsukuba 305-8572, Japan. Phone and fax: 81-29-853-773. E-mail: ezura@gene.tsukuba.ac.jp.

[∇] Published ahead of print on 29 February 2008.




FIG. 1. Construction of the ACC deaminase-producing *A. tumefaciens* strain. (A) Plasmid construction for expression of ACC deaminase in *A. tumefaciens*. An HindIII and XbaI fragment (1 kb) containing the ACC deaminase gene from *Pseudomonas* sp. strain ACP was ligated into the HindIII and XbaI sites of the broad-host-range plasmid pBBR1MCS-5, resulting in pBBRacdS. The expression of the ACC deaminase gene *acdS* was under the control of the *lac* promoter. MCS stands for multiple cloning sites. (B) Detection of ACC deaminase activity in *A. tumefaciens*. The α -ketobutirate (α -KB) accumulation in the reaction buffer was measured according to the method of Honma and Shimomura (7). The triangles and circles indicate the lysates from *A. tumefaciens* C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) and C58C1Rif^R(pBBRacdS, pIG121-Hm), respectively. Closed and open symbols represent samples with and without ACC in the reaction buffer, respectively. Bars indicate standard deviations ($n = 3$).

nase enhanced the ability for gene transfer from *A. tumefaciens* (Fig. 2B).

Seeds of *Arabidopsis thaliana* (Columbia) were sterilized and grown at 22°C for 7 days with 16 h of light per day after 4 days of vernalization. Intact *A. thaliana* plants were dipped into a suspension of *A. tumefaciens* C58 or A136 (10^7 cells ml⁻¹). A136 lacks the Ti plasmid and the T-DNA region and was used as a control. The inoculated seedlings were blotted onto sterile filter paper to remove excess suspension material and cocultivated on MS for 7




FIG. 2. Effect of ACC deaminase activity on ethylene evolution and gene transfer. (A) Measurement of ethylene evolution. The accumulation of ethylene in the headspace was measured on a gas chromatograph. 121-Hm, 121-Hm/AVG, MCS-5, and ACDS indicate samples inoculated with *A. tumefaciens* C58C1Rif^R(pIG121-Hm), those inoculated with *A. tumefaciens* C58C1Rif^R(pIG121-Hm) and treated with the addition of 1 μ M AVG to the cocultivation medium, and samples inoculated with C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) and C58C1Rif^R(pBBRacdS, pIG121-Hm), respectively. Bars represent standard deviations ($n = 3$). The characters *a* and *b* show statistically significant differences (*t* test; $P < 0.05$). fw, fresh weight. (B) Quantification of gene transfer by GUS assay. Melon cotyledon segments were cocultivated with three different *A. tumefaciens* strains for 3 days. 121-Hm, 121-Hm/AVG, MCS-5, and ACDS indicate samples inoculated with *A. tumefaciens* C58C1Rif^R(pIG121-Hm), those inoculated with *A. tumefaciens* C58C1Rif^R(pIG121-Hm) and treated with the addition of 1 μ M AVG to the cocultivation medium, and samples inoculated with C58C1Rif^R(pBBR1MCS-5, pIG121-Hm) and C58C1Rif^R(pBBRacdS, pIG121-Hm), respectively. Bars represent standard deviations ($n = 3$). Different letters indicate statistically significant differences (*t* test; $P < 0.05$).

days with 16 h of light per day. After cocultivation, to eliminate the bacteria, the plants were washed in sterilized water and then incubated on MS containing 375 mg of amoxicillin-clavulanic acid (Augmentin) liter⁻¹ for 3 weeks. Four weeks after inoculation with C58, C58(pBBR1MCS-5), and C58(pBBRacdS), green tumors had formed on the stems (Fig. 3A). The sizes of tumors among the different infections were almost the same (Fig. 3A). There were no tumors observed on plants inoculated with A136 (Fig. 3). This result indicated that the tumor formation was induced by stable transformation (22). To estimate the genetic transformation efficiency, the numbers of *A. thaliana* plants forming green tumors were determined and the percentages were calculated. Fifteen intact *A. thaliana* seedlings were used in each




FIG. 3. Estimation of the frequency of genetic transformation by *acdS*-expressing *A. tumefaciens*. (A) Photographs of tumor formation on *A. thaliana* Columbia (Col) stems. (B) Frequency of genetic transformation. A136, C58, MCS5, and ACDS indicate samples inoculated with *A. tumefaciens* strains A136, C58, C58(pBBRMCS-5), and C58(pBBR*acdS*), respectively. Bars indicate standard deviations ($n = 3$). The characters *a* and *b* represent statistically significant differences based on chi-square testing ($P < 0.05$).

experiment, and there were three independent repetitions. The percentages of plants that formed tumors were $8.1\% \pm 2.3\%$, $10.6\% \pm 4.1\%$, and $27.2\% \pm 2.4\%$, respectively, of those inoculated with *A. tumefaciens* C58, C58(pBBRMCS-5), and C58(pBBR*acdS*) (Fig. 3B). The tumor incidence was higher among plants inoculated with the ACC deaminase-producing strain. This result indicated that ACC deaminase activity increased the ability for stable transformation with *A. tumefaciens* (Fig. 3B).

Genetic transformation is a key technology for plant molecular breeding. Among several techniques of genetic transformation, *Agrobacterium*-mediated gene transfer is the most frequently used. Although great efforts have been made to establish efficient protocols of genetic transformation for plants of interest, species and genotypes recalcitrant to genetic transformation, such as cotton (11) and soybeans (9), still exist. We succeeded in producing an *Agrobacterium* strain with improved potential for gene transfer by providing the ability to

reduce the ethylene level of the plant during cocultivation. The knowledge obtained in this study will provide a clue to overcome such problems in plant molecular breeding as producing transgenic plants of recalcitrant species and genotypes.

We thank the members of the Ezura laboratory for helpful discussions. We are grateful to Shohael Abdullah for his advice on the English in this article.

This work was supported by the 21st Century Centers of Excellence Program and a grant-in-aid for scientific research, category B (no. 15380002), from the Ministry of Education, Science, Sports, and Technology of Japan to H.E.

REFERENCES

- Davis, M. E., A. R. Miller, and R. D. Lineberger. 1992. Studies on the effects of ethylene on transformation of tomato cotyledons (*Lycopersicon esculentum* Mill.) by *Agrobacterium tumefaciens*. *J. Plant Physiol.* **139**:309–312.
- Deblaere, R., B. Bytebier, H. De Greve, F. Deboeck, J. Schell, M. van Montagu, and J. Leemans. 1985. Efficient octopine Ti plasmid-derived vectors for *Agrobacterium*-mediated gene transfer to plants. *Nucleic Acids Res.* **13**:4777–4788.
- Ezura, H., K. Yuhashi, T. Yasuta, and K. Minamisawa. 2000. Effect of ethylene on *Agrobacterium tumefaciens*-mediated gene transfer to melon. *Plant Breed.* **119**:75–79.
- Glick, B. R., D. M. Penrose, and J. Li. 1998. A model of the lowering of plant ethylene concentrations by plant growth-promoting bacteria. *J. Theor. Biol.* **190**:63–68.
- Han, J. S., C. K. Kim, S. H. Park, K. D. Hirschi, and I. G. Mok. 2005. *Agrobacterium*-mediated transformation of bottle gourd (*Lagenaria siceraria* Standl.). *Plant Cell Rep.* **23**:692–698.
- Hiei, Y., S. Ohta, T. Komari, and T. Kumashiro. 1994. Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**:271–282.
- Honma, S., and T. Shimomura. 1978. Metabolism of 1-aminocyclopropane-1-carboxylic acid. *Agric. Biol. Chem.* **42**:1825–1831.
- Jefferson, R. K., T. A. Kavanagh, and M. W. Bevan. 1987. GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**:3901–3907.
- Ko, T. S., S. S. Korban, and D. A. Somers. 2006. Soybean (*Glycine max*) transformation using immature cotyledon explants. *Methods Mol. Biol.* **343**:397–405.
- Kovach, M. E., P. H. Elzer, D. S. Hill, G. T. Robertson, M. A. Farris, R. M. Roop II, and K. M. Peterson. 1995. Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **166**:175–176.
- Leelavathi, S., V. G. Sunnichan, R. Kumria, G. P. Vijaykannth, R. K. Bhatnagar, and V. S. Reddy. 2004. A simple and rapid *Agrobacterium*-mediated transformation protocol for cotton (*Gossypium hirsutum* L.): embryogenic calli as a source to generate large numbers of transgenic plants. *Plant Cell Rep.* **22**:465–470.
- Madhaiyan, M., S. Poonguzhali, J. Ryu, and T. Sa. 2006. Regulation of ethylene levels in canola (*Brassica campestris*) by 1-aminocyclopropane-1-carboxylate deaminase-containing *Methylobacterium fujisawaense*. *Planta* **224**:268–278.
- Murashige, T., and F. Skoog. 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant.* **15**:473–497.
- Newell, C. A. 2000. Plant transformation technology. *Developments and applications.* *Mol. Biotechnol.* **16**:53–65.
- Ohta, S., S. Mira, T. Hattori, and K. Nakamura. 1990. Construction and expression in tobacco of a β -glucuronidase (GUS) reporter gene containing an intron within the coding sequence. *Plant Cell Physiol.* **31**:805–813.
- Penrose, D. M., and B. R. Glick. 2001. Levels of 1-aminocyclopropane-1-carboxylic acid (ACC) in exudates and extracts of canola seeds treated with plant growth-promoting bacteria. *Can. J. Microbiol.* **47**:368–372.
- Sciaky, D., A. L. Montoya, and M. D. Chilton. 1978. Fingerprints of *Agrobacterium* Ti plasmids. *Plasmid* **1**:238–253.
- Sheehy, R. E., M. Honma, M. Yamada, T. Sasaki, B. Martineau, and W. R. Hiatt. 1991. Isolation, sequence, and expression in *Escherichia coli* of the *Pseudomonas* sp. strain ACP gene encoding 1-aminocyclopropane-1-carboxylate deaminase. *J. Bacteriol.* **173**:5260–5265.
- Shen, W. J., and B. G. Forde. 1989. Efficient transformation of *Agrobacterium* spp. by high voltage electroporation. *Nucleic Acids Res.* **17**:8353.
- Sun, H. J., S. Uchii, S. Watanabe, and H. Ezura. 2006. A highly efficient transformation protocol for Micro-Tom, a model cultivar of tomato functional genomics. *Plant Cell Physiol.* **47**:426–431.
- Tanaka, Y., Y. Katsumoto, F. Brugliera, and J. Mason. 2005. Genetic engineering in floriculture. *Plant Cell Tissue Organ Cult.* **80**:1–24.
- Yuan, Z. C., M. P. Edlind, P. Liu, P. Saenkham, L. M. Banta, A. A. Wise, E. Ronzone, A. N. Binns, K. Kerr, and E. W. Nester. 2007. The plant signal salicylic acid shuts down expression of the *vir* regulon and activates quorum-quenching genes in *Agrobacterium*. *Proc. Natl. Acad. Sci. USA* **104**:1179–11795.