

Regulation of Urokinase-Type Plasminogen Activator Gene Transcription by Macrophage Colony-Stimulating Factor

KATRYN J. STACEY,¹ LINDSAY F. FOWLES,¹ MICHAEL S. COLMAN,²
MICHAEL C. OSTROWSKI,² AND DAVID A. HUME^{1*}

*Centre for Molecular and Cellular Biology, University of Queensland, Brisbane, Australia 4072,¹ and
Department of Microbiology, Duke University School of Medicine,
Durham, North Carolina 27710²*

Received 9 December 1994/Returned for modification 3 February 1995/Accepted 8 March 1995

The mouse urokinase-type plasminogen activator (uPA) gene was used as a model macrophage colony-stimulating factor 1 (CSF-1)-inducible gene to investigate CSF-1 signalling pathways. Nuclear run-on analysis showed that induction of uPA mRNA by CSF-1 and phorbol myristate acetate (PMA) was at the transcriptional level in bone marrow-derived macrophages. CSF-1 and PMA synergized strongly in the induction of uPA mRNA, showing that at least some components of CSF-1 action are mediated independently of protein kinase C. Promoter targets of CSF-1 signalling were investigated with NIH 3T3 cells expressing the human CSF-1 receptor (*c-fms*). uPA mRNA was induced in these cells by treatment with CSF-1, and a PEA3/AP-1 element at -2.4 kb in the uPA promoter was involved in this response. Ets transcription factors can act through PEA3 sequences, and the involvement of Ets factors in the induction of uPA was confirmed by use of a dominant negative Ets-2 factor. Expression of the DNA binding domain of Ets-2 fused to the *lacZ* gene product prevented CSF-1-mediated induction of uPA mRNA in NIH 3T3 cells expressing the CSF-1 receptor. Examination of *ets-2* mRNA expression in macrophages showed that it was also induced synergistically by CSF-1 and PMA. In the macrophage cell line RAW264, the uPA PEA3/AP-1 element mediated a response to both PMA and cotransfected Ets-2. uPA promoter constructs were induced 60- to 130-fold by Ets-2 expression, and the recombinant Ets-2 DNA binding domain was able to bind to the uPA PEA3/AP-1 element. This work is consistent with a proposed pathway for CSF-1 signalling involving sequential activation of *fms*, *ras*, and Ets factors.

Macrophage colony-stimulating factor (CSF-1) acts primarily on cells of the mononuclear phagocyte lineage to control growth, differentiation, and survival of precursors as well as functional modulation of mature macrophages (59). The cell surface receptor for CSF-1 (CSF-1R) is a tyrosine kinase, the product of the *c-fms* proto-oncogene (reviewed in references 59 and 68), first identified by its homology to the viral oncogene *v-fms* from the McDonough feline sarcoma virus (69). Binding of CSF-1 is required for the kinase activity of the normal CSF-1R, whereas the *v-fms* product is constitutively active. The CSF-1R is most closely related to the platelet-derived growth factor receptor and the *c-kit* gene product (59).

When CSF-1 binds to its receptor, a range of cellular proteins and the receptor itself are phosphorylated (22). Like the platelet-derived growth factor receptor, CSF-1R binds and activates the 85-kDa subunit of phosphatidylinositol 3' kinase (55), but the role of phosphatidylinositol 3' kinase in signal transduction is not known. Other proteins which become tyrosine phosphorylated in response to CSF-1 include *src* family kinases (17), tyrosine phosphatase PTP-1C (84), Shc proteins, and an uncharacterized 150-kDa protein potentially involved in the *ras* signalling pathway (43), although these may not all be direct substrates of the receptor. The *ras* GTPase-activating protein (GAP)-associated proteins p62 and p190, and to a small extent GAP itself, have been reported to be phosphorylated in response to activated CSF-1R in fibroblasts (31, 55), but this has not been observed with macrophages (55).

Some of the actions of CSF-1 are mimicked by the protein

kinase C (PKC) agonist phorbol myristate acetate (PMA) (75), leading to speculation about the role of PKC in CSF-1 signalling. Evidence for an increase in production of diacylglycerol (an activator of PKC) after CSF-1 treatment of monocytes and bone marrow-derived macrophages (BMM) (34, 76) and for an increase in membrane-bound PKC activity (34, 66) has been presented, but recent experiments fail to support these findings (35). The generation of diacylglycerol is commonly the result of activation of phospholipase C- γ , an enzyme which cleaves phosphatidylinositol-4,5-bisphosphate to diacylglycerol and inositol-1,4,5-trisphosphate. Unlike the platelet-derived growth factor receptor, CSF-1R does not activate phospholipase C- γ (21), and there is no associated increase in phosphatidylinositol turnover (34, 81). An alternative route for diacylglycerol production may be by activation of a phosphatidylcholine-specific phospholipase C (83).

Evidence from a number of groups points to CSF-1-activated mitogenic signal pathways involving activation of *fms*, *ras*, *raf*, Ets, and *myc* factors (7, 9, 39, 60). This may not be a simple linear pathway, as recent experiments with a macrophage cell line showed that activation of Raf-1 was independent of *ras* (10). The involvement of p21^{ras} is well established in experiments with fibroblasts. CSF-1 treatment of *c-fms*-expressing NIH 3T3 cells and BAC1.2F5 macrophages increased the amount of active GTP-bound p21^{ras} (10, 26, 31). Microinjection of anti-p21^{ras} antibodies into *v-fms*-transformed fibroblasts caused reversion of the transformed phenotype (70), as did overexpression of the catalytic domain of GAP (7) and expression of a dominant negative Ets protein (39). Raf-1 was activated by CSF-1 treatment, and expression of oncogenic *v-raf* in a macrophage cell line led to CSF-1-independent growth and constitutive expression of *ets-2* and *c-myc* mRNAs (9). Mutation of the CSF-1R at tyrosine 809, an autophosphor-

* Corresponding author. Mailing address: Centre for Molecular and Cellular Biology, University of Queensland, Brisbane, Australia 4072. Phone: 61-7-3654493. Fax: 61-7-3654388. Electronic mail address: D.Hume@mailbox.uq.oz.au.

ylation site, prevented the mitogenic response to CSF-1 and impaired induction of *c-myc*, while induction of *c-fos* and *junB* was unaffected (60, 64). The mitogenic response could be restored by either *c-myc* (60), *ets-2* (39), or *ets-1* (61) expression.

Progress in defining promoter targets of CSF-1 action has been made in studies of the long terminal repeat of NVL3, a member of the VL30 family of mouse retrotransposons. A ras-responsive promoter element which also mediated a response to CSF-1 in NIH 3T3 fibroblasts expressing *c-fms* was found (7). The sequence element is a compound PEA3/AP-1-like site (54). Additional evidence for the role of p21^{ras} in the CSF-1 signalling pathway was obtained by showing that the catalytic domain of GAP inhibited *fms*-mediated activation of this element (7). Expression of a dominant negative Ets protein (Ets-2 DNA binding domain-LacZ fusion protein), which presumably binds the PEA3 sequence, also prevented CSF-1 activation of this element (39). This work provided good evidence for the roles of p21^{ras} and Ets factors in both CSF-1-mediated mitogenesis and induction of the NVL3 gene. More recent data have implicated Ets factors in *c-myc* induction by CSF-1 (61), in regulation of the CSF-1-responsive macrophage scavenger receptor gene (82), and in CSF-1-mediated induction of the *c-fos* gene (33).

The ultimate aim of studies of growth factor-activated signal transduction is to find all the molecular links from ligand binding at the cell surface to transcription factor activation and gene induction. Studies of nuclear targets of CSF-1 action in macrophages have been hampered by a lack of well-characterized genes which are induced by CSF-1 in transfectable cell lines, as primary macrophages and macrophage-like cell lines are difficult to transfect efficiently (73). This paper addresses the regulation of urokinase-type plasminogen activator (uPA), which is acutely regulated by CSF-1 in mature macrophages.

uPA is a serine protease which cleaves the proenzyme plasminogen to plasmin, a protease of broad specificity (reviewed in references 18 and 52). Plasmin directly catalyzes the degradation of the extracellular matrix and also acts indirectly by activating other proenzymes such as procollagenase. uPA activity is associated with processes which require tissue remodelling or cell movement, such as wound healing and involution of the mammary glands, and is frequently expressed at the invading front of cancerous tissue. Since macrophage function requires migration through tissues, uPA may be required to allow the necessary extracellular matrix degradation. uPA activity was first shown to be induced by CSF-1 in bone marrow-derived macrophages (BMM) (42). Accumulation of uPA mRNA in response to CSF-1 has been confirmed to occur in BMM (30) and in the macrophage-like cell line RAW264 (72). The level of macrophage or monocyte uPA mRNA has also been found to be increased by PMA (30) and gamma interferon (15) and decreased by dexamethasone and cyclic AMP (cAMP) agonists such as cholera toxin (15).

In this work we have examined the regulation of the uPA gene by CSF-1 and PMA. These agents synergistically induced uPA mRNA in macrophages, suggesting divergent pathways of activation. The difficulty of transfecting primary macrophages has meant that other cell types have had to be used for transient-transfection analysis. A PEA3/AP-1-like element at -2430 of the mouse uPA promoter was shown to be important in the induction of the gene by PMA in RAW264 cells and by CSF-1 in NIH 3T3 cells expressing *c-fms*. This element was first described by Rørth et al. (56) as being required for induction of uPA transcription by PMA and epidermal growth factor in keratinocytes. The element is conserved in mouse, pig, and human promoters (12) and is required for induction by PMA in a pig kidney epithelial cell line (41) and by PMA and basic

fibroblast growth factor in NIH 3T3 cells (5). Hence, diverse signalling pathways can converge on this one element. Here we provide evidence that Ets-2 can bind and activate transcription through this element.

MATERIALS AND METHODS

Materials. Recombinant human CSF-1 was kindly provided by Chiron Corp. PMA was obtained from Sigma.

Cell culture. BALB/c BMM were differentiated in culture with CSF-1 as previously described (73). The RAW264 macrophage-like cell line (53) is derived from tumor cells of an Abelson leukemia virus-infected mouse. This cell line was routinely grown in bacteriological petri dishes in RPMI 1640 medium with 5% fetal calf serum. NIH 3T3 fibroblasts, NIH 3T3 cells expressing human *c-fms* (63), and NIH 3T3 cells expressing both *c-fms* and the human Ets-2 DNA binding region fused to *lacZ* (39) were cultured in Dulbecco modified Eagle medium supplemented with 10% fetal calf serum.

Northern (RNA) analysis. Total RNA was extracted from cells by the method of Chomczynski and Sacchi (13). Total RNA (10 µg) was run on MOPS (morpholinepropanesulfonic acid)-formaldehyde denaturing gels, transferred to Hybond N nylon membranes, and hybridized with ³²P-labelled DNA probes. The probes used were mouse uPA, a 1.2-kb *Pst*I cDNA fragment (4); mouse Ets-2, excised from the plasmid Ets.pECE (obtained from R. Maki, La Jolla Cancer Research Foundation); 18S rRNA, a 24-bp oligonucleotide complementary to murine 18S rRNA (5'-CAT GGT AGG CAC GGC GAC TAC CAT3'); and human *K-*rev**-1 (rap 1A), a gene fragment obtained by PCR.

Nuclear run-on analysis. The plasmids used were pGEM2 (Promega) (as a control); uPA 5', a genomic fragment of mouse uPA from +8 to +398 (including exon 1, intron 1, and the beginning of exon 2) in pBluescript; uPA3', a fragment encompassing bp 424 to 1643 of the 2.3-kb mouse uPA cDNA in pGEM2; β-actin, a 1.8-kb chicken β-actin cDNA fragment in pGEM2; *c-fms*, p755 containing the 3.7-kb mouse *c-fms* cDNA, obtained from L. Rohrschneider (Fred Hutchinson Cancer Research Center, Seattle, Wash.); lysozyme, pMLC which contains a 780-bp fragment of mouse lysozyme m cDNA, obtained from R. Renkawitz (Max Planck Institute, Munich, Germany).

Plasmids were linearized, phenol-chloroform extracted, ethanol precipitated, and resuspended in 10 mM Tris-HCl (pH 8)–1 mM EDTA. The plasmid (1 µg) was slot blotted onto Hybond N membranes. Cells (2×10^7 to 3×10^7 per treatment group) were harvested and rinsed twice with ice-cold phosphate-buffered saline (PBS). The cells were lysed with 5 ml of ice-cold lysis buffer (10 mM Tris HCl [pH 7.4], 10 mM NaCl, 3 mM MgCl₂, 0.5% Nonidet P-40), pelleted (500 × g, 5 min), and then resuspended in 1 ml of lysis buffer. Nuclei were pelleted (500 × g, 5 min) and resuspended in 100 µl of glycerol storage buffer (50 mM Tris HCl [pH 8.3], 40% glycerol, 5 mM MgCl₂) with 40 U of RNasin (Boehringer Mannheim). To 100 µl of nuclei were added 100 µl of reaction buffer (10 mM Tris HCl [pH 8.0], 5 mM MgCl₂, 0.3 M KCl, 1 mM ATP, 1 mM CTP, 1 mM GTP, 5 mM dithiothreitol) and 50 µCi of [α-³²P]UTP (3,000 Ci/mmol; Amersham). Nuclei were incubated at 30°C for 30 min with shaking. RNA was then extracted by the method of Chomczynski and Sacchi (13) with the addition of 20 µg of carrier yeast tRNA (Boehringer Mannheim).

Filters were prehybridized and hybridized in 5 × SSPE (1 × SSPE is 0.18 M NaCl, 10 mM NaH₂PO₄, and 1 mM EDTA [pH 7.7]), 5 × Denhardt's solution, 0.5% sodium dodecyl sulfate (SDS), 50 µg of denatured herring sperm DNA per ml, and 200 µg of yeast tRNA per ml at 42°C in small glass scintillation vials inside rolling hybridization bottles. The prehybridization solution was removed after at least 6 h of incubation, and equivalent counts of RNA from each sample in 0.5 ml of hybridization solution were added. Hybridization proceeded for 36 h. The filters were washed twice for 20 min each time with 2 × SSC (1 × SSC is 0.15 M NaCl plus 0.015 M sodium citrate) at 42°C, twice for 20 min each time with 1 × SSC–0.1% SDS at 55°C, and once for 30 min with 2 × SSC with 10 µg of RNase A at 37°C and finally were rinsed several times in 2 × SSC. The filters were exposed for 2 days to 2 weeks.

Transfection analysis. RAW264 cells were transfected by electroporation with 10 µg of plasmid DNA as described previously (73). We found that the voltage needed to be optimized for the individual electroporator in use, as delivered voltages differed from the readout values. puPACAT2 plasmids containing the mouse uPA promoter linked to the chloramphenicol acetyltransferase (CAT) gene (see Fig. 3E) have been described (12). Plasmid pPR99, constructed by Rørth et al. (56), contains a 90-bp *Hae*III fragment of the mouse uPA promoter (-2446 to -2356) linked to the uPA proximal promoter (-114 to +398) and the *Escherichia coli* CAT gene. The related plasmids pPR119 and pPR127 contain point mutations in the PEA3/AP-1 site of the *Hae*III fragment (56) (see Fig. 3E). pPR126 contains only the proximal promoter (-114 to +398) and the CAT gene. Plasmid pGLuPA-6.6 contains the same promoter region (-6616 to +398) as puPACAT2, inserted into the *Hind*III site of the luciferase reporter plasmid pGL2-Basic (Promega). Plasmid pGL99 is equivalent to pPR99 but drives the luciferase reporter gene. The mouse Ets-2 expression vector Ets.pECE and PU.1 expression vector PU.pECE were obtained from R. Maki (La Jolla Cancer Research Foundation). Mouse Ets-2 expression vector was obtained from Martine Roussel (St. Jude Children's Research Hospital, Memphis, Tenn.). Mouse PEA3 cDNA (a gift from J. Hassell, McMaster University, Hamilton, Ontario,




FIG. 1. Northern analysis of uPA and *ets-2* mRNA expression in RAW264 cells and BMM. RAW264 cells and 6-day-old BMM were plated overnight (16 h) with or without 10^4 U of CSF-1 per ml. Cells were then either harvested or treated for a further 4 h with 200 ng of PMA per ml. Cell treatments were as follows: control, 16 h without CSF-1; +CSF-1, 16 h with CSF-1; +PMA, PMA added to medium for 4 h after incubation without CSF-1 for 16 h; and +CSF-1 +PMA, PMA added to medium for 4 h after incubation with CSF-1 for 16 h. Total RNA was extracted and hybridized with cDNA probes for uPA and *Ets-2*. An oligonucleotide probe for 18S rRNA is shown as a control for loading.

Canada) was inserted into a cytomegalovirus promoter-driven expression plasmid. The expression vector pECE (23) was used as a control in cotransfection experiments. CAT assays were performed as described previously (12), and luciferase assays were performed as described in Promega protocols. Total protein was assayed with a Bio-Rad Bradford assay kit.

NIH 3T3 fibroblasts were transfected by calcium phosphate coprecipitation and assayed for secreted human placental alkaline phosphatase as described previously (49). The *v-fms* expression plasmid pSMFeSV (62) was used in cotransfections.

Electrophoretic mobility shift analysis (EMSA). Extracts were prepared from approximately 3×10^7 BMM nuclei by a modification of the method of Osborn et al. (48). Cells were harvested, washed with PBS, and resuspended in 5 ml of hypotonic wash buffer (HWB) (10 mM HEPES [*N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid] [pH 7.9], 10 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA [ethylene glycol-bis(β -aminoethyl ether)-*N,N,N',N'*-tetraacetic acid]). After the extracts were pelleted and resuspended in 0.5 ml of HWB, 1 ml of HWB containing 0.05% Nonidet P-40 was added. Nuclei were spun down ($500 \times g$, 5 min, 4°C; Heraeus Biofuge) and resuspended in 50 to 100 μ l of nuclear extraction buffer (20 mM HEPES [pH 7.9], 0.42 M NaCl, 20% glycerol, 0.2 mM EDTA, 0.2 mM EGTA) and left on ice for 10 min. Nuclei were spun down ($15,000 \times g$, 10 min, 4°C; Heraeus Biofuge) and the supernatant was removed and stored at -20°C . All solutions were ice cold and contained 1 mM phenylmethylsulfonyl fluoride, 1 mM dithiothreitol, 10 μ g of leupeptin per ml, and 10 μ g of aprotinin per ml. Protein extract (1 to 5 μ g in 2 μ l of nuclear extraction buffer) was incubated with 0.027 pmol of end-labelled double-stranded oligonucleotide probe in a total volume of 10 μ l containing 20 mM HEPES (pH 7.9), 0.5 mM dithiothreitol, 12% glycerol, 50 mM NaCl, 0.5 μ g of poly[d(I-C)], and 1 mM EDTA. After the addition of protein, samples were incubated at room temperature for 30 min and run on 5 to 6% acrylamide gels (acrylamide-to-bis ratio, 29:1) at 100 V in 0.5 \times Tris-borate-EDTA buffer. The gels were dried and visualized by autoradiography.

Mouse *Ets-2* DNA binding domain (amino acids 334 to 466) was expressed in *E. coli* as a glutathione *S*-transferase (GST) fusion protein with the vector pGEX2T (Pharmacia). Protein was purified with glutathione-agarose (Sigma), and 10 ng of protein was used in EMSA as described above, with the omission of poly[d(I-C)] from the binding reaction. Expressed and purified GST protein was used as a control.

RESULTS

Northern analysis of uPA in macrophages. The effects of CSF-1 and PMA on uPA mRNA levels were examined with BMM and the CSF-1-independent macrophage-like cell line RAW264 (Fig. 1). Cells were plated overnight either with or without CSF-1 and then either harvested or treated for a further 4 h with PMA. The level of uPA mRNA was approximately 40-fold higher in CSF-1-treated BMM than in CSF-1-starved cells. The decline in uPA mRNA level with overnight

CSF-1 starvation is fully reversible with the readdition of CSF-1 (71), so the decrease in mRNA level is not due to cell death. RAW264 cells also responded to CSF-1, giving an induced level of uPA mRNA that was approximately 25% of that in CSF-1-treated BMM. In six experiments, RAW264 cells responded to CSF-1 with three- to sixfold inductions of uPA mRNA during 8- to 24-h incubations (71, 72). In experiments examining the time course of induction of uPA mRNA by CSF-1 in both BMM and RAW264 cells, mRNA levels were increased after 1 to 2 h, reached a maximum at 8 to 16 h, and remained elevated above basal levels thereafter, provided that CSF-1 concentrations remained saturating (25, 30, 72).

The tumor-promoting phorbol ester PMA is believed to exert most of its effects by activation of PKC. PMA alone had a moderate effect on uPA levels in BMM but gave a very large increase in uPA levels in RAW264 cells (Fig. 1). It has been suggested that similarities in the actions of CSF-1 and PMA in macrophages are due to a common activation of PKC (75). If PMA and CSF-1 both induce uPA by activation of PKC, then treatment with optimal concentrations of the two agents together should not have an additive effect. Adding PMA to CSF-1-pretreated BMM increased uPA mRNA levels 400-fold above the level for CSF-1-starved cells (Fig. 1). Since CSF-1 and PMA added alone gave 40- and 20-fold inductions, respectively, this effect is much greater than additive. The synergism of the two factors added at saturating concentrations indicates that CSF-1 does not induce uPA by merely activating PKC in a manner similar to that of PMA. Although CSF-1 and PMA may activate some of the same pathways, they do so in distinct manners such that synergy is possible. In RAW264 cells the effect of CSF-1 and PMA together was also more than additive. The effect was less striking than in BMM because of the large level of induction by PMA alone.

Ets family transcription factors are implicated in CSF-1 induction of the *NVL3* and *c-myc* genes and in mitogenesis in fibroblasts (39, 61). *Ets* factors are a family defined by homology to the *v-Ets* oncogene product of E26 erythroblastosis virus and include the products of the *ets-1*, *ets-2*, *erg*, *fli-1*, *spi-1* (PU.1), *elf-1*, and *PEA3* genes (78). PU.1 is a macrophage- and B-cell-specific transcription factor (57), but PU.1 expression and interaction with its target sequence are not affected by CSF-1 (58). *Ets-2* protein and mRNA were found to be induced in chicken BMM by the chicken myelomonocytic growth factor and PKC activators (8). In the mouse macrophage cell line BAC1.2F5, *ets-2* mRNA was induced by CSF-1 (9). Examination of *ets-2* mRNA expression in macrophages showed induction by PMA and CSF-1 closely paralleling that of uPA mRNA, with apparent synergism between the two stimuli (Fig. 1). A time course of the response to CSF-1 in BMM showed a peak of *ets-2* mRNA expression at 1 to 2 h, preceding the major induction of uPA (25). In RAW264 cells, the response of *ets-2* mRNA to CSF-1 was barely detectable even at 2 h (25), although the message was induced by PMA (Fig. 1). The possibility that transcription factor *Ets-2* can regulate uPA expression is addressed later in this paper. If induction of *Ets-2* is a prerequisite for uPA induction, then it would be anticipated that blockage of protein synthesis would prevent uPA induction. Investigations of such events are difficult, as both *ets-2* mRNA and uPA mRNA are induced in macrophages by cycloheximide (14, 25).

Effects of PMA and CSF-1 on rates of transcription. Nuclear run-on analyses were performed to establish whether the increases in mRNA levels observed with PMA and CSF-1 were due to transcriptional activation. Figure 2A shows results of an analysis of the rates of transcription of a number of genes in BMM treated with CSF-1 and PMA. Two mouse uPA clones




FIG. 2. Nuclear run-on analysis of transcription of the uPA, β -actin, *c-fms*, and lysozyme genes in BMM and RAW264 cells. uPA5' is a genomic fragment of uPA including exon 1 and intron 1. uPA3' is a cDNA clone encompassing bp 424 to 1643 of the 2.3-kb mouse uPA cDNA. pGEM2 plasmid DNA was used as a control. (A) Six-day-old BMM were starved of CSF-1 for 15 h (control), grown in the continued presence of CSF-1 (+CSF-1), or starved of CSF-1 for 15 h prior to the addition of PMA for 1 h (+PMA) before the isolation of nuclei. (B) RAW264 cells were grown without additions (control), with CSF-1 for 12 h (+CSF-1 12 hr), with CSF-1 for 4 h (+CSF-1 4 hr), or with PMA for 1 h (+PMA). Seven-day-old BMM grown in the continued presence of CSF-1 (+CSF-1) were assayed in the same experiment. These assays were performed with duplicate cell samples, and results are representative of two independent run-on analyses. CSF-1 was used at 10^4 U/ml, and PMA was used at 200 ng/ml.

were used. uPA 5' is a genomic clone containing the noncoding first exon, the first intron, and the beginning of exon 2. uPA 3' is a cDNA clone containing bp 424 to 1643 (exons 5 to 11) of the 2.3-kb uPA mRNA (4, 20). These two clones were used in order to distinguish between effects on initiation of transcription and effects on elongation. Earlier results have shown transcriptional termination or pausing in the second intron of the uPA gene in both fibroblasts and thioglycolate-elicited peritoneal macrophages (3). The results here do not suggest that release of a transcriptional blockage is a major form of regulation of the uPA gene in BMM or RAW264 cells. Both CSF-1 and PMA gave a clear increase in the rate of transcription of uPA in BMM (Fig. 2A). In RAW264 cells, PMA treatment increased the rate of transcription, but there was no detectable change when cells were treated with CSF-1 for either 4 or 12 h (Fig. 2B). This suggests that the moderate level of induction of uPA mRNA by CSF-1 in RAW264 cells (Fig. 1) occurs at a posttranscriptional level.

The rates of uPA transcription in BMM and RAW264 cells were compared by using equivalent counts per minute of run-on transcripts from the two cell types (Fig. 2B). Such an approach assumes that the total nascent transcripts produced per cell per unit of time is independent of the cell type. This is unlikely to be the case, as RAW264 cells proliferate more rapidly than BMM. However, even if it is assumed that RAW264 nuclei incorporated twice as much radiolabel as BMM nuclei, the rate of transcription of uPA in BMM treated with CSF-1 was still apparently much greater than in RAW264

cells with or without CSF-1 (Fig. 2B). The results of Northern and nuclear run-on analyses suggest that the CSF-1 responsiveness of the uPA gene is impaired in RAW264 cells compared with that in BMM. CSF-1 and PMA treatments also affected the transcription of other genes studied. In both RAW264 cells and BMM, transcription of *c-fms* and the lysozyme gene, which are macrophage marker genes, was reduced by CSF-1 and PMA. Both of these agonists are known to down-regulate *c-fms* mRNA in these cells (85).

Promoter studies in RAW264 cells. In previous work, we showed that RAW264 cells could be effectively transfected by electroporation but that BMM could not be transfected by the same method because of DNA-dependent cell death (73). Unfortunately no macrophage cell line which is readily transfectable and gives a good transcriptional response to CSF-1 has yet been identified. Previously reported analysis of the mouse uPA promoter in RAW264 cells (12) showed that maximal expression required the first intron and 6.6 kb of promoter sequence. The full-length (6.6-kb) uPA promoter was induced sixfold by PMA, but no induction was seen with CSF-1 (72). Basal expression of the pig promoter transfected into RAW264 cells was dependent on the PEA3/AP-1 element first described by Rørth et al. (12, 56).

Rørth et al. (56) prepared plasmid pPR99 containing a 90-bp *Hae*III fragment with the PEA3/AP-1 element (−2446 to −2356) linked to the uPA-proximal promoter region (−114 to +398) and the CAT reporter gene (Fig. 3E). The activity of this construct in RAW264 cells was compared with that of the full-length promoter in puPACAT2 (−6.6 kb to +398) and that of the promoter truncated to −2657. Truncation of the promoter to −2657 reduced the basal level of expression (12) but gave little change in PMA inducibility (Fig. 3A). Thus, regions upstream of −2657 have enhancer activity (12) but are not required for the PMA response. The PMA-induced level of expression from pPR99 is seven to eight times that of the full-length promoter in puPACAT2. The higher level of inducibility may be due to the deletion of inhibitory regions, such as those found in the human promoter between −660 and −338 (11) and between −1824 and −1572 (77), both of which are 3' to the PEA3/AP-1 element.

Mutating either the PEA3 half (plasmid pPR119) or the AP-1 half (plasmid pPR127) (56) of the PEA3/AP-1 site greatly reduced the PMA response (Fig. 3B). As the basal level of expression from these plasmids was very close to the experimental background level, it was not possible to accurately determine the fold induction by PMA. Using analogous constructs in a luciferase vector in which low-level activity is more easily detectable, we showed that the PEA3 and AP-1 mutations caused only a 50% drop in activity (25). Hence, the reductions in PMA-induced activity in pPR119 and pPR127 are not likely to be due to a drop in basal expression. The PEA3/AP-1 element can mediate a response to PMA in RAW264 cells, and both halves of the site are required for maximal induction. A plasmid with only the uPA-proximal promoter (−114 to +398), pPR126, was not PMA responsive (Fig. 3B).

Figure 1 shows that the expression of *ets-2* mRNA closely paralleled that of uPA. The *ets-2* and uPA genes may be regulated together through common pathways, or Ets-2 levels may control uPA mRNA levels. Since the peak of *ets-2* mRNA expression precedes most of the uPA mRNA induction (25) and PEA3 sequences are bound by Ets family transcription factors (79), the second possibility is an attractive hypothesis. The relatively impaired response of RAW264 cells to CSF-1 may be a consequence of the failure to induce Ets-2. For this reason, the effect of Ets-2 expression on promoter activity in




FIG. 3. (A) PMA response of uPA promoter constructs in transiently transfected RAW264 cells. The cells were electroporated with 10 μ g of puPACAT2 and molar equivalent amounts of puPACAT2-2657 and pPR99. The cells were divided into two groups for treatment with or without 200 ng of PMA per ml at 24 h after transfection. The cells were harvested after a further 24 h in culture. CAT activity was normalized to that of puPACAT2 without PMA treatment. (B) Involvement of the PEA3/AP-1 site in the PMA response. RAW264 cells were electroporated with 10 μ g of uPA promoter construct, and the cells were divided into two groups for treatment with or without 100 ng of PMA per ml at 24 h after transfection. The cells were harvested after a further 24 h in culture. CAT activity was normalized to the results for pPR99 treated with PMA. Results shown are the averages of four separate experiments (error bars show standard deviations). Plasmid constructs are shown in Fig. 3E. pPR126 contains only the uPA proximal promoter region (-114 to +398) linked to the CAT gene (56). (C) Involvement of the PEA3/AP-1 site in the response to Ets-2 expression. RAW264 cells were electroporated with 10 μ g of uPA promoter construct and 1 μ g of either an Ets-2 expression plasmid (Ets.pECE) or control expression vector (pECE). Cells were harvested 48 h after transfection. CAT activity was normalized to results for pPR99 with cotransfected Ets.pECE. Results shown are the averages from four separate experiments (error bars show standard deviations). (D) Transactivation of the uPA promoter by various Ets family members. RAW264 cells were electroporated with 10 μ g of pGLuPA-6.6 or a molar equivalent amount of pGL99 together with 2 μ g of Ets family expression plasmids or a control plasmid, pECE. Cells were harvested 24 h after transfection. Results were normalized to those for cotransfection of each of the reporter plasmids with pECE. Absolute values obtained for pGL99 were approximately 25% of those for pGLuPA-6.6. Results from three separate experiments (averages of duplicates) are indicated by the solid, hatched, and shaded bars. (E) Diagram of uPA promoter constructs used in transfection experiments. The uPA exonic sequences included in constructs are noncoding sequences (20). Derivatives of plasmid pPR99 contained the mutations shown in the PEA3/AP-1 site (56).

RAW264 cells was examined. Figure 3C shows the results of the cotransfection of an Ets-2 expression plasmid (Ets.pECE) with pPR99 and mutated constructs into RAW264 cells. Ets-2 expression dramatically increased the activity of the uPA promoter in pPR99. Mutations of the PEA3 site (pPR119) and the

AP-1 site (pPR127) both reduced the response to cotransfected Ets-2. It is likely that Ets-2 activates transcription directly through the PEA3/AP-1 site, but the possibility that Ets-2 induces another factor involved in transactivation cannot be ruled out. The fact that the AP-1 mutation decreases the

response to Ets-2 emphasizes the compound nature of this element; both halves of the element are required for full enhancer activity. The uPA proximal promoter region alone in pPR126 did not respond to Ets-2. The residual Ets-2 response seen with the PEA3/AP-1 mutant constructs pPR119 and pPR127 may be due to another region within the 90-bp *Hae*III fragment which has a potential Ets factor binding sequence (GGAT at -2396 bp in the mouse promoter). This sequence is not perfectly conserved among mouse, pig, and human promoters (12), and whether it has an enhancer function has not been established.

Transactivation of the uPA promoter is not a property of all Ets factors. Figure 3D shows results of cotransfection of uPA promoter constructs with expression vectors for the Ets family members Ets-2, Elf-1, PU.1, and PEA3. The promoter constructs used were pGLuPA-6.6, which contains the same promoter region as puPACAT2 (-6616 to +398) linked to the luciferase reporter gene, and pGL99, which is a luciferase version of pPR99 (Fig. 3E). For both plasmids, the induction by Ets-2 was 60- to 130-fold. It should be noted that the level of luciferase activity from pGL99 was approximately 25% of that from pGLuPA-6.6. Expression of PU.1 and Elf-1 had little or no effect on promoter activity, and PEA3 gave only a modest increase compared with that of Ets-2. Although native PU.1 is expressed relatively highly in RAW264 cells (57), we still observe some transactivation by coexpressed PU.1 on several other promoters. Our previous work showed that PU.1 from RAW264 nuclear extracts did not bind the uPA PEA3/AP-1 element under EMSA conditions (12). An Ets-1 expression vector gave the same magnitude of response as Ets-2 (result not shown). Ets-1 is not a good candidate for regulating expression in macrophages, as it is expressed strongly in lymphoid cells but not in macrophage or myeloid cell lines (37).

Consistent with the results of nuclear run-on (Fig. 2), which show little effect of CSF-1 on uPA gene transcription in RAW264 cells, no activation of the uPA promoter by CSF-1 was detected in transfection analyses with this cell line (72). Stably integrated full-length (6.6-kb) promoter was also not responsive to CSF-1 (25). Results of transient and stable transfection therefore support the idea that the three- to sixfold induction of uPA mRNA in RAW264 cells over an 8- to 16-h treatment with CSF-1 (Fig. 1) (71, 72) is largely a posttranscriptional effect.

Regulation of the uPA gene by CSF-1 in NIH 3T3 cells. As transcriptional regulation of uPA by CSF-1 in RAW264 cells could not be demonstrated, the regulation of uPA was examined with a heterologous system, i.e., with NIH 3T3 fibroblasts expressing the human CSF-1R, *c-fms* (referred to herein as NIH 3T3/*c-fms*). The kinetics of stimulation of uPA mRNA expression by the addition of recombinant human CSF-1 to NIH 3T3/*c-fms* are depicted in Fig. 4. The level of uPA mRNA was rapidly stimulated by CSF-1 in these cells, increasing by eightfold within 1 h. The level thereafter declined but at 2 and 4 h remained threefold higher than in unstimulated cells. This level of expression was maintained even after 24 h of CSF-1 stimulation (16). The kinetics of induction are in contrast to the results found for CSF-1 stimulation of uPA mRNA in BMM (71) or RAW264 cells (72), in which expression peaks at 8 to 16 h. Similarly, the NVL3 retrotransposon in NIH 3T3/*c-fms* is maximally induced 16 to 24 h after CSF-1 stimulation (7).

To examine the role of Ets proteins in the induction of uPA, we next examined the effect of a dominant negative repressor of Ets action. Stable expression of the DNA binding domain of human Ets-2 fused to the *lacZ* gene product (39) with *c-fms* in NIH 3T3 cells prevented CSF-1 induction of uPA mRNA (Fig.




FIG. 4. Effects of CSF-1 and Ets-2 DNA binding domain on uPA mRNA expression in NIH 3T3 cells. NIH 3T3 cells were stably transfected with a construct expressing Ets-2 DNA binding domain fused to the *lacZ* gene (pAPretZ-neo) and a retroviral vector expressing the CSF-1R (39). Control cells were transfected with the parental expression vector (39) and the CSF-1 expression vector. RNA was prepared from NIH 3T3 cells expressing human *c-fms* (open circles) and NIH 3T3 cells expressing both *c-fms* and the human Ets-2 DNA binding domain fused to the *lacZ* gene product (solid circles). The cells were serum starved for 24 h and then treated with 5×10^4 U of CSF-1 per ml for the indicated times (in hours). Northern blot analysis was performed with a uPA-specific probe and a K-rev-specific probe to control for RNA loading. The data shown are the uPA hybridization/K-rev-1 hybridization ratios as determined by scanning of autoradiographs with a model SL-504-XL laser scanning densitometer (Biomed Instruments).

4). This is similar to the result obtained for CSF-1 stimulation of NVL3 expression in NIH 3T3/*c-fms*. The NVL3 gene contains a PEA3/AP-1-like element which behaves as both a ras-responsive element and a CSF-1-responsive element (7). The possibility that the uPA PEA3/AP-1 element can behave as a CSF-1-responsive element was addressed by transient transfections into NIH 3T3 cells.

uPA promoter activity in NIH 3T3/*c-fms*. Promoter elements involved in the CSF-1 response of NIH 3T3/*c-fms* were analyzed by transient transfection. Plasmid pPR99 and the mutant plasmids pPR119 and pPR127 (56) were transfected into NIH 3T3 cells with and without cotransfected *v-fms* expression plasmid, which encodes a constitutively active form of the CSF-1R (62) (Fig. 5A). The basal level of activity of pPR99 was low in NIH 3T3 cells, but *v-fms* expression activated this reporter by an average of 25-fold in three separate experiments. In contrast, the activities of reporters that contained point mutations in either the PEA3 site (pPR119) or the AP-1 site (pPR127) were not greatly stimulated by *v-fms*. In a second set of experiments, reporters were transfected into NIH 3T3/*c-fms* cells and stimulation by CSF-1 was determined (Fig. 5B). In these experiments, pPR99 activity was stimulated three- to fourfold, while the activities of pPR119 and pPR127 were not significantly affected by the ligand-activated CSF-1R. The CSF-1-mediated increase in uPA mRNA (Fig. 4) is therefore likely to involve activation of the PEA3/AP-1 element and requires Ets family transcription factors.

Proteins binding the PEA3/AP-1 element. The transient-transfection analyses described above implicate AP-1 or cAMP response element (CRE)-binding and Ets family transcription factors in the action of CSF-1 and PMA. EMSA was performed in order to determine whether AP-1 transcription factors are able to bind the octameric AP-1 site of the uPA element and whether any proteins binding the PEA3 site are detectable.

In EMSA the uPA PEA3/AP-1 element (uPA wt oligonucleotide) formed two major complexes with nuclear proteins from CSF-1-treated BMM (Fig. 6A). These complexes were abolished by mutation of the AP-1 half of the element (uPA




FIG. 5. *fms* response of uPA promoter-CAT constructs transfected into NIH 3T3 cells. (A) NIH 3T3 cells were transiently transfected by calcium phosphate coprecipitation with 3 μ g of uPA-CAT reporter construct in the presence (+v-fms) or absence (-) of 2.5 μ g of the *v-fms* expression plasmid pSMFeSV (62). (B) NIH 3T3/*c-fms* cells were transfected with 3 μ g of uPA promoter-CAT reporter constructs. After the precipitate was removed, the cells were either grown in the presence of 10^4 U of CSF-1 per ml for 48 h and restimulated with CSF-1 at 2 h before harvest (+CSF-1) or grown in the absence of CSF-1 (-). CAT assays were done with equivalent amounts of protein as determined by protein concentration. A plasmid expressing a secreted form of human placental alkaline phosphatase was used as an internal control for transfection efficiency. Thin-layer chromatography plates were autoradiographed and then analyzed with a Betascope 603 blot analyzer (Betagen, Waltham, Mass.). Error bars in both panels show the standard deviations of results from three separate experiments.

M5). Mutation of the PEA3 site (uPA M7) or use of a truncated oligonucleotide completely lacking the PEA3 site (uPA CRE) did not affect the formation of these complexes. Thus, the major proteins detectable under the conditions used bind only to the AP-1 half of the element. The possibility that phosphorylation may be necessary for factor binding to the PEA3 half was considered, but inclusion of phosphatase inhibitors in extraction and binding buffers did not alter the pattern of binding (25).

Incubation of CSF-1-treated BMM nuclear extract with an oligonucleotide able to bind AP-1 yielded one band (Fig. 6B). Cold uPA wt oligonucleotide was able to compete for proteins binding to the AP-1 site but was a little less efficient than self-competition with cold AP-1 oligonucleotide. Mutation of the AP-1 half of the uPA element (M8) abolished this competition, whereas mutation of the PEA3 half (M5) did not. Thus, the octameric site in the uPA PEA3/AP-1 element (TGAGGTCA) can bind some of the same factors which bind to a conventional AP-1 site (TGAGTCA).

Treatment of BMM with CSF-1 or PMA increased the levels of proteins binding to the AP-1 oligonucleotide (Fig. 6C). With CSF-1 treatment, levels were highest at 1 h and subsequently




FIG. 6. (A) EMSA of proteins binding to the uPA PEA3/AP-1 element and mutated oligonucleotides. Nuclear extract was prepared from BMM starved of CSF-1 for 16 h and then treated with 10^4 U of CSF-1 per ml for 3 h. Nuclear protein (4.5 μ g) was incubated with four different oligonucleotide probes labelled to achieve the same specific activity. Binding reactions were run on a 5% acrylamide gel. Lane wt probe, no protein extract. Arrows indicate protein-DNA complexes. (B) Competition for proteins binding to a labelled AP-1 oligonucleotide. AP-1 oligonucleotide was mixed with the indicated fold molar excess of cold competitor before the addition of 1 μ g of nuclear protein. Nuclear extract was made from BMM grown in the presence of CSF-1. Binding reactions were run on a 6% polyacrylamide gel. Free probe has been run off the bottom of the gel. Lane no comp., no competing cold oligonucleotide. (C) Effect of treatment of BMM with CSF-1 and PMA on AP-1 binding. BMM were either starved of CSF-1 overnight (-CSF-1 O/N) and subsequently treated with CSF-1 for 1 or 3 h or with PMA for 1 h or left with continued CSF-1 stimulation (+CSF-1 O/N) and subsequently treated with PMA for 1 h (+CSF-1 O/N +PMA 1hr). Nuclear protein (1 μ g) was incubated with labelled AP-1 oligonucleotide before being run on a 5% polyacrylamide gel. Oligonucleotides used were as follows: AP-1, 5'-CGCTTGATGAGTCAGCCGGAA3' (Promega); uPA wt, 5'-GGCCAGGAG GAAATGAGGTCATCTTGCT3' (uPA PEA3/AP-1); uPA M5, 5'-GGCCAG GAGGAAATGAGGATCTTGCT3' (AP-1 mutant); uPA M7, 5'-GGCCAG TACCAAATGAGGTCATCTTGCT3' (PEA3 mutant); uPA M8, 5'-GGCCAG GACCAAATGAGGTCATCTTGCT3' (PEA3 mutant); and uPA CRE, 5'-GA AATGAGGTCATCTTGCT3' (AP-1 half-site).

declined. However, CSF-1 and PMA did not synergize to increase AP-1 binding activity. In fact, CSF-1 pretreatment prevented induction of AP-1 binding activity by PMA. The same pattern of inductions was found with labelled uPA wt oligonucleotide, although the degree of induction was less pronounced (result not shown). The low level of protein binding in extracts from cells treated with CSF-1 and PMA together indicates that uPA promoter activity does not correlate directly with the induction of AP-1 binding activity.

DNA binding of Ets-1 and Ets-2 is known to be impeded by a repressive domain under the conditions of EMSA (67). High-affinity binding may require either posttranslational modification or protein-protein interaction to relieve this repressive




FIG. 7. Binding of the Ets-2 DNA binding domain to the uPA PEA3/AP-1 element. (A) Purified bacterially expressed GST-Ets-2 fusion protein and control GST protein were incubated with labelled uPA wt oligonucleotide. When cold competitors were used, they were preincubated with the protein for 15 min before the addition of the labelled probe. Samples were loaded onto a 6% polyacrylamide gel 15 min after the addition of labelled probe. (B) Competition for GST-Ets-2 fusion protein binding to the labelled uPA wt oligonucleotide by either cold uPA wt or Ets-2 consensus oligonucleotides. Labelled probe and the indicated molar excesses of cold competitor were mixed together before the addition of protein. After 30 min, samples were loaded onto a 6% polyacrylamide gel. The Ets-2 consensus oligonucleotide (5'CTAGGACCGGAAGTGG GAGT3') is based on sequences established by R. Maki (La Jolla Cancer Research Foundation), using recombinant Ets-2 DNA binding domain and random oligonucleotides. Other oligonucleotides used are as indicated in the legend to Fig. 6.

effect. Perhaps because of this problem and the lack of good supershifting antibodies specific for Ets-2, to our knowledge native Ets-2 binding activity in nuclear extracts has not been clearly identified. Even using a consensus sequence determined for binding of recombinant Ets-2 DNA binding domain (44), we could not detect a PMA- or CSF-1-inducible protein in macrophages (result not shown), although *ets-2* mRNA is clearly induced (Fig. 1). To examine whether Ets-2 can bind the uPA PEA3/AP-1 element, the DNA binding domain alone was expressed as a GST fusion protein. Figure 7A shows that the Ets-2 DNA binding domain binds the uPA wt oligonucleotide and can be competed for by the AP-1 mutant (uPA M5) and an Ets-2 consensus oligonucleotide but not by the PEA3 mutant (uPA M8). On the basis of competition with increasing concentrations of cold oligonucleotides, the uPA wt sequence was evidently a much lower-affinity site for Ets-2 binding than the Ets-2 consensus sequence (Fig. 7B).

	PEA3	AP-1
Polyoma virus	AGCAGGAAG	-TGAC-TAACT
mouse uPA	AGGAGGAAA	-TGAGGTCATC
mouse NVL3	CACAGGATA	-TGAC-TCTTT
human TGF- β 1	ACGGAAGGAGAG	-TCAGG
human scavenger receptor	GAAAGGAAA	-TGAC-ACATT
	PEA3	AP-1
human collagenase	GAGGATGTTATAAAGCAT	TGAGTCA
human keratin 18	AAGCGGATGTGGCTAAGGCT	TGAGTCA

FIG. 8. Functional PEA3/AP-1 elements identified in a number of genes. Ets core binding sequences (GGAA or GGAT) (78) are underscored, and AP-1-like sequences are indicated by dashed underlining.

DISCUSSION

A PEA3/AP-1 element was first identified in the polyoma-virus enhancer. It mediated responses to serum, PMA, and the *v-src*, *Ha-ras*, *v-mos*, *v-raf*, and polyomavirus middle-T-antigen oncogenes (80) and has been termed a tetradecanoyl phorbol acetate- and oncogene-responsive unit (28). Similar elements have been found in other genes such as collagenase (28), transforming growth factor β (50, 54), the NVL3 retrotransposon (54), keratin 18 (51), uPA (56), and the macrophage scavenger receptor (82) (Fig. 8). These elements have all been characterized as ras-responsive elements. Although Rørth et al. (56) mention that ras transformation of keratinocytes did not activate the uPA PEA3/AP-1 element, the element responded strongly to activated *ras* in NIH 3T3 cells (16) and RAW264 macrophages (25). The collagenase and keratin 18 PEA3 and AP-1 elements are more widely spaced (Fig. 8), but it is interesting that the insertion of 10 bp in both cases corresponds to one turn of the DNA helix and may give a similar orientation of binding factors.

As noted in the introduction, the uPA PEA3/AP-1 site at -2430 in the mouse promoter mediates induction by signals such as epidermal growth factor, PMA, and basic fibroblast growth factor. In this work we have found that the uPA PEA3/AP-1 element is involved in the transcriptional response to CSF-1 in NIH 3T3/*c-fms*. Both halves of the element were required for efficient activation of transcription by either *v-fms* or *c-fms* with CSF-1 (Fig. 5). This may reflect the fact that the PEA3 and AP-1 halves are comparatively weak binding sites for Ets and AP-1 proteins (Fig. 6 and 7). Induction of uPA mRNA by CSF-1 was prevented by both a dominant negative Ets protein (Fig. 4) and the catalytic domain of GAP (16). uPA induction by CSF-1 in this cell line therefore appears to involve ras signalling and activation of Ets transcription factors. This makes uPA regulation similar to that of both the NVL3 retrotransposon and *c-myc* (6, 7, 39, 61). Work on the induction of the NVL3 retrotransposon by CSF-1 in NIH 3T3/*c-fms* identified a PEA3/AP-1-like element mediating the CSF-1 response (7, 54). This element had been previously identified as a ras-responsive element (49). Another CSF-1-inducible gene, the macrophage scavenger receptor gene, contains a similar ras- and PMA-responsive element, although this element has not yet been directly shown to be activated by CSF-1 (82). Comparison of the NVL3, uPA, and scavenger receptor PEA3/AP-1 elements (Fig. 8) shows similar spacings of elements but a number of sequence differences. Wu et al. (82) found that Ets-2, but not Ets-1, was able to transactivate the scavenger receptor element and to cooperate with c-Jun. This differs from results for the uPA element in RAW264 cells, in which Ets-2 and Ets-1 have similar activities.

Introduction of a mutation at Tyr-809 in the CSF-1R prevents induction of *c-myc*, the NVL3 gene, and the mitogenic response in NIH 3T3/*c-fms* treated with CSF-1 (39, 60, 64). These effects are mimicked by the expression of dominant negative Ets-2 (39). CSF-1-induced mitogenesis can be restored to cells with the Tyr-809 mutation by the expression of *ets-1*, *ets-2*, or *c-myc* (39, 60, 61), and induction of *c-myc* appears to involve Ets factors (61). This implicates an Ets factor with binding specificity and activity similar to those of Ets-1 and Ets-2 in the CSF-1 signalling pathway in these cells. The identification of the NVL3 and uPA PEA3/AP-1 sequences as promoter targets of CSF-1 is an important step towards finding the specific transcription factors involved in signalling.

The 3' half of the uPA PEA3/AP-1 site is octameric and more closely resembles a CRE (TGACGTCA) than it does the consensus AP-1 site (TGA[C/G]TCA). The distinction between CRE and AP-1 sites has become blurred by the findings that AP-1 factors can dimerize with the CRE-binding factors (CREB/ATF family) (29) and that AP-1 factors themselves can also bind to octameric sequences (1, 65). The wide range of Ets family proteins and AP-1 and CREB/ATF proteins makes the determination of which factors bind these elements a challenge. The octameric site in uPA is not responsive to cAMP and functions similarly to an AP-1 site (40). In extracts from pig kidney epithelial cells, c-Jun has been found to bind the pig uPA PEA3/AP-1 element (41). In another study, a c-Jun/ATF2 heterodimer has been shown to bind the AP-1 half of the uPA PEA3/AP-1 element (19). Macrophage AP-1-binding factors will also clearly bind to the octameric uPA AP-1 sequence (Fig. 6), although with slightly lower affinity than to a conventional AP-1 site. When either a labelled AP-1 oligonucleotide (Fig. 6C) or the uPA PEA3/AP-1 element (25) was used, treatment with CSF-1 or PMA increased levels of AP-1 factor binding. However, CSF-1 pretreatment prevented induction of AP-1 factors by PMA. Hence, the synergy seen in induction of uPA mRNA (Fig. 1) is not due to synergistic induction of general AP-1 binding activity.

The effect of PEA3 site mutation in transient-expression analysis (Fig. 3) showed that this element was functionally important. Despite this, with EMSA we were unable to detect any strong binding of macrophage nuclear factors to the PEA3 half of the uPA element. The two major detectable bands were both accounted for by proteins binding to the AP-1 site (Fig. 6). Conditions used in making nuclear protein extracts or in EMSA may be inadequate for detecting factors binding to the PEA3 site. In EMSA with extract from pig kidney LLC-PK₁ cells, only one complex was observed; this result was explainable by c-Jun binding to the AP-1 half of the pig uPA PEA3/AP-1 element (41). However, *in vivo* footprinting showed the uPA PEA3 site to be constitutively occupied regardless of enhancer activity (46). Owen and Ostrowski (50) found a 120-kDa protein in NIH 3T3 which can bind to the NVL3 PEA3 site and does not correspond to known Ets factors. In studies of the scavenger receptor element, large amounts of protein were required before evidence of binding to the PEA3 half of the element was observed (82).

Mutation of the PEA3 site reduced the high level of transactivation by Ets-2 (Fig. 3C), suggesting that Ets-2 acts through this site. Bacterially expressed Ets-2 DNA binding domain was clearly capable of binding to this element, but it is a lower-affinity site than the consensus Ets-2 binding site (Fig. 7). It is impossible to determine what represents sufficient affinity to be of biological relevance, especially as the uPA site is a compound element and cooperative binding could increase the effective affinity. The evidence that Ets-2 can transactivate through this site, is capable of binding to it, and is induced by

CSF-1 makes it a good candidate for a normal regulator of uPA *in vivo*.

The macrophage cell line RAW264 can be readily transfected and responds to CSF-1 with a three- to sixfold induction of uPA mRNA (Fig. 1) (71, 72). In contrast to that of BMM, this induction did not seem to be mediated at the transcriptional level (Fig. 2). The rate of transcription of uPA in RAW264 cells was low compared with that in CSF-1-stimulated BMM (Fig. 2), leading us to propose that in RAW264 cells there is a deficiency in the CSF-1 signal transduction pathway leading to uPA gene transcription. Recent results comparing protein tyrosine phosphorylation levels in RAW264 cells and BMM show similar levels of induction of phosphorylation by CSF-1 in the two cell types, including phosphorylation of mitogen-activated protein kinases (25). Thus, there appears to be no fundamental defect in CSF-1R function in RAW264 cells. RAW264 cells are transformed by Abelson leukemia virus, which presumably encodes an active *v-abl* tyrosine kinase. This tyrosine kinase activity may lead to the down-regulation of a factor in the CSF-1 signalling pathway.

A simple hypothesis to account for the poor level of induction of uPA by CSF-1 in RAW264 cells is that uPA transcription is regulated by Ets-2 and CSF-1 fails to induce *ets-2* mRNA efficiently in the cell line (Fig. 1). If Ets-2 is involved in the normal regulation of the uPA gene, then the synergistic induction of uPA mRNA by CSF-1 and PMA could be at least partly accounted for by synergistic effects of the two agents on *ets-2* mRNA levels (Fig. 1).

An alternative interpretation is that *ets-2* and uPA are regulated in parallel, via the activation of another Ets protein. However, the time course of *ets-2* mRNA induction and decay is quite different from that for uPA (25). The *ets-2* gene promoter itself contains an Ets factor binding site involved in expression of the gene, and protein binding to an adjacent promoter element was competed for, although not efficiently, by an AP-1 oligonucleotide (45). Whether this element functions in the same way as the uPA element remains to be established. Other Ets factors expressed in macrophages include PU.1 (57), Elf-1 (32), and Fli-1 (36). Neither PU.1 nor Elf-1 was able to effectively transactivate the uPA promoter (Fig. 3D), although the uPA PEA3 element is similar to a published Elf-1 binding site (AGGAGGAAAA [74]) and an Elf-1 binding site is closely associated with an AP-1 site in the human interleukin-3 gene (27).

Although analysis of the CSF-1 response in transfected RAW264 cells was not possible, the PEA3/AP-1 element was identified as a PMA-responsive element in this cell line (Fig. 3B). The same element mediated a response to PMA in keratinocytes (56) and in the pig kidney epithelial cell line LLC-PK₁ (41). In HepG2 hepatoma cells, the cooperation between the PEA3/AP-1 site and a downstream AP-1 site was necessary for induction by PMA (47).

Although both PMA and CSF-1 are able to activate the PEA3/AP-1 element, the evidence does not support the hypothesis that the pathways of activation are coincident. Both the synergy between CSF-1 and PMA in induction of uPA mRNA in BMM (Fig. 1) and the differential response of RAW264 cells to CSF-1 and PMA in transient transfections suggest that CSF-1 and PMA induce uPA by distinct mechanisms. PMA does not mimic the action of CSF-1. The two agonists may activate separate pathways or may have synergistic effects on the one signalling pathway. Although some evidence that CSF-1 activates PKC has been presented (34, 66), a recent study suggests that PKC activation is not significant, at least not in early signalling by CSF-1 (35). Like CSF-1 (2), PMA and PKC can activate Raf-1 (38). It will be interesting to

ascertain whether PMA can increase Raf-1 activation when added to CSF-1-treated cells.

PMA is frequently used in promoter studies without indication of its physiological relevance. There may be natural stimuli which act through PKC and strongly enhance the induction of uPA by CSF-1. Transforming growth factor β induces uPA in RAW264 cells and may act through a PKC-dependent mechanism (24). It is also possible that PMA provides a combination of stimuli to cells which does not resemble the action of any physiological molecule.

In this work we have provided evidence that the CSF-1 response of the uPA gene is distinct from the response to PMA, is mediated through a PEA3/AP-1 element, and requires activation of an Ets transcription factor, possibly Ets-2. Earlier evidence pointed to bifurcating signalling pathways from the CSF-1 receptor, one through ras, raf, Ets, and myc factors and another leading to induction of fos and jun proteins (7, 9, 39, 60). Although ras is generally considered an activator of Raf-1, recent work showed that ras and Raf-1 were activated independently by CSF-1 in the macrophage cell line BAC1.2F5 (10) and are therefore likely to be part of cooperating but separate pathways. Since expression of *v-raf* in BAC1.2F5 macrophages leads to constitutive expression of *ets-2* mRNA (9), induction of *ets-2* by CSF-1 may be due to activation of a pathway involving Raf-1. We have found that coexpressed Ets-2 and activated ras in RAW264 cells synergistically increase uPA promoter activity (25). This is consistent with a model in which a Raf-1-activated pathway induces Ets-2 expression and ras either activates Ets-2 or induces cooperating factors.

Many studies of the CSF-1 signal transduction pathway have been performed with fibroblasts expressing the CSF-1R. This is partly due to the difficulties of transfection and manipulation of macrophages. In this study we have used NIH 3T3 cells expressing *c-fms*, as the transfectable macrophage cell lines available to us did not respond adequately to CSF-1 with transcriptional activation of the uPA gene. Differences in CSF-1 signalling between fibroblasts and macrophages have been detected (43), and the challenge remains to confirm with macrophages the results of numerous studies performed with fibroblasts.

ACKNOWLEDGMENTS

This work was supported by grants from the National Health and Medical Research Council of Australia and the Queensland Cancer Fund to D.A.H. and National Cancer Institute grant CA5327 to M.C.O. Michael Colman was supported by a National Cancer Institute training grant in viral oncology.

CSF-1 was generously provided by Chiron Corp.

REFERENCES

1. Angel, P., and M. Karin. 1991. The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim. Biophys. Acta* **1072**:129-157.
2. Baccarini, M., D. M. Sabatini, H. App, U. R. Rapp, and E. R. Stanley. 1990. Colony stimulating factor-1 (CSF-1) stimulates temperature dependent phosphorylation and activation of the RAF-1 proto-oncogene product. *EMBO J.* **9**:3649-3657.
3. Belin, D., and J.-D. Vassalli. 1993. Transcription of the murine uPA gene, p. 36. *In* F. Blasi, K. Danø, J. Quigley, D. Rifkin, and S. Strickland (ed.), *Molecular and cellular biology of plasminogen activation*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
4. Belin, D., J.-D. Vassalli, C. Combépine, F. Godeau, Y. Nagamine, E. Reich, H. P. Kocher, and R. M. Duvoisin. 1985. Cloning, nucleotide sequencing and expression of cDNAs encoding mouse urokinase-type plasminogen activator. *Eur. J. Biochem.* **148**:225-232.
5. Besser, D., and Y. Nagamine (Friedrich Miescher-Institut, Basel, Switzerland). 1994. Personal communication.
6. Bortner, D. M., M. Ulivi, and M. C. Ostrowski. 1992. Constitutive expression of c-myc in NIH3T3 cells which co-express the CSF-1 receptor and the catalytic domain of the GTPase-activating protein restores CSF-1 dependent mitogenic growth. *Int. J. Oncol.* **1**:221-225.
7. Bortner, D. M., M. Ulivi, M. F. Roussel, and M. C. Ostrowski. 1991. The carboxy-terminal catalytic domain of the GTPase-activating protein inhibits nuclear signal transduction and morphological transformation mediated by the CSF-1 receptor. *Genes Dev.* **5**:1777-1785.
8. Boulukos, K. E., P. Pognonec, E. Sariban, M. Bailly, C. Lagrou, and J. Ghysdael. 1990. Rapid and transient expression of Ets2 in mature macrophages following stimulation with cMGF, LPS, and PKC activators. *Genes Dev.* **4**:401-409.
9. Büscher, D., P. Dello Sbarba, R. A. Hipskind, U. R. Rapp, E. R. Stanley, and M. Baccarini. 1993. *v-raf* confers CSF-1 independent growth to a macrophage cell line and leads to immediate early gene expression without MAP-kinase activation. *Oncogene* **8**:3323-3332.
10. Büscher, D., R. A. Hipskind, S. Krautwald, T. Reimann, and M. Baccarini. 1995. Ras-dependent and -independent pathways target the mitogen-activated protein kinase network in macrophages. *Mol. Cell. Biol.* **15**:466-475.
11. Cannio, R., P. S. Rennie, and F. Blasi. 1991. A cell-type specific and enhancer-dependent silencer in the regulation of the expression of the human urokinase plasminogen activator gene. *Nucleic Acids Res.* **19**:2303-2308.
12. Cassady, A. I., K. J. Stacey, K. A. Nimmo, K. M. Murphy, D. von der Ahe, D. Pearson, F. Botteri, Y. Nagamine, and D. A. Hume. 1991. Constitutive expression of the urokinase plasminogen activator gene in murine RAW264 macrophages involves distal and 5' non-coding sequences that are conserved between mouse and pig. *Nucleic Acids Res.* **19**:6839-6847.
13. Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**:156-159.
14. Collart, M. A., D. Belin, J.-D. Vassalli, S. de Kossodo, and P. Vassalli. 1986. γ interferon enhances macrophage transcription of the tumour necrosis factor/cachectin, interleukin 1, and urokinase genes, which are controlled by short lived repressors. *J. Exp. Med.* **164**:2113-2118.
15. Collart, M. A., D. Belin, J.-D. Vassalli, and P. Vassalli. 1987. Modulations of functional activity in differentiated macrophages are accompanied by early and transient increase or decrease in *c-fos* gene transcription. *J. Immunol.* **139**:949-955.
16. Colman, M. S., and M. C. Ostrowski. Unpublished data.
17. Courtneidge, S. A., R. Dhand, D. Pilat, G. M. Twamley, M. D. Waterfield, and M. F. Roussel. 1993. Activation of Src family kinases by colony stimulating factor-1, and their association with its receptor. *EMBO J.* **12**:943-950.
18. Danø, K., P. A. Andreasen, J. Grøndahl-Hansen, P. Kristensen, and L. S. Nielsen. 1985. Plasminogen activators, tissue degradation and cancer. *Adv. Cancer Res.* **44**:139-266.
19. De Cesare, D., D. Vallone, G. Cirillo, and P. Verde. 1993. A cell specific c-jun/ATF2 heterodimer binds to the essential AP-1 site of the human urokinase enhancer, p. 24. *In* F. Blasi, K. Danø, J. Quigley, D. Rifkin, and S. Strickland (ed.), *Molecular and cellular biology of plasminogen activation*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
20. Degen, S. J. F., J. L. Heckel, E. Reich, and J. L. Degen. 1987. The murine urokinase-type plasminogen activator gene. *Biochemistry* **26**:8270-8279.
21. Downing, J. R., B. L. Margolis, A. Zilberstein, R. A. Ashmun, A. Ullrich, C. J. Sherr, and J. Schlessinger. 1989. Phospholipase C- γ , a substrate for PDGF receptor kinase, is not phosphorylated on tyrosine during the mitogenic response to CSF-1. *EMBO J.* **8**:3345-3350.
22. Downing, J. R., C. W. Rettenmier, and C. J. Sherr. 1988. Ligand-induced tyrosine kinase activity of the colony-stimulating factor 1 receptor in a murine macrophage cell line. *Mol. Cell. Biol.* **8**:1795-1799.
23. Ellis, L., E. Clauser, D. O. Morgan, M. Edery, R. A. Roth, and W. J. Rutter. 1986. Replacement of insulin receptor tyrosine residues 1162 and 1163 compromises insulin-stimulated kinase activity and uptake of 2-deoxyglucose. *Cell* **45**:721-732.
24. Falcone, D. J., T. A. McCaffrey, A. Haimovitz-Friedman, and M. Garcia. 1993. Transforming growth factor- β 1 stimulates macrophage urokinase expression and release of matrix-bound basic fibroblast growth factor. *J. Cell. Physiol.* **155**:595-605.
25. Fowles, L. F., and K. J. Stacey. Unpublished data.
26. Gibbs, J. B., M. S. Marshall, E. M. Scolnick, R. A. F. Dixon, and U. S. Vogel. 1990. Modulation of guanine nucleotides bound to ras in NIH3T3 cells by oncogenes, growth factors, and the GTPase activating protein (GAP). *J. Biol. Chem.* **265**:20437-20442.
27. Gottschalk, L. R., D. M. Giannola, and S. G. Emerson. 1993. Molecular regulation of the human IL-3 gene: inducible and T cell-restricted expression requires intact AP-1 and Elf-1 nuclear protein binding sites. *J. Exp. Med.* **178**:1681-1692.
28. Gutman, A., and B. Wasylyk. 1990. The collagenase gene promoter contains a TPA and oncogene-responsive unit encompassing the PEA3 and AP-1 binding sites. *EMBO J.* **9**:2241-2246.
29. Hai, T., and T. Curran. 1991. Cross-family dimerization of transcription factors Fos/Jun and ATF/CREB alters DNA binding specificity. *Proc. Natl. Acad. Sci. USA* **88**:3720-3724.
30. Hamilton, J. A., G. Vairo, K. R. Knight, and B. G. Cocks. 1991. Activation and proliferation signals in murine macrophages. *Biochemical signals con-*

- trolling the regulation of macrophage urokinase-type plasminogen activator activity by colony-stimulating factors and other agents. *Blood* **77**:616-627.
31. **Heidaran, M. A., C. J. Molloy, M. Pangelinan, G. G. Choudhury, L.-M. Wang, T. P. Fleming, A. Y. Sakaguchi, and J. H. Pierce.** 1992. Activation of the colony-stimulating factor 1 receptor leads to the rapid tyrosine phosphorylation of GTPase-activating protein and activation of cellular p21ras. *Oncogene* **7**:147-152.
 32. **Hilfinger, J. M., N. Clark, M. Smith, K. Robinson, and D. M. Markovitz.** 1993. Differential regulation of the human immunodeficiency virus type 2 enhancer in monocytes at various stages of differentiation. *J. Virol.* **67**:4448-4453.
 33. **Hipskind, R. A., D. Büscher, A. Nordheim, and M. Baccharini.** 1994. Ras/ MAP kinase-dependent and -independent signaling pathways target distinct ternary complex factors. *Genes Dev.* **8**:1803-1816.
 34. **Imamura, K., A. Dianoux, T. Nakamura, and D. Kufe.** 1990. Colony-stimulating factor 1 activates protein kinase C in human monocytes. *EMBO J.* **9**:2423-2429.
 35. **Jaworowski, A., S. Argyriou, P. Yusoff, and J. A. Hamilton.** 1994. Phospholipase D is activated by phorbol ester but not CSF-1 in murine bone marrow-derived macrophages. *Biochem. Biophys. Res. Commun.* **201**:733-739.
 36. **Klemsz, M. J., R. A. Maki, T. Papayannopoulou, J. Moore, and R. Hromas.** 1993. Characterization of the *ets* oncogene family member, *fli-1*. *J. Biol. Chem.* **268**:5769-5773.
 37. **Kola, I., S. Brookes, A. R. Green, R. Garber, M. Tymms, T. Papas, and A. Seth.** 1993. The *Ets1* transcription factor is widely expressed during murine embryo development and is associated with mesodermal cells involved in morphogenetic processes such as organ formation. *Proc. Natl. Acad. Sci. USA* **90**:7588-7592.
 38. **Kolch, W., G. Heldecker, G. Kochs, R. Hummel, H. Vahidl, H. Mischak, G. Finkenzeller, D. Marmé, and U. Rapp.** 1993. Protein kinase C α activates RAF-1 by direct phosphorylation. *Nature (London)* **364**:249-252.
 39. **Langer, S. J., D. M. Bortner, M. F. Roussel, C. J. Sherr, and M. C. Ostrowski.** 1992. Mitogenic signaling by colony-stimulating factor 1 and *ras* is suppressed by the *ets-2* DNA-binding domain and restored by *myc* overexpression. *Mol. Cell. Biol.* **12**:5355-5362.
 40. **Lee, J. S., B. Favre, B. A. Hemmings, B. Kiefer, and Y. Nagamine.** 1994. Okadaic acid-dependent induction of the urokinase-type plasminogen activator gene associated with stabilization and autoregulation of c-Jun. *J. Biol. Chem.* **269**:2887-2894.
 41. **Lee, J. S., D. von der Ahe, B. Kiefer, and Y. Nagamine.** 1993. Cytoskeletal reorganization and TPA differently modify AP-1 to induce the urokinase-type plasminogen activator gene in LLC-PK1 cells. *Nucleic Acids Res.* **21**:3365-3372.
 42. **Lin, H.-S., and S. Gordon.** 1979. Secretion of plasminogen activator by bone marrow-derived mononuclear phagocytes and its enhancement by colony-stimulating factor. *J. Exp. Med.* **150**:231-245.
 43. **Lioubin, M. N., G. M. Myles, K. Carlberg, D. Bowtell, and L. R. Rohrschneider.** 1994. *Sch, Grb2, Sos1*, and a 150-kilodalton tyrosine-phosphorylated protein form complexes with Fms in hematopoietic cells. *Mol. Cell. Biol.* **14**:5682-5691.
 44. **Maki, R. (La Jolla Cancer Research Foundation).** 1994. Personal communication.
 45. **Mavrothalassitis, G. J., and T. S. Papas.** 1991. Positive and negative factors regulate the transcription of the *ETS2* gene via an oncogene-responsive-like unit within the *ETS2* promoter region. *Cell Growth Differ.* **2**:215-224.
 46. **Nerlov, C., P. Busk, and M. Johnsen.** 1993. Molecular characterization of the structure and function of the uPA enhancer, p. 23. *In* F. Blasi, K. Danø, J. Quigley, D. Rifkin, and S. Strickland (ed.), *Molecular and cellular biology of plasminogen activation*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 47. **Nerlov, C., D. De Cesare, F. Pergola, A. Caracciolo, F. Blasi, M. Johnsen, and P. Verde.** 1992. A regulatory element that mediates co-operation between a PEA3-AP-1 element and an AP-1 site is required for phorbol ester induction of urokinase enhancer activity in HepG2 hepatoma cells. *EMBO J.* **11**:4573-4582.
 48. **Osborn, L., S. Kunkel, and G. J. Nabel.** 1989. Tumor necrosis factor alpha and interleukin 1 stimulate the human immunodeficiency virus enhancer by activation of the nuclear factor κ B. *Proc. Natl. Acad. Sci. USA* **86**:2336-2340.
 49. **Owen, R. D., D. M. Bortner, and M. C. Ostrowski.** 1990. *ras* oncogene activation of a VL30 transcriptional element is linked to transformation. *Mol. Cell. Biol.* **10**:1-9.
 50. **Owen, R. D., and M. C. Ostrowski.** 1990. Transcriptional activation of a conserved sequence element by *ras* requires a nuclear factor distinct from c-fos or c-jun. *Proc. Natl. Acad. Sci. USA* **87**:3866-3870.
 51. **Pankov, R., A. Umezawa, R. Maki, C. J. Der, C. A. Hauser, and R. G. Oshima.** 1994. Oncogene activation of human keratin 18 transcription via the Ras signal transduction pathway. *Proc. Natl. Acad. Sci. USA* **91**:873-877.
 52. **Pöllänen, J., R. W. Stephens, and A. Vaheri.** 1991. Directed plasminogen activation at the surface of normal and malignant cells. *Adv. Cancer Res.* **57**:273-328.
 53. **Raschke, W. C., S. Baird, P. Ralph, and I. Nakoinz.** 1978. Functional macrophage cell lines transformed by Abelson leukemia virus. *Cell* **15**:261-267.
 54. **Reddy, M. A., S. J. Langer, M. S. Colman, and M. C. Ostrowski.** 1992. An enhancer element responsive to *ras* and *fms* signalling pathways is composed of two distinct nuclear factor binding sites. *Mol. Endocrinol.* **6**:1051-1060.
 55. **Reedijk, M., X. Liu, and T. Pawson.** 1990. Interactions of phosphatidylinositol kinase, GTPase-activating protein (GAP), and GAP-associated proteins with the colony-stimulating factor 1 receptor. *Mol. Cell. Biol.* **10**:5601-5608.
 56. **Rørth, P., C. Nerlov, F. Blasi, and M. Johnsen.** 1990. Transcription factor PEA3 participates in the induction of urokinase plasminogen activator transcription in murine keratinocytes stimulated with epidermal growth factor or phorbol-ester. *Nucleic Acids Res.* **18**:5009-5017.
 57. **Ross, I. L., T. L. Dunn, Y. Xie, S. Roy, C. J. K. Barnett, and D. A. Hume.** 1994. Comparison of the expression and function of the transcription factor PU.1 (SPI-1 proto-oncogene) between murine macrophages and B lymphocytes. *Oncogene* **9**:121-132.
 58. **Ross, I. L., and D. A. Hume.** Unpublished data.
 59. **Roth, P., and E. R. Stanley.** 1992. The biology of CSF-1 and its receptor. *Curr. Top. Microbiol. Immunol.* **181**:141-167.
 60. **Roussel, M. F., J. L. Cleveland, S. A. Shurtleff, and C. J. Sherr.** 1991. *Myc* rescue of a mutant CSF-1 receptor impaired in mitogenic signalling. *Nature (London)* **353**:361-363.
 61. **Roussel, M. F., J. N. Davis, J. L. Cleveland, J. Ghysdael, and S. W. Hiebert.** 1994. Dual control of *myc* expression through a single DNA binding site targeted by *ets* family proteins and E2F-1. *Oncogene* **9**:405-415.
 62. **Roussel, M. F., C. W. Rettenmier, A. T. Look, and C. J. Sherr.** 1984. Cell surface expression of v-*fms*-coded glycoproteins is required for transformation. *Mol. Cell. Biol.* **4**:1999-2009.
 63. **Roussel, M. F., and C. J. Sherr.** 1989. Mouse NIH3T3 cells expressing human colony-stimulating factor 1 (CSF-1) receptors overgrow in serum-free medium containing human CSF-1 as their only growth factor. *Proc. Natl. Acad. Sci. USA* **86**:7924-7927.
 64. **Roussel, M. F., S. A. Shurtleff, J. R. Downing, and C. J. Sherr.** 1990. A point mutation at tyrosine-809 in the human colony-stimulating factor 1 receptor impairs mitogenesis without abrogating tyrosine kinase activity, association with phosphatidylinositol 3-kinase, or induction of *c-fos* and *junB* genes. *Proc. Natl. Acad. Sci. USA* **87**:6738-6742.
 65. **Rybeck, R. P., and R. Bravo.** 1991. c-JUN, JUN B, and JUN D differ in their binding affinities to AP-1 and CRE consensus sequences: effect of FOS proteins. *Oncogene* **6**:533-542.
 66. **Sariban, E., K. Imamura, M. Sherman, V. Rothwell, P. Pantazis, and D. Kufe.** 1989. Downregulation of *c-fms* gene expression in human monocytes treated with phorbol esters and colony-stimulating factor 1. *Blood* **74**:123-129.
 67. **Schneikert, J., Y. Lutz, and B. Wasyluk.** 1992. Two independent activation domains in c-Ets-1 and c-Ets-2 located in non-conserved sequences of the *ets* gene family. *Oncogene* **7**:249-256.
 68. **Sherr, C. J.** 1990. Colony-stimulating factor-1 receptor. *Blood* **75**:1-12.
 69. **Sherr, C. J., C. W. Rettenmier, R. Sacca, M. F. Roussel, A. T. Look, and E. R. Stanley.** 1985. The *c-fms* proto-oncogene product is related to the receptor for the mononuclear phagocyte growth factor, CSF-1. *Cell* **41**:665-676.
 70. **Smith, M. R., S. J. DeGudicibus, and D. W. Stacey.** 1986. Requirement for *c-ras* proteins during viral oncogene transformation. *Nature (London)* **320**:540-543.
 71. **Stacey, K. J.** 1993. Ph.D. thesis. University of Queensland, Brisbane, Australia.
 72. **Stacey, K. J., A. I. Cassady, K. Nimmo, K. M. Murphy, D. von der Ahe, D. Pearson, F. Botteri, Y. Nagamine, and D. A. Hume.** 1992. The regulation of urokinase plasminogen activator gene expression in macrophages, p. 233-240. *In* R. van Furth (ed.), *Mononuclear phagocytes: biology of monocytes and macrophages*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
 73. **Stacey, K. J., I. L. Ross, and D. A. Hume.** 1993. Electroporation and DNA-dependent cell death in murine macrophages. *Immunol. Cell Biol.* **71**:75-85.
 74. **Thompson, C. B., C.-Y. Wang, I.-C. Ho, P. R. Bohjanen, B. Petryniak, C. H. June, S. Miesfeldt, L. Zhang, G. J. Nabel, B. Karpinski, and J. M. Leiden.** 1992. *cis-acting* sequences required for inducible interleukin-2 enhancer function bind a novel Ets-related protein, Elf-1. *Mol. Cell. Biol.* **12**:1043-1053.
 75. **Vairo, G., and J. A. Hamilton.** 1991. Signalling through CSF receptors. *Immunol. Today* **12**:362-369.
 76. **Weis, N., and J. A. Hamilton.** 1991. Colony stimulating factor-1 stimulates diacylglycerol generation in murine bone marrow-derived macrophages, but not in resident peritoneal macrophages. *J. Cell. Physiol.* **147**:298-305.
 77. **Verde, P., S. Boast, A. Franze, F. Robbiati, and F. Blasi.** 1988. An upstream enhancer and a negative element in the 5' flanking region of the human urokinase plasminogen activator gene. *Nucleic Acids Res.* **16**:10699-10716.
 78. **Wasyluk, B., S. L. Hahn, and A. Giovane.** 1993. The Ets family of transcription factors. *Eur. J. Biochem.* **211**:7-18.
 79. **Wasyluk, B., C. Wasyluk, P. Flores, A. Begue, D. Leprince, and D. Stehelin.** 1990. The *c-ets* proto-oncogenes encode transcription factors that cooperate with c-Fos and c-Jun for transcriptional activation. *Nature (London)* **346**:191-193.
 80. **Wasyluk, C., P. Flores, A. Gutman, and B. Wasyluk.** 1989. PEA3 is a nuclear

- target for transcription activation by non-nuclear oncogenes. *EMBO J.* **8**:3371-3378.
81. **Whetton, A. D., P. N. Monk, S. D. Consalvey, and C. P. Downes.** 1986. The haemopoietic growth factors interleukin-3 and colony stimulating factor-1 stimulate proliferation but do not induce inositol lipid breakdown in murine bone-marrow derived macrophages. *EMBO J.* **5**:3281-3286.
82. **Wu, H., K. Moulton, A. Horvai, S. Parik, and C. K. Glass.** 1994. Combinatorial interactions between AP-1 and ets domain proteins contribute to the developmental regulation of the macrophage scavenger receptor gene. *Mol. Cell. Biol.* **14**:2129-2139.
83. **Xu, X.-X., T. G. Tessner, C. O. Rock, and S. Jackowski.** 1993. Phosphatidylcholine hydrolysis and *c-myc* expression are in collaborating mitogenic pathways activated by colony-stimulating factor 1. *Mol. Cell. Biol.* **13**:1522-1533.
84. **Yeung, Y.-G., K. Berg, F. J. Pixley, R. H. Angeletti, and E. R. Stanley.** 1992. Protein tyrosine phosphatase-1C is rapidly phosphorylated in tyrosine in macrophages in response to colony stimulating factor-1. *J. Biol. Chem.* **267**:23447-23450.
85. **Yue, X., P. Favot, T. L. Dunn, A. I. Cassady, and D. A. Hume.** 1993. Expression of mRNA encoding the macrophage colony-stimulating factor receptor (*c-fms*) is controlled by a constitutive promoter and tissue-specific transcription elongation. *Mol. Cell. Biol.* **13**:3191-3201.