

# Identification and Characterization of Ral-Binding Protein 1, a Potential Downstream Target of Ral GTPases

SHARON B. CANTOR, TAKESHI URANO, AND LARRY A. FEIG\*

Department of Biochemistry, Tufts University School of Medicine, Boston, Massachusetts 02111

Received 19 April 1995/Returned for modification 23 May 1995/Accepted 26 May 1995

**Ral proteins constitute a distinct family of Ras-related GTPases. Although similar to Ras in amino acid sequence, Ral proteins are activated by a unique nucleotide exchange factor and inactivated by a distinct GTPase-activating protein. Unlike Ras, they fail to promote transformed foci when activated versions are expressed in cells. To identify downstream targets that might mediate a Ral-specific function, we used a *Saccharomyces cerevisiae*-based interaction assay to clone a novel cDNA that encodes a Ral-binding protein (RalBP1). RalBP1 binds specifically to the active GTP-bound form of RalA and not to a mutant Ral with a point mutation in its putative effector domain. In addition to a Ral-binding domain, RalBP1 also contains a Rho-GTPase-activating protein domain that interacts preferentially with Rho family member CDC42. Since CDC42 has been implicated in bud site selection in *S. cerevisiae* and filopodium formation in mammalian cells, Ral may function to modulate the actin cytoskeleton through its interactions with RalBP1.**

Two closely related Ral proteins (RalA and RalB; 85% identical) constitute a distinct family within the Ras superfamily of GTPases (5). They are among the closest relatives of Ras proteins, sharing 58% sequence identity, and have similar overall structural features (15). For example, their affinities for guanine nucleotides and intrinsic GTPase activities are comparable. Moreover, mutations in Ral that are comparable to oncogenic mutations in Ras also suppress the intrinsic GTPase activity in Ral and make the protein resistant to its GTPase-activating protein (GAP), Ral-GAP (10). Finally, both Ras and Ral are prenylated at the carboxy-terminal CAAX box that targets them to the membrane fraction of cells (21).

Ras and Ral are distinguishable in many respects, however. Unlike Ras, Ral does not produce transformed foci of cells when it is locked in its active GTP state (13). This difference is likely due to the fact that the effector domain of Ral is distinct from that of Ras. This 10- to 15-amino-acid (aa) (aa 26 to 45 in Ras) region present in Ras family members plays an important role in interactions with downstream targets and GAPs. Ras and Ral are also regulated by distinct GAPs (10) and guanine nucleotide exchange factors that promote the replacement of bound GDP with GTP on these GTPases (12). Interestingly, the Ral exchange factor, Ral-GDS, has been shown to bind preferentially to the active forms of Ras, Rap1A, and R-Ras in the *Saccharomyces cerevisiae* two-hybrid system and in vitro (17, 20, 34). This raises the possibility that Ral activation constitutes a distinct downstream pathway from Ras.

The localization of Ral in cells is also different from that of Ras. Whereas Ras is found almost exclusively on the internal face of the plasma membrane, Ral has a more diverse distribution. Ral can be found in plasma membrane fractions, but it is present primarily in cytoplasmic vesicles, including clathrin-coated vesicles and secretory vesicles (2, 9, 36).

To understand the role of Ral function in cells, we used the yeast two-hybrid system to screen for proteins that can bind preferentially to the active GTP-bound form of RalA. Here we describe some properties of one such protein, termed Ral-binding protein 1 (RalBP1).

## MATERIALS AND METHODS

**Cell culture and transient expression.** COS-7 cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% iron-enriched calf serum (Hyclone) at 37°C in 5% CO<sub>2</sub>. Cells were plated at a density of ~5 × 10<sup>5</sup>/60-mm-diameter plate 1 day before transfection. Adenovirus major late promoter-based expression vector pMT3 (35) was used for transient expression of the cDNAs in COS-7 cells by the DEAE-dextran method.

**Interaction cloning with the yeast two-hybrid system.** For two-hybrid screening, rala (72L, 203S) was subcloned into pAS-CYH2 (Trp marker) (8). This vector directs the expression of a fusion protein between the DNA-binding domain of GAL4 and the entire RalA protein. A rat brain cDNA library, cloned into pACT (4) (kindly provided by I. Macara, University of Vermont), contained a total of 5 × 10<sup>6</sup> primary recombinants with an average insert size of 1.5 kb. *S. cerevisiae* Y190 (*MATa gal4 gal80 his3 trp1-901 ade2-101 ura3-52 leu2-3,112 URA3::GAL-lacZ LYS2::GAL-HIS3 cyh'*), expressing the GAL4 DNA-binding domain-Ral fusion protein, was transformed with the pACT library (Leu marker) with salmon sperm DNA as the carrier, and 2.3 × 10<sup>6</sup> primary transformants were selected for growth on medium lacking leucine, tryptophan, and histidine and containing 30 mM 3-aminotriazole. The plates were incubated at 30°C for 3 days. Surviving yeast colonies were transferred to nitrocellulose and laid onto minimal medium plates containing 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) and screened for expression of β-galactosidase by incubation at 30°C for 1 to 2 days. Blue coloration of a colony was indicative of a positive interaction. Of the 200 His<sup>+</sup> colonies, 15 were also LacZ<sup>+</sup>. The 15 His<sup>+</sup> LacZ<sup>+</sup> colonies were rescued from the plate and grown in selective medium. Plasmid DNA was recovered and introduced by electroporation into leucine-deficient *Escherichia coli* KC8. Transformants were plated on minimal medium lacking leucine so that only transformants carrying the library plasmid grew.

Positives were tested for target specificity by retransformation into reporter strain Y190 alone or in conjunction with the Ral-Gal4 DNA-binding domain fusion or with different Gal4 DNA-binding domain fusions. Only library plasmids that did not activate marker expression in the presence of SNF1, Ras, or Ral49N were analyzed further.

**Plasmid construction.** pASCYH-Ral72L,203S was constructed as follows. The 771-nucleotide *EcoRI-HindIII* fragment was excised from plasmid ptaCRal72L (5). *BamHI* linkers were added, and the resulting fragment was subcloned into the *BamHI* site of pGEX3X. The Cys at position 203 was changed to a Ser (C203S), and a 3' *SalI* site was introduced by PCR. A *BamHI-SalI* RalA fragment was then cloned into vector pASCYH. An effector domain mutation was created by using overlap PCR (19) to change the Asp at position 49 to an Asn (D49N). pASCYH-Ras61L186C was constructed as follows. pXCR61L (14) was used as the template for site-directed PCR mutagenesis, which changed Cys-186 to a Ser (C186S) and generated both 5' and 3' *BamHI* sites. The resulting product was excised by *BamHI* digestion and subcloned into vector pASCYH.

**Isolation of the complete RalBP1 cDNA and sequencing.** Additional rat RalBP1 cDNAs were isolated from an oligo(dT)-primed and randomly primed brain library (Clontech) by using the 1.6-kb RalBP1 clone isolated from the two-hybrid system as a probe. The RalBP1 cDNA sequence was determined on both strands by the dideoxy-chain termination technique with multiple cDNA clones and subclones.

**Northern (RNA) blot analysis.** A multiple rat tissue poly(A) RNA blot (Clon-

\* Corresponding author. Phone: (617) 636-6956. Fax: (617) 636-6409.




FIG. 1. Specificity of the interaction between RalBP1 and Ral in *S. cerevisiae*. The 1.6-kb cDNA clone (RalBP1) isolated from the two-hybrid screen as a fusion with the Gal4 activation domain was used to cotransform reporter strain *S. cerevisiae* Y190 along with various cDNAs fused to the Gal4 DNA-binding domain. Segments: A, RalA72L; B, SNF1; D, Ras61L; E, Ral49N effector mutant. Cotransformation of the DNA-binding domain of Gal4 fused to SNF1 and the activation domain of Gal4 fused to SNF4 served as a positive control (segment C). To assay for protein interaction-induced growth, cells were streaked on permissive medium (+ His) or on selective medium (– His) containing 3-aminotriazole. The plates were incubated for 2 days and photographed.

tech) was probed for expression of the RalBP1 gene by using randomly primed,  $^{32}\text{P}$ -labeled fragments from the 1.6-kb RalBP1 cDNA originally isolated from the expression library.

**Expression of Ral and RalBP1.** Recombinant Ral and Ral49N were subcloned into pGEX2T and transformed into *E. coli* BL21. The bacteria were induced to express protein with 0.1 mM isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG) for 4 h. The fusion proteins were isolated by affinity chromatography on glutathione-Sepharose, eluted with elution buffer (50 mM Tris [pH 8.0], 120 mM NaCl, 20 mM glutathione), and dialyzed into phosphate-buffered saline. Full-length RalBP1 was inserted as a *Bam*HI-*Eco*RI restriction fragment into an altered version of mammalian expression vector pMT3 (35) that contained a modified Glu epitope (RMEFMPME) 5' to the cloning site. RalBP1 (positions 86 to 626) was excised from pACT by *Xho*I digestion, and 5' *Bam*HI and 3' *Eco*RI cleavage sites were added by PCR. The resulting fragment was subcloned into the same modified pMT3 vector. Glu-tagged RalBP1 was affinity purified from transiently transfected COS-7 cells by using Sepharose A beads coated with anti-Glu immunoglobulin G (kindly provided by S. Powers, Onyx Pharm). The RalBP1 cDNA sequence encoding codons 375 to 626 was amplified by PCR with primers that incorporated *Bam*HI and *Eco*RI restriction sites at the 5' and 3' ends, respectively. The amplified products were subcloned into pGEX2T and transformed into *E. coli* BL21. RalBP1 (positions 86 to 415), a subfragment of RalBP1 (positions 86 to 626), was isolated by digestion with *Bam*HI and *Bgl*II. It was subcloned into pGEX-2T and transformed into *E. coli* BL21. Recombinant RalA and RasH proteins were expressed in and purified from *E. coli* as described previously (11).

**Binding of Ral to RalBP1 in vitro.** In vitro binding assays were performed with GST-RalA or recombinant Ral and either Glu epitope tagged-RalBP1 or glutathione-S-transferase (GST)-RalBP1. Glu-tagged RalBP1 (~0.2  $\mu\text{g}$ ) bound to Sepharose A beads or RalBP1 fragments (~1.0  $\mu\text{g}$ ) bound to glutathione-Sepharose were incubated at 4°C for 3 h with either 0.5  $\mu\text{g}$  of the nucleotide-bound GST-RalA fusion protein or purified RalA, respectively, in 20 mM Tris-HCl (pH 7.5)–10 mM  $\text{MgCl}_2$ –150 mM NaCl–0.5% Nonidet P-40. Ral proteins were pre-loaded with guanine nucleotides by incubation with 1 mM GDP $\beta$ S or GTP $\gamma$ S for 20 min at 37°C in 20 mM Tris-HCl (pH 7.5)–1 mM dithiothreitol (DTT)–2 mM EDTA–25 mM NaCl–40  $\mu\text{g}$  of bovine serum albumin per ml. The reactions were stopped by addition of  $\text{MgCl}_2$  to a final concentration of 10 mM. The beads were washed six times with 1 ml of 20 mM Tris-HCl (pH 7.5)–10 mM  $\text{MgCl}_2$ –20 mM NaCl–0.5% Nonidet P-40 and denatured in Laemmli sample buffer. After separation in a sodium dodecyl sulfate (SDS)–10% polyacrylamide gel and transfer to nitrocellulose, the Ral protein retained on the beads was detected by immunoblotting with anti-Ral antibody. The anti-RalA antibody was prepared by injection of rabbits with the GST-RalA fusion protein, followed by affinity purification with RalA. Detection was done by chemiluminescence (Amersham) with horseradish peroxidase-conjugated anti-rabbit immunoglobulin. As a control, equivalent amounts of recombinant Ras, bound to either GDP $\beta$ S or GTP $\gamma$ S, were incubated with GST-RalBP1 (positions 375 to 627) as described above for Ral proteins. Additionally, nucleotide-free Ras was incubated with the catalytic domain of Ras-guanine nucleotide-releasing factor (33) bound to glutathione-Sepharose. Ras proteins retained on the beads were visualized by immunoblotting with anti-Ras antibody (Upstate Biotechnology, Inc.).

**Immunoblot analysis of RalBP1.** GST-RalBP1 (positions 86 to 415), described above, was purified with glutathione-Sepharose and eluted with 50 mM Tris (pH 8.0)–120 mM NaCl–20 mM glutathione–1 mM DTT and used to immunize

rabbits. The resulting antiserum and preimmune serum control were used for immunoblotting at a dilution of 1:100. Rat brains were fractionated into soluble or particulate fractions as follows. The tissue was homogenized in lysis buffer (25 mM Tris-HCl (pH 7.5), 1 mM  $\text{MgCl}_2$ , 1 mM ethylene glycol-bis( $\beta$ -aminoethyl ether)-*N,N,N',N'*-tetraacetic acid [EGTA], 1 mM DTT, 1% aprotinin) with 10 strokes with a Dounce homogenizer. Unbroken cells were removed by three successive centrifugations at 10,000  $\times g$  for 15 min each time at 4°C. The sample was then separated into soluble and particulate fractions by centrifugation at 100,000  $\times g$  and 4°C for 60 min. The protein concentration of each fraction was determined by Lowry assay (25). A 75- $\mu\text{g}$  sample of protein from each fraction was denatured in Laemmli sample buffer, separated on an SDS-polyacrylamide gel, and transferred to nitrocellulose. RalBP1 was detected by immunoblotting with an anti-RalBP1 serum described above.

**Measurement of GTP hydrolysis rate of p21s.** The GTPase activities of RalA, RhoA, Rac1, and CDC42Hs were measured by a nitrocellulose filtration assay as follows. A 2- $\mu\text{g}$  sample of each p21 was incubated with either  $10^{-7}$  M [ $\gamma$ - $^{32}\text{P}$ ]GTP or  $10^{-7}$  M [ $\alpha$ - $^{32}\text{P}$ ]GTP (650 Ci/mmol; ICN) in the presence of 20 mM Tris-HCl (pH 7.5)–1 mM DTT–5 mM EDTA–40  $\mu\text{g}$  of bovine serum albumin per ml for 10 min at 30°C in a total volume of 50  $\mu\text{l}$ . The exchange reaction was stopped by addition of  $\text{MgCl}_2$  (10 mM final concentration). An 8- $\mu\text{l}$  sample of the p21-[ $\gamma$ - $^{32}\text{P}$ ]GTP complex was added to 12  $\mu\text{l}$  of exchange buffer (1 mM GTP, 40  $\mu\text{g}$  of bovine serum albumin per ml, 20 mM Tris-HCl [pH 7.5], 1 mM DTT, 25 mM NaCl, 10 mM  $\text{MgCl}_2$ ) in the absence or presence of the RalBP1 GAP domain (positions 86 to 415). The assay was incubated at 30°C for RhoA, RalA, and CDC42Hs and at 15°C for Rac1. Samples of 15  $\mu\text{l}$  were removed from each incubation over a 10-min time course and filtered through nitrocellulose. The filters were washed twice with ice-cold wash buffer (20 mM Tris-HCl [pH 7.5], 150 mM NaCl, 10 mM  $\text{MgCl}_2$ ), and the amount of [ $\gamma$ - $^{32}\text{P}$ ]GTP still bound to the filters was measured by scintillation counting. RhoA, Rac, and CDC42Hs were expressed as GST fusion proteins and obtained in purified form after thrombin cleavage.

**Nucleotide sequence accession number.** The nucleotide sequence reported here has been assigned GenBank accession no. U28830.

## RESULTS

**Identification of a novel Ral-binding protein.** A yeast two-hybrid system was used to identify potential downstream targets of the RalA GTPase (8). A Ral-GAL4 DNA-binding domain fusion was constructed in vector pAS-CYH2. A mutant Ral protein with a Gln-to-Leu substitution at position 72 was used. This protein should accumulate in the active GTP-bound state in *S. cerevisiae* because it has decreased intrinsic GTPase activity (15) and fails to respond to Ral-GAP (10). Cys-203 in the CAAX box of RalA was also mutated to Ser to prevent C-terminal geranyl-geranylation. Such a modification is known to target Ral to membrane fractions and may thus prevent nuclear translocation of the fusion protein required in this assay system. Expression of the fusion protein was confirmed

1 gaa ttc cgg ggc taa gtg cga gga ggc ggc tgc agc gga gag gac ggt cgc tgg cgc  
 61 gca ggc act ggt gta atg gat agg taa cag aag acc act tcc ctt cct cct cag ggc ttc  
 121 agc atg act gag tgc ttc ctg ccc ccc acc agc agc cct agt gaa cac cgc agg gcc gag  
 1 M T E C F L P P T S S P S E H R R A E  
 181 cac ggc agc ggg ctg acc cgg acc ccc agc tcc gag gag atc agc cct aca aaa ttt cct  
 20 H G S G L T R T P S S E E I S P T K F P  
 241 gaa ttg tac cgg acg ggt gag cct tca cct ccc cat gac atc ctc cac gag cct cct gat  
 40 G L Y R T G E P S P H D I L H E P P D  
 301 atc gtg tca gat gat gag aaa gac cat gga aag aaa aaa gga aaa ttt aag aea aag gag  
 60 I V S D D E K D H G K K K G K P K K E  
 361 aaa agc agt gaa ggc tac gtc gcc ttc cag gaa gac agc tct gga gat gag cct gaa agt  
 80 K R T E G Y A A F Q E D S S G D E A E S  
 421 cct tcc aag atg aag agt tcc aag ggc atc atg ttc aaa aag ccc agc ttc tct aaa  
 100 P S K M K R S K G I H V F K K P S F S K  
 481 aag aag gaa aag gat ttt aaa atc aaa gaa aag ccc aaa gaa aag ccc aca aaa gaa gaa  
 120 K K E K D F K I K E K P K E E K H K E E  
 541 aag cat aaa gag gaa aag cat aaa gaa aag aaa tgt aaa gac ttc acg gcc gcc gat gtt  
 140 K H K E E K H K E K K C K D L T A A D V  
 601 gtt aaa cag tgg aag gaa aag aaa aag aag aag cca act cag gag cca gag gtg cct  
 160 V K Q W K E K K K K P T Q E P E V P  
 661 cag acg gac gcc ccc agt ctc aga ccc att ttc ggc gct ccc ttc gct gaa gca gta gag  
 180 Q T D A P S L R P I P G A P L A D A V E  
 721 agc acc atg atg tac gac gcc atc cgg ctg coa gct gtc ttc cgg gag tgc gtg gac tac  
 200 R T M M Y D G I R L P A V F R E C V D Y  
 781 atg gag aag cac ggc atg aaa tgt gaa ggc atc tac cga gtg tca gga atc aaa tca aag  
 220 M E K H G M K C E G T Y R V S G I K S K  
 841 gta gat gag ctc aaa gca gcc tat gac cga gag gag tct cca aac ttg gaa gaa tat gaa  
 240 V D E L K A A Y D R E R S P N L E R Y E  
 901 cct aac act gta gcc agc tgg ctg aag cag tat ctg cga gac ctt ccc gag aat ttg ctt  
 260 P N T V A S L L K Q Y L R D L P E N L L  
 961 acc aaa gag ctt atg ccc cga ttt gaa gag gct tgt ggg agg acc aca gag gtg gag aaa  
 280 T K E L L M P R F E E A C G R T T E V E K  
 1021 gtg cag gag ttc cag cgc ttg ctc agg gaa ctg ccg gag tac aac cat ctt ctg ctc tcc  
 300 V Q E F Q R L L R F L P E V N H L L L S  
 1081 tgg ctc att gtc cag atg gac cac gtc atc ggc aag gag ctg gaa acc aag atg aac atc  
 320 M T L H M H H K A N T L A K T L A  
 1141 cag aac act tcc ata tgc ctg agc cca acc gtc cag atc agc aac cgg ttc ctg tat gtc  
 340 Q N I S T V L S P P T V Q I S N P V L Y V  
 1201 ctt ttc aca cac gtg caa gag ctc ttt ggc acc gtc ctc ctg aag caa gta acg aga cct  
 360 L P T H V Q E L F G T V V L K Q V T R P  
 1261 ctg cgt tgg tcc aac atg gcc aca atg cct aca cta gac acc caa gaa gac atc aag  
 380 L R W S N M A T M P T L P E T Q A G I K  
 1321 gaa gaa atc agc aga cag gac ttc ctt ttg aac tgt tta cat cga gat ctg cag ggc ggg  
 400 E E I R R Q E F L L N C L H R D L Q G G  
 1381 ata aag gac ttc tot aaa gaa gaa ata tgg gaa gta cag agt atc ctg act gcc ctc  
 420 I K D L S K E E R L W E V Q R I L T A C  
 1441 aag aga aag ctg agg gaa gct aaa aga caa gag tgt gag acc aag att gca cag gat ata  
 440 K R K L R E A K R Q A E C E T K I A Q E I  
 1501 gcc agc ctt tcy aag gag gat gtt tcc aaa gaa gaa acy aac gaa aac gag gag gtc ata  
 460 A S L S K E D S K E E T M E N E E V I  
 1561 aat atc ctc ctt gcc cag gag aat gag atc ctg act gag cag gag ctc ctg gcc atg  
 480 N I L L A Q E N E T L T E Q E L L A M  
 1621 gag cag ttt ctg cgt cgt cag atc gcc tcy gag aaa gaa gag att gac cgc ctc cga gct  
 500 E Q F L R R Q I A S E K E E I D R L R A K  
 1681 gag att gcg gag atc cag agt gcc cag cat ggc agc gag aca cad gag tac tcc tct  
 520 E I A E I Q S R Q H G R S E T E E Y S S  
 1741 gac agc gag agc agc gag gat gag gag gat gag cag atc att ctg gaa gac tta cag  
 540 D S E S E S E D E E E L Q I I L E D L Q  
 1801 agc cag aat gaa gaa tta gaa ata aag aac aat cac ttg aac cag gca gtc cat gaa gag  
 560 R Q N E E L E I K N N H L N Q A V H E E  
 1861 cgg gag gcc atc gtt gag ctc cga gtc cag ctc cgg ctg ctc cag atg ctg gac acc aag  
 580 R E A I V E L R V Q L R L L Q M L R A K  
 1921 tct gag cag cag ctc cag gag gag gag ccc gag agy cga ggg ggc aca ggc cca atc  
 600 S E Q Q L Q E E E F E R R G G T G P L  
 1981 ccc tgt gaa ggt gtc ctc gag gtc gaa gcc aag gag cag gag aag ccc tgg ccc agc  
 620 P C E G V L E V R A A K E Q A F P S P S  
 2041 aaa gac cgg aag gag acg ccc atc TGA act gcc ccg gcc ctg tgc atc gta ctg taa gcg  
 640 K D R K E T P I \*

FIG. 2. Nucleotide and deduced amino acid sequences of the open reading frame encoding RalBP1. The start and stop codons bordering the open reading frame are in capitals, and stop codons preceding the first methionine are underlined. Nucleotide and amino acid sequence numbers are at the left. The sequence homologous to Rho-GAP family members (aa 210 to 352) is underlined, as is a highly basic region predicted to be  $\alpha$ -helical.

by immunoblot analysis of yeast extracts with anti-Ral serum (data not shown). To identify proteins that interact with Ral, the yeast reporter strain, containing pAS-RAL72L,203S, was transformed along with a rat brain cDNA library expressed as fusions to the GAL4 activation domain in pACT (4). From  $2.5 \times 10^6$  primary transformants, approximately 200 survived the initial histidine selection. Fifteen colonies were also positive for the secondary screen, expression of  $\beta$ -galactosidase. For seven of these clones, histidine prototrophy and  $\beta$ -galactosidase activity depended on the presence of both plasmids. The cDNA library plasmids were transferred into *E. coli* and subjected to DNA sequencing, which revealed that six clones (1.6 kb) were identical and one contained an independent clone of the same gene.

To determine the specificity of this protein-protein interaction, yeast cells were cotransformed with the 1.6-kb partial cDNA for this Ral-binding protein, RalBP1, along with either Ral or one of two control fusion proteins, the SNF1 kinase and constitutively active Ras. As a positive control, we cotrans-




FIG. 3. Tissue distribution of RalBP1 expression. Northern blot analysis of poly(A) mRNAs isolated from a variety of tissues probed with RalBP1 cDNA encoding aa 86 to 626. Lanes: a, heart; b, brain; c, spleen; d, lung; e, liver; f, skeletal muscle; g, kidney; h, testis. The numbers at the left are molecular sizes in kilobases.

formed SNF1 and SNF4, which are known to interact in the two-hybrid system (Fig. 1). All cotransformants grew well under nonselective conditions in medium containing histidine, but only Ral and RalBP1 and SNF1 and SNF4 grew well under selective conditions in medium lacking histidine. Next, the role of the putative effector domain of Ral (aa 37 to 56) in this binding reaction was evaluated. The analogous region in the mammalian Ras protein (residues 26 to 45) interacts with downstream targets such as Raf, Ras-GAP, and RalGDS (6, 17, 20, 30, 34). Substitution of Asn for Asp at position 38 of Ras blocks its interaction with many of these proteins. Thus, we introduced an equivalent substitution into Ral (49N) and observed that it abolished the protein's interaction with RalBP1 in *S. cerevisiae* (Fig. 1). These results show that RalBP1 interacts specifically with Ral and that this interaction requires a functional effector domain. Thus, RalBP1 may function as a downstream target of Ral.

The 1.6-kb partial cDNA isolated by the two-hybrid system was used as a probe to isolate the entire coding sequence of the gene. A single open reading frame was found that coded for a 647-aa protein with an estimated molecular mass of ~75 kDa (Fig. 2). The cDNA clone was also used to probe Northern blots containing RNAs from a variety of tissues. RalBP1 was found to be expressed in all of the tissues examined as a major transcript of ~4.6 kb (Fig. 3).

Antiserum to RalBP1 was generated by injecting rabbits with a portion of RalBP1 (aa 86 to 415) purified from bacteria as a fusion protein with GST. When the antiserum was used to immunoblot total-cell lysates from rat brain tissue, a predominant band of ~95 kDa was detected. This band likely represented RalBP1 because it was absent when preimmune serum was used (Fig. 4, compare lanes a and b). When brain extracts were fractionated into cytoplasmic and particulate fractions, a similar band was present in both (lanes c and d). The size of this 95-kDa band was significantly greater (~20 kDa) than that predicted from the cDNA. When the full-length cDNA for RalBP1 was expressed transiently in COS-7 cells, it also migrated as a protein of ~95 kDa (Fig. 4, lane e). Furthermore, when the carboxyl-terminal region of RalBP1 (aa 375 to 626; see Fig. 6) was expressed as a fusion protein with GST in *E. coli*, it also migrated to the position of a protein ~20 kDa larger than that predicted (Fig. 4, lane f). Endogenous RalBP1 in COS-7 cells was below the level of detection in this gel (lane g). The discrepancy in apparent size may be due to the abnormal mobility of RalBP1 in SDS-polyacrylamide gels.

Examination of the amino acid sequence of RalBP1 revealed that it contained a domain between residues 210 and 352 that




FIG. 4. Immunoblot detection of RalBP1 in cell extracts. Whole-cell lysates (lanes a and b) and cytosolic (lane c) and particulate (lane d) fractions of rat brain tissue were run on SDS-polyacrylamide gels and immunoblotted with antiserum against RalBP1 (positions 86 to 415) or preimmune serum (lane a). Alternatively, whole-cell extracts from COS-7 cells (lane g), from COS-7 cells transiently transfected with RalBP1 (lane e), or from *E. coli* expressing GST-RalBP1 (positions 375 to 626) (lane f) were used. Antibody interaction was detected by a chemiluminescence assay (Amersham) with horseradish peroxidase-conjugated anti-rabbit immunoglobulin. The numbers on the left are molecular sizes in kilodaltons. The arrow indicates full-length RalBP1.




FIG. 6. Linear map of RalBP1 with putative functional domains and fragments expressed in recombinant form. Functional and putative functional domains of RalBP1 are marked. Amino acid sequences of RalBP1 that were expressed in recombinant form are indicated: a, positions 86 to 626; b, positions 375 to 626; c, positions 86 to 415.

is homologous to a family of GAPs for the Rho family of GTPases (Fig. 2 and 5). The closest homology was with the Rho-GAP domain of BCR (35% identity). Other interesting regions of RalBP1 that warrant further study are an extremely basic region between aa 113 and 172 that contains ~35% lysines (Fig. 2). Secondary-structure analysis predicted that this region forms an  $\alpha$ -helix. Another region predicted to be enriched in  $\alpha$ -helix (acidic) is between aa 400 and 600. A linear map of the protein is presented in Fig. 6.

**Binding of RalBP1 to Ral in vitro.** The yeast two-hybrid system indicated that RalBP1 bound specifically to Ral and that binding was dependent upon a functional Ral effector

domain. These results suggested that RalBP1 could be a downstream target of Ral. If this is so, RalBP1 should bind preferentially to the active, GTP-bound form of Ral. To confirm this hypothesis and to map the region of RalBP1 that binds to Ral, *in vitro* binding assays were developed.

To measure the nucleotide dependence of binding, the original 1.6-kb cDNA (encoding aa 86 to 626; Fig. 6) of RalBP1 was expressed in COS-7 cells with a Glu epitope added to the N terminus. This form of RalBP1 was immunoprecipitated with anti-Glu antibodies and then incubated with GST-Ral bound to either GTP $\gamma$ S or GDP $\beta$ S. After washing away of unbound protein, the immune complexes were run on SDS-polyacrylamide gels and immunoblotted with antiserum to RalA (Fig. 7A). As expected, only active GTP-bound Ral bound tightly to RalBP1 (positions 86 to 626). Moreover, the binding of RalBP1 to Ral was blocked by a mutation in the putative effector domain of Ral.

To localize the binding site for Ral on RalBP1, GST fusion constructs containing RalBP1 aa 86 to 626, 86 to 415, and 375 to 626 were generated (Fig. 6). RalBP1 (positions 86 to 626) contained the same amino acid sequences as those used for Fig. 7A. RalBP1 (positions 86 to 415) contained the basic  $\alpha$ -helix and the Rho-GAP domain, whereas RalBP1 (positions 375 to 626) contained the remaining carboxyl-terminal sequences of the protein. The encoded proteins were then expressed in bacteria, purified on glutathione beads, and incubated with Ral bound to either GTP $\gamma$ S or GDP $\beta$ S. After washing, the beads were run on SDS-polyacrylamide gels and immunoblotted with anti-Ral serum (Fig. 7B). Ral-binding activity was found to be associated with the acidic  $\alpha$ -helix-rich carboxyl terminus of the protein (lane c) and not with the Rho-GAP domain (lane d). RalBP1 failed to bind Ras complexed to either GTP $\gamma$ S or GDP $\beta$ S (Fig. 7C, lanes b and c). The Ras protein was functional because in its nucleotide-free bound state, it bound to Ras exchange factor Ras-GRF (Fig. 7C, lane a).

**GAP activity of RalBP1.** RalBP1 positions 86 to 415 and 375 to 626, described above, were tested for GAP activity against Rho family members CDC42, Rac1, and RhoA, as well as RalA. The proteins were preloaded with [ $\gamma$ -<sup>32</sup>P]GTP and then incubated with RalBP1 for various lengths of time. [ $\gamma$ -<sup>32</sup>P]GTP remaining bound to the proteins was measured to assess the rate of GTP hydrolysis (Fig. 8). GTP-binding proteins prebound to [ $\alpha$ -<sup>32</sup>P]GTP were used as negative controls to assess nonspecific losses of GTP. RalBP1 (positions 86 to 415) clearly enhanced the GTP hydrolysis activity of CDC42 (panel A). In

	210				
Ralbp1	PAVFRCEVDY	M. EKHGKCE	GIYRVSGIKS	KVDELKAAVD	RE
Bcr	PYIVRQCVEE	I. ERRGMEEV	GIYRVSGVAT	DIQALKAAPD	VN
Nchim	PMVVMDCIRE	I. ESRGLNSE	GLYRVSGFSD	LIEDVKMAPD	RD
B2chim	PMVVDICIRE	I. EARGLKSE	GLYRVSGFTE	HIEDVKMAPD	RD
P190	PIFIERCIEY	I. EATGLSTE	GIYRVSGNKS	EMESLQRQPD	QD
Rhogap	PIVLRETVAY	L. QAHALTTE	GIFRRSANTQ	VVREVQQKYN	MG
P122 RhoGAP	PQSIQQAMRY	L. RNHCLDQV	GLFRKSGVKS	RIQALRQMNE	SA
myr5	PIVLEKLLRH	V. EHHLYTE	GLYRKSGAAN	RTALRQALQ	TD
Consensus	P .	. .	G . R S .	. . .	
Ralbp1	PNTVASLLKQ	YLRDLPEPLL	TKELMPRFEE	ACGRTTEVEK	VQE
Bcr	VNAIAGTLKL	YFRELPEPLF	TDEFYPNFAE	GIALSDPVAK	ESC
Nchim	INIITGALKL	YFRDLPIPLI	TYDAYPKFIE	SAKIMDPDEQ	LET
B2chim	INIITGALKL	YFRDLPIFVI	TYDYTSKFDI	AAKISNADER	LEA
P190	VNTVAGAMKS	FFSELDPPLV	PYSMQIDLVE	AHKINDREQK	LHA
Rhogap	LHLPAVILKLT	FLRELPEPLL	TFDLYPHVVG	FLNIDESQRV	PAT
P122 RhoGAP	V. . . ADMIKQ	YFRDLPEPLM	TNKLSETFLQ	IYQYVVKDQR	LQA
myr5	IHAITGVLLKQ	WLRLEPEPLM	TPAQYGDFLR	AVELPEKQEQ	LAA
	. * K	. . . * LP . .	. . . .	. . . .	
					352
Ralbp1	LPEYNHLLLS	WLIHVMDHVI	AKELETKMNI	QNISIVLSPT	VQI
Bcr	LPEANLITFL	FLLDHLKRVA	EKEAVNKMSL	HNLATVFGPT	LLR
Nchim	LPPAHCTELR	YLMHLKRVV	LHEKENLMNA	ENLGIIVFGPT	LMR
B2chim	LPPAHYETLR	YLMHLKRVV	MNEKDNFMNA	ENLGIIVFGPT	LMR
P190	FPKENHEVFK	VYVSHLNKVS	HNNKVNLMTS	ENLSICFWPT	LMR
Rhogap	LPEENYQVLR	FLTARFLQIS	AHSDQNKMTN	TNLAVVFGPN	LLW
P122 RhoGAP	LPDENREVLQ	TLLYFLSDVT	AAVKENQMTF	TNLAVCLAPS	LFH
myr5	LPEANHTSLE	RLLPHLVKVA	LLEDVNRMSF	GALAIIFAPC	LLR
	. * P	. . . .	. M . . . .	. . . . P .	

FIG. 5. Alignment of RalBP1 with other Rho-GAP family members with detected GAP activity against Rho GTPases. Sequence comparisons were generated with the PILE-UP program (version 7.2; Wisconsin Genetics Computer Group). Identical amino acids are indicated, and highly conserved residues are marked in a consensus sequence.




FIG. 7. Ral binding to RalBP1 in vitro. (A) Epitope-tagged RalBP1 (positions 86 to 626) was transiently transfected into COS-7 cells. The protein was then immunoprecipitated from cell lysates and incubated with either GST-RalA or effector mutant GST-RalA49N preloaded with either GTP $\gamma$ S or GDP $\beta$ S. RalBP1-containing beads were then washed, loaded onto SDS-polyacrylamide gels, and immunoblotted with anti-Ral serum. S, supernatant from washes; IP, washed beads containing RalBP1. (B) GST-RalBP1 positions 86 to 626 (lanes b), 375 to 626 (lanes c), and 86 to 415 (lanes d) were purified from bacterial lysates on glutathione beads and incubated with RalA bound to either GTP $\gamma$ S or GDP $\beta$ S. The beads were then washed, run on SDS-polyacrylamide gels, and immunoblotted with anti-Ral serum. Recombinant RalA was run as a size standard (lane a). (C) GST-RalBP1 (positions 375 to 626) bound to glutathione beads was incubated with RasH loaded with either GTP $\gamma$ S (lane b) or GDP $\beta$ S (lane c). The beads were washed, loaded into SDS-polyacrylamide gels, and immunoblotted with anti-Ras serum. As a positive control, nucleotide-free RasH was incubated with GST-GRF. The beads were washed and run on the same gel (lane a) for immunoblotting with anti-Ras serum.

contrast, the GAP domain of RalBP1 had little, if any, effect on Rac1 (panel B) and no observable effect on RhoA (panel C) or RalA (data not shown). RalBP1 (positions 375 to 626), containing the Ral-binding activity, also had no detectable GAP

activity on any of these proteins (data not shown), demonstrating that RalBP1 is not likely to be the previously detected Ral-GAP (10).

## DISCUSSION

Interest in the function of Ral GTPases has increased recently, with the discovery that Ral GDS, the exchange factor that activates Ral, binds to the active GTP-bound form of Ras in the *S. cerevisiae* two-hybrid system and in vitro (17, 20, 34). Thus, it is possible that activation of Ral represents a distinct downstream pathway from Ras.

To identify proteins that may mediate Ral function in cells, we used the *S. cerevisiae* two-hybrid system. This report describes the cloning and characterization of a novel protein, RalBP1, that has many properties expected of such a downstream target of Ral proteins. First, RalBP1 binds specifically to the activated GTP-bound form of RalA. Second, this binding interaction is inhibited by a mutation (D49N) in the putative effector domain of Ral. In Ras, an analogous Asp-to-Asn mutation is known to block interactions with downstream targets such as Raf. Since RalB differs from RalA in amino acid sequences outside known effector domains, it too likely interacts with RalBP1. Finally, like Ral proteins, RalBP1 displays a ubiquitous tissue expression pattern.

The first clue to the function of RalBP1 came from the observation that in addition to a Ral-binding domain, the protein also contains a Rho-GAP domain. Rho-GAPs are a family of proteins that inactivate Rho GTPases by enhancing their intrinsic GTPase activity (23). By interacting preferentially with the GTP-bound form of Rho proteins, some Rho-GAPs may also function to transmit signals from these Rho proteins to downstream targets. A property common to Rho family members is the ability to influence the actin cytoskeleton (16). For example, Rho proteins mediate growth factor alterations of focal adhesions and actin stress fibers, while Rac proteins link receptors to actin polymerization associated with plasma membrane ruffling (16, 28, 29). Moreover, CDC42 has been implicated in bud site selection in *S. cerevisiae* (1) and filopodium formation in mammalian cells (22, 26). Interestingly, these GTPases may form a cascade of GTPase cycles in which CDC42 leads to Rac activation, which then leads to Rho activation (26).

At least eight functional mammalian Rho-GAPs have been detected. They are a diverse group of proteins with differing specificities toward Rho family members. They also contain a variety of additional signaling domains that may endow them with distinct cellular functions. For example, p190 is a GAP for all three of the best-characterized Rho subfamilies, Rho, Rac, and CDC42 (31). p190 binds to Ras-GAP upon cell stimulation and may help coordinate Ras and Rho signaling pathways. The BCR protein contains a Rho-GAP domain that is functional on Rac and CDC42 but not on Rho (7). BCR also contains a Ser-Thr kinase domain whose function is poorly understood. Rho-GAPs such as N-chimerin and B-chimerin are specific for Rac. Distinguishing features of these GAPs are their regulation by lipids and their tissue-specific expression (7, 24). More recently, a Rho-GAP containing a phospholipase *c*-activating domain has been cloned. It is a GAP for Rho but not Rac (18). Finally, a GAP specific for Rho that contains a myosin domain has been characterized (27). This is of particular interest since it could directly link Rho proteins to actin.

RalBP1 needs to be added to this growing family of regulatory proteins. RalBP1 can be distinguished from these other family members in that it displays the greatest specificity toward CDC42, at least in vitro. Microinjection of CDC42 into




FIG. 8. GAP activity of RalBP1. GST RalBP1 (positions 86 to 415) was incubated with [ $\gamma$ - $^{32}$ P]GTP-CDC42Hs (A), [ $\gamma$ - $^{32}$ P]GTP-Rac1 (B), or [ $\gamma$ - $^{32}$ P]GTP-RhoA (C) for the indicated amounts of time. The samples were then passed through nitrocellulose filters, and the amounts of radioactivity remaining associated with CDC42, Rac1, and RhoA were determined by filtering the free radioactivity through nitrocellulose filters. GAP activity is indicated by loss of radioactivity from the GTPases. As controls, GST-RalBP1 (positions 86 to

cells has recently been shown to promote actin microspikes and filopodium formation (22, 26). Filopodia are actin-containing structures which have been proposed to have a sensory function in fibroblasts and neural growth cones. Interestingly, a similar phenotype has also been noted in cells stimulated with the neuropeptide bradykinin (22). In fact, inhibition of endogenous CDC42 activation by injection of dominant negative CDC42 blocked these effects of bradykinin. Thus, by binding to a CDC42-GAP, Ral may influence these actin-mediated cellular activities. Since, at least in some cells, CDC42 may regulate the activity of Rac and Rho proteins (26), RalBP1 may also indirectly influence plasma membrane ruffling and actin stress fiber formation.

How might Ral binding to RalBP1 influence CDC42 function in cells? Ral could attract RalBP1 to membrane compartments where Ral is known to reside. This would lead to local inactivation of CDC42. Ral has been detected in plasma membrane fractions. However, most of the protein was found in intracellular vesicles, including clathrin-coated vesicles and secretory vesicles, suggesting that Ral is rapidly internalized from the cell surface (2, 9, 36). Thus, Ral could target RalBP1 to the plasma membrane to influence CDC42 function in filopodium formation. Alternatively, Ral could target RalBP1 to intracellular vesicles, where CDC42 could theoretically influence the interaction between vesicles and the cytoskeleton. This model is consistent with the existence of many GAPs in cells that have the potential to alter CDC42 function at discrete cellular locations. This mechanism is also consistent with the emerging theme that Ras family members function as regulatable localizing devices for other signaling molecules.

By binding selectively to the active GTP-bound form of CDC42, RalBP1 could also function to transmit signals downstream from this GTPase. A body of evidence supports such a role for Ras-GAP (3). Again, one would predict that this is a localized effect at the site of Ral-RalBP1 interactions. Finally, Ral binding might inhibit the GAP activity of RalBP1. If RalBP1 is normally in the vicinity of its target CDC42, activation of Ral could then lead to the activation of CDC42.

RalBP1 may mediate additional Ral functions through activities encoded by other regions of RalBP1, such as the basic  $\alpha$ -helix near its amino terminus. Ral also is known to interact with a distinct Ral-GAP that could be responsible for transmitting other Ral-induced cellular effects (10).

It is becoming clear that although members of the Ras superfamily display distinct functions in cells, they form a complicated network of GTPase cycles. For example, the Ras GTPase cycle may influence the Ral GTPase cycle (17, 20, 34), and here we show that the Ral GTPase cycle has the potential to influence the GTPase cycle of CDC42. Ras has already been connected to Rho family members through the Ras-GAP binding partner, p190 Rho-GAP (32). In addition, Ras nucleotide exchange factors Ras-GRF and SOS have putative exchange domains for Rho family members (12). Finally, within the Rho family, new data have revealed multiple interactions between the activity cycles of Rho, Rac, and CDC42 proteins (22, 26). Undoubtedly, these complicated cascades are necessary for these GTPases to help coordinate the complex phenotypes displayed by eukaryotic organisms.

415) was incubated with CDC42 and Rac1 bound to [ $\alpha$ - $^{32}$ P]GTP. GAP activity should not promote the loss of counts associated with the GTPases labeled in this fashion. Symbols:  $\circ$  and  $\bullet$ , [ $\gamma$ - $^{32}$ P]GTP;  $\triangle$  and  $\blacktriangle$ , [ $\alpha$ - $^{32}$ P]GTP. Filled symbols, RalBP1 added; open symbols, no RalBP1 added.

## ACKNOWLEDGMENTS

Sharon Cantor and Takeshi Urano contributed equally to this work.

We thank Steve Elledge for components of the two-hybrid system, Ian Macara for the two-hybrid rat brain cDNA expression library, and Scott Powers for anti-Glu antibodies.

This work was supported by a Public Health Service grant (GM47717) from the National Institute of General Medical Sciences and an American Cancer Society faculty research award to L.A.F. and a Human Frontiers Science Program long-term fellowship to T.U.

## REFERENCES

- Adams, A. E. M., D. I. Johnson, R. M. Longnecker, B. F. Sloat, and J. R. Pringle. 1990. CDC42 and CDC42, two additional genes involved in budding and the establishment of cell polarity in the yeast *Saccharomyces cerevisiae*. *J. Cell Biol.* **111**:131–142.
- Bielinski, D. F., N. Y. Pyun, K. Linko-Stentz, I. Macara, and R. E. Fine. 1993. Ral and Rab3a are major GTP binding proteins of axonal rapid transport vesicles of synaptic vesicles and do not redistribute following depolarization stimulated synaptosomal exocytosis. *Biochim. Biophys. Acta* **1151**:246–256.
- Boguski, M. S., and F. McCormick. 1993. Proteins regulating Ras and its relatives. *Nature* (London) **366**:643–654.
- Brondyk, W. H., C. J. McKiernan, K. A. Fortner, P. Stabila, R. W. Holz, and I. G. Macara. 1995. Interaction cloning of Rabin3, a novel protein that associates with the Ras-like GTPase Rab3A. *Mol. Cell Biol.* **15**:1137–1143.
- Chardin, P., and A. Tavitian. 1986. The ral gene: a new ras-related gene isolated by the use of a synthetic probe. *EMBO J.* **5**:2203–2208.
- Chuang, E., D. Barnard, L. Hettich, X.-F. Zhang, J. Avruch, and M. S. Marshall. 1994. Critical binding and regulatory interactions between Ras and Raf occur through a small, stable N-terminal domain of Raf and specific Ras effector residues. *Mol. Cell Biol.* **14**:5318–5325.
- Diekmann, D., S. Brill, M. D. Garrett, N. Totty, J. Hsuan, C. Monfries, C. Hall, L. Lim, and A. Hall. 1991. Bcr encodes a GTPase-activating protein for p21Rac. *Nature* (London) **351**:400–402.
- Durfee, T., K. Becherer, P.-L. Chen, S.-H. Yeh, Y. Yang, A. E. Kilourn, W.-H. Lee, and S. J. Elledge. 1993. The retinoblastoma protein associates with the protein phosphatase type 1 catalytic subunit. *Genes Dev.* **7**:555–569.
- Emkey, R., and L. A. Feig. Subcellular distribution of Ral and its GTPase activating protein, Ral-GAP. Submitted for publication.
- Emkey, R., S. Freedman, and L. A. Feig. 1991. Characterization of a GTPase activating protein for the Ras-related Ral protein. *J. Biol. Chem.* **266**:9703–9706.
- Farnsworth, C. L., M. S. Marshall, J. B. Gibbs, D. W. Stacey, and L. A. Feig. 1991. Preferential inhibition of the oncogenic form of RasH by mutations in the GAP binding/"effector" domain. *Cell* **64**:625–633.
- Feig, L. A. 1994. Guanine-nucleotide exchange factors: a family of positive regulators of Ras and related GTPases. *Curr. Opin. Cell Biol.* **6**:204–211.
- Feig, L. A., and R. Emkey. 1993. Ral gene products and their regulation, p. 247–258. In J. C. Lacal and F. McCormick (ed.), *The ras superfamily of GTPases*. CRC Press, Inc., Boca Raton, Fla.
- Feig, L. A., B. T. Pan, T. M. Roberts, and G. M. Cooper. 1986. Isolation of ras GTP-binding mutants using an in situ colony-binding assay. *Proc. Natl. Acad. Sci. USA* **83**:4607–4611.
- Frech, M., I. Schlichting, A. Wittinghofer, and P. Chardin. 1990. Guanine nucleotide binding properties of the mammalian RalA protein produced in *Escherichia coli*. *J. Biol. Chem.* **265**:6353–6359.
- Hall, A. 1992. Ras-related GTPases and the cytoskeleton. *Mol. Biol. Cell* **3**:475–479.
- Hofer, F., S. Fields, C. Schneider, and G. S. Martin. 1994. Activated Ras interacts with the Ral guanine nucleotide dissociation stimulator. *Proc. Natl. Acad. Sci. USA* **91**:11089–11093.
- Homma, Y., and Y. Emori. 1995. A dual functional signal mediator showing RhoGAP and phospholipase C- $\gamma$  stimulating activities. *EMBO J.* **14**:286–291.
- Horton, R. M., H. D. Hunt, S. N. Ho, J. K. Pullen, and L. R. Pease. 1989. Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene* **77**:61–68.
- Kikuchi, A., S. D. Demo, Z. N. Ye, Y. W. Chen, and L. T. Williams. 1994. RalGDS family members interact with the effector domain of Ras p21. *Mol. Cell Biol.* **14**:7483–7491.
- Kinsella, B. T., and W. A. Maltese. 1991. Carboxyl-terminal isoprenylation of ras-related GTP binding proteins rac1, rac2 and ralA. *J. Biol. Chem.* **266**:9786–9794.
- Kozma, R., S. Ahmed, A. Best, and L. Lim. 1995. The Ras-related protein CDC42 and bradykinin promote formation of peripheral actin microspikes and filopodia in Swiss 3T3 fibroblasts. *Mol. Cell Biol.* **15**:1942–1952.
- Lamarche, N., and A. Hall. 1994. GAPs for rho-related GTPases. *Trends Genet.* **10**:436–440.
- Leung, T., B.-E. How, E. Manser, and L. Lim. 1993. Germ cell beta chimerin, a new GTPase activation protein for p21 Rac, is specifically expressed during the acrosome assembly stage in rat testis. *J. Biol. Chem.* **268**:3813–3816.
- Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**:265–275.
- Nobes, C. D., and A. Hall. 1995. Rho, Rac and CDC42 GTPases regulate the assembly of multimolecular focal complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**:53–62.
- Reinhard, J., A. A. Scheel, D. Diekmann, A. Hall, C. Ruppert, and M. Bahler. 1995. A novel type of myosin implicated in signalling by rho family GTPases. *EMBO J.* **14**:697–704.
- Ridley, A., and A. Hall. 1992. The small GTP-binding protein rho regulates the assembly of focal adhesion and actin stress fibers in response to growth factors. *Cell* **70**:389–399.
- Ridley, A. J., H. F. Paterson, C. L. Johnston, D. Diekmann, and A. Hall. 1992. The small GTP binding protein rac regulates growth factor-induced membrane ruffling. *Cell* **70**:401–410.
- Schaber, M. D., V. M. Garsky, D. B. Boylon, W. S. Hill, E. M. Scolnick, M. S. Marshall, I. S. Sigal, and J. B. Gibbs. 1989. Ras interactions with the GTPase-activating protein (GAP). *Protein Struct. Funct. Genet.* **6**:306–315.
- Settleman, J., C. F. Albright, L. C. Foster, and R. A. Weinberg. 1992. Association between GTPase activators for Rho and Ras families. *Nature* (London) **359**:153–154.
- Settleman, J., V. Narasimhan, L. C. Foster, and R. A. Weinberg. 1992. Molecular cloning of cDNAs encoding the GAP-associated protein p190: implications for a signalling pathway from Ras to the nucleus. *Cell* **69**:539–549.
- Shou, C., C. L. Farnsworth, B. G. Neel, and L. A. Feig. 1992. Molecular cloning of cDNAs encoding a guanine-nucleotide releasing factor for Ras p21. *Nature* (London) **358**:351–354.
- Spaargaren, M., and J. R. Bischoff. 1994. Identification of the guanine nucleotide dissociation stimulator for Ral as a putative effector molecule of R-Ras, N-Ras, K-Ras and Rap. *Proc. Natl. Acad. Sci. USA* **91**:12609–12613.
- Swick, A. G., M. Janicot, T. Cheneval-Kastelic, J. C. McLenithan, and M. D. Lane. 1992. Promoter-cDNA-directed heterologous protein expression in *Xenopus laevis* oocytes. *Proc. Natl. Acad. Sci. USA* **89**:1812–1816.
- Volkmand, W., J. Pevsner, L. A. Elferink, and R. H. Scheller. 1993. Association of three small GTP-binding proteins with cholinergic synaptic vesicles. *FEBS Lett.* **317**:53–56.