

Research article

Open Access

A web-based Alcohol Clinical Training (ACT) curriculum: Is in-person faculty development necessary to affect teaching?

Daniel P Alford*^{1,2}, Jessica M Richardson¹, Sheila E Chapman^{1,2}, Catherine E Dubé³, Robert W Schadt⁴ and Richard Saitz^{1,2,5,6}

Address: ¹Clinical Addiction Research and Education (CARE) Unit, Section of General Internal Medicine, Department of Medicine, Boston Medical Center, Boston, MA, USA, ²Boston University School of Medicine, Boston, MA, USA, ³Department of Community Health, Brown University, Providence, RI, USA, ⁴Department of Educational Technology, Boston University School of Public Health, Boston, MA, USA, ⁵Youth Alcohol Prevention Center, Boston University School of Public Health, Boston, MA, USA and ⁶Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

Email: Daniel P Alford* - dalford@bu.edu; Jessica M Richardson - jesricha@montefiore.org; Sheila E Chapman - schapman@bu.edu; Catherine E Dubé - Catherine_Dube@Brown.edu; Robert W Schadt - rschadt@bu.edu; Richard Saitz - rsaitz@bu.edu

* Corresponding author

Published: 6 March 2008

Received: 13 June 2007

BMC Medical Education 2008, 8:11 doi:10.1186/1472-6920-8-11

Accepted: 6 March 2008

This article is available from: <http://www.biomedcentral.com/1472-6920/8/11>

© 2008 Alford et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Physicians receive little education about unhealthy alcohol use and as a result patients often do not receive efficacious interventions. The objective of this study is to evaluate whether a free web-based alcohol curriculum would be used by physician educators and whether in-person faculty development would increase its use, confidence in teaching and teaching itself.

Methods: Subjects were physician educators who applied to attend a workshop on the use of a web-based curriculum about alcohol screening and brief intervention and cross-cultural efficacy. All physicians were provided the curriculum web address. Intervention subjects attended a 3-hour workshop including demonstration of the website, modeling of teaching, and development of a plan for using the curriculum. All subjects completed a survey prior to and 3 months after the workshop.

Results: Of 20 intervention and 13 control subjects, 19 (95%) and 10 (77%), respectively, completed follow-up. Compared to controls, intervention subjects had greater increases in confidence in teaching alcohol screening, and in the frequency of two teaching practices – teaching about screening and eliciting patient health beliefs. Teaching confidence and teaching practices improved significantly in 9 of 10 comparisons for intervention, and in 0 comparisons for control subjects. At follow-up 79% of intervention but only 50% of control subjects reported using any part of the curriculum ($p = 0.20$).

Conclusion: In-person training for physician educators on the use of a web-based alcohol curriculum can increase teaching confidence and practices. Although the web is frequently used for dissemination, in-person training may be preferable to effect widespread teaching of clinical skills like alcohol screening and brief intervention.

Background

Practice guidelines of leading professional societies rec-

ommend alcohol screening and behavioral counseling interventions in primary care settings [1-3]. Valid, brief,

practical screening tools exist for the detection of unhealthy alcohol use in primary care settings [4], and brief interventions by physicians can reduce drinking and improve health outcomes when delivered to primary care patients with unhealthy alcohol use [5-8]. However, unhealthy alcohol use in primary care is often unrecognized and untreated, as reported in studies performed well after research demonstrating efficacy and national guidelines were published [9-14]. Although physicians recognize their responsibility in identifying and addressing alcohol problems [15], it often does not occur using effective patient-centered techniques [16]. Physician avoidance of and discomfort with brief alcohol counseling have been identified as important barriers [17].

Physician education can improve screening and brief intervention skills resulting in decreased patient drinking [7,18-22]. Some education and training programs aimed at improving physician attitudes and clinical practice around substance abuse issues have been effective [23-32]. However, despite the existence of numerous curricula [33], they are not being widely used [34]. Only half of internal medicine residency training programs have training on initial diagnosis and management of substance use disorders [34].

Web-based training can be an innovative and efficient way to connect with many individuals, while allowing learning at a convenient time for the learner. Adult learning principles [35] suggest that physicians' use of information sources outside the local sphere, such as journals, conferences, and the Internet, are essential to the enhancement and acceleration of information diffusion throughout the medical community. Although journals and books are the most common mechanism by which research findings are disseminated, they are not always read by practicing physicians [36]. The Internet can provide flexible, adaptable, tailored and sustainable access to current information [37-41] allowing for self-directed and individualized learning. Physicians have come to rely on the Internet for accessing clinical information [42] and for continuing medical education (CME) [43]. Internet-based CME has been shown to improve physician knowledge and change physician behavior [44-48]. However, little data is available on the use of web-based curricular materials by physician educators or the effectiveness of faculty development programs aimed at increasing physician use of the Internet curriculum resources. Further, although the train-the-trainer model is an efficient and widely accepted mechanism of curriculum dissemination, it is not known whether and to what degree such efforts enhance physician use of web-based curriculum tools.

To enhance dissemination of alcohol skills training to physicians, we developed an easily transportable curricu-

lum that meets the general requirements of successful web-based courses [49] and adult learning theories [35] that could be actively distributed, easily integrated into existing curricula and used by internal medicine faculty educators. In this study, we tested whether in-person faculty development training is associated with a) use of a free web-based Alcohol Clinical Training (ACT) curriculum among physician educators, b) increased alcohol-related teaching confidence, and c) increased specific alcohol-related teaching practices.

Methods

The ACT curriculum

The Alcohol Clinical Training (ACT) curriculum is a federally funded, web-based curriculum created specifically for general internist educators to teach improved clinical and communication skills (screening, assessment and brief intervention) important in addressing unhealthy alcohol use in primary care settings. The ACT curriculum is based on the U.S. National Institute on Alcohol Abuse and Alcoholism (NIAAA) *Helping Patients Who Drink Too Much: A Clinician's Guide* [50]. With a special focus on health disparities, curricular topics include the spectrum of alcohol use, selected health consequences of alcohol use, epidemiology of unhealthy alcohol use, alcohol problems frequently missed, effects of physician culture on doctor/patient communication, screening, and brief intervention. The ACT curriculum was developed by and for general internists and is designed for teaching faculty, residents and medical students in a variety of teaching settings including small group conferences and large group didactic sessions. It consists of PowerPoint slides with case-based video vignettes, as well as speaker notes and audio, and learner evaluation materials. The curriculum is designed to be flexible and modifiable (i.e. slide content can be changed and videos are available as streaming or downloadable files) and can be taught using all the components together in a 3 hour workshop or by using various components separately in 45 minute sessions (i.e. preclinic conference or attending rounds).

Pilot studies

Pilot testing was conducted to fine tune the Alcohol Clinical Training (ACT) curriculum based on input from learners in real practice settings caring for diverse (economically and culturally) patient populations. Pilot testing was performed with 3 types of physicians including residents in internal medicine, practicing community clinicians, and faculty physician educators.

Study design

In this controlled educational study, we analyzed baseline and 3-month follow-up survey data collected from applicants to a satellite workshop conducted at an American College of Physicians (ACP) national meeting. The study

was approved by the Institutional Review Board at Boston Medical Center.

Subjects were physician educators in the U.S. who applied to attend a workshop on the use of the web-based ACT curriculum. The workshop was advertised on the ACP website and newspaper and via electronic mail to members of several medical professional organizations (e.g. Society of General Internal Medicine, Association of Program Directors in Internal Medicine). The workshop was limited to 20 participants to facilitate an interactive format and because workshop space and resources were limited. Due to this limitation, the first 20 eligible applicants were invited to attend the workshop (intervention subjects), while all eligible applicants who applied after the workshop filled were asked to enroll as control subjects (to complete assessments and have access to the curriculum website but not attend the workshop). The control group was limited by the number of additional applications received, beyond the 20 accepted for the workshop. When intervention subjects attended the workshop, control subjects were sent a letter including a description of, and web address for, the online ACT curriculum [51]. Upon completion of the study, intervention subjects received reimbursement up to \$500 for travel costs or a \$500 honorarium for completing the baseline assessment, attending the workshop and completing the 3-month follow-up survey; control subjects received \$100 for completing the baseline assessment and 3-month follow-up survey. All subjects provided informed consent.

Faculty development workshop

The workshop consisted of 3 hours of in-person, interactive teaching on the effective use of the ACT curriculum, including demonstration of navigating and using website materials, modeling of teaching by expert faculty, and creating an individual action plan: a teaching project focused on using the ACT curriculum within 2 months of the workshop. Participants were required to develop an action plan objective, and to identify the target audience, setting, available resources, potential barriers, and plan for evaluation. Attendees received continuing medical education credits from Boston University.

Assessments

Both intervention and control subjects completed baseline surveys with their applications to attend the workshop. Follow-up surveys were mailed to all subjects 3 months after the workshop. Because up to 5 months separated completion of the baseline survey and workshop attendance, intervention subjects repeated the baseline survey directly preceding the workshop to assess whether baseline results changed (e.g. due to secular trend or in response to being selected to attend the workshop) (Figure 1: Participation Summary).




Figure 1
Participation Summary.

Baseline surveys included questions on respondent characteristics such as: demographics (gender, race, ethnicity, first language [English: yes/no], age, number of fluent languages other than English), residency completion year, primary teaching settings and expertise in the diagnosis and management of alcohol problems (yes/no) with any affirmative response to "Do you have expertise in the diagnosis and management of alcohol problems through: American Society of Addiction Medicine (ASAM) certification, past faculty fellowship(s), practice in an addiction specialty setting or other specified being counted as substance abuse "expertise". Additionally, baseline surveys assessed the settings in which subjects had taught about alcohol problems (i.e., resident conferences/seminars, medical student courses or conferences, continuing medical education (CME) courses, grand rounds, morning report, inpatient attending rounds, teaching while providing clinical care, other). Note that although the term "unhealthy alcohol use" better encompasses the spectrum of use of clinical interest than the term "alcohol problems," we use the latter in describing our methods and results because it was the term in use at the time of the study (consistent with the contemporaneous NIAAA guideline) [52].

Based on work by D'Onofrio and colleagues [21], the subjects were asked at both baseline and follow-up to rate, on

a 5-point Likert Scale, their teaching confidence (from "not at all" to "very") and specific teaching practices (from "rarely" to "always") in the following 5 domains: alcohol screening, assessment of readiness to change, counseling about alcohol problems, eliciting patient health beliefs, and assuring patients that they are understood [21] Follow-up surveys assessed which components (slides, notes, audio, any, none) and in which settings the ACT curriculum was used in the prior 3-months. Intervention subjects were provided a copy of their action plan and asked how much of it had been completed (none, some, or all).

Outcomes

The primary outcomes are baseline to follow-up change in self-reported teaching confidence (5 domains) and specific teaching practices (5 domains). Secondary outcomes included curriculum use, type of teaching settings, and frequency of alcohol-related advice sought. Degree of action plan completion was an outcome for intervention subjects only.

Statistical analyses

All data was analyzed with SAS/STAT software, Version 8.2 [53]. Initial analyses consisted of descriptive statistics (means, standard deviations, medians, interquartile ranges, and proportions). Comparisons were performed with 2 sample t-tests for continuous variables and chi square tests for categorical variables. Reported p-values are two-tailed, and a p-value less than 0.05 was considered statistically significant.

For the primary outcomes, we compared mean change from baseline to follow-up (calculated as follow-up score minus baseline score) in the 5 domains of teaching confidence and 5 domains of specific teaching practices both between and within groups. We also compared between-group differences in secondary outcomes. Lastly, for intervention subjects, we compared responses on the baseline

application to those from the pre-workshop repeat baseline surveys.

Results

Of the 35 physicians who completed baseline surveys, 1 was not U.S.-based and 1 refused to participate in the control group; thus 33 were enrolled (Figure 1). Of 20 intervention and 13 control subjects, 19 (95%) and 10 (77%), respectively, completed follow-up. There were no statistically significant differences in baseline characteristics by group (Table 1) including self-reported counseling and teaching others to counsel patients with alcohol problems regarding their alcohol use. One subject was missing all baseline data, with the exception of gender.

Of the 5 domains of teaching confidence and the 5 domains of specific teaching practices evaluated, compared to controls, intervention subjects increased significantly more in their confidence in teaching alcohol screening (mean change, intervention + 1.24 vs. control + 0.11, p = 0.006) and in the frequency of teaching about alcohol screening (mean change, intervention +0.56 vs. control -0.56, p = 0.02) (Table 2). Intervention subjects also increased significantly more than controls in the frequency of teaching learners to elicit patient health beliefs (mean change, intervention +0.81 vs. control -0.33, p = 0.03). Within group changes from baseline to follow-up in teaching confidence and frequency of specific teaching practices were significant for 9 of the 10 comparisons in intervention subjects, and 0 of the 10 comparisons in control subjects. The intervention subjects' pre-workshop repeat baseline surveys significantly increased compared with the baseline survey in only 1 domain – confidence in teaching to assure patients that they're understood, which did not fully explain the difference between baseline and follow-up scores (baseline to preworkshop repeat baseline mean change +0.51 vs. baseline to follow-up mean change +1.47).

Table 1: Characteristics of the 33 enrolled physician educators

	Intervention Group (N = 20) [†]	Control Group (N = 13)	p-value
Male (%)	79	62	0.43
Race (%)			0.85
Asian	37	31	
Black/African American	11	8	
White	37	54	
Other	16	8	
Hispanic (%)	5	8	1.00
English First Language (%)	58	54	1.00
Has Substance Abuse Expertise (%)	50	54	1.00
Mean Age	41	45	0.14
Mean # Fluent Languages	2	1	0.37
Mean # Years Since Residency	10	11	0.56

[†] 1 subject missing on all characteristics except gender

Table 2: Baseline to follow-up change in 5 domains of teaching confidence and specific teaching practices

	Intervention (N = 18) [†]	Control (N = 9) [†]	Between- group p-value
Teaching confidence[§]			
Alcohol screening	+ 1.24**	+ 0.11	0.006
Assessment of readiness to change	+ 1.00**	+ 0.11	0.06
Counseling about alcohol problems	+ 1.18**	+ 0.44	0.12
Eliciting patient health beliefs	+ 1.29**	+ 0.67	0.23
Assuring patients that they are understood	+ 1.47**	+ 0.56	0.07
Specific teaching practice frequency[¶]			
Alcohol screening	+ 0.56*	- 0.56	0.02
Assessment of readiness to change	+ 0.44	- 0.44	0.09
Counseling about alcohol problems	+ 0.67*	- 0.22	0.08
Eliciting patient health beliefs	+ 0.81**	- 0.33	0.03
Assuring patients that they are understood	+ 0.94*	+ 0.11	0.18

*p < .05; **p < .01; in within-group comparisons of baseline to follow-up change

[†]Baseline data were missing for one subject with follow-up data in each group (1 of 19 in the intervention group and 1 of 10 in the control group)

[§]5-point Likert scale, where 1 = Not at all Confident and 5 = Very Confident

[¶]5-point Likert scale, where 1 = Rarely and 5 = Always

At follow-up, there was more curriculum use among intervention subjects than control subjects, though the difference was not significant (79% vs. 50%, p = 0.20) (Table 3). The most commonly used component of the curriculum was the slides, whereas use of the audio component was nearly nonexistent.

Although not intended for this purpose, the curriculum was used for self-learning by the majority of subjects (71%) with no difference between intervention and control groups (Table 4). For teaching, the curriculum was used in a variety of settings, the most common of which were while providing clinical care (61%) and resident teaching conferences (43%).

Among intervention subjects, 84% (16/19) completed at least part of their action plan including 8 participants who completed their entire action plan. Two of the three intervention subjects who did not complete any of their action plan also did not use any part of the curriculum.

Discussion

In-person training for physician educators on the use of a web-based Alcohol Clinical Training (ACT) curriculum is associated with increases in confidence in teaching about alcohol screening and specific teaching practices – more frequent teaching about alcohol screening and eliciting patient health beliefs. Given the small sample size, non-significant increases are also noteworthy, including the increases associated with in-person training in confidence in teaching about assessment of readiness to change and assuring patients that they are understood, and more frequent teaching about assessment of readiness to change and counseling about alcohol problems. Also notable are the within-group findings demonstrating that intervention group confidence and teaching frequency increased significantly in 9 of 10 comparisons, which were not a result of improvements prior to the workshop. In comparison, the control group never improved significantly, and in fact, worsened in some cases. These findings suggest that in-person training of, and not only access to, this web-based curriculum can lead to improvements in alcohol-related teaching confidence and practice.

Table 3: Proportion with curriculum use at follow-up

	Intervention Group (N = 19) N (%)	Control Group (N = 10) N (%)	p-value
Any curriculum use	15 (79)	5 (50)	0.20
Slide Use	11 (58)	4 (40)	0.17
Notes Use	7 (37)	2 (20)	0.26
Audio Use	0 (0)	1 (10)	0.39
Video Use	3 (16)	1 (10)	1.00

Table 4: ACT curriculum teaching settings at follow-up

	Intervention Group (N = 18†) N (%)	Control Group (N = 10) N (%)	p-value
For my own learning	14 (78)	6 (60)	0.42
Resident teaching conferences	9 (50)	3 (30)	0.43
Medical student teaching conferences	7 (39)	1 (10)	0.19
Continuing Medical Education courses	0 (0)	0 (0)	N/A
Grand rounds	2 (11)	0 (0)	0.52
Morning report	5 (28)	1 (10)	0.37
Inpatient attending rounds	9 (50)	1 (10)	0.04*
Teaching while providing clinical care (e.g., precepting)	12 (67)	5 (50)	0.44
Other	2 (11)	0 (0)	0.52

*p < .05

† Data missing for 1 subject

Although not significant, a greater proportion of educators who had in-person training used the curriculum. Even among those without in-person training, a substantial proportion of subjects reported using the web-based curriculum. Although physicians used the curriculum, none used the audio portion and very few used the video portion. While the curriculum was used for its intended purpose, as an education tool, surprisingly, it was most commonly used for self-study, even by educators self-selected as having an interest in teaching about alcohol.

Curricular topics for generalist physician educators are expanding in number and scope while residency duration remains the same. Considering the ineffectiveness of medical residency programs in training for alcohol screening and management [34,54], target audiences for this curriculum include both the physicians who will use the curriculum to train others, and physicians being trained. The intent of developing and making this alcohol education curriculum available is to provide faculty with a variety of educational materials (i.e. video, slides) that they can take "off the shelf," modify if desired and use in a variety of settings.

The ACT curriculum was developed by and for the same group – general internists. This approach is in keeping with the emphasis on specialty specific teaching in physicians' (adult) learning principles. This approach uses the internist teacher as a role model with credibility specifically applicable to the learner's specialty [55].

Diffusion of alcohol skills training is enhanced or impeded by fundamental characteristics of the training mechanism, such as its complexity and accessibility [56]. Many previously created alcohol curricula for physicians are less easy to access, less tailored to their audiences, and less focused. In the 1990s, NIAAA developed two multi-module curricula, which include materials available for purchase on diskette [57-59]. The Project ADEPT (*Alcohol and Drug Education for Physician Training*) curriculum is a

comprehensive substance abuse curriculum for primary care physicians, which includes 7 modules, each with approximately 300 pages of instructional material. Several other non-web-based curricula are similarly lengthy, ranging from 9-hour sessions to 4-day workshops, and often address substance abuse in general, rather than being alcohol-specific, despite the fact that guidelines recommend universal screening for alcohol, but not other drugs. Many require payment or are no longer available. On the other hand, web-based materials are more easily available, and evaluations of web-based physician education have shown significant changes in both non-behavioral measures (e.g. knowledge, attitude, confidence and satisfaction) [44,45,60] as well as behavioral changes [61,62] that impact patient care [44]. With the increasing number of physicians using the Web for continuing medical education [42], it is not surprising that a number of organizations are making curricula available on the web. The NIAAA's *Helping Patients Who Drink Too Much: A Clinician's Guide*, upon which the ACT curriculum is based, is freely available on the Web and in print [50], however, it does not include the audio, video or evaluation features offered with ACT. Project Cork, Clinical Tools, Inc., the University of Florida Division of Addiction Medicine, and likely others, also provide Internet-based alcohol curricula, some of which are focused on screening and brief intervention and are geared toward physicians in general [63,64] while unlike any others, the ACT curriculum is specifically tailored to internist educators.

Several important limitations of this study evaluating the ACT curriculum should be considered. The small sample size makes it difficult to identify differences between groups and caution must be exercised when interpreting the results of non-significant findings. However, some results did reach significance. Second, the nonrandomized nature of the study could have led to confounding. For example, because enrollment in the intervention group was based on early workshop application, intervention subjects might have been more highly motivated

than control subjects. However, we believe individuals in both groups were highly motivated since they all applied to attend a workshop that involved a considerable time commitment. Further, selection bias could lead to difficulty generalizing these results from this group of physician educators volunteered to travel to attend a course to a representative sample of physician educators. Nevertheless, our original intention was to study physicians with an interest in alcohol use and not to generalize beyond that population. Further, it is possible that the workshop learners considered the workshop instructors to be opinion leaders or field experts. Relatedly, subjects' specific teaching practices were not directly measured; as such it is possible that some of the findings may be attributed to social desirability bias, reporting favorable behaviors to researchers evaluating the course they attended.

Despite these limitations, these study results suggest that posting a Web-based curriculum tailored for internist educators can lead to its use, and to improvements in teaching confidence and frequency of teaching practices that are further improved when the curriculum is demonstrated in person. Furthermore, although intended for use by educators to train others, such a curriculum can be used for self-study. More sophisticated enhancements, such as audio and video components, might require more substantial faculty development efforts, and additional research is needed on both how to better disseminate these curricula, and on practice and patient-level outcomes. Nonetheless, this educational tool has the potential, perhaps in conjunction with other efforts [65,66], to improve clinical practice in an area recognized as needing substantial improvement [9].

Conclusion

Leading professional societies recommend that alcohol screening and behavioral counseling interventions be implemented in primary care settings. But physician education to support this implementation has not been effectively or widely disseminated. This study demonstrates that a free web-based alcohol clinical training curriculum will be used by physician educators and that in-person training on the use of the curriculum can further increase teaching confidence and practices.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

DPA and RS led the creation of the study concept and design. All authors made substantial contributions to acquisition, analysis and interpretation of data and were involved in drafting and revising the manuscript for

important intellectual content. All authors have read and approved the manuscript for publication.

Acknowledgements

The authors appreciate the data management assistance of Erika Edwards, MPH, and Michael Winter, MPH, at the Data Coordinating Center, Boston University School of Public Health, Boston, MA, USA. Support from this study came from the US National Institute of Alcohol Abuse and Alcoholism (R25 AA13822). Preliminary results were presented at the annual national meetings of the Society of General Internal Medicine (SGIM), April 26–29, 2006 in Los Angeles, CA, USA, the Research Society on Alcoholism, June 24–28, 2006 in Baltimore, MD, USA, and the Association for Medical Education and Research in Substance Abuse (AMERSA), November 2–4, 2006 in Washington, DC, USA.

References

1. **United States Preventive Services Task Force and Screening and behavioral counseling interventions in primary care to reduce alcohol misuse. Recommendation Statement** [<http://www.ahrq.gov/clinic/3rduspstf/alcohol/alcomisrs.htm>]
2. United States Preventive Services Task Force: *Guide to Clinical Preventive Services* 2nd edition. Washington, DC: Office of Disease Prevention and Health Promotion; 1996.
3. Anderson P, Gual A, Colom J: **Alcohol and Primary Health Care: Clinical Guidelines on Identification and Brief Interventions**. Barcelona, Spain, Department of Health of the Government of Catalonia; 2005.
4. Saitz R: **Unhealthy alcohol use**. *N Engl J Med* 2005, **352**:596-607.
5. Fleming MF, Barry KL, Manwell LB, Johnson K, London R: **Brief physician advice for problem alcohol drinkers: a randomized controlled trial in community-based primary care practices**. *JAMA* 1997, **277**:1039-45.
6. Kristenson H, Ohlin H, Hulten-Nosslin MB, Trell E, Hood B: **Identification and intervention of heavy drinking in middle-aged men: results and follow-up of 24–60 months of long-term study with randomized controls**. *Alcohol Clin Exp Res* 1983, **7**:203-9.
7. Ockene JK, Adams A, Hurley TG, Wheeler EV, Hebert JR: **Brief physician- and nurse practitioner-delivered counseling for high-risk drinkers: does it work?** *Arch Intern Med* 1999, **159**:2198-205.
8. Allen JP, Maisto SA, Connors GJ: **Self-report screening tests for alcohol problems in primary care**. *Arch Intern Med* 1995, **155**:1726-30.
9. McGlynn EA, Asch SM, Adams J, Keesey J, Hicks J, DeCristofaro A, Kerr EA: **The quality of health care delivered to adults in the United States**. *N Engl J Med* 2003, **348**:2635-45.
10. The National Center on Addiction and Substance Abuse at Columbia University: *Missed Opportunity: National Survey of Primary Care Physicians and Patients on Substance Abuse*. Conducted by the Survey Research Laboratory, University of Illinois at Chicago New York, NY: Josiah Macy, Jr. Foundation; 2000.
11. Buchsbaum DG, Buchanan RG, Poses RM, Schnoll SH, Lawton MJ: **Physician detection of drinking problems in patients attending a general medicine practice**. *J Gen Intern Med* 1992, **7**:517-21.
12. Coulehan JL, Zettler-Segal M, Block M, McClelland M, Schulberg HC: **Recognition of alcoholism and substance abuse in primary care patients**. *Arch Intern Med* 1987, **147**:349-52.
13. Rydon P, Redman S, Sanson-Fisher RW, Reid AL: **Detection of alcohol-related problems in general practice**. *J Stud Alcohol* 1992, **53**:197-202.
14. Saitz R, Mulvey KP, Plough A, Samet JH: **Physician unawareness of serious substance abuse**. *Am J Drug Alcohol Abuse* 1997, **23**:343-54.
15. Rush B, Ellis K, Crowe T, Powell L: **How general practitioners view alcohol use. Clearing up the confusion**. *Can Fam Physician* 1994, **40**:1570-1579.
16. Makoul G, Dhurandhar A, Goel MS, Scholtens D, Rubin AS: **Communication about behavioral health risks: a study of videotaped encounters in 2 internal medicine practices**. *J Gen Intern Med* 2006, **21**:698-703.
17. McCormick KA, Cochran NE, Back AL, Merrill JO, Williams EC, Bradley KA: **How primary care providers talk to patients about alcohol: a qualitative study**. *J Gen Intern Med* 2006, **21**:966-72.

18. Adams A, Ockene JK, Wheller EV, Hurley TG: **Alcohol counseling: physicians will do it.** *J Gen Intern Med* 1998, **13**:692-98.
19. Ockene JK, Wheeler EV, Adams A, Hurley TG, Hebert J: **Provider training for patient-centered alcohol counseling in a primary care setting.** *Arch Intern Med* 1997, **157**:2334-41.
20. Saitz R, Sullivan LM, Samet JH: **Training community-based clinicians in screening and brief intervention for substance abuse problems: Translating evidence into practice.** *Subst Abus* 2000, **21**:21-31.
21. D'Onofrio G, Nadel ES, Degutis LC, Sullivan LM, Casper K, Bernstein E, Samet JH: **Improving emergency medicine residents' approach to patients with alcohol problems: a controlled educational trial.** *Ann Emerg Med* 2002, **40**:50-62.
22. Kaner EF, Lock CA, McAvoy BR, Heather N, Gilvarty E: **A RCT of three training and support strategies to encourage implementation of screening and brief alcohol intervention by general practitioners.** *British Journal of General Practice* 1999, **49**:699-703.
23. Karam-Hage M, Nerenberg L, Brower KJ: **Modifying residents' professional attitudes about substance abuse treatment and training.** *Am J Addict* 2001, **10**:40-47.
24. Rose AJ, Stein MR, Arnsten JH, Saitz R: **Teaching internal medicine resident physicians about alcoholics anonymous: a pilot study of an educational intervention.** *Subst Abus* 2006, **27**:5-11.
25. Friedmann PD, Rose J, Hayaki J, Ramsey S, Charuvastra A, Dube C, Herman D, Stein MD: **Training primary care clinicians in maintenance care for moderated alcohol use.** *J Gen Intern Med* 2006, **21**:1269-75.
26. Levin FR, Owen P, Stinchfield R, Rabinowitz E, Pace N: **Use of standardized patients to evaluate the physicians in residence program: a substance abuse training approach.** *J Addict Dis* 1999, **18**:39-50.
27. Humair JP, Cornuz J: **A new curriculum using active learning methods and standardized patients to train residents in smoking cessation.** *J Gen Intern Med* 2003, **18**:1023-27.
28. Ury WA, Rahn M, Tolentino V, Pignotti MG, Yoon J, McKeegney P, Sulmasy DP: **Can a pain management and palliative care curriculum improve the opioid prescribing practices of medical residents?** *J Gen Intern Med* 2002, **17**:625-31.
29. Amodeo M, Litchfield L: **Integrating Substance Abuse Content into Social Work Courses: Effects of Intensive Faculty Training.** *Subst Abus* 1999, **20**:5-16.
30. Fleming M, Barry K, Davis A, Kropp S, Kahn R, Rivo M: **Medical education about substance abuse: changes in curriculum and faculty between 1976 and 1992.** *Acad Med* 1994, **69**:362-69.
31. Lewis DC, Niven RG, Czechowicz D, Trumble JG: **A review of medical education in alcohol and other drug abuse.** *JAMA* 1987, **257**:2945-48.
32. Alford DP, Clark TW, Samet JH: **Promoting substance abuse education among generalists: the Chief Resident Immersion Training (CRIT) program.** *Journal of General Internal Medicine* 2004, **19**(Suppl):93.
33. Gual A, Anderson P, Sequera L, Colom J: **Alcohol and Primary Health Care: Training Programme on Identification and Brief Interventions.** Barcelona, Spain, Department of Health of the Government of Catalonia; 2005.
34. Isaacson JH, Fleming M, Kraus M, Kahn R, Mundt M: **A national survey of training in substance use disorders in residency programs.** *J Stud Alcohol* 2000, **61**:912-15.
35. Spencer JA, Jordan RK: **Learner centred approaches in medical education.** *BMJ* 1999, **318**:1280-1283.
36. Brown BS: **Reducing impediments to technology transfer in drug abuse programming.** *NIDA Res Monogr* 1995, **155**:169-85.
37. Curran VR, Fleet L: **A review of evaluation outcomes of web-based continuing medical education.** *Med Educ* 2005, **39**:561-67.
38. Cook DA, Dupras DM, Thompson WG: **An online core curriculum in primary care medicine for internal medicine residents.** *Med Educ* 2003, **37**:1043.
39. Wellbery C, Gooch R: **A Web-based multimedia medical humanities curriculum.** *Fam Med* 2005, **37**:165-67.
40. Peterson MW, Galvin JR, Dayton C, D'Alessandro MP: **Realizing the promise: delivering pulmonary continuing medical education over the Internet.** *Chest* 1999, **115**:1429-36.
41. Godin P, Hubbs R, Woods B, Tsai M, Nag D, Rindfleisch T, Dev P, Melmon KL: **New paradigms for medical decision support and education: the Stanford Health Information Network for Education.** *Top Health Inf Manage* 1999, **20**:1-14.
42. Bennett NL, Casebeer LL, Kristofco RE, Strasser SM: **Physicians' Internet information-seeking behaviors.** *J Contin Educ Health Prof* 2004, **24**:31-38.
43. Sargeant J, Curran V, Jarvis-Selinger S, Ferrier S, Allen M, Kirby F, Ho K: **Interactive on-line continuing medical education: physicians' perceptions and experiences.** *J Contin Educ Health Prof* 2004, **24**:227-36.
44. Fordis M, King JE, Ballantyne CM, Jones PH, Schneider KH, Spann SJ, Greenberg SB, Greisinger AJ: **Comparison of the instructional efficacy of Internet-based CME with live interactive CME workshops: a randomized controlled trial.** *JAMA* 2005, **294**:1043-51.
45. Proude EM, Conigrave KM, Haber PS: **Effectiveness of skills-based training using the Drink-less package to increase family practitioner confidence in intervening for alcohol use disorders.** *BMC Med Educ* 2006, **6**:8.
46. Casebeer L, Kristofco RE, Strasser S, Reilly M, Krishnamoorthy P, Rabin A, Zheng S, Karp S, Myers L: **Standardizing evaluation of on-line continuing medical education: physician knowledge, attitudes, and reflection on practice.** *J Contin Educ Health Prof* 2004, **24**:68-75.
47. Short LM, Surprenant ZJ, Harris JM Jr: **A community-based trial of an online intimate partner violence CME program.** *Am J Prev Med* 2006, **30**:181-85.
48. Harris JM Jr, Kutob RM, Surprenant ZJ, Maiuro RD, Delate TA: **Can Internet-based education improve physician confidence in dealing with domestic violence?** *Fam Med* 2002, **34**:287-92.
49. Wong G, Greenhalgh T, Russell J, Boynton P, Toon P: **Putting your course on the Web: lessons from a case study and systematic literature review.** *Med Educ* 2003, **37**:1020-1023.
50. National Institute on Alcohol Abuse and Alcoholism: *U.S. Department of Health and Human Services. Helping Patients Who Drink Too Much: A Clinician's Guide. Updated 2005 Edition. NIH Publication No. 07-3769* Rockville, MD; 2007.
51. **Alcohol Screening and Brief Intervention Curriculum** [<http://www.mdalcoholtraining.org/>]
52. National Institute on Alcohol Abuse and Alcoholism: *Helping Patients with Alcohol Problems. A Health Practitioner's Guide* Bethesda, MD: National Institutes of Health; 2003.
53. **SAS/STAT Software, Version 8.2 of the SAS System for Windows.** SAS Institute Inc. Cary, NC; 1999.
54. Stimmel B, Cohen D, Colliver J, Swartz M: **An Assessment of House Staff's Knowledge of Alcohol and Substance Abuse Utilizing Standardized Patients.** *Subst Abus* 2000, **21**:1-7.
55. Slotnick HB: **How doctors learn: education and learning across the medical-school-to-practice trajectory.** *Acad Med* 2001, **76**:1013-26.
56. Rogers EM: *Diffusion of Innovations.* New York, NY 5th edition. 2003.
57. Fleming M, Murray M: *A Medical Education Model for the Prevention and Treatment of Alcohol Use Disorders* National Institute of Alcohol Abuse and Alcoholism; 1998.
58. Sirica C, Lewis DC: **Training about alcohol and substance abuse for all primary care physicians.** *A conference sponsored by the Josiah Macy, Jr. Foundation, Phoenix, Arizona, Josiah Macy, Jr. Foundation, New York, NY* 1994.
59. Murray M, Fleming M: **Prevention and treatment of alcohol-related problems: an international medical education model.** *Acad Med* 1996, **71**:1204-10.
60. Harris JM, Salasche SJ, Harris RB: **Can Internet-based continuing medical education improve physicians' skin cancer knowledge and skills?** *J Gen Intern Med* 2001, **16**:50-56.
61. Allison JJ, Kiefe CI, Wall T, Casebeer L, Ray MN, Spettell CM, Hook EW III, Oh MK, Person SD, Weissman NW: **Multicomponent Internet continuing medical education to promote chlamydia screening.** *Am J Prev Med* 2005, **28**:285-90.
62. Stewart M, Marshall JN, Ostbye T, Feightner JW, Brown JB, Harris S, Galajda J: **Effectiveness of case-based on-line learning of evidence-based practice guidelines.** *Fam Med* 2005, **37**:131-38.
63. **ProjectCork** [<http://www.projectcork.org/>]
64. **Alcohol Research and Education** [<http://webapps.health.ufl.edu/aec/jsp/02CourseList.jsp>]. Division of Addiction Medicine, University of Florida
65. Garnick DW, Lee MT, Chalk M, Gastfriend D, Horgan CM, McCorry F, McLellan AT, Merrick EL: **Establishing the feasibility of per-**

formance measures for alcohol and other drugs. *J Subst Abuse Treat* 2002, **23**:375-85.

66. Garnick DW, Horgan CM, Chalk M: **Performance measures for alcohol and other drug services.** *Alcohol Res Health* 2006, **29**:19-26.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1472-6920/8/11/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

