

Shankar Prasad Kanaujia,^a
Chellamuthu Vasuki Ranjani,^a
Jeyaraman Jeyakanthan,^b Seiki
Baba,^{b,c} Chizu Kuroishi,^b Akio
Ebihara,^b Akeo Shinkai,^b Seiki
Kuramitsu,^{b,c} Yoshitsugu Shiro,^b
Kanagaraj Sekar^{a,d,*} and
Shigeyuki Yokoyama^{b,e,f,*}

^aBioinformatics Centre (Centre of Excellence in Structural Biology and Biocomputing), Indian Institute of Science, Bangalore 560 012, India, ^bRIKEN SPring-8 Center, Harima Institute, 1-1-1 Kouto, Sayo, Hyogo 679-5148, Japan, ^cDepartment of Biology, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan, ^dSupercomputer Education and Research Centre, Indian Institute of Science, Bangalore 560 012, India, ^eRIKEN Genomic Sciences Center, 1-7-22 Suehiro-cho, Tsurumi, Yokohama 230-0045, Japan, and ^fDepartment of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Correspondence e-mail:
sekar@serc.iisc.ernet.in,
yokoyama@biochem.s.u-tokyo.ac.jp

Received 30 October 2006
Accepted 30 December 2006



© 2007 International Union of Crystallography
All rights reserved

Cloning, expression, purification, crystallization and preliminary X-ray crystallographic study of DHNA synthetase from *Geobacillus kaustophilus*

The aerobic Gram-positive bacterium *Geobacillus kaustophilus* is a bacillus species that was isolated from deep-sea sediment from the Mariana Trench. 1,4-Dihydroxy-2-naphthoate (DHNA) synthetase plays a vital role in the biosynthesis of menaquinone (vitamin K₂) in this bacterium. DHNA synthetase from *Geobacillus kaustophilus* was crystallized in the orthorhombic space group C222₁, with unit-cell parameters $a = 77.01$, $b = 130.66$, $c = 131.69$ Å. The crystal diffracted to a resolution of 2.2 Å. Preliminary studies and molecular-replacement calculations reveal the presence of three monomers in the asymmetric unit.

1. Introduction

Interest in thermophilic microorganisms that live and thrive at high temperatures of up to or above 373 K has been increasing in recent decades (Andrade *et al.*, 1999; Singleton & Amelunxen, 1973). One thermophilic organism of immense research interest is *Geobacillus kaustophilus*, a bacillus species isolated from deep-sea sediment from the Mariana Trench (Takami *et al.*, 1997). It is an aerobic endospore-forming Gram-positive bacterium that grows optimally at 333 K with an upper temperature limit of 347 K (Takami *et al.*, 2004). Vitamin K₂ (menaquinone) plays a vital role as an electron carrier in this bacterium (Bentley & Meganathan, 1982; Johnston *et al.*, 2005). Quinones serve as a source of energy and are of fundamental importance to all life forms (Bentley & Meganathan, 1982). They occur as ubiquinone in the mitochondria of animals, plastoquinone in the chloroplasts of plants and menaquinone in bacteria. A key reaction in the biosynthesis of menaquinone involves the conversion of *O*-succinylbenzoyl-CoA to 1,4-dihydroxy-2-naphthoyl-CoA (Truglio *et al.*, 2003). 1,4-Dihydroxy-2-naphthoate synthetase (MenB), abbreviated as DHNA synthetase (EC 4.1.3.36), is believed to be responsible for cyclization in the reaction (Meganathan & Bentley, 1981; Sharma *et al.*, 1992).

Menaquinone has been found to be an important product of metabolism in many bacterial species (Bentley & Meganathan, 1982). Inactivation of the genes encoding the menaquinone-biosynthetic enzymes has been found to produce a demand for menaquinone in the growth medium (Truglio *et al.*, 2003). Thus, the enzymes involved in the menaquinone-biosynthesis pathway play a significant role in the metabolism of bacterial species. Understanding the three-dimensional crystal structures of enzymes from the Gram-positive bacterium *G. kaustophilus* will be of great significance in the field of thermophile research. Intensive research needs to be performed to study the thermostability of macromolecules and to screen new thermophiles for biotechnological processes. The recent focus has been on the study of metabolic pathways and mechanisms involved in the regulation of thermophilic proteins. Furthermore, insight into the crystal structure may shed light on the design of a potent inhibitor for the protein.

2. Materials and methods

2.1. Cloning, expression and purification of the protein

The DHNA gene (GK2873) was amplified by PCR using *G. kaustophilus* HTA426 genomic DNA as the template. The

Table 1

X-ray diffraction data statistics of DHNA synthetase.

Values in parentheses are for the last resolution bin.

Wavelength (Å)	1.0
Space group	C222 ₁
Temperature (K)	100
Unit-cell parameters (Å)	$a = 77.01, b = 130.66, c = 131.69$
Matthews coefficient (Å ³ Da ⁻¹)	1.83 (3 subunits per ASU)
Solvent content (%)	32.8
Resolution range (Å)	50.0–2.20 (2.28–2.20)
No. of observed reflections	1320862
No. of unique reflections	33813 (3181)
Completeness (%)	98.9 (94.2)
R_{sym}^{\dagger} (%)	4.1 (8.1)
Multiplicity	9.8 (6.5)
Average $I/\sigma(I)$	39 (14)

$$\dagger R_{\text{sym}} = \sum I_{hkl} - \langle I \rangle / \sum I_{hkl}$$

amplified fragment was cloned under the control of the T7 promoter of the *Escherichia coli* expression vector pET-HisTEV, with the tobacco etch virus (TEV) protease-recognition site, Glu-Asn-Leu-Tyr-Phe-Gln-Gly, instead of the thrombin-recognition site of the pET-15b vector (Novagen, Madison, WI, USA). The expression vector was introduced into *E. coli* BL21-CodonPlus(DE3)-RIL strain (Stratagene, La Jolla, CA, USA) and the recombinant strain was cultured in 2.5 l LB broth supplemented with 50 µg ml⁻¹ ampicillin. The cells (9.6 g) were collected by centrifugation, washed with 50 ml buffer A (20 mM Tris-HCl pH 8.0) containing 0.5 M NaCl, 5 mM 2-mercaptoethanol and 1 mM phenylmethylsulfonyl fluoride and resuspended in 25 ml of the same buffer. The cells were then disrupted by sonication in a chilled water bath and the cell lysate was incubated at 343 K for 13 min. The sample was centrifuged at 15 000g for 30 min and the supernatant was applied onto a HisTrap Chelating HP5 Superdex 200 column (GE Healthcare Bioscience Corp., Piscataway, NJ, USA) pre-equilibrated with buffer A containing 0.5 M NaCl and 20 mM imidazole, which was eluted with a linear gradient of 0.02–0.5 M imidazole. The sample containing DHNA synthetase was then loaded onto a HiLoad 16/60 Superdex 200 column (GE Healthcare Bioscience Corp.) pre-equilibrated with buffer A containing 0.5 M NaCl and 20 mM imidazole. The eluted fractions containing DHNA synthetase were collected and treated with TEV protease at 303 K for 1 h. The protein sample was applied onto a HisTrap Chelating HP5-Superdex 200 column (GE Healthcare Bioscience Corp.) pre-equilibrated with buffer A containing 0.5 M NaCl and 20 mM



Figure 1
Crystal of DHNA synthetase from *G. kaustophilus* (C-centred orthorhombic).

imidazole. The flowthrough fraction was collected and the purified protein was loaded onto a HiPrep 26/10 desalting column, which was eluted with buffer A containing 0.2 M NaCl.

2.2. Crystallization experiments

The protein was concentrated using a Vivaspin 20 concentrator (10 kDa molecular-weight cutoff, Sartorius, AG, Göttingen, Germany). The protein concentration was determined by measuring the absorbance at 280 nm (Kuramitsu *et al.*, 1990). The concentration of the purified protein was ~10 mg ml⁻¹ in 20 mM Tris-HCl pH 8.0 and 200 mM NaCl. Preliminary crystallization screening was carried using the Index kit (Hampton Research). Sitting-drop vapour-diffusion experiments were set up by pipetting drops consisting of 1 µl protein solution and 1 µl well solution. The well solution contained 30% (v/v) pentaerythritol ethocylate, 0.05 M ammonium sulfate and 0.05 M bis-tris pH 6.5. Crystals were obtained when 1 µl protein solution was mixed with 1 µl well solution and allowed to equilibrate against 100 µl well solution at 293 K. 50% (v/v) Paratone-N and 50% (v/v) paraffin oil were used as a cryoprotectant. Crystals (Fig. 1) appeared in around 20 d.

2.3. Data collection and processing

The diffraction data were collected from a single crystal (approximately 0.2 × 0.2 × 0.2 mm in size) at 100 K using the RIKEN Structural Genomics Beamline II (BL26B2) at SPring-8 (Hyogo, Japan) with a Jupiter 210 CCD detector (Rigaku MSC Co., Tokyo, Japan). The crystal-to-detector distance was maintained at 230 mm. The program DENZO (Otwinowski, 1993) was used for data processing. The crystal diffracted to 2.2 Å resolution. The details of the diffraction data statistics are given in Table 1.

3. Results and discussion

Analysis of the diffracted intensities indicates that DHNA synthetase from *G. kaustophilus* belongs to the C-centred orthorhombic space group C222₁, with unit-cell parameters $a = 77.01, b = 130.66, c = 131.69$ Å. The calculated Matthews coefficient ($V_M = 1.83$ Å³ Da⁻¹), corresponding to a solvent content of 32.8% (Matthews, 1968), indicates the presence of three monomers in the asymmetric unit. The X-ray diffraction data statistics are summarized in Table 1. The DHNA enzyme consists of 272 amino-acid residues with a subunit molecular weight of 30 kDa.

3.1. Molecular replacement

The crystal structure of DHNA synthetase from *G. kaustophilus* was solved by molecular-replacement calculations using the program CNS v.1.1 (Brünger *et al.*, 1998). The coordinates of the search model used were those of MenB from *Mycobacterium tuberculosis* (PDB code 1rjn), which has 50% amino-acid sequence identity to DHNA synthetase from *G. kaustophilus*. Three monomers were located in the asymmetric unit and were subsequently subjected to rigid-body refinement using the program CNS v.1.1 (Brünger *et al.*, 1998). 5% of the reflections were used for R_{free} calculation (Brünger, 1992). The crystallographic R_{work} and R_{free} values of the partially refined structure were 36.6% and 38.1%, respectively.

SPK, CVR and KS thank the Bioinformatics Centre, the Interactive Graphics Based Molecular Modelling Facility and the Super-computer Education and Research Centre. We thank Drs Satoru Watanabe and Ryoichi Arai for construction of the pET-HisTEV

vector and Nagisa Takemoto for purification of the protein. We also thank Dr Koji Takio and Naoko Takahashi for the N-terminal amino-acid sequence analysis of the recombinant DHNA synthase. This work was supported by the RIKEN Structural Genomic/Proteomics Initiative (RSGI), the National Project on Protein Structural and Functional Analyses, Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Andrade, C. M. M. C., Pereira, N. Jr & Antranikian, G. (1999). *Rev. Microbiol.* **30**, 287–298.
- Bentley, R. & Meganathan, R. (1982). *Microbiol. Rev.* **46**, 241–280.
- Brünger, A. T. (1992). *Nature (London)*, **355**, 472–474.
- Brünger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W., Jiang, J.-S., Kuszewski, J., Nilges, M., Pannu, N. S., Read, R. J., Rice, L. M., Simonson, T. & Warren, G. L. (1998). *Acta Cryst.* **D54**, 905–921.
- Johnston, J. M., Arcus, V. L. & Baker, E. N. (2005). *Acta Cryst.* **D61**, 1199–1206.
- Kuramitsu, S., Hiromi, K., Hayashi, H., Morino, Y. & Kagamiyama, H. (1990). *Biochemistry*, **29**, 5469–5476.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Meganathan, R. & Bentley, R. (1981). *Biochemistry*, **20**, 5336–5340.
- Otwinowski, Z. (1993). *Proceedings of the CCP4 Study Weekend. Data Collection and Processing*, edited by L. Sawyer, N. Isaacs & S. Bailey, pp. 56–62. Warrington: Daresbury Laboratory.
- Sharma, V., Suvarna, K., Meganathan, R. & Hudspeth, M. E. (1992). *J. Bacteriol.* **174**, 5057–5062.
- Singleton, R. Jr & Amelunxen, R. E. (1973). *Bacteriol. Rev.* **37**, 320–342.
- Takami, H., Inoue, A., Fuji, F. & Horikoshi, K. (1997). *FEMS Microbiol. Lett.* **152**, 279–285.
- Takami, H., Nishi, S., Lu, J., Shimamura, S. & Takaki, Y. (2004). *Extremophiles*, **8**, 351–356.
- Truglio, J. J., Theis, K., Feng, Y., Gajda, R., Machutta, C., Tonge, P. J. & Kisker, C. (2003). *J. Biol. Chem.* **278**, 42352–42360.