

Akira Nakamura,^a Chieko Wada^b
and Kunio Miki^{a,c,*}

^aDepartment of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan, ^bGraduate School of Biostudies, Kyoto University, Sakyo-ku, Kyoto 606-8315, Japan, and ^cRIKEN SPring-8 Center at Harima Institute, Koto 1-1-1, Sayo, Hyogo 679-5148, Japan

Correspondence e-mail:
miki@kuchem.kyoto-u.ac.jp

Received 6 December 2006
Accepted 19 March 2007

Expression and purification of F-plasmid RepE and preliminary X-ray crystallographic study of its complex with operator DNA

The replication initiator factor RepE of the F plasmid in *Escherichia coli* is an essential protein that stringently regulates the F-plasmid copy number. The RepE protein has a dual function: its monomer functions as a replication initiator, while its dimer acts as a transcriptional repressor of the *repE* gene. The wild-type dimeric RepE protein was expressed as an N-terminal histidine-tagged protein, purified under native conditions with a high salt concentration and crystallized in complex with the *repE* operator DNA using the sitting-drop vapour-diffusion technique. The crystals diffracted to a resolution of 3.14 Å after the application of dehydration and crystal annealing and belong to space group $P2_1$, with unit-cell parameters $a = 60.73$, $b = 99.32$, $c = 95.00$ Å, $\beta = 108.55^\circ$.

1. Introduction

The mini-F plasmid, which is a derivative of the F factor in *Escherichia coli*, replicates under strict control, maintaining a copy number of 1–2 per host chromosome (Kline, 1985). The plasmid-encoded initiator protein RepE (251 amino acids, 29 kDa) plays an important role in the regulation of mini-F replication. Two distinct oligomeric states, monomeric and dimeric, of RepE have been found and their functions are known to be different (Ishiai *et al.*, 1994). The RepE monomer is a replication initiator that recognizes a specific 19 bp sequence DNA (iteron; Masson & Ray, 1986, 1988; Tokino *et al.*, 1986). Within the replication origin (*ori2*), four directly repeated iterons (DR) are present and are bound by RepE molecules, which could cause local DNA melting at the 13-mer region with the assistance of a histone-like protein HU and subsequent initiation of mini-F replication (Kawasaki *et al.*, 1996; Komori, Matsunaga *et al.*, 1999). In contrast, the RepE dimer is an autogenous repressor that binds to the *repE* promoter/operator region located just upstream of the *repE* gene (Rokeach *et al.*, 1985; Masson & Ray, 1986, 1988; Tokino *et al.*, 1986). The operator of *repE* includes an inverted-repeat (IR) sequence, which shares 8 bp with the iterons. The RepE dimer recognizes the common 8 bp of the IR and inhibits the transcription of *repE* itself. Since RepE exists primarily as a dimer, in order to activate RepE as an initiator, conversion from the dimer to the monomer is accomplished by the molecular chaperones DnaK, DnaJ and GrpE (Kawasaki *et al.*, 1990, 1991; Wada *et al.*, unpublished results). The crystal structure of the active initiator form of RepE has been reported previously (Komori, Matsunaga *et al.*, 1999) and suggested that the C-terminal domain recognizes the specific DNA sequence and that the N-terminal domain interacts with the DNA backbone. Based on the RepE monomer structure, a marked conformational change is required for dimerization. Structural information on the repressor form of RepE, however, remains unknown. Because most initiator proteins generally aggregate easily, wild-type and full-length initiators have not been targeted for structural studies. We have successfully expressed and purified wild-type RepE and here we report the crystallization and preliminary X-ray diffraction analysis of RepE bound to DNA containing an IR




Table 1

Data-collection and processing statistics of the His₆-RepE-DNA1 complex.

Values in parentheses refer to the highest resolution shell.

Wavelength (Å)	1.0000
Space group	<i>P2₁</i>
Unit-cell parameters (Å, °)	<i>a</i> = 60.73, <i>b</i> = 99.32, <i>c</i> = 95.00, β = 108.55
Resolution range (Å)	50.0–3.14 (3.25–3.14)
Observed reflections	42887
Unique reflections	16535
Completeness (%)	88.3 (73.6)
<i>I</i> / σ (<i>I</i>)	11.0 (3.9)
<i>R</i> _{sym} † (%)	6.7 (15.0)

† $R_{sym} = \frac{\sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl) \rangle|}{\sum_{hkl} \sum_i I_i(hkl)}$, where *I_i(hkl)* is the *i*th intensity measurement of reflection *hkl* and $\langle I(hkl) \rangle$ is the average value over multiple measurements.

sequence in order to elucidate the crystal structure of dimeric RepE in complex with *repE* operator DNA.

2. Materials and methods

2.1. Expression and purification

E. coli strain M15 harbouring pREP4 plasmid (Qiagen, Hilden, Germany) was transformed with pKV7202 plasmid carrying an N-terminal His-tagged *repE* gene (Matsunaga *et al.*, 1995) and grown at 310 K in Luria broth medium containing 50 µg ml⁻¹ ampicillin and 25 µg ml⁻¹ kanamycin. When the culture reached mid-log growth phase (optical density at 600 nm = 0.6), the culture was rapidly cooled on ice and protein expression was induced by the addition of isopropyl β-D-thiogalactopyranoside (IPTG) to a final concentration of 0.1 mM. After further growth for 22 h at 293 K, the cells were harvested by centrifugation at 2110g for 30 min at 277 K. All of the following purification steps of N-terminal His-tagged RepE (His₆-RepE) were performed under cooled conditions at 277 K. The bacterial pellets were resuspended in lysis buffer [50 mM NaH₂PO₄ pH 8.0, 1 M KCl, 20% (v/v) glycerol and 10 mM imidazole] supplemented with 1 mg ml⁻¹ lysozyme and 0.5 mM 4-(2-aminoethyl)-benzenesulfonyl fluoride (*p*-ABSF) and disrupted by sonication on ice. The cell lysate was centrifuged at 27 000g for 60 min and the supernatant was then filtered using a membrane filter with 0.45 µm pore size. After Ni²⁺-nitrilotriacetic acid (Ni-NTA) Superflow resin (Qiagen) equilibrated with lysis buffer was added to the supernatant of the cell lysate, the mixture was stirred for 60 min and loaded onto an Ni-NTA Superflow column. The column was then washed with a stepwise gradient of a wash buffer [50 mM sodium phosphate pH 8.0, 1 M KCl and 20% (v/v) glycerol] containing 30, 50 and 100 mM imidazole. His₆-RepE was eluted with a wash buffer containing 200 mM imidazole. Several cycles of purification by means of Ni-NTA immobilized metal-affinity chromatography (IMAC) were performed. For a stock solution, glycerol was added to a final concentration of 35% (v/v) and purified His₆-RepE was stored in small aliquots at 193 K.

2.2. Preparation of RepE–DNA complex

For RepE–DNA complex formation, oligonucleotides containing the inverted-repeat sequence in the *repE* operator recognized by the RepE dimer were prepared by Hokkaido System Science (Sapporo, Japan). Their sequences were designed based on the expectation that DNA duplexes form a pseudo-continuous helix in crystal packing (Joachimik & Sigler, 1991), as shown in Fig. 1. DNA duplexes were prepared essentially as described by Komori, Sasai *et al.* (1999), except that the final concentrations were 2.4 mM. Each DNA duplex

was added to His₆-RepE solution in a 1.2-fold molar excess relative to the RepE dimer and the mixture was subsequently dialyzed against crystallization buffer [100 mM potassium citrate pH 6.2, 100 mM KCl, 10 mM MgCl₂, 10% (v/v) glycerol and 2 mM dithiothreitol (DTT)] at 277 K and concentrated to approximately 5 mg ml⁻¹ His₆-RepE concentration. Before crystallization, the formation of the protein–DNA complex was confirmed by native polyacrylamide gel electrophoresis (PAGE), in which a band corresponding to the protein–DNA complex was stained with both Coomassie Brilliant Blue (CBB) and Mupid-Blue (Advance, Tokyo, Japan).

2.3. Crystallization

The crystallization conditions for the His₆-RepE–operator complexes were screened by the sitting-drop vapour-diffusion method at 277 K. Initial conditions were found using a Hydra II Plus One (Matrix Technologies Corporation, Hudson, NH, USA) with commercially available screening kits (Hampton Research, Aliso Viejo, CA, USA; Sigma–Aldrich, St Louis, MO, USA). A drop was prepared by mixing equal volumes (0.3 µl for initial screening and 1.0 µl for optimization) of the His₆-RepE–operator complex solution and the reservoir solution, allowing the mixture to equilibrate against the reservoir solution (80 µl for initial screening and 100 µl for optimization). The presence of the complex in the crystal component was identified by native PAGE using the washed and dissolved crystals.

2.4. X-ray diffraction analysis

A His₆-RepE–DNA1 (Fig. 1) crystal mounted in a nylon loop was transferred and soaked for 36 h at 277 K in a dehydration solution containing a 1.2-fold higher concentration of each component of the crystallization condition. The dehydrated crystal was briefly immersed into the dehydration solution supplemented with 25–30% (v/v) glycerol as a cryoprotectant and then frozen by flash-cooling in a stream of nitrogen gas at 100 K. Prior to X-ray irradiation, we performed crystal annealing (Harp *et al.*, 1998; Heras & Martin, 2005) by blocking the cryostream twice for 8 s each time. X-ray diffraction data were collected using an ADSC Quantum 315 CCD detector on SPring-8 beamline BL41XU at a wavelength of 1.0000 Å. A total of 180 frames were recorded with an oscillation angle of 1.0°, an exposure time of 3.0 s per frame and a crystal-to-detector distance of 350 mm. The diffraction intensities were integrated with the *HKL-2000* (Otwinowski & Minor, 1997) program

DNA name	Sequence	Crystals obtained
DNA1	TT AGTGTGACAACTCTAAAAACTTGTGCACACT TC AA TCACACTGTTAGATTTTTTGAACAGTGTGA AG	Diffract to 3.14 Å
DNA2	T AGTGTGACAACTCTAAAAACTTGTGCACACT TC AA TCACACTGTTAGATTTTTTGAACAGTGTGA A	Very small
DNA3	T AGTGTGACAACTCTAAAAACTTGTGCACACT TC GA TCACACTGTTAGATTTTTTGAACAGTGTGA A	Diffract to ~8 Å
DNA4	C AGTGTGACAACTCTAAAAACTTGTGCACACT GT TG TCACACTGTTAGATTTTTTGAACAGTGTGA C	Not obtained

Figure 1
DNA duplexes used in complex formation and crystallization trials. The common 8 bp sequences between iteron and operator are underlined and their directions are indicated by the arrows.

suite. The data-collection and processing statistics are summarized in Table 1.

3. Results and discussion

Wild-type RepE proteins tend to aggregate and more than 80% of the RepE expressed in *E. coli* was collected as inclusion bodies. RepE protein has therefore previously been purified under denaturing conditions using guanidine hydrochloride (Kawasaki *et al.*, 1992; Kline *et al.*, 1992; Ishiai *et al.*, 1994; Matsunaga *et al.*, 1997). However, in order to keep RepE proteins with their native fold for subsequent crystallization, we attempted to purify His₆-RepE without denaturing. For bacterial culture conditions, the temperature was shifted from 310 to 293 K in order to decrease the protein-expression level during expression of the His₆-RepE protein. Furthermore, during RepE purification a high salt concentration of 1 M was maintained in the buffers to prevent protein aggregation. Owing to these modifications, a significant improvement in the yield of purified RepE protein was achieved (from less than 1 mg to 25 mg of His₆-RepE per litre of culture). After several cycles of purification by the IMAC method, His₆-RepE attained over 90% purity as estimated by SDS-PAGE followed by staining with CBB. The result of gel-filtration chromatography for His₆-RepE using a wash buffer containing 200 mM imidazole [eluted at 76 ml from Hi-Load 16/60 Superdex 200 (GE Healthcare Biosciences, Tokyo, Japan) at a flow rate of 0.3 ml min⁻¹] indicates the presence of an ~80 kDa protein but not of an ~30 kDa protein, suggesting that His₆-RepE forms dimers even under high-salt conditions (1 M).

The complexes of His₆-RepE with each of four DNA duplexes containing IR (31 or 33 bp; Fig. 1) were prepared and their complexation was confirmed by native PAGE as described above. Under normal conditions (pH 8.8), His₆-RepE is unable to diffuse into a native polyacrylamide gel because the theoretical pI value is 9.5, whereas the complexes are able to move into the gel owing to the effect of the acidic oligonucleotides. As a result of staining the gels with CBB and Mupid-Blue independently to confirm the presence of His₆-RepE and the DNA, respectively, specific sharp bands corresponding to the complexes were indeed detected at the same position in both gels (Fig. 2). No corresponding band was found in the lanes in which His₆-RepE and DNA had been loaded independently onto the native gel (data not shown). The complex His₆-RepE-DNA1 was




Figure 2 Native PAGE of the crystals of the His₆-RepE-DNA1 complex. Samples were loaded onto a 15% polyacrylamide gel (37.5:1 acrylamide:bisacrylamide) and electrophoresed. The gels were stained with either Mupid-Blue (left) or CBB (right) for the detection of DNA1 and His₆-RepE, respectively. The arrows labelled Complex and Free DNA indicate the positions of the His₆-RepE-DNA1 complex and free DNA1, respectively. Lane 1, solution used in crystallization; lane 2, crystallization mother liquor; lanes 3–4, washing solution; lane 5, dissolved crystals.

estimated to have a molecular weight of ~100 kDa by gel-filtration chromatography. In addition, it has been reported that the *repE* operator is preferentially bound by the dimeric form of RepE compared with the monomeric form (Ishiai *et al.*, 1994). Therefore, we concluded that these complexes are constituted of the RepE dimer and the DNA duplex.

The crystallization conditions for the four complexes were screened by the sparse-matrix method. Although crystals of three complexes (complexes with DNA1, DNA2 and DNA3) were obtained, only one crystal (for the complex with DNA1) was suitable for X-ray analysis. After the presence of both His₆-RepE and DNA1 in the obtained crystals had been confirmed (Fig. 2), the optimization of the crystallization conditions for His₆-RepE-DNA1 afforded a final condition of 100 mM HEPES-NaOH pH 7.2, 200 mM NaCl, 10% (w/v) PEG 4000 and 5 mM SmCl₃. Prismatic crystals grew in two weeks (Fig. 3; 0.12 × 0.04 × 0.02 mm). Prior to X-ray exposure, both a dehydration treatment (Schick & Jurnak, 1994; Heras & Martin,




Figure 3 Crystal of the His₆-RepE-DNA1 complex. The scale bar is 100 μm in length.




Figure 4 Pseudo-precession image of a His₆-RepE-DNA1 crystal representing the *h0l* zone of reciprocal space. The circle indicates a resolution of 4 Å. The picture was generated with *HKLVIEW* (Collaborative Computational Project, Number 4, 1994).

2005) and a process of crystal annealing were indispensable to improve the resolution limit of X-ray diffraction from His₆-RepE-DNA1 crystals (from approximately 8 to 3 Å resolution). Diffraction intensities were recorded to a maximum resolution of 3.14 Å using synchrotron radiation, but the diffraction patterns were anisotropic, showing diffraction to only 4.2 Å in the weakest direction (Fig. 4). The diffraction symmetry and the systematic absences of the diffraction data indicated that the His₆-RepE-DNA1 crystals belong to the monoclinic space group *P*2₁, with unit-cell parameters *a* = 60.73, *b* = 99.32, *c* = 95.00 Å, β = 108.55°. Assuming the presence of one His₆-RepE-DNA1 complex in the asymmetric unit, the *V*_M value (Matthews, 1968) was calculated to be 3.3 Å³ Da⁻¹, which corresponds to a solvent content of 63%. Although diffraction data were also collected at a wavelength of 1.6000 Å using one crystal, no significant peaks derived from Sm atoms were found either in the anomalous difference Patterson map or in the isomorphous difference Patterson map. Structure determination is currently in progress.

The authors would like to thank Drs M. Kawamoto and N. Shimizu and the other staff at beamline BL41XU, SPring-8, Japan for their kind assistance with X-ray diffraction data collection. This work was supported by Grants-in-Aid for Scientific Research (to CW and KM) and by a grant from the National Project on Protein Structural and Function Analysis from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to KM).

References

- Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.
- Harp, J. M., Timm, D. E. & Bunick, G. J. (1998). *Acta Cryst.* **D54**, 622–628.
- Heras, B. & Martin, J. L. (2005). *Acta Cryst.* **D61**, 1173–1180.
- Ishiai, M., Wada, C., Kawasaki, Y. & Yura, T. (1994). *Proc. Natl Acad. Sci. USA*, **91**, 3839–3843.
- Joachimiak, A. & Sigler, P. B. (1991). *Methods Enzymol.* **208**, 82–99.
- Kawasaki, Y., Matsunaga, F., Kano, Y., Yura, T. & Wada, C. (1996). *Mol. Gen. Genet.* **253**, 42–49.
- Kawasaki, Y., Wada, C. & Yura, T. (1990). *Mol. Gen. Genet.* **220**, 277–282.
- Kawasaki, Y., Wada, C. & Yura, T. (1991). *J. Bacteriol.* **173**, 1064–1072.
- Kawasaki, Y., Wada, C. & Yura, T. (1992). *J. Biol. Chem.* **267**, 11520–11524.
- Kline, B. C. (1985). *Plasmid*, **14**, 1–16.
- Kline, B. C., Sandhu, G. S., Eckloff, B. W. & Aleff, R. A. (1992). *J. Bacteriol.* **174**, 3004–3010.
- Komori, H., Matsunaga, F., Higuchi, Y., Ishiai, M., Wada, C. & Miki, K. (1999). *EMBO J.* **18**, 4597–4607.
- Komori, H., Sasai, N., Matsunaga, F., Wada, C. & Miki, K. (1999). *J. Biochem. (Tokyo)*, **125**, 24–26.
- Masson, L. & Ray, D. S. (1986). *Nucleic Acids Res.* **14**, 5693–5711.
- Masson, L. & Ray, D. S. (1988). *Nucleic Acids Res.* **16**, 413–424.
- Matsunaga, F., Ishiai, M., Kobayashi, G., Uga, H., Yura, T. & Wada, C. (1997). *J. Mol. Biol.* **274**, 27–38.
- Matsunaga, F., Kawasaki, Y., Ishiai, M., Nishikawa, K., Yura, T. & Wada, C. (1995). *J. Bacteriol.* **177**, 1994–2001.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Rokeach, L. A., Sogaard-Andersen, L. & Molin, S. (1985). *J. Bacteriol.* **164**, 1262–1270.
- Schick, B. & Jurnak, F. (1994). *Acta Cryst.* **D50**, 563–568.
- Tokino, T., Murotsu, T. & Matsubara, K. (1986). *Proc. Natl Acad. Sci. USA*, **83**, 4109–4113.