

## Minireview

# Host epigenetic modifications by oncogenic viruses

JM Flanagan<sup>\*,1</sup>

<sup>1</sup>CR-UK Viral Oncology Group, Wolfson Institute for Biomedical Research, Gower Street, University College London, London WC1E 6BT, UK

Epigenetic alterations represent an important step in the initiation and progression of most human cancers, but it is difficult to differentiate the early cancer causing alterations from later consequences. Oncogenic viruses can induce transformation via expression of only a small number of viral genes. Therefore, the mechanisms by which oncogenic viruses cause cancer may provide clues as to which epigenetic alterations are critical in early carcinogenesis.

*British Journal of Cancer* (2007) **96**, 183–188. doi:10.1038/sj.bjc.6603516 www.bjcancer.com

Published online 19 December 2006

© 2007 Cancer Research UK

**Keywords:** oncogenic virus; epigenetics; methylation; histone modifications; p300; CBP; KSHV; HHV8

Epigenetics describes the regulation of gene expression and genomic stability by heritable, but potentially reversible, changes in DNA methylation and chromatin structure. Significant progress has been made towards understanding the complementary functions between the various epigenetic regulators, including DNA methyltransferases, methylated DNA-binding proteins and potential DNA demethylases (such as cytosine deaminases). Histone modifying enzymes interact with other regulators of chromatin structure, such as ATP-dependent chromatin remodeling complexes and the opposing effects of the polycomb group (PcG) and trithorax group (TxG) genes (Esteller, 2006). The mediation of transcriptional repression by microRNAs is also gaining popularity as an epigenetic mechanism potentially disrupted in cancer (Esquela-Kerscher and Slack, 2006).

Transcriptionally active genes in a normal cell are marked by unmethylated promoter CpG islands, histone hyperacetylation and particular histone modifications, such as H3 lysine 4 (H3K4) di- and tri-methylation and H3K79 methylation. Transcriptionally repressed genes are marked by promoter CpG island methylation, histone hypoacetylation and H3K9 and H3K27 methylation. The epigenetic landscape of the genome is markedly different in cancers. A general genome-wide hypomethylation is associated with repetitive DNA sequences in all cancer cells, alongside specific promoter hypermethylation of tumour suppressor genes and hypomethylation of oncogenes (Esteller, 2006). In many cancer types, the most commonly hypermethylated genes are involved in DNA repair (hMLH1, MGMT), the cell cycle (p16INK4a, p15INK4b, p14ARF, SFN), apoptosis (DAPK), cell adherence (CDH1, CDH13) and detoxification (GSTP1) (Esteller, 2006). In addition to DNA methylation alterations, a number of critical histone modifications are considered common epigenetic alterations in cancer, including loss of H4K16 monoacetylation and H4K20 trimethylation and gains in H3K4 di- and tri-methylation, H3K79 methylation and H3 and H4 hyperacetylation (Feinberg *et al*, 2006).

As with many genetic changes in cancer, it is difficult to differentiate cause from consequence (Baylin and Ohm, 2006). Moreover, it is unclear whether genetic or epigenetic changes occur first and, if epigenetic, whether these involve DNA methylation or histone modifications. It is very difficult to study early cancer events in tumours or tumour-derived cell lines, because they represent end products of transformation and likely bear little resemblance to the cellular environment of early carcinogenic events (Baylin and Ohm, 2006). The increasingly popular idea that cancers originate from disrupted or 'cancer-primed' progenitor cells is supported by the hypothesis that epigenetic disruption of key genes occurs at the earliest stage of cancer development (Feinberg *et al*, 2006). Some of the most convincing evidence for epigenetic disruption of progenitor cells derive from the ubiquitous nature of genome-wide hypomethylation in almost all cancers, common hypermethylation of genes, such as p16INK4a, in many cancer types (and, occasionally, in surrounding normal tissues) and 'germline epimutations', such as heightened loss of imprinting (LOI) of insulin-like growth factor 2 in patients at risk for colorectal cancer (Feinberg *et al*, 2006). Such observations suggest the exciting prospect that an epigenetic model could elucidate the early carcinogenic events common to all cancers.

The identification and isolation of the subpopulation of cancer progenitor stem cells, most recently in glioblastoma, represents one potential strategy to elucidate the earliest carcinogenic changes (Singh *et al*, 2004). Identification of genetic and epigenetic differences between cancer progenitor cells and the bulk of other cancer cells may establish a causal link to their phenotypic differences. To date, however, the identification of cancer progenitor stem cells has met with limited success for most cancer types.

Another strategy to investigate the early events in cancer is to manipulate and dissect the process by which oncogenic viruses infect cells and initiate transformation *in vitro* and *in vivo*. Viruses are small and tend to encode only the most important genes required for their survival, for example genes for viral persistence and replication, immune regulation and key genes required to trigger transformation. Oncogenic viruses include retroviruses, such as human T-cell lymphotropic viruses 1 and 2 (HTLV1/

\*Correspondence: Dr JM Flanagan; E-mail: j.flanagan@ucl.ac.uk

Received 14 September 2006; revised 6 November 2006; accepted 9 November 2006; published online 19 December 2006

HTLV2), the RNA flavivirus, hepatitis C virus (HCV) and DNA viruses, such as human papilloma virus (HPV), hepatitis B viruses (HBV), polyomaviruses (JC virus (JCV), BK virus (BKV) and simian virus 40 (SV40)) and the  $\gamma$ herpesviruses, Epstein–Barr virus (EBV) and Kaposi sarcoma-associated herpesvirus (KSHV). *In vitro*, oncogenic viruses can induce cellular transformation, producing cells capable of uncontrolled proliferation and often tumorigenic in athymic mice. *In vivo*, viral oncogenesis also requires additional cellular alterations that enable the transformed cell to escape host responses, such as the immune system and apoptosis. Furthermore, viruses exploit epigenetic mechanisms, such as DNA methylation of viral genes, that would otherwise illicit an immune response or the increased expression of DNA methyltransferases that results in increased methylation of cellular genes (Tsai *et al*, 2002; Tao and Robertson, 2003). The frequency of hypermethylation of the cell cycle regulating gene p16INK4a in HBV-, EBV-, KSHV- and HPV-related tumours suggests the importance of its inactivation in virally induced cancers (Platt *et al*, 2002; Li *et al*, 2005). This Minireview describes the interactions between viral proteins and epigenetic regulators that occur in the host cell to mediate the methylation and histone alterations required by the virus. The potential involvement of virally induced epigenetic changes in initiating carcinogenesis is discussed alongside the viral genes potentially responsible. If virally induced epigenetic defects mimic the first alterations in cancer, then these would obviously represent the most effective targets for cancer drug design.

## HOST EPIGENETIC CHANGES OWING TO VIRUSES AND VIRUS-ASSOCIATED CANCERS

The cause or consequence conundrum in cancer epigenetics is equally relevant to the epigenetics of viral infection. It is difficult to

differentiate an epigenetic change that is directly due to viral infection, due to the host antiviral response or due to a subsequent downstream effect of the transformation process. Do these viruses happen to infect cancer progenitor cells that are already committed to cancer development and are, thus, just along for the ride? Or are cancer progenitor cells more susceptible to viral infection? Important cancer causing changes may be separated from their consequences via the identification of direct interactions between viral proteins and epigenetic regulators (Table 1).

### KSHV as an example

Kaposi sarcoma-associated herpesvirus is an oncogenic  $\gamma$ -herpesvirus identified as the causative agent of the endothelial tumour Kaposi–sarcoma (KS) and associated with the lymphoproliferative disorders, multicentric Castleman's disease (MCD) and primary effusion lymphoma (PEL) (reviewed in Jenner and Boshoff, 2002). Kaposi sarcoma-associated herpesvirus has both a latent and lytic phase of its life cycle with a more restricted set of genes expressed during latency and in the majority of tumour cells. Particularly in MCD and to a lesser extent in KS and PEL, a small percentage of cells still express lytic genes and maintain lytic replication, which may serve to further spread the virus and maintain important paracrine growth effects of lytic genes such as vGPCR (Jenner and Boshoff, 2002). Kaposi sarcoma-associated herpesvirus contains over 80 open reading frames (ORFs), some of which are pirated from the host genome in order to control proliferation, immune regulation and cell signalling (Jenner and Boshoff, 2002). The virus also employs epigenetic mechanisms, for example chromatin remodelling and demethylation of the lytic switch gene Rta (ORF50) promoter, to control its entry into the lytic phase and hyperacetylation of the latent replication origin to control latent cycle replication (Lu *et al*, 2006). Given this use of cellular epigenetic machinery to regulate its own genome during latency

**Table 1** Epigenetic interactions of oncogenic viral proteins

Virus	Human Neoplasms	Viral Proteins	Epigenetic interactions	References
<i>DNA viruses</i>				
KSHV	KS	LANA	<ul style="list-style-type: none"> <li>● Activates DNMT3a</li> <li>● Interacts with SUV39H1, MeCP2, mSin3, HP1</li> </ul>	(Li <i>et al</i> , 2000; Shamay <i>et al</i> , 2006)
	PEL		<ul style="list-style-type: none"> <li>● Binds and inhibits p300/CBP HAT activity.</li> </ul>	
EBV	MCD	vIRFs	<ul style="list-style-type: none"> <li>● Activates DNMTs 1,3a,3b</li> </ul>	(Wang <i>et al</i> , 2000; Tsai <i>et al</i> , 2002; Knight <i>et al</i> , 2003)
	BL, NPC, HD	LMP1		
HPV	Gastric Cancer	EBNA2	<ul style="list-style-type: none"> <li>● Binds p300 to activate transcription</li> </ul>	(Patel <i>et al</i> , 1999; Longworth and Laimins, 2004; Lin <i>et al</i> , 2005; Burgers <i>et al</i> , 2006)
	PTLD	EBNA3c	<ul style="list-style-type: none"> <li>● Interacts with HDACs</li> <li>● DNMT3b protein is increased by HPV in females only</li> </ul>	
HBV	Papilloma, carcinomas	?		(Lee <i>et al</i> , 2005)
		E7	<ul style="list-style-type: none"> <li>● Binds DNMT1 to increase DNA methyltransferase activity</li> <li>● Binds HDACs and Nurd ATP-dependent remodelling complex</li> </ul>	
SV40	HCC	E6	<ul style="list-style-type: none"> <li>● Binds and inhibits p300/CBP HAT activity</li> </ul>	(Soejima <i>et al</i> , 2003)
	? Osteosarcoma	HBx	<ul style="list-style-type: none"> <li>● Activates DNMT1</li> </ul>	
BKV	? Mesothelioma	Large T-Ag	<ul style="list-style-type: none"> <li>● Activates DNMT3b</li> </ul>	(McCabe <i>et al</i> , 2006)
	? Brain tumours		<ul style="list-style-type: none"> <li>● May induce methylator phenotype in CRC</li> </ul>	
JCV	? Gliomas	T-Ag		(Goel <i>et al</i> , 2006)
	? Medulloblastoma			
Adenovirus	? CRC			(Ghosh and Harter, 2003; Tumell <i>et al</i> , 2005; Burgers <i>et al</i> , 2006; McCabe <i>et al</i> , 2006)
	None	E1A	<ul style="list-style-type: none"> <li>● Binds DNMT1 to increase DNA methyltransferase activity</li> <li>● Binds E2F promoters to demethylate H3K9</li> <li>● Binds and perturbs p300/CBP HAT activity</li> </ul>	
<i>RNA viruses</i>				
HTLV1/2	ATL	Tax	<ul style="list-style-type: none"> <li>● Binds with p300/CBP to repress transcription</li> </ul>	(Kwok <i>et al</i> , 1996)


ATL = adult T cell leukemia; BKV = BK virus; BL = Burkitts Lymphoma; CRC = colorectal cancer; EBV = Epstein–Barr Virus; HBV = hepatitis B virus; HCC = hepatocellular carcinoma; HD = Hodgkins Disease; HPV = human papilloma virus; HTLV = human lymphotropic virus; JCV = JC virus; KS = Kaposi Sarcoma; KSHV = Kaposi Sarcoma-associated herpesvirus; MCD = multicentric Castleman's Disease; NPC = Nasopharyngeal Carcinoma; PEL = primary effusion lymphoma; PTLD = posttransplant lymphoproliferative disease; SV40 = simian virus 40.

and lytic replication, the virus may also interact with the host epigenome as a deliberate mechanism to alter the cellular environment. There exists clear evidence that KSHV reprogrammes cellular gene expression in both lymphatic and blood vessel endothelial cells, although the mechanism of reprogramming remains unclear (Wang *et al*, 2004). One possibility is that the virus encodes proteins that are involved in DNA methylation, histone modifications, chromatin remodelling or microRNA processing to ensure complete control of cellular gene expression (Figure 1).

Regarding cellular methylation by KSHV, only *p16INK4a* hypermethylation has been identified in the majority of KSHV-infected PEL cell lines and, potentially, in primary PEL samples (Platt *et al*, 2002). This is not surprising, given that *p16INK4a* is one of the most commonly hypermethylated genes in nearly all cancers. This suggests that, in B cells, at least, KSHV is able to invoke DNA methyltransferase activity and may inactivate other cellular genes via methylation. A direct link between LANA and the *de novo* methyltransferase DNMT3a has been proposed in a study, that reveals that LANA recruits DNMT3a to the chromatin and targets repression of approximately 80 cellular genes, some of which are known targets of epigenetic inactivation in various cancers (Shamay *et al*, 2006). In addition, LANA interacts with the DNA methyl binding protein MeCP2, the mSin3 transcriptional

repression complex and the histone methyltransferase SUV39H1, thus enabling numerous roles in epigenetic gene regulation (see Shamay *et al* (2006) for individual references). Regarding histone modification of cellular genes, a number of studies have identified a direct link between the KSHV encoded interferon regulatory factors (viral interferon regulatory factor (IRF) 1, 2 and 3) and the histone acetyltransferase complex p300/CBP (Li *et al*, 2000). The binding of cellular IRF3 to CBP is thought to be a requirement for transcriptional activation of the antiviral cytokine interferon- $\beta$  (IFN- $\beta$ ). Accordingly, one may assume that KSHV would evolve a mechanism to disrupt IFN- $\beta$  activation, which it has by direct interaction between vIRF1 and either p300 or CBP. In effect, these interactions inhibit histone acetyltransferase activity and lead to histone hypoacetylation, altered chromatin structure and reduced expression of cytokine genes and, presumably, other genes activated by p300/CBP (Lin *et al*, 2001). Increased DNA methylation and decreased histone acetylation are both associated with gene silencing, which could account for the many genes down-regulated after KSHV infection.

The importance of epigenetic regulation for gene expression and the reprogramming of cellular gene expression during viral transformation suggests that other viral genes (latent or lytic) may also be involved in epigenetic control of gene expression. Kaposi sarcoma-associated herpesvirus is relatively unique in that



**Figure 1** Viral control of the host epigenome. Epigenetic control of gene expression occurs at four different levels starting with chromatin packaging into higher order chromatin structures controlled by ATP-dependent chromatin remodelling complexes (ATP-DRs), PcG and TxG genes. DNA is wrapped around nucleosomes that are assembled from dimers of the histones H2A, H2B, H3 and H4, all of which contain tails that can be either acetylated (HATs) and demethylated (HDMTs) during active transcription or deacetylated (HDACs) and highly methylated (HMTs) during repressed transcription. At the nucleotide level, epigenetic control via DNA methylation is mediated by the DNA methylating enzymes (DNMTs), methyl-DNA-binding proteins (e.g. MeCBPs) and cytosine deaminases that could act as demethylating enzymes. Finally, transcriptional repression is also mediated by microRNAs. Oncogenic viruses target DNA methyltransferase activity and p300/CBP histone acetyltransferase activity, but may also target other epigenetic mechanisms to induce carcinogenesis.

it contains numerous genes with cellular homologues and, by investigating the epigenetic functions of the cellular genes, one may also infer interactions or functions related to epigenetic control by the viral genes. Using this strategy, a number of viral genes appear likely epigenetic regulators, including viral interleukin-6 (IL-6), viral G-protein coupled receptor and the 12 KSHV microRNAs (Table 2). Speculation on viral gene function based on the functions of cellular homologues can provide interesting leads and has proved correct for the viral IRFs, but there exist cases wherein the viral gene does not function in the manner of its cellular homologue. For instance, vIL-6 does not bind to the cellular IL-6 receptor. Rather, it binds only to the gp130 receptor. At present, the potential cellular or viral targets of the KSHV microRNAs exist only as *in silico* predictions, among which are the cytokine signalling regulator *SOCS3*, DNA repair gene *RAD21*, the tumour necrosis factor- $\alpha$  (*TNF- $\alpha$* ) and the kinase *MAP3K8*, which is also involved in production of *TNF- $\alpha$*  (Cai *et al.*, 2005). Whether or not there are cellular homologues of the viral microRNAs has yet to be explored. Accordingly, whether or not the viral genes described here are actually involved in epigenetic regulation must be investigated experimentally.

### Other oncogenic viruses

Increased DNA methylation activity and decreased histone acetylation activity are not unique properties of KSHV, as other oncogenic viruses also share these properties (Table 1). The related  $\gamma$ -herpesvirus, EBV, activates DNA methyltransferase activity by increasing the expression of the maintenance methyltransferase DNMT1 and both *de novo* methyltransferases, DNMT3a and DNMT3b (Tsai *et al.*, 2002). The LMP 1 protein is one of the key oncogenic viral proteins of EBV and it is not surprising that this is the protein that alters the DNA methyltransferase activity. The hepatitis B virus HBx protein, BK polyoma virus T-antigen and the adenovirus oncogene E1A all increase the activity of DNMT1 alone (Lee *et al.*, 2005; McCabe *et al.*, 2006). In both EBV and HBV-related epithelial cancers, this results in increased methylation and suppression of E-cadherin, thereby increasing cell migration (Tsai *et al.*, 2002; Lee *et al.*, 2005). The HPV acts similar to KSHV in that it appears to increase the *de novo* methyltransferase, DNMT3b protein, in nonsmoking lung cancers, curiously only in females (Lin *et al.*, 2005). Although no HPV protein has been identified as the culprit for this increase in DNMT3b, a recent study has now shown that HPV E7-protein increases DNA methyltransferase enzymatic activity by directly interacting with DNMT1 (Burgers *et al.*, 2006). The link between JCV and increased DNA methylation is more speculative with an association between the JCV T-antigen expression and the methylator phenotype in colorectal cancer (Goel *et al.*, 2006). The simian SV40 virus increases DNA

methylation by upregulation of DNMT3b via the large T antigen (Soejima *et al.*, 2003). The association between SV40 and human cancers is still contentious; however, this does provide an interesting insight into the possibility that activation of methylation may be conserved among other mammalian viruses. Although methylation of p16INK4a occurs in hepatocellular carcinoma associated with the HCV, no direct interactions have yet been identified.

The ability to alter histone modifications and chromatin structure is also common to many oncogenic viruses including EBV, HPV, adenovirus and HTLV1. The EBV nuclear antigens EBNA 2 and 3c alter histone acetylation by interacting with the p300/CBP complex or with histone deacetylase (HDAC), respectively (Wang *et al.*, 2000; Knight *et al.*, 2003). Human papilloma virus oncoprotein E6 binds and inhibits the histone acetyltransferase activity of the p300/CBP complex similar to KSHV (Patel *et al.*, 1999). This transcriptional repressive activity by HPV is further supported by E7 protein interaction with HDAC1 and the Nurd ATP-dependent chromatin remodelling complex that are also involved in repression (Longworth and Laimins, 2004). The adenovirus transforming protein E1A also interacts with the p300/CBP transcriptional complex to perturb or alter the normal functioning during the cell cycle (Turnell *et al.*, 2005). This has been described as one of the key events in the induction of E1A-induced cellular transformation. Finally, the retrovirus HTLV1 Tax protein also interacts with the p300/CBP complex to mediate transcriptional repression (Kwok *et al.*, 1996).

It is interesting to note that the EBV proteins that interact with epigenetic regulators (EBNA2, EBNA3c and LMP1) are all latent genes and are not typically expressed in Burkitt's lymphoma, gastric cancer and most nasopharyngeal carcinomas. This may lead to the conclusion that the role of virus-induced host epigenetic changes may be limited in these particular cancers. However, it cannot be ruled out that the initial cancer precursor cells may have harboured the virus in a latent phase in which the host epigenome could be altered. Epigenetic fingerprints such as DNA methylation and histone modifications are mitotically heritable and, thus, even if the progeny cancer cell no longer expresses these latent genes, the epigenetic history of the cell may remain. It is also interesting to observe that the proteins that interact with epigenetic mechanisms, LMP1, LANA, E6 and E7, large T antigen, E1A and Tax, are often described as the viral 'oncoproteins' as they are often either oncogenic on their own or essential components of viral transformation. This typically leads to the conclusion that the functions of these proteins, in this case the epigenetic functions, are indeed essential for viral-induced transformation.

Returning to the hypothesis that viral-induced epigenetic changes might mimic the early events in cancers, we can observe from these studies described that disruption or alteration of

**Table 2** Potential epigenetic functions of selected KSHV proteins and microRNAs

KSHV Gene	Cellular Homolog	Activities	Epigenetic functions of cellular homologues	References
K2 (vIL6)	IL6	B-cell growth factor, angiogenesis, hematopoiesis	<ul style="list-style-type: none"> <li>● IL6 regulates expression of DNMT1</li> <li>● IL6 supports the maintenance of p53 promoter methylation</li> <li>● IL6 supports methylation of the RAD23B DNA repair gene</li> <li>● IL6 induces polycomb group gene EZH2</li> </ul>	(Croonquist and Van Ness, 2005; Hodge <i>et al.</i> , 2005; Peng <i>et al.</i> , 2005)
ORF74 (vGPCR)	GPCR/ IL-8R	Angiogenesis; transformation	<ul style="list-style-type: none"> <li>● GPCR signaling regulates histone acetylation and gene transcription</li> </ul>	(Kang <i>et al.</i> , 2005)
KSHV miRNAs (mirK1-mirK12)	None yet identified	Transcriptional repression. Some predicted cellular targets include TNF, SOCS3, RAD21 and MAP3K8	<ul style="list-style-type: none"> <li>● miRNAs (Oncomirs) such as the let-7 family regulate expression of oncogenes</li> </ul>	(Cai <i>et al.</i> , 2005; Esquela-Kerscher and Slack, 2006)

p300/CBP histone acetyltransferase activity is common to many oncogenic viruses, which suggests that it may be one of the critical early events in viral-induced carcinogenesis. Further evidence for an early involvement of p300/CBP in nonviral cancers can be observed in the increased predisposition to childhood malignancies in Rubinstein–Taybi syndrome, which is characterised by germline mutation of CBP as well as numerous somatic mutations in colorectal, breast and gastric carcinomas (Iyer *et al*, 2004). This suggests that abrogation or perturbation of the histone acetyltransferase activity of p300/CBP may also be one of the critical early events in all cancers. Similarly, an increase in DNA methyltransferase activity might also be a crucial early event. Disruption of either the maintenance methyltransferase, DNMT1, or the *de novo* methyltransferases, DNMT3a or 3b, suggests that each oncogenic virus may have evolved different routes to the same goals (methylation of key tumour suppressor genes) or perhaps that different viruses require inactivation of different genes via different methyltransferases. Interestingly, ranid herpesvirus 1 (RaHV-1), which is the aetiological agent of kidney cancers in the North American leopard frog *Rana pipiens* encodes its own DNA (cytosine-5) methyltransferase, further suggesting that alteration of methylation in the host genome is a highly conserved function in oncogenic viruses (Davison *et al*, 1999). Again, in nonviral cancers, this is also a common finding with an overall increase in DNA methyltransferase activity in many tumours compared with normal tissues (Esteller, 2006). Both of these activities, disruption of histone acetylation and increased DNA methylation, are related to repression of transcription, which provides support for the idea that the critical early carcinogenic events begin with the inactivation of tumour-suppressor genes (Feinberg *et al*, 2006). A more complete understanding of the interactions between viral proteins and epigenetic regulators may be a very important step in understanding the earliest alterations in cancer. In addition, identification of the particular changes induced by viruses might provide novel, and more effective, targets for cancer therapies.

## CONCLUSIONS

It is apparent from current knowledge that KSHV and other oncogenic viruses increase activity of DNA methyltransferases and can decrease p300/CBP-mediated histone acetylation, which are both likely requirements for the inactivation of tumour-suppressor genes. The fact that both of these events also occur in many other

## REFERENCES

- Baylin SB, Ohm JE (2006) Epigenetic gene silencing in cancer – a mechanism for early oncogenic pathway addiction? *Nat Rev Cancer* **6**: 107–116
- Burgers WA, Blanchon L, Pradhan S, de Launoit Y, Kouzarides T, Fuks F (2006) Viral oncoproteins target the DNA methyltransferases. *Oncogene* Sept 18 [Epub ahead of print] PMID: 16983344
- Cai X, Lu S, Zhang Z, Gonzalez CM, Damania B, Cullen BR (2005) Kaposi's sarcoma-associated herpesvirus expresses an array of viral microRNAs in latently infected cells. *Proc Natl Acad Sci USA* **102**: 5570–5575
- Croonquist PA, Van Ness B (2005) The polycomb group protein enhancer of zeste homolog 2 (EZH 2) is an oncogene that influences myeloma cell growth and the mutant ras phenotype. *Oncogene* **24**: 6269–6280
- Davison AJ, Sauerbier W, Dolan A, Addison C, McKinnell RG (1999) Genomic studies of the Lucké tumor herpesvirus (RaHV-1). *J Cancer Res Clin Oncol* **125**: 232–238
- Esquela-Kerscher A, Slack FJ (2006) Oncomirs – microRNAs with a role in cancer. *Nat Rev Cancer* **6**: 259–269
- Esteller M (2006) Epigenetics provides a new generation of oncogenes and tumour-suppressor genes. *Br J Cancer* **94**: 179–183
- Feinberg AP, Ohlsson R, Henikoff S (2006) The epigenetic progenitor origin of human cancer. *Nat Rev Genet* **7**: 21–33

nonviral cancers suggests that these may indeed be some of the earliest alterations in carcinogenesis. Identification of cancer progenitor cells with increased DNA methyltransferase activity and abrogated p300/CBP histone acetyltransferase activity would further support this hypothesis. Furthermore, there exists the potential for other viral genes in epigenetic control, as proposed in this review for KSHV, and thus there is scope for identification of novel epigenetic alterations as early carcinogenic events.

There is gathering momentum for the use of epigenetic related therapies, such as HDAC inhibitors, for example suberoylanilide hydroxamic acid (SAHA), which is currently in Phase III trials and has shown reasonable efficacy in haematological and solid tumours. At present, five trials are investigating the use of HDAC inhibitor valproic acid (VPA) in Kaposi sarcoma, EBV-related nasopharyngeal carcinoma and non-Hodgkin's lymphoma, as well as in HIV-infected patients (<http://clinicaltrials.gov/>). Some circumspection is required, however, because histone deacetylase inhibitors, including valproate, may induce lytic replication, as has been shown for KSHV and EBV (Klass *et al*, 2005; Feng and Kenney, 2006; Lu *et al*, 2006). For an effective drug, it would be preferable to induce lytic replication and induce apoptosis without increasing the viral load. A recent study has shown that valproate can induce entry into the lytic cycle in KSHV-infected PEL cells and can significantly increase production of the virus, leading to cell death by apoptosis (Klass *et al*, 2005). With the addition of ganciclovir and PFA, viral DNA replication was blocked without blocking the lytic cascade and apoptosis, which advocates the use of HDAC inhibitors in combination therapies rather than as single agents in virus-associated cancers. Other HDAC inhibitors (depsipeptide, SAHA, MS-275 and trichostatin A) have also been tested in PEL cells with minimal (3–14%) reactivation (Niedermeier *et al*, 2006). Given our present understanding of the epigenetic control required by oncogenic viruses in the transformation process, the use of HDAC inhibitors, or other epigenetically targeted therapies, is warranted in virus associated tumours.

## ACKNOWLEDGEMENTS

I would like to apologise to the authors of many of the important original studies that have not been cited owing to space limitations. I would like to thank Chris Boshoff, Nessa Carey and Matthew Trotter for critical review of this manuscript. JMF is a Cellcentric funded Research Fellow.

- Feng WH, Kenney SC (2006) Valproic acid enhances the efficacy of chemotherapy in EBV-positive tumors by increasing lytic viral gene expression. *Cancer Res* **66**: 8762–8769
- Ghosh MK, Harter ML (2003) A viral mechanism for remodeling chromatin structure in G0 cells. *Mol Cell* **12**: 255–260
- Goel A, Li MS, Nagasaka T, Shin SK, Fuerst F, Ricciardiello L, Wasserman L, Boland CR (2006) Association of JC virus T-antigen expression with the methylator phenotype in sporadic colorectal cancers. *Gastroenterology* **130**: 1950–1961
- Hodge DR, Peng B, Cherry JC, Hurt EM, Fox SD, Kelley JA, Munroe DJ, Farrar WL (2005) Interleukin 6 supports the maintenance of p53 tumor suppressor gene promoter methylation. *Cancer Res* **65**: 4673–4682
- Iyer NG, Ozdag H, Caldas C (2004) p300/CBP and cancer. *Oncogene* **23**: 4225–4231
- Jenner RG, Boshoff C (2002) The molecular pathology of Kaposi's sarcoma-associated herpesvirus. *Biochim Biophys Acta* **1602**: 1–22
- Kang J, Shi Y, Xiang B, Qu B, Su W, Zhu M, Zhang M, Bao G, Wang F, Zhang X, Yang R, Fan F, Chen X, Pei G, Ma L (2005) A nuclear function of beta-arrestin1 in GPCR signaling: regulation of histone acetylation and gene transcription. *Cell* **123**: 833–847

- Klass CM, Krug LT, Pozharskaya VP, Offermann MK (2005) The targeting of primary effusion lymphoma cells for apoptosis by inducing lytic replication of human herpesvirus 8 while blocking virus production. *Blood* **105**: 4028–4034
- Knight JS, Lan K, Subramanian C, Robertson ES (2003) Epstein–Barr virus nuclear antigen 3C recruits histone deacetylase activity and associates with the corepressors mSin3A and NCoR in human B-cell lines. *J Virol* **77**: 4261–4272
- Kwok RP, Laurance ME, Lundblad JR, Goldman PS, Shih H, Connor LM, Marriott SJ, Goodman RH (1996) Control of cAMP-regulated enhancers by the viral transactivator Tax through CREB and the co-activator CBP. *Nature* **380**: 642–646
- Lee JO, Kwun HJ, Jung JK, Choi KH, Min do S, Jang KL (2005) Hepatitis B virus X protein represses E-cadherin expression via activation of DNA methyltransferase 1. *Oncogene* **24**: 6617–6625
- Li HP, Leu YW, Chang YS (2005) Epigenetic changes in virus-associated human cancers. *Cell Res* **15**: 262–271
- Li M, Damania B, Alvarez X, Ogryzko V, Ozato K, Jung JU (2000) Inhibition of p300 histone acetyltransferase by viral interferon regulatory factor. *Mol Cell Biol* **20**: 8254–8263
- Lin R, Genin P, Mamane Y, Sgarbanti M, Battistini A, Harrington Jr WJ, Barber GN, Hiscott J (2001) HHV-8 encoded vIRF-1 represses the interferon antiviral response by blocking IRF-3 recruitment of the CBP/p300 coactivators. *Oncogene* **20**: 800–811
- Lin TS, Lee H, Chen RA, Ho ML, Lin CY, Chen YH, Tsai YY, Chou MC, Cheng YW (2005) An association of DNMT3b protein expression with P16INK4a promoter hypermethylation in non-smoking female lung cancer with human papillomavirus infection. *Cancer Lett* **226**: 77–84
- Longworth MS, Laimins LA (2004) The binding of histone deacetylases and the integrity of zinc finger-like motifs of the E7 protein are essential for the life cycle of human papillomavirus type 31. *J Virol* **78**: 3533–3541
- Lu F, Day L, Gao SJ, Lieberman PM (2006) Acetylation of the latency-associated nuclear antigen regulates repression of Kaposi's sarcoma-associated herpesvirus lytic transcription. *J Virol* **80**: 5273–5282
- McCabe MT, Low JA, Imperiale MJ, Day ML (2006) Human polyomavirus BKV transcriptionally activates DNA methyltransferase 1 through the pRb/E2F pathway. *Oncogene* **25**: 2727–2735
- Niedermeier A, Talanin N, Chung EJ, Sells RE, Borris DL, Orenstein JM, Trepel JB, Blauvelt A (2006) Histone deacetylase inhibitors induce apoptosis with minimal viral reactivation in cells infected with Kaposi's sarcoma-associated herpesvirus. *J Invest Dermatol* **126**: 2516–2524
- Patel D, Huang SM, Baglia LA, McCance DJ (1999) The E6 protein of human papillomavirus type 16 binds to and inhibits co-activation by CBP and p300. *EMBO J* **18**: 5061–5072
- Peng B, Hodge DR, Thomas SB, Cherry JM, Munroe DJ, Pompeia C, Xiao W, Farrar WL (2005) Epigenetic silencing of the human nucleotide excision repair gene, hHR23B, in interleukin-6-responsive multiple myeloma KAS-6/1 cells. *J Biol Chem* **280**: 4182–4187
- Platt G, Carbone A, Mittnacht S (2002) p16INK4a loss and sensitivity in KSHV associated primary effusion lymphoma. *Oncogene* **21**: 1823–1831
- Shamay M, Krithivas A, Zhang J, Hayward S (2006) Recruitment of the de novo DNA methyltransferase Dnmt3a by Kaposi's sarcoma-associated herpesvirus LANA. *Proc Natl Acad Sci USA* **103**: 14554–14559
- Singh SK, Clarke ID, Hide T, Dirks PB (2004) Cancer stem cells in nervous system tumors. *Oncogene* **23**: 7267–7273
- Soejima K, Fang W, Rollins BJ (2003) DNA methyltransferase 3b contributes to oncogenic transformation induced by SV40T antigen and activated Ras. *Oncogene* **22**: 4723–4733
- Tao Q, Robertson KD (2003) Stealth technology: how Epstein–Barr virus utilizes DNA methylation to cloak itself from immune detection. *Clin Immunol* **109**: 53–63
- Tsai CN, Tsai CL, Tse KP, Chang HY, Chang YS (2002) The Epstein–Barr virus oncogene product, latent membrane protein 1, induces the downregulation of E-cadherin gene expression via activation of DNA methyltransferases. *Proc Natl Acad Sci USA* **99**: 10084–10089
- Turnell AS, Stewart GS, Grand RJ, Rookes SM, Martin A, Yamano H, Elledge SJ, Gallimore PH (2005) The APC/C and CBP/p300 cooperate to regulate transcription and cell-cycle progression. *Nature* **438**: 690–695
- Wang HW, Trotter MW, Lagos D, Bourbouliou D, Henderson S, Makinen T, Elliman S, Flanagan AM, Alitalo K, Boshoff C (2004) Kaposi sarcoma herpesvirus-induced cellular reprogramming contributes to the lymphatic endothelial gene expression in Kaposi's sarcoma. *Nat Genet* **36**: 687–693
- Wang L, Grossman SR, Kieff E (2000) Epstein–Barr virus nuclear protein 2 interacts with p300, CBP, and PCAF histone acetyltransferases in activation of the LMP1 promoter. *Proc Natl Acad Sci USA* **97**: 430–435