

# 11q13 amplification status and human papillomavirus in relation to p16 expression defines two distinct etiologies of head and neck tumours

CCR Ragin<sup>1,2,3</sup>, E Taioli<sup>2,3</sup>, JL Weissfeld<sup>2,3</sup>, JS White<sup>1,2,3</sup>, KM Rossie<sup>4</sup>, F Modugno<sup>2,3</sup> and SM Gollin<sup>\*,1,3</sup>

<sup>1</sup>Department of Human Genetics, University of Pittsburgh Graduate School of Public Health, 130 DeSoto Street, Pittsburgh, PA 15261, USA;

<sup>2</sup>Department of Epidemiology, University of Pittsburgh Graduate School of Public Health, Pittsburgh, PA 15261, USA; <sup>3</sup>The University of Pittsburgh Cancer Institute, Pittsburgh, PA 15232, USA; <sup>4</sup>Department of Oral Medicine and Pathology, School of Dental Medicine, Pittsburgh, PA 15232, USA

Two distinct etiologies of head and neck squamous cell carcinoma (HNSCC) have been proposed, DNA damage owing to tobacco and alcohol exposure and human papillomavirus (HPV) oncogene-mediated transformation. Common genetic alterations in HNSCC include *TP53* mutations, 11q13 amplification (amp) and *CDKN2A/p16* mutations or promoter methylation. However, in HPV + HNSCC it is frequent to observe wild-type *TP53* and expression of p16. The relationship of this unusual pattern with 11q13 amp has not been tested. In a retrospective study on 125 HNSCC patients, only 17% (five out of 30) of HPV + vs 44% (39 out of 89) of HPV – tumours expressed 11q13 amp (adjusted odds ratio (OR) = 0.2, 95% confidence interval (CI) = 0.1–0.6). A subpopulation of tumours ( $n = 69$ ) were classified according to the three molecular markers, *TP53*, p16 and 11q13 amp. In addition to wild-type *TP53*, and p16 expression, HPV + tumours were more likely not to be amplified at 11q13 (OR = 6.5, 95% CI = 1.8–23.9). As HPV + HNSCC lack the genetic alterations which are common in other tumours, we hypothesise that HPV infection may represent an early event in the HNSCC carcinogenic process, thus suggesting a distinct molecular pathway.

British Journal of Cancer (2006) 95, 1432–1438. doi:10.1038/sj.bjc.6603394 www.bjancer.com

Published online 26 September 2006

© 2006 Cancer Research UK

**Keywords:** HPV; 11q13; gene amplification; *CDKN2A/p16*; head and neck cancer; *TP53*

Although the majority of cases of head and neck squamous cell carcinoma (HNSCC) are attributed to tobacco and alcohol consumption (Blot *et al*, 1988; Merletti *et al*, 1989; Franceschi *et al*, 1990; Franceschi *et al*, 1999), morphologic signs of human papillomavirus (HPV) infection and detection of the viral antigens in oral squamous cell carcinomas were first reported in 1983 (Syrjanen *et al*, 1983). Increasing evidence from large case–control studies of HPV prevalence in head and neck tumours suggests that this virus may be an additional independent risk factor for a subset (~25%) of HNSCC (Smith *et al*, 1998; Gillison *et al*, 2000; Herrero *et al*, 2003).

Head and neck squamous cell carcinoma frequently carries numerous genetic alterations, such as mutations, chromosomal losses, gains and amplifications (amp) (Gollin, 2001). Common genetic alterations in HNSCC involve *TP53*, *CDKN2A/p16* and chromosomal band 11q13. Amplification of 11q13 is among the most common sites of gene amplification, observed in a number of cancers (Schraml *et al*, 1999), and in approximately 45% of HNSCC

(Lese *et al*, 1995; Schuurin, 1995; Gollin, 2001). A number of putative oncogenes map to this region; among them, *CCND1* (cyclin D1) is thought to play a role in tumorigenesis. Cyclin D1 is an important cell cycle protein that binds cyclin-dependent kinase (CDK) 4 or 6 to phosphorylate and inactivate the RB1 protein thus driving the cell from G1 into S-phase. Therefore, over-expression of cyclin D1 may lead to deregulate cell cycling from G1 to S-phase and may result in a selective growth advantage for the affected cell.

*CDKN2A/p16* opposes cyclin D1 function by negatively regulating RB1 phosphorylation and a genetic (loss of heterozygosity (LOH)) or epigenetic (promoter methylation) alteration in this gene is frequently observed, resulting in a lack of p16 expression in most HNSCC (Weber *et al*, 2003). *TP53*, also is a frequently mutated tumour-suppressor gene in a number of cancers, including about 45% of HNSCC (Sidransky, 1995).

In contrast to most HNSCC, HPV-positive tumours are more likely to have wild-type *TP53*. Human papillomavirus-infected cells accomplish G<sub>1</sub> to S-phase transition through the targeted degradation of pRb by the viral E7 oncoprotein and consequently, *CDKN2A/p16* is overexpressed. Correlations between genetic alterations at *TP53*, 11q13 and p16 expression with HPV have not been evaluated. We report a retrospective study examining the association of HPV with these molecular risk factors (*TP53* mutation, p16 protein expression and 11q13 amp) in patients with first primary head and neck tumours.

\*Correspondence: Dr SM Gollin; Department of Human Genetics, University of Pittsburgh Graduate School of Public Health, 130 Desoto Street, Pittsburgh, PA 15261, USA;

E-mail: sgollin@hgen.pitt.edu

Received 2 June 2006; accepted 29 August 2006; published online 26 September 2006



statistical software package for analysis. Statistical analyses were performed using the Intercooled STATA (version 8.2) software (StataCorp. LP, College Station TX, USA). Factors associated with HPV status were selected based on cross-tabulations and univariate logistic regression analyses. Cross-tabulations were analysed using the  $\chi^2$  test or Fisher's exact test, where appropriate. Logistic regression was used to perform the analyses of associations between demographic, clinical and pathologic variables with HPV status. Crude odds ratio (OR) were first calculated. Univariate analyses for the association of selected variables with 11q13 amp status were performed in the subset of HPV-negative subjects to identify possible confounding factors of association. Potential confounders were identified based on *P*-values  $\leq 0.1$ . Logistic regression was used to calculate the relative odds of HPV status with 11q13 amp after adjusting for the potential confounders (age, tumour stage and site).

## RESULTS

### Study population demographics

Table 1 summarises the patient demographics. Approximately 64% (80 out of 125) of the patients were males and 36% (45 out of 125) were females. The majority of patients were current smokers ( $n=76$ , 61%). The majority of the study population (65%) was dead by the last follow-up (May 2004). Our study population ranged in age from 18 to 92 years; the median age at diagnosis was 63 years, and the median age at death was 67 years.

### Human papillomavirus prevalence and genotype in tumour specimens

Twenty-four percent (30 out of 125) of the tumours were positive for HPV. All HPV-positive tumours contained HPV16 whereas two out of 30 (7%) were coinfecting with other HPV types. One had an additional low-risk HPV type 11 and the other had an additional high-risk HPV type 33 and possibly 52 (a positive hybridisation signal for cross-reactive HPV33 probe on the HPV linear array genotyping strip could not rule out the presence of HPV52) (Table 1).

### Distribution of tumour human papillomavirus status according to patient characteristics

The majority of tumours occurred in the oral cavity (91 out of 125, 73%). Human papillomavirus-positive tumours occurred at various sites throughout the oral cavity and pharynx (data not shown). Approximately 20% (18 out of 91) of tumours from the oral cavity were found to be HPV-positive, whereas a slightly higher proportion of HPV-positive tumours were observed in the oropharynx (35% (12 out of 34)). When stratified by smoking status, seven HPV-positive tumours from the oropharynx of current smokers occurred in the base of tongue (3), soft palate (2), tonsil (1) and an overlapping lesion of the oropharynx (1). In contrast, the five HPV-positive tumours from the oropharynx of nonsmokers and former smokers occurred in the tonsil (3) and base of tongue (2) only.

Analyses of factors associated with a positive HPV status are summarised in Table 2. Human papillomavirus-positive tumours were less likely to occur in older individuals ( $<55$  years, 40% were HPV-positive vs  $\geq 55$  years, 19% were HPV-positive: adjusted OR = 0.3, 95% confidence interval (CI) = 0.1–0.8), and were less likely to carry amplified genes in chromosomal band 11q13. After adjusting for age, tumour site and stage, 11q13 amp status was still statistically associated with HPV status, with only 17% (five out of 30) of HPV-positive tumours vs 44% (39 out of 89) of HPV-negative tumours having 11q13 amp (adjusted OR = 0.2, 95% CI = 0.1–0.6).

**Table 1** Patient characteristics ( $n=125$ )

Characteristic	N	Percent (%)
<b>Age (years)</b>		
<35	3	2.40
36–45	9	7.20
46–55	18	14.40
56–65	43	34.40
66–75	37	29.60
76–85	13	10.40
>85	2	1.60
<b>Gender</b>		
Male	80	64.00
Female	45	36.00
<b>Race</b>		
Caucasian	117	93.60
Other	8	6.40
<b>Vital status</b>		
Alive	44	35.20
Dead	81	64.80
<b>Tobacco exposure</b>		
Never	13	10.40
Former	36	28.80
Current	76	60.80
<b>Alcohol duration/years</b>		
Never	18	14.40
Social	3	2.40
2–10	6	4.80
11–30	33	26.40
31–50	53	42.40
>50	12	9.60
<b>HPV status</b>		
Negative	95	76.00
HPV16	28	22.40
HPV16 & 11	1	0.80
HPV16, 33 & (52) <sup>a</sup>	1	0.80
<b>Family history cancer</b>		
No	62	49.60
Yes	63	50.40
<b>Tumour stage</b>		
I	21	16.80
II	28	22.40
III	27	21.60
IV	49	39.20
<b>Subsequent tumour(s)</b>		
No	61	48.80
Yes	64	51.20
Total	125	100.00

Abbreviation: HPV = human papillomavirus. <sup>a</sup>Unable to rule out HPV52 owing to cross-reactivity of the HPV33 probe.

We evaluated the pattern of tobacco and alcohol risk behaviours by HPV status. Although the majority of patients from both the HPV-positive and negative groups were ever smokers and ever drinkers (77 and 80%, respectively), these patients were no more likely to be HPV-positive than those who never smoked or consumed alcohol (OR = 0.8, 95% CI = 0.1–7.8). However, no statistical interaction between alcohol and drinking habits with HPV-positivity was observed.

Individuals with HPV-positive tumours were less likely to have subsequent tumours, either recurrences, metastases or new

**Table 2** Association of patient/tumour characteristics with HPV status

Characteristics	Any HPV subtype			
	No. positive (n = 30)	No. negative (n = 95)	Unadjusted OR (95% CI)	Adjusted OR <sup>a</sup> (95% CI)
Age (years)				
<55	12	18	1.0 (referent)	
≥55	18	77	0.4 (0.1–0.9)	0.3 (0.1–0.8)
Gender				
Male	23	57	1.0 (referent)	
Female	7	38	0.5 (0.2–1.1)	0.5 (0.2–1.3)
Race				
Caucasian	29	88	1.0 (referent)	
Other	1	7	0.4 (0.1–3.7)	0.2 (0.02–2.2)
Tobacco exposure				
Never	4	9	1.7 (0.5–6.1)	1.7 (0.4–6.7)
Former	10	26	1.4 (0.6–3.6)	2.4 (0.8–6.6)
Current	16	60	1.0 (referent)	
Alcohol exposure				
Never	4	14	1.0 (referent)	
Ever	26	81	1.1 (0.3–3.7)	0.9 (0.3–3.2)
Alcohol duration				
Never user	4	14	1.0 (referent)	
Short-term user	3	6	1.8 (0.3–10.3)	1.3 (0.2–8.4)
Long-term user	23	75	1.1 (0.3–3.6)	0.9 (0.3–3.2)
Combined tobacco and alcohol exposure				
Ever/Ever	23	76	1.0 (referent)	
Never/Ever	3	5	2.0 (0.4–8.9)	1.8 (0.4–8.6)
Ever/Never	3	10	1.0 (0.3–3.9)	1.3 (0.3–5.6)
Never/Never	1	4	0.8 (0.1–7.8)	0.8 (0.1–8.3)
Family cancer history				
No	13	49	1.0 (referent)	
Yes	17	46	1.4 (0.6–3.2)	1.3 (0.6–3.2)
Tumour site				
Oral cavity	18	73	1.0 (referent)	
Oropharynx	12	22	2.2 (0.9–5.3)	2.4 (1.0–5.8)
11q13 amplification status				
Not amplified	25	50	1.0 (referent)	
Amplified	5	39	0.3 (0.1–0.7)	0.2 (0.1–0.6)
Not performed	0	6	—	
Tumour grade				
Well	8	24	1.0 (referent)	
Moderate	17	52	1.0 (0.4–2.6)	1.0 (0.3–2.8)
Poor	3	11	0.8 (0.2–3.7)	0.7 (0.1–3.2)
Unknown	2	8	—	—
Tumour stage				
I & II	10	39	1.0 (referent)	
III & IV	20	56	1.4 (0.6–3.3)	1.4 (0.6–3.5)
Recurrent, metastatic or new primary tumours				
No	21	41	1.0 (referent)	
Yes	9	54	0.4 (0.2–0.9)	0.4 (0.2–1.0)

Abbreviations: CI, confidence interval; HPV, human papillomavirus; OR, odds ratio. <sup>a</sup>Adjusted for age, site and stage, where appropriate.

primaries (adjusted OR = 0.4, 95% CI = 0.2–1.0). Subsequent tumours that developed in HPV-positive patients (33%, 10 out of 30) were confined to head and neck sites. In contrast, among patients with HPV-negative tumours, 57% (54 out of 95) developed recurrences, metastases or new primaries at head and neck sites, as well as other sites throughout the body.

A higher proportion of HPV-positive tumours occurred in the oropharynx. Forty percent (12 out of 30) of HPV-positive tumours compared to 23% (22 out of 95) of HPV-negative tumours occurred in the oropharynx (adjusted OR = 2.4, 95% CI = 1.0–5.8). A high tumour stage (stage III & IV: adjusted OR = 1.4, 95% CI = 0.6–3.5) and family history of cancer (adjusted OR = 1.3, 95%

CI = 0.6–3.2) were also more likely to be associated with HPV-positive tumours, but none of these associations were statistically significant.

### Human papillomavirus and 11q13 amplification status according to tumour site

Amplification status at chromosomal band 11q13 was available for 119 tumours from our study population. Approximately 37% (44 out of 119) were amplified at 11q13. Figure 1 shows the frequency of 11q13 amp stratified by tumour site and HPV status. Overall, we detected a lower prevalence of 11q13 amp in the HPV-positive tumours than in the HPV-negative group (17 vs 44%, respectively, Fisher's exact  $P=0.009$ ). In the oral cavity, 22% of the HPV-positive tumours were amplified at 11q13, whereas 39% of the HPV-negative tumours had 11q13 amp (Fisher's exact  $P=0.270$ ). In the oropharynx, a significantly smaller proportion of HPV-positive tumours had 11q13 amp when compared with HPV-negative tumours (8% HPV-positive vs 60% HPV-negative, Fisher's exact  $P=0.008$ ).

### TP53 mutation status, p16 expression and 11q13 amplification status in human papillomavirus-positive tumours

In a subset of tumours ( $n=69$ ) for which data on all three molecular markers were available (i.e. *TP53* mutation status, p16 expression and 11q13 amp status), their associations with HPV status was assessed (Table 3). *TP53* mutations were less likely to occur in the HPV-positive HNSCC (adjusted OR = 0.4, 95% CI = 0.0–3.1). As expected, HPV-positive tumours were more likely to express p16 (adjusted OR = 3.0, 95% CI = 0.9–9.7). Sixty percent (nine out of 15) of the HPV-positive tumours exhibited high expression of p16 whereas 33% (18 out of 54) of HPV-negative tumours expressed high levels of p16.

We classified the subpopulation according to combined p16 expression, *TP53* mutation status and 11q13 amp status. Class II included tumours that carried wild-type *TP53*, expressed p16 and was not amplified at 11q13, class III included tumours that also carried wild-type *TP53* and expressed p16 but were amplified at chromosomal band 11q13 and class I tumours were those with all other combinations of *TP53* mutation status, p16 expression and 11q13 amp status. Human papillomavirus-positive tumours were more likely not to be amplified at 11q13 and simultaneously overexpress p16 and contain wild-type *TP53* (class II: adjusted OR = 6.5, 95% CI = 1.8–23.9). The tumours that contained wild-type *TP53* overexpressed p16, but those with 11q13 amp were less likely to be HPV-positive than those without 11q13 amp and were no different from all other tumours (class III: OR = 0.9, 95% CI = 0.1–8.9).

### DISCUSSION

This retrospective study of 125 patients with first primary head and neck tumours evaluated the association between tumour HPV status and patient demographics, clinical risk factors and amplification status at chromosomal band 11q13. We further evaluated the relationship between genetic alterations: *TP53* mutation status, 11q13 amp and p16 protein expression in a subpopulation of 69 unselected tumours. Our analysis shows that along with wild-type *TP53* and p16 overexpression, HPV-positive tumours were less likely to carry gene amplification of chromosomal band 11q13, which further defines the characteristic of HPV-positive head and neck tumours and suggests a distinct molecular pathway for HNSCC development.

The relationship between LOH at *TP53*, 11q13 amp and HPV has been shown in one study including only 37 HNSCC (Rodrigo et al,




Figure 1 Frequency of 11q13 amp according to HPV status and tumour site.

Table 3 Association of molecular markers with HPV status

Molecular characteristics	Any HPV		
	No. positive (n = 15)	No. negative (n = 54)	Adjusted <sup>a</sup> OR (95% CI)
<i>11q13 amplification status</i>			
Not amplified	13	26	1.0 (referent)
Amplified	2	28	0.1 (0.0–0.7)
<i>TP53 mutation status</i>			
Wild-type	14	45	1.0 (referent)
Mutant	1	9	0.4 (0.0–3.1)
<i>CDKN2A/p16 expression</i>			
Negative	6	36	1.0 (referent)
Positive	9	18	3.0 (0.9–9.7)
<i>Three molecular<sup>b</sup> markers combined</i>			
Class I	6	39	1.0 (referent)
Class II	8	8	6.5 (1.8–23.9)
Class III	1	7	0.9 (0.1–8.9)

Abbreviations: CI, confidence interval; HPV, human papillomavirus; OR, odds ratio. <sup>a</sup>Adjusted for age, site and stage, where appropriate. <sup>b</sup>Three molecular markers combined were classified as: class II = *TP53*wt, p16-positive and no 11q13 amplification. Class III = *TP53*wt, p16-positive and 11q13 amplification. Class I = all other possible combination of *TP53* mutation status, p16 expression and 11q13 amplification status.

2002). Loss of heterozygosity at *TP53*, 11q13 amp and HPV status were reported on only 11 (30%) of their samples and p16 overexpression was not evaluated. The study reported a lower frequency of 11q13 amp in HPV-positive tumours (HPV+ with 11q13 amp = 2 out of 4 (50%); HPV- with 11q13 amp = 5 out of 7 (71%) but their HPV-positive tumours were more likely to have LOH at *TP53* (three out of four (75%)). Earlier studies utilising immunohistochemistry have shown that the overexpression of p53 and p16 and reduced expression of cyclin D1 were associated with HPV-positive tumours but 11q13 amp was not evaluated and these studies only included tumours arising in the tonsils (Andl et al, 1998; Li et al, 2004). One other study evaluated the combined relationship between the three molecular markers and HPV status using immunohistochemistry (Wilczynski et al, 1998), but similar to the earlier mentioned studies, their analysis was limited to tumours that arose in the tonsils. In addition, the study included tumours that were first primaries, recurrences, lymph node metastases and lung metastases. Despite the differences in sample size and tumour site included in these analyses, the results from these three studies are in agreement with our findings. In our study, the combined characteristics of *TP53* mutation status, p16 protein expression (a protein which is overexpressed as a consequence of HPV E7 expression) and 11q13 amp (a genetic alteration frequently observed in head and neck tumours) were

evaluated to further characterise the role of HPV in head and neck tumorigenesis. Tumours that were HPV-positive were more likely to carry all three markers (wild-type *TP53*, express p16 and no amplification at 11q13). We included in our analyses tumours that were first primaries from various sites within the oral cavity and pharynx and demonstrate that HPV-positive tumours with this distinct molecular phenotype can arise not only in the oropharynx (50%, four out of eight) but also in the oral cavity (50%, four out of eight).

Our results shed further light on the multiple genetic alterations that are commonly found in head and neck tumours (Gollin, 2001). A genetic progression model for HNSCC has been proposed by Califano *et al* (Califano *et al*, 1996, 2000) and describes a series of ordered genetic changes which may occur during tumour development. It is thought that, for the most part, early genetic alterations include loss of chromosomal bands 9p21, 3p, 17p13 and amplification of 11q13. The candidate genes involved in these genetic alterations are thought to be *CDKN2A/p16*, the tumour suppressor gene *FHIT*, *TP53* and the *CCND1* oncogene, respectively. Examination of hyperplastic, dysplastic, carcinoma *in situ* and invasive carcinoma lesions revealed increasing LOH which corresponded with histopathological progression (Califano *et al*, 1996). Loss of heterozygosity at 9p21 was the most frequent alteration in benign hyperplastic lesions, followed by LOH at 3p21 and 17p13. In addition, dysplastic lesions revealed an increased incidence of allelic imbalance at 11q13 (benign hyperplasia: (6%), dysplasia: (29%), carcinoma *in situ*: 40%, invasive carcinoma (61%)). Our results show that HPV-positive tumours are less likely to be amplified at 11q13 and at the same time p16 is overexpressed, providing indirect evidence that the infection may have occurred at an earlier time point in the carcinogenic process.

The loss of functional *TP53* by mutation seems not to be necessary in HPV-positive HNSCC, as p53 protein loss results from increased degradation in these tumours (Wiest *et al*, 2002; Hafkamp *et al*, 2003; Braakhuis *et al*, 2004). The HPV E7 oncoprotein binds and degrades the RB1 tumour suppressor protein (Boyer *et al*, 1996), which in turn causes the release of E2F (a transcriptional regulator of cell proliferation genes) from pRb/E2F complexes, permitting E2F to transactivate S-phase-related genes. The functional inactivation of pRb by E7 leads to overexpression of the cyclin-dependent kinase inhibitor p16 (Khleif *et al*, 1996). The detection of p16 expression therefore is considered to be a surrogate marker for HPV infection, this observation was also confirmed in our studies.

Amplification of chromosomal band 11q13 has been reported in a number of carcinomas (head and neck, breast, lung, pancreatic, prostate, ovarian, bladder and so on.) (Schwab, 1998). Approximately 45% of HNSCC have amplification of 11q13 (Schuurin, 1995; Lese *et al*, 1995). The amplification status of chromosomal band 11q13 was determined in our study, using a sensitive methodology, FISH with probes for *FGF3/INT2*, *FGF4/HST1* or *CCND1* (Lese *et al*, 1995; Shuster *et al*, 2000). A significantly smaller proportion of patients with 11q13 amp were observed in the HPV-positive group (13%, two out of 15) compared to the HPV-negative group (54%, 26 out of 48). One consequence of

11q13 amp is *CCND1* overexpression, which is thought to play a direct role in this disease (Callender *et al*, 1994). Other proto-oncogenes which map to the 11q13 core region are amplified and overexpressed (Huang *et al*, 2002), but their role in head and neck tumour development has not yet been fully delineated. The *CCND1* gene product, cyclin D1 associates with cyclin-dependent kinase 4 and 6, and this complex promotes RB1 phosphorylation and like the HPV E7 oncoprotein, their interaction leads to the dissociation of pRb from the transcription factor E2F. This results in transition of the cell from G<sub>1</sub> into S-phase. One possible explanation for the lack of 11q13 amp in HPV-positive tumours may be that amplification of chromosomal band 11q13 might be unnecessary, as the resulting interaction of the HPV E7 oncoprotein with pRb might allow the cells to be less dependent on *CCND1* for cell cycle progression. We therefore believe that the low frequency of 11q13 amp observed in our HPV-positive head and neck tumours may be explained by a distinct mechanism for HPV carcinogenesis.

Several studies report that patients with HPV-positive head and neck tumours have an improved prognosis (Ritchie *et al*, 2003; Schlecht, 2005), whereas amplification of 11q13 has been associated with a more rapid and frequent recurrence of disease (Fujii *et al*, 2001) and poorer survival (Akervall *et al*, 1997; Rodrigo *et al*, 2000; Namazie *et al*, 2002). Weinberger *et al* (2006) has shown that patients with HPV-positive tumours which express high levels of p16 protein and low levels of p53 protein present with a favourable prognosis. We have shown in our study that HPV-positive tumours have high levels of p16 expression and wild-type *TP53* and in addition, a low frequency of amplification at 11q13. Further analyses are warranted to determine whether this lack of 11q13 amp also contributes to the improved prognosis observed in patients with HPV-positive HNSCC.

## ACKNOWLEDGEMENTS

The authors are grateful to Drs Jonas T Johnson, Eugene N Meyers and Jennifer R Grandis for their support of this study, and Ms Robin Wagner, Ms Jennifer Vates and Ms Jennifer Ridge-Hetrick for assistance with clinical correlates. We thank Drs Jennifer Grandis and Robert Ferris for critical reading of the manuscript. We thank Jane Dudek and David Reed from Roche Diagnostics for allowing us to evaluate the LightCycler<sup>®</sup> 2.0 system for use in HPV testing and for making available the HPV Linear Array reagents (an RUO product) for the HPV genotyping. We also thank Drs Ilyas Kamboh and Robert Redner for the use of their instrumentation; Ryan Minster and Dr David Sloan (Biotage) for their assistance in pyrosequencing. This work was supported in part by NIH Grants R01DE10513, R01DE12008 and R01DE14729 to SMG, P50CA097190 (SPOR: Project 1) to ET and JLW and K07CA80668 to FM. CCRR and JSW were supported by the UPCI Cancer Education and Career Development Grant R25CA089507 to William Bigbee. The FISH studies were carried out in the University of Pittsburgh Cancer Institute Cytogenetics Facility, funded in part by P30CA47904 to Ronald B Herberman.

## REFERENCES

- Akervall JA, Michalides RJ, Mineta H, Balm A, Borg A, Dictor MR, Jin Y, Loftus B, Mertens F, Wennerberg JP (1997) Amplification of cyclin D1 in squamous cell carcinoma of the head and neck and the prognostic value of chromosomal abnormalities and cyclin D1 overexpression. *Cancer* **79**: 380–389
- Andl T, Kahn T, Pfuhl A, Nicola T, Erber R, Conradt C, Klein W, Helbig M, Dietz A, Weidauer H, Bosch FX (1998) Etiological involvement of oncogenic human papillomavirus in tonsillar squamous cell carcinomas lacking retinoblastoma cell cycle control. *Cancer Res* **58**: 5–13

- Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, Bernstein L, Schoenberg JB, Stemhagen A, Fraumeni Jr JF (1988) Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res* **48**: 3282–3287
- Boyer SN, Wazer DE, Band V (1996) E7 protein of human papilloma virus-16 induces degradation of retinoblastoma protein through the ubiquitin-proteasome pathway. *Cancer Res* **56**: 4620–4624
- Braakhuis BJ, Snijders PJ, Keune WJ, Meijer CJ, Ruijter-Schippers HJ, Leemans CR, Brakenhoff RH (2004) Genetic patterns in head and neck

- cancers that contain or lack transcriptionally active human papillomavirus. *J Natl Cancer Inst* **96**: 998–1006
- Califano J, van der Riet P, Westra W, Nawroz H, Clayman G, Piantadosi S, Corio R, Lee D, Greenberg B, Koch W, Sidransky D (1996) Genetic progression model for head and neck cancer: implications for field cancerization. *Cancer Res* **56**: 2488–2492
- Califano J, Westra WH, Meininger G, Corio R, Koch WM, Sidransky D (2000) Genetic progression and clonal relationship of recurrent premalignant head and neck lesions. *Clin Cancer Res* **6**: 347–352
- Callender T, El-Naggar AK, Lee MS, Frankenthaler R, Luna MA, Batsakis JG (1994) PRAD-1 (CCND1)/cyclin D1 oncogene amplification in primary head and neck squamous cell carcinoma. *Cancer* **74**: 152–158
- de Roda Husman AM, Walboomers JM, van den Brule AJ, Meijer CJ, Snijders PJ (1995) The use of general primers GP5 and GP6 elongated at their 3' ends with adjacent highly conserved sequences improves human papillomavirus detection by PCR. *J Gen Virol* **76**(Part 4): 1057–1062
- Franceschi S, Levi F, La VC, Conti E, Dal ML, Barzan L, Talamini R (1999) Comparison of the effect of smoking and alcohol drinking between oral and pharyngeal cancer. *Int J Cancer* **83**: 1–4
- Franceschi S, Talamini R, Barra S, Baron AE, Negri E, Bidoli E, Serraino D, La VC (1990) Smoking and drinking in relation to cancers of the oral cavity, pharynx, larynx, and esophagus in northern Italy. *Cancer Res* **50**: 6502–6507
- Fujii M, Ishiguro R, Yamashita T, Tashiro M (2001) Cyclin D1 amplification correlates with early recurrence of squamous cell carcinoma of the tongue. *Cancer Lett* **172**: 187–192
- Gillison ML, Koch WM, Capone RB, Spafford M, Westra WH, Wu L, Zahurak ML, Daniel RW, Viglione M, Symer DE, Shah KV, Sidransky D (2000) Evidence for a causal association between human papillomavirus and a subset of head and neck cancers. *J Natl Cancer Inst* **92**: 709–720
- Gollin SM (2001) Chromosomal alterations in squamous cell carcinomas of the head and neck: window to the biology of disease. *Head Neck* **23**: 238–253
- Gravitt PE, Peyton CL, Alessi TQ, Wheeler CM, Coutlee F, Hildesheim A, Schiffman MH, Scott DR, Apple RJ (2000) Improved amplification of genital human papillomaviruses. *J Clin Microbiol* **38**: 357–361
- Hafkamp HC, Speel EJ, Haesevoets A, Bot FJ, Dinjens WN, Ramaekers FC, Hopman AH, Manni JJ (2003) A subset of head and neck squamous cell carcinomas exhibits integration of HPV 16/18 DNA and overexpression of p16INK4A and p53 in the absence of mutations in p53 exons 5–8. *Int J Cancer* **107**: 394–400
- Herrero R, Castellsague X, Pawlita M, Lissowska J, Kee F, Balaram P, Rajkumar T, Sridhar H, Rose B, Pintos J, Fernandez L, Idris A, Sanchez MJ, Nieto A, Talamini R, Tavani A, Bosch FX, Reidel U, Snijders PJ, Meijer CJ, Viscidi R, Munoz N, Franceschi S (2003) Human papillomavirus and oral cancer: the International Agency for Research on Cancer multicenter study. *J Natl Cancer Inst* **95**: 1772–1783
- Huang X, Gollin SM, Raja S, Godfrey TE (2002) High-resolution mapping of the 11q13 amplicon and identification of a gene, TAOS1, that is amplified and overexpressed in oral cancer cells. *Proc Natl Acad Sci USA* **99**: 11369–11374
- Khleif SN, DeGregori J, Yee CL, Otterson GA, Kaye FJ, Nevins JR, Howley PM (1996) Inhibition of cyclin D-CDK4/CDK6 activity is associated with an E2F-mediated induction of cyclin kinase inhibitor activity. *Proc Natl Acad Sci USA* **93**: 4350–4354
- Law JC, Whiteside TL, Gollin SM, Weissfeld J, El-Ashmawy L, Srivastava S, Landreneau RJ, Johnson JT, Ferrell RE (1995) Variation of p53 mutational spectra between carcinoma of the upper and lower respiratory tract. *Clin Cancer Res* **1**: 763–768
- Lese CM, Rossie KM, Appel BN, Reddy JK, Johnson JT, Myers EN, Gollin SM (1995) Visualization of INT2 and HST1 amplification in oral squamous cell carcinomas. *Genes Chromosomes Cancer* **12**: 288–295
- Li W, Thompson CH, Cossart YE, O'Brien CJ, McNeil EB, Scolyer RA, Rose BR (2004) The expression of key cell cycle markers and presence of human papillomavirus in squamous cell carcinoma of the tonsil. *Head Neck* **26**: 1–9
- Merletti F, Boffetta P, Ciccone G, Mashberg A, Terracini B (1989) Role of tobacco and alcoholic beverages in the etiology of cancer of the oral cavity/oropharynx in Torino, Italy. *Cancer Res* **49**: 4919–4924
- Namazie A, Alavi S, Olopade OI, Pauletti G, Aghamohammadi N, Aghamohammadi M, Gornbein JA, Calcaterra TC, Slamoni DJ, Wang MB, Srivatsan ES (2002) Cyclin D1 amplification and p16(MTS1/CDK41) deletion correlate with poor prognosis in head and neck tumors. *Laryngoscope* **112**: 472–481
- Ritchie JP, Smith EM, Summersgill KF, Hoffman HT, Wang D, Klusmann JP, Turek LP, Haugen TH (2003) Human papillomavirus infection as a prognostic factor in carcinomas of the oral cavity and oropharynx. *Int J Cancer* **104**: 336–344
- Rodrigo JP, Garcia LA, Ramos S, Lazo PS, Suarez C (2000) EMS1 gene amplification correlates with poor prognosis in squamous cell carcinomas of the head and neck. *Clin Cancer Res* **6**: 3177–3182
- Rodrigo JP, Gonzalez MV, Lazo PS, Ramos S, Coto E, Alvarez I, Garcia LA, Suarez C (2002) Genetic alterations in squamous cell carcinomas of the hypopharynx with correlations to clinicopathological features. *Oral Oncol* **38**: 357–363
- Saiki RK, Bugawan TL, Horn GT, Mullis KB, Erlich HA (1986) Analysis of enzymatically amplified beta-globin and HLA-DQ alpha DNA with allele-specific oligonucleotide probes. *Nature* **324**: 163–166
- Schlecht NF (2005) Prognostic value of human papillomavirus in the survival of head and neck cancer patients: an overview of the evidence. *Oncol Rep* **14**: 1239–1247
- Schraml P, Kononen J, Bubendorf L, Moch H, Bissig H, Nocito A, Mihatsch MJ, Kallioniemi OP, Sauter G (1999) Tissue microarrays for gene amplification surveys in many different tumor types. *Clin Cancer Res* **5**: 1966–1975
- Schuuring E (1995) The involvement of the chromosome 11q13 region in human malignancies: cyclin D1 and EMS1 are two new candidate oncogenes – a review. *Gene* **159**: 83–96
- Schwab M (1998) Amplification of oncogenes in human cancer cells. *Bioessays* **20**: 473–479
- Shuster MI, Han L, Le Beau MM, Davis E, Sawicki M, Lese CM, Park NH, Colicelli J, Gollin SM (2000) A consistent pattern of RIN1 rearrangements in oral squamous cell carcinoma cell lines supports a breakage-fusion-bridge cycle model for 11q13 amplification. *Genes Chromosomes Cancer* **28**: 153–163
- Sidransky D (1995) Molecular genetics of head and neck cancer. *Curr Opin Oncol* **7**: 229–233
- Smith EM, Hoffman HT, Summersgill KS, Kirchner HL, Turek LP, Haugen TH (1998) Human papillomavirus and risk of oral cancer. *Laryngoscope* **108**: 1098–1103
- Syrjanen K, Syrjanen S, Lamberg M, Pyrhonen S, Nuutinen J (1983) Morphological and immunohistochemical evidence suggesting human papillomavirus (HPV) involvement in oral squamous cell carcinogenesis. *Int J Oral Surg* **12**: 418–424
- Weber A, Wittekind C, Tannapfel A (2003) Genetic and epigenetic alterations of 9p21 gene products in benign and malignant tumors of the head and neck. *Pathol Res Pract* **199**: 391–397
- Weinberger PM, Yu Z, Haffty BG, Kowalski D, Harigopal M, Brandsma J, Sasaki C, Joe J, Camp RL, Rimm DL, Psyrri A (2006) Molecular classification identifies a subset of human papillomavirus-associated oropharyngeal cancers with favorable prognosis. *J Clin Oncol* **24**: 736–747
- Wiest T, Schwarz E, Enders C, Flechtenmacher C, Bosch FX (2002) Involvement of intact HPV16 E6/E7 gene expression in head and neck cancers with unaltered p53 status and perturbed pRb cell cycle control. *Oncogene* **21**: 1510–1517
- Wilczynski SP, Lin BT, Xie Y, Paz IB (1998) Detection of human papillomavirus DNA and oncoprotein overexpression are associated with distinct morphological patterns of tonsillar squamous cell carcinoma. *Am J Pathol* **152**: 145–156