

Research article

Open Access

The calcium channel $\beta 2$ (CACNB2) subunit repertoire in teleosts

Alicia M Ebert¹, Catherine A McAnelly¹, Ashok Srinivasan²,
Rachel Lockridge Mueller¹, David B Garrity¹ and Deborah M Garrity*¹

Address: ¹Department of Biology, Colorado State University, Fort Collins, CO 80523, USA and ²NIH/NHLBI, 10 Center Drive, Rm. 6C/104, Bethesda, MD 20892, USA

Email: Alicia M Ebert - amebert@lamar.colostate.edu; Catherine A McAnelly - cmcanell@holly.colostate.edu;
Ashok Srinivasan - srinivasana@mail.nih.gov; Rachel Lockridge Mueller - rachel.mueller@colostate.edu;
David B Garrity - david.garrity@colostate.edu; Deborah M Garrity* - deborah.garrity@colostate.edu

* Corresponding author

Published: 17 April 2008

Received: 24 December 2007

BMC Molecular Biology 2008, 9:38 doi:10.1186/1471-2199-9-38

Accepted: 17 April 2008

This article is available from: <http://www.biomedcentral.com/1471-2199/9/38>

© 2008 Ebert et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Cardiomyocyte contraction is initiated by influx of extracellular calcium through voltage-gated calcium channels. These oligomeric channels utilize auxiliary β subunits to chaperone the pore-forming α subunit to the plasma membrane, and to modulate channel electrophysiology [1]. Several β subunit family members are detected by RT-PCR in the embryonic heart. Null mutations in mouse $\beta 2$, but not in the other three β family members, are embryonic lethal at E10.5 due to defects in cardiac contractility [2]. However, a drawback of the mouse model is that embryonic heart rhythm is difficult to study in live embryos due to their intra-uterine development. Moreover, phenotypes may be obscured by secondary effects of hypoxia. As a first step towards developing a model for contributions of β subunits to the onset of embryonic heart rhythm, we characterized the structure and expression of $\beta 2$ subunits in zebrafish and other teleosts.

Results: Cloning of two zebrafish $\beta 2$ subunit genes ($\beta 2.1$ and $\beta 2.2$) indicated they are membrane-associated guanylate kinase (MAGUK)-family genes. Zebrafish $\beta 2$ genes show high conservation with mammals within the SH3 and guanylate kinase domains that comprise the "core" of MAGUK proteins, but $\beta 2.2$ is much more divergent in sequence than $\beta 2.1$. Alternative splicing occurs at the N-terminus and within the internal HOOK domain. In both $\beta 2$ genes, alternative short ATG-containing first exons are separated by some of the largest introns in the genome, suggesting that individual transcript variants could be subject to independent cis-regulatory control. In the *Tetraodon nigroviridis* and *Fugu rubripes* genomes, we identified single $\beta 2$ subunit gene loci. Comparative analysis of the teleost and human $\beta 2$ loci indicates that the short 5' exon sequences are highly conserved. A subset of 5' exons appear to be unique to teleost genomes, while others are shared with mammals. Alternative splicing is temporally and spatially regulated in embryo and adult. Moreover, a different subset of spliced $\beta 2$ transcript variants is detected in the embryonic heart compared to the adult.

Conclusion: These studies refine our understanding of $\beta 2$ subunit diversity arising from alternative splicing, and provide the groundwork for functional analysis of $\beta 2$ subunit diversity in the embryonic heart.

Background

Voltage-dependent L-type Ca^{2+} channels (V-LTCC) are essential for the initiation and regulation of excitation-contraction coupling in cardiac muscle. In addition, Ca^{2+} entry through V-LTCC channels can also serve as a second messenger to modulate regulatory protein kinases, calmodulin and beta-adrenergic responses [3]. Genetic mutation of key Ca^{2+} homeostasis proteins is frequently associated with defects in both cardiac cell differentiation and contractility, suggesting Ca^{2+} signaling is essential for normal cardiac development [4-7]. Ca^{2+} signals can also contribute to the control of gene expression [8]. Ca^{2+} influx through V-LTCC can dramatically affect the phosphorylation, activity, or expression of many genes [5,8-10]. These data support the hypothesis that Ca^{2+} signaling through V-LTCC impacts not only cardiac contraction but also embryonic cardiac growth and morphology, physiology, and gene expression.

V-LTCCs are oligomeric proteins composed of a pore-forming $\alpha 1$ subunit and up to four auxiliary subunits, termed $\alpha 2$, β , γ , and δ , associated in a 1:1:1:1 stoichiometry [11,12]. Mammals encode 10 distinct $\alpha 1$ subunit genes (four of which form V-LTCC) and four distinct β subunit genes ($\beta 1 - \beta 4$). Particular $\alpha 1$ and β subtypes are able to associate non-exclusively to form heterogeneous channels [13-17]. V-LTCC that contain the $\alpha 1c$ subunit ($\text{Ca}_v1.2$) constitute the vast majority of L-type channels expressed in adult cardiac muscle [17-22].

The V-LTCC β subunit genes have important functions in cardiac, skeletal and smooth muscle, the central nervous system and the retina [2,23-28]. β subunits modulate Ca^{2+} channel function in two distinct ways. First, they facilitate transport of the $\alpha 1$ to the plasmalemma. In addition, they interact with $\alpha 1$ to modulate both the kinetics and voltage-sensitivity of the channels [29-31]. As members of the membrane-associated guanylate kinase (MAGUK) family, the V-LTCC β subunit genes include conserved SH3 and guanylate kinase (GK) domains connected by a bridging region termed the HOOK domain [32]. The larger MAGUK protein family appears to be specific to metazoans, and its core structure, including the SH3 and GK domains, can be traced back to early metazoan history [33]. All known β genes undergo alternative splicing of 5' exons, and some exhibit alternative splicing of internal exons as well [34-36]. Alternative splicing in internal exons can create truncated proteins, some of which have been shown to mediate calcium channel-independent functions in the cell [1,36,37].

In the mouse, the earliest documented expression of a cardiac β protein is a Western blot showing $\beta 2$ protein in the linear heart tube (E8.5) [2]. By E9.5 (chamber morphogenesis stages), $\beta 4$ and $\beta 2$ mRNA and protein are clearly

present in the embryonic myocardium [2,10,38]. $\beta 2$ and $\beta 4$ proteins show detectable differences in spatial distribution as early as E9.5, which become more pronounced in mid- to late embryogenesis (E10.5 - E15.5) and in later fetal development [38]. In rat postnatal development (4.5 weeks), RT-PCR studies indicate all four β subunits are expressed in cardiac tissue, though they differ in abundance [39]. Significantly, by 4.5 weeks RNA from atria and ventricles show different β subunit RT-PCR expression profiles [39]. In brain and heart, newborn and juvenile animals express different subsets of β genes (including different transcript variants) than the adult [39,40]. The physiological significance of these observations is not known. However, a recent report describing the targeted mutagenesis of mouse $\beta 2$ indicated that homozygous loss of this gene ($\beta 2^{-/-}$) is embryonic lethal at E10.5, significantly earlier than the $\alpha 1c$ -deficient mouse [2,19]. No up-regulation of other β subunits was observed in $\beta 2^{-/-}$ mutants. In $\beta 2^{-/-}$ mutant mice, cardiac contractility is severely compromised and V-LTCC currents diminished. Hearts in mutant mice exhibit severe bradycardia and vascular patterning defects, but these phenotypes may be secondary to severe morphological defects that develop in the heart or to the overall reduced embryonic growth [2]. In contrast to $\beta 2$, mice with null mutations in $\beta 1$, $\beta 3$ and $\beta 4$ have no reported cardiac phenotypes [23,25-28]. However, the zebrafish $\beta 1^{-/-}$ mutant *relaxed* develops pericardial edema and reduced circulation at 4 dpf (days post-fertilization), suggesting a possible compromise in cardiac function [41]. Thus, while multiple β subunits are clearly present in the embryonic heart, much remains to be learned about their functional roles in development.

As a model for the study of β subunit developmental function in the heart, zebrafish has several advantages, including the external fertilization of embryos and rapid, transparent development of the larvae, which allows experimental access to the developing heart and circulatory system. Moreover, for about the first 6 days of development the small zebrafish larvae subsist on oxygen acquired via diffusion, enabling them to survive without a functional heart [42]. As a first step in establishing a zebrafish model, we cloned two $\beta 2$ subunit gene homologs from zebrafish. We find one $\beta 2$ subunit is much more similar than the other to known vertebrate $\beta 2$ subunits. Significant alternative splicing occurs involving several ATG-containing first exons in both zebrafish genes, as well as two other teleost species. The separation of these short exons by extremely large introns suggests potential independent cis-regulatory control of alternatively spliced transcripts. Unexpectedly, both $\beta 2$ subunits were maternally expressed, consistent with possible early roles for this gene in morphogenetic movements or patterning. Additionally, both genes showed evidence of tis-

sue-, stage- and transcript-specific regulatory control in embryos and adults.

Results

Identification of zebrafish and pufferfish calcium channel $\beta 2$ subunit (CACNB2) genes

TBLASTN searches of the GenBank database at NCBI using human $\beta 2$ subunit sequences suggested that zebrafish encode two $\beta 2$ homologues [43]. We therefore designed primers to highly conserved sequences within the SH3 and GK protein domains, and performed 5' and 3' RACE-PCR on RNA extracted from embryos aged 1–3 dpf. Using RACE and reverse-transcriptase PCR (RT-PCR), we isolated cDNAs representing two zebrafish $\beta 2$ genes, termed $\beta 2.1$ and $\beta 2.2$ (see Figs. 1, 2 and Additional File 1). The $\beta 2.1$ gene showed a near perfect match with genomic sequences located within Genbank zebrafish chromosome 22 (NC_007133.1), whereas $\beta 2.2$ matched genomic

sequences found on chromosome 2 (NC_007113.1) [43–45]. Alternative splicing occurred at the N-terminus for both genes and within the HOOK domain for $\beta 2.1$ (see Figs. 1 and 2D for descriptions of transcript variants). In the initial RACE analysis for $\beta 2.1$, we recovered four $\beta 2.1_{tv1}$ clones, and five $\beta 2.1_{tv6}$ clones, suggesting these may be the most abundant transcript variants. One clone each was found for the $\beta 2.1_{tv2}$, 3, 4, 5, 7 and 8 transcripts, suggesting they may be less abundant. For $\beta 2.2$, we recovered two $\beta 2.2_{tv1}$ and three $\beta 2.2_{tv2}$ 5' RACE clones. In additional RT-PCR analysis using primers closely flanking the HOOK domain, we confirmed that $\beta 2.1$ transcript variants containing more than one exon between exons 6 and 10 (i.e. $\beta 2.1_{tv3}$, 4, 7 and 8) could be detected in the RNA samples (see Additional file 2). Conversely, we confirmed that no alternative splicing occurred in the $\beta 2.2$ HOOK domain (see Additional File 2). Several 3' RACE clones were of a single variant for $\beta 2.1$




Figure 1
Structure of zebrafish $\beta 2.1$ and $\beta 2.2$ MAGUK genes. Exons are shown as black boxes; introns are not drawn to scale. Alternative splicing is indicated in the N-terminus and HOOK domain-encoding regions. * indicates the site of a premature stop codon in $\beta 2.1$ transcripts that include exon 7. Labeled arrows indicate the locations and names of primers used in RACE and RT-PCR. Human exon structure was adapted from [57]. See Additional File 2A for primer sequences.




Figure 2
Comparative gene structure in teleost $\beta 2$ subunits. A) The exons identified for each of four teleost $\beta 2$ genes are listed vertically with indication of their size in bp. Exons on each horizontal line of the table are homologous in sequence (see later figures for alignments). Exons are color coded to indicate what portion of the 5-domain MAGUK protein (diagrammed in B) they comprise. ^{ATG}, indicates the presence of an initiation codon in the exon. *, indicates the exon contains a premature in-frame termination codon. C) Percent amino acid identity among vertebrate $\beta 2$ subunits within the SH3 through GK sequences. D) The transcript variant numbers assigned to the alternatively spliced transcripts are listed on the left; transcript variant composition is depicted by exon numbers.

and a single variant for $\beta 2.2$, suggesting that no alternative splicing occurred in the GK or C-terminal regions of the genes.

To better assess the conservation of the new genes, we performed a comparative search for orthologous $\beta 2$ sequences in the pufferfish *Tetraodon nigroviridis* and *Fugu rubripes* genomes [45-47]. We identified a single $\beta 2$ subunit gene in each pufferfish genome (Fig. 2A). All four teleost $\beta 2$ genes contained highly conserved SH3 and GK domains characteristic of MAGUK family proteins (Fig. 2B). The size and number of exons comprising these domains are nearly identical in teleosts and mammals (Fig. 2A, yellow and green sections). Pairwise comparisons of amino acid sequences within these core (SH3 through GK) domains, using the most similar transcript variants available, indicated the zebrafish $\beta 2.1$, Fugu and Tetraodon $\beta 2$ genes shared ~87% identity with other vertebrate $\beta 2$ genes (Figs. 2C; see Fig S3 for alignment of core

regions). We extended this analysis to five teleost species by including ESTs or proteins derived from genomic sequences. Three crystallographic studies on mammalian CACNB genes have identified a total of 21 amino acid residues as critical for interaction of the β subunit with the α subunit AID (alpha interaction domain). In our five different teleost species, we observed that all 21 residues were nearly 100% conserved with mammals (see Additional File 3) [48-50].

In contrast, the more divergent zebrafish $\beta 2.2$ core region shares only ~62% amino acid identity (see Additional File 4 for alignment of core regions). The most extreme sequence divergence in the zebrafish $\beta 2.2$ gene occurs at the 5' end of the gene, both 5' of and within the SH3 domain, although high levels of divergence exist throughout the entire protein. We identified a single EST, termed DW608729, from three-spined stickleback as a possible ortholog to the zebrafish $\beta 2.2$ gene. Within the core

domain, DW608729 is 74% identical to zebrafish $\beta 2.2$ (EU301442) but only 52% identical to zebrafish $\beta 2.1$ (EU301434). In addition, DW608729 maps to a genomic contig, AANH01005391.1, which contains sequences homologous to zebrafish $\beta 2.2$ exon 1 (encoding MFC-CGLGHWREQSTY) and $\beta 2.2$ exon 2 (encoding MPP-KKK)(Fig. 3). In the $\beta 2.2$ gene(s), regions of high divergence encompass sequences both within and outside of inferred secondary structures [32]. Nevertheless, 19 of

21 β subunit residues noted for interaction with the α subunit AID are conserved in zebrafish $\beta 2.2$ (see Additional File 4) [48-50]. High sequence divergence in $\beta 2.2$ relative to other vertebrate β genes is reflected in the branch lengths on the phylogeny (Fig. 6). This pattern contrasts with that seen in the $\beta 4$ group, where the two zebrafish paralogs have experienced similar rates of amino acid substitution.

N-terminal domain alignments (Percent identity)

A) Danio $\beta 2.1_{tv1}$ ($\geq 89\%$)

TN $\beta 2$ (BK006355)	1	MYHSCSEESDLGSADSYTSRPSDSVSL	3	EEEDKEAVRREAERQAQAQLDKAKTKP	54
FR $\beta 2$ (BK006351)	1	MYHSCSEESDLGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	54
GA $\beta 2$ (DN723626)	1	MYRSGSEESDLGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	54
DR $\beta 2.1$ (EU301434)	1	MYLSTSEESDLGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKSKP	54

B) Danio $\beta 2.1_{tv5}$ (like Human $\beta 2e$) ($\geq 82\%$)

FR $\beta 2$ (BK006352)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL	3	EEEDKEAVRREAERQAQAQLDKAKTKP	65
TN $\beta 2$ (BK006356)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	65
OL $\beta 2$ (BAAF04030879.1*)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	65
HS $\beta 2$ (NP_963864.1)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	65
XT $\beta 2$ (fgeneshl...)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	65
MM $\beta 2$ (BAD001474)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	65
GG $\beta 2$ (XP_425981.1)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	64
DR $\beta 2.1$ (EU301438)	1	MKATWIRLLKRAKGGHVKSSDACGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKSKP	65

C) Danio $\beta 2.2_{tv1}$ (like Human $\beta 2a$) ($\geq 44\%$)

FR $\beta 2$ (BK006350)	1	MPCCGLGHRRAQLPYGSADSYTSRPSDSVSL	3	EEEDKEAVRREAERQAQAQLDKAKTKP	59
TN $\beta 2$ (BK006354)	1	MPCCGLGHRRAQLPYGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	59
OM $\beta 2$ (ABA46357.1)	1	MPCCGLGHRRAQLPYGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	59
HS $\beta 2$ (NP_000715.2)	1	MCCCLVHRRRVRVSYGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	59
RN $\beta 2$ (NP_446303.1)	1	MCCCLVHRRRVRVSYGSADSYTSRPSDSVSL		EEEDKEAVRREAERQAQAQLDKAKTKP	59
DR $\beta 2.2$ (EU301442)	1	MFCCGLGHWRRRQSTYGSADRQDT		EEEDKEAVRREAERQAQAQLDKAKTKP	45
GA $\beta 2.2$ (AANH01005391.1*)	1	MYCCGLGHWRRVQSTNG--ERQDS		EEEDKEAVRREAERQAQAQLDKAKTKP	39

D) Danio $\beta 2.2_{tv2}$ ($\geq 68\%$)

GA $\beta 2.2$ (DW608729)	1	MPPKKKG--ERQDSVVEE--NAQVQLEKAKYKP	31
DR $\beta 2.2$ (EU301443)	1	MPPKKKGSADRQDTGATEEKEKNAQVQLDKAKYKP	35

E) Additional teleost $\beta 2$ partial transcripts (like Human $\beta 2d$ and $\beta 2c$) ($\geq 90\%$)

HS $\beta 2$ (NP_963887.2)	1A: 1	MVQRDMSKSPPTAAAQAQIQLMELLENVAPAGALGAAAQ	40
HS $\beta 2$ (NP_963865)	1B: 1	MNQSGLDLLKI	12
TN $\beta 2$ (BK006353)	1	SHSRGARRKGRFKGSDGSTSSDITNSL	7
FR $\beta 2$ (BK006349)	1	SHGRGARRKGRFKGSDGSTSSDITNSL	7
OM $\beta 2$ (CA369627)	1	SHGRGARRKGRFKGSDGSTSSDITNSL	7
HS $\beta 2$ (NP_963887.2)	1	SYGKGRARRKGRFKGSDGSTSSDITNSL	7

Figure 3

Sequences and alignments of alternatively spliced 5' regions. (A -E) Alignments show the entire N-terminal portion of the protein (sequences prior to the SH3 domain) for vertebrate $\beta 2$ sequences culled from public databases. An overall percent amino acid identity, calculated for the various pairwise combinations, is listed in parentheses for each peptide alignment. Where appropriate, homology to previously described human transcript variants is indicated [57]. E) In humans, exon 1A or exon 1B is spliced to exon 2A. No exon 1A or 1B-like exons could be identified in the current databases for teleosts, but an exon homologous to human 2A is present in several species. DR, *Danio rerio*; FR, *Fugu rubripes*; GA, *Gasterosteus aculeatus* (three-spined stickleback); GG, *Gallus gallus*; HS, *Homo sapiens*; MM, *Mus musculus*; OL, *Oryzias latipes* (Medaka killifish); OM, *Onchorhynchus mykiss* (trout); RN, *Rattus norvegicus*, TN, *Tetraodon nigroviridis*; XT, *Xenopus tropicalis*. In this and other alignments, conceptual translations were used if protein accession numbers were not available. * denotes a single genomic contig which contains the predicted exons shown. 6 denotes an exon border, numbered with reference to (A-D) zebrafish or (E) pufferfish exons. ##, indicates the location of two conserved cysteine residues which are palmitoylated in human $\beta 2$ proteins.

Alignment of residues in the C terminus of the β subunit genes shows that sequences 3' to the GK domain are not highly conserved even among the teleost species (see Additional File 5). It has previously been noted among multiple vertebrate β subunit genes that the C terminal regions are not highly conserved. Specific functions for C terminal domains are poorly defined, although studies of truncated $\beta 2$ proteins suggest the C terminus could contribute to protein function [35].

Alternative splicing at the N terminus and within the HOOK domain

The zebrafish $\beta 2.1$ and $\beta 2.2$ genes undergo alternative splicing within the N-terminus and within the internal HOOK domain (Figs. 1, 2A, and 2D). The $\beta 2.1$ and $\beta 2.2$ genes each encode two mutually exclusive N termini (Fig. 3). Zebrafish $\beta 2.1$ and $\beta 2.2$, as well as the *Tetraodon* and *Fugu* $\beta 2$ genes, share some 5' exons in common with mammals (Fig. 2A, lines 4 and 8; Fig. 3). In contrast, the $\beta 2.1$ exon 1 sequence could not be found in any mammalian, *Xenopus* or chick databases, but was present in the *Tetraodon* and *Fugu* genomes, suggesting this exon may be specific to teleosts (Fig. 2A, line 6). The $\beta 2.2$ exon 2 was

found only in one other sequence, the stickleback EST DW608729, although the small size of this exon (6 amino acids) could make it difficult to recognize if it had diverged in other species (Fig. 2A, line 5). Additional 5' exons occur in mammals that were not observed in any teleost cDNAs or in genomic sequences available to date, although the limitation of this analysis is that complete genomic sequences are not available for all teleost species examined, and that our data relies in part on genome mining rather than extensive cDNA analysis. However, our data support the hypothesis that each of the teleost $\beta 2$ loci have evolved to contain a unique combination of 5' exons, each of which is predicted to encode an alternative protein N-terminus.

Alternative splicing also occurs internally for one zebrafish $\beta 2$ subunit gene. $\beta 2.1$ encodes three alternatively spliced HOOK domain exons (exons 7, 8 and 9; Figs. 1 and 4). The high conservation of $\beta 2.1$ exons 8 and 9 with mammalian counterparts (Fig. 2A, lines 15 and 16), and exon 8 with several other teleost genes (Fig. 4), suggests that these internal sequences could have functional relevance. The $\beta 2.1$ exon 7, which appears to be

HOOK-domain exons (Percent identity)

A) Danio $\beta 2.1$ exon 7

DR $\beta 2.1$ (EU301437)	164	ADPLPAANQYWETPIHTQSHTLIDYGQFSFLNSLIAHV FV .	
		FAVPCCKTVCLLR LDRW . S . HRSCGCWPELHVSFTHMKLP SDYL	247

B) Danio $\beta 2.1$ exon 8 ($\geq 60\%$)

FN $\beta 2$ (BK006352)	176	ALDIDATDLDPEDNELPVNLRSPKASPNTVMSPLSKEKRM PFFKK	221
TN $\beta 2$ (BK006356)	176	SLDIDATDLDPEDNELPVHLRSPKANPNTVMS SLSREKRM PFFKK	221
GA $\beta 2$ (DN723626)	164	ALDIDATDLDPEDNEFPVNL RSTKASPNTVMSPLAKEKRC P S FRR	209
HS $\beta 2$ (NP_963884)	169	AIDIDATCLDAEENDIPANHRSPKPSANSVTS PHSKEKRM PFFKK	213
DR $\beta 2.1$ (EU301434)	164	AVDIDAAGLDAAEESD PPLHORS PKASPNTVI S PLAREKRM PFFKK	209

C) Danio $\beta 2.1$ exon 9 (86%)

DR $\beta 2.1$ (EU301439)	175	AKQWOKS	181
HS $\beta 2$ (NP_963887.2)	227	AKQKOKS	230

D) Danio $\beta 2.2$ exon 7 (75%)

DR $\beta 2.2$ (EU301442)	151	VQQVKKKS	162
GA $\beta 2$ (DW608729)	137	NQQVKKKP	145

Figure 4
Sequences and alignments of alternatively spliced exons contributing to the HOOK domain. Four distinct exons occur in zebrafish $\beta 2$ transcript variants which differentially join to exon 6 to encode the HOOK domain (see Fig. 1C). (A-C) Three of these exons are alternatively spliced in $\beta 2.1$, whereas the sequence in (D) was the sole sequence found in all $\beta 2.2$ transcripts. Species names are abbreviated as in Figure 3.

unique to zebrafish, includes a premature in-frame stop codon expected to truncate the protein in the HOOK domain (Fig. 2A, line 14). In $\beta 2.1$ transcripts that contain both exon 8 and 9, the reading frame is altered such that a premature in-frame stop codon truncates the protein in exon 10.

No alternative splicing was observed for $\beta 2.2$ transcripts in the HOOK domain; instead, all variants encode a short exon (exon 7) specific to fish that contains several positively charged residues (Fig. 2A, line 13). Likewise, the *Tetraodon* and *Fugu* genomes contain only one recogniza-

bly homologous exon (exon 9), similar to human exon 11 (Fig. 2B, line 15). The variety of $\beta 2$ transcript variants is expected to encode an array of different $\beta 2$ subunit proteins in teleosts.

The large size of 5' introns

Remarkably, the introns that separate the first few 5' exons of the $\beta 2$ loci are among the largest introns known in the zebrafish, pufferfish or human genomes. Each *Tetraodon* and *Fugu* $\beta 2$ locus contains one intron over 10 kb, a size that ranks within the top 5% of the largest introns in pufferfish [51]. In addition, the two pufferfish genes, the stick-

Human $\beta 2$




Figure 5

Large intron sizes in $\beta 2$ genomic loci. Boxes denote the location of exons that encode the 5' region of $\beta 2$ subunit genes. The intervening regions denote the relative sizes of introns, labeled in bp. The scale is expanded for the teleost gene diagrams relative to human. The various N-terminal (grey) exons splice to the solid black exon, which begins with amino acids GSAD... The following genomic contigs were used for the analysis: Human $\beta 2$ (NT_008705.15), *Danio* $\beta 2.2$ (NC_007113), *Danio* $\beta 2.1$ (NC_007133), *Tetraodon* $\beta 2$ (CAAE01015017.1), *Fugu* $\beta 2$ (CAAB01000004.1), *Gasterosteus* $\beta 2$ (AANH01005391.1, using the EST DW608729, which resembles zebrafish $\beta 2.2$).

leback $\beta 2.2$ -like gene, and both zebrafish $\beta 2.2$ and $\beta 2.1$ each contain introns of > 5000 bp that separate 5'-most exons (Fig. 5). In pufferfish, the modal value for intron size is 79 bp, with 75% of introns < 425 bp in length [51]. Given the compact nature of the pufferfish genomes [52], it was not surprising that introns of the human $\beta 2$ locus exceeded the size of those in fish. Nevertheless, the trend of megasized introns in the N-terminus of $\beta 2$ loci extends to the human genome. Three giant introns ranging in size from 60–100 kb each separate the human $\beta 2$ exons in the N-terminus. Thus, when human exon 3 is spliced to exon 7 (beginning GSAD...), the splicing machinery must exclude over 250 kb of intronic sequence. In contrast, introns in the remaining part of the human $\beta 2$ gene averaged 3261 bp in size. In humans, the mean size of introns adjoining coding sequences is 3749 bp, but 75% of introns are smaller than 2609 bp [51,53,54]. A recent analysis of the genomes of *Arabidopsis thaliana*, *Drosophila*, mouse and human indicated that the median sizes of introns separating 5' UTR (non-coding) sequence are significantly larger than introns separating coding sequence [54]. Even so, the median size of human 5' UTR introns (8223 bp, [54]) is still much smaller than the human 5' $\beta 2$ introns, which separate coding sequences.

Divergence of the calcium channel $\beta 2$ genes in zebrafish

We used CLUSTALW software to align the $\beta 2$ subunit core domain amino acid sequences and TreePuzzle and PAUP* to construct phylogenetic trees using maximum likelihood and maximum parsimony, respectively [55]. By using only core domains (~SH3-GK), we minimized differences due to alternative splicing (Fig. 6). The *Fugu* and *Tetraodon* $\beta 2$ subunit genes, zebrafish $\beta 2.1$, zebrafish $\beta 2.2$, and other vertebrate $\beta 2$ genes form a monophyletic group (MLQP = 87%, MPBP < 50%). The sequence of $\beta 2.2$ is substantially more diverged from other vertebrate $\beta 2$ genes than is $\beta 2.1$, both in amino acid replacements (see branch lengths in Fig. 6) and insertions/deletions. Portions of the gene were sufficiently divergent to be essentially random in sequence with respect to all other taxa, and the region 5' of the SH3 domain also contained an 18-amino acid deletion. No convincing region of synteny could be established between any of the teleost $\beta 2$ genes and human $\beta 2$, despite the fact that human chromosome 10 and mouse chromosome 2 share a region of synteny inclusive of the $\beta 2$ genes. Thus, a syntenic approach was not useful to confirm or refute the orthology of zebrafish $\beta 2.1$ or $\beta 2.2$ with mammalian $\beta 2$ genes.

Heterogeneity of calcium channel $\beta 2$ expression in the embryo and adult

To determine whether $\beta 2$ genes are expressed in a stage- or tissue-specific manner in early embryogenesis, we performed RT-PCR on RNA isolated from embryos of several stages in early development. To track the expression pat-

terns of specific transcript variants (Fig. 2D), we used forward primers specific to the 5' exons 1 or 2, and reverse primers in exon 10 (the GK domain) in RT-PCR experiments (Fig. 7). Surprisingly, amplification of $\beta 2.1$ and $\beta 2.2$ transcripts occurred in embryos as young as the 4-cell and 1000-cell stages (Fig. 7A). Since zebrafish zygotic transcription does not initiate until the 10th cell division (~the 1000-cell stage) [56], the presence of mRNA in 4-cell embryos indicates that the transcripts are of maternal origin. The $\beta 2.2$ transcript variant 2 was expressed steadily from the 4-cell stage through 72 hours post-fertilization (hpf). In contrast, $\beta 2.2$ transcript variant 1 showed a pulse of expression in early epiboly stages. $\beta 2.1$ transcript variant 6 was robustly detected from 26 hpf through at least 3 dpf. Other $\beta 2.1$ transcript variants (1 and 2) were detected more sporadically or were undetectable in this assay, consistent with their rare recovery in RACE reactions. Thus, both $\beta 2.1$ and $\beta 2.2$ are expressed from the earliest stages of embryogenesis, but show significant heterogeneity in patterns of transcript variant expression throughout the first three days of embryogenesis.

To determine which transcript variants of the zebrafish $\beta 2$ genes are expressed in the embryonic heart, we performed RT-PCR on RNA extracted from cardiac tissue at 72 hpf (Fig. 7B). These data indicate that only a subset of $\beta 2.1$ and $\beta 2.2$ transcript variants are expressed in the embryonic heart relative to the adult heart. In humans, at least seven $\beta 2$ transcript variants are expressed in the heart, including $\beta 2a$ and $\beta 2e$ [34,35,39]. In zebrafish, the $\beta 2.1$ transcript variant 6 (resembling $\beta 2e$ at the N-terminus) was expressed only in the embryo, whereas the $\beta 2.2$ transcript variant 1 (resembling human $\beta 2a$ at the N-terminus) was expressed in the adult but not embryonic heart.

The expression of $\beta 2$ subunits in several adult organs or tissues was assayed. In most adult tissues, we observed transcript variant-specific patterns of expression for the $\beta 2.1$ and $\beta 2.2$ genes (Fig. 7C). Mutually exclusive expression of $\beta 2.1$ transcript variants 1 and 6 occurred in the heart and brain, respectively. Some other tissues express more than one $\beta 2.1$ transcript variant. Many adult tissues express both $\beta 2.2$ transcript variants 1 and 2, but a few tissues (muscle, gill and skin) expressed only $\beta 2.2$ transcript variant 2. Thus, adult tissues also show significant heterogeneity in expression of $\beta 2$ subunit transcript variants.

Discussion

Teleost calcium channel β subunits and embryonic development

We have identified four new calcium channel $\beta 2$ subunits in teleosts, including two in zebrafish and one gene each in the *Fugu* and *Tetraodon* genomes. Like the mammalian calcium channel β subunits, the four teleost $\beta 2$ subunit genes encode characteristic MAGUK family proteins, with




Figure 6
Phylogeny of $\beta 2$ subunit genes. Phylogenetic tree showing the relationships among $\beta 2$ subunit core domain (SH3 – GK) sequences. Numbers above the nodes indicate maximum likelihood quartet puzzling support values; numbers below the nodes are maximum parsimony bootstrap proportions. "----" indicates a node that was unresolved in the maximum parsimony analysis. The long branch associated with zebrafish $\beta 2.2$ reflects an elevated rate of amino acid substitution throughout the core domain, particularly at the 5' end. See Methods for accession numbers of sequences used.




Figure 7

Expression of $\beta 2$ subunit transcript variants in the embryo and adult. RT-PCR analysis using transcript variant-specific primers (located in the 5' exons 1 or 2 and exon 10) was performed on RNA samples from A) whole embryos at various developmental stages, B) cardiac tissue dissected from *cmic2:GFP* embryos or from adult fish, and C) adult organs and tissues. Expression of a housekeeping gene, *EF1 α* , was used as a control for RNA integrity. In B, 72 hpf or adult RNA reactions were run on single gels, subsequently subdivided to multiple panels for clarity in presentation. Transcript variant numbers are listed to the right of panels; refer to Fig. 1D.

SH3 and GK domains that are highly conserved with other vertebrates. We show that several alternatively spliced transcripts arise from both $\beta 2.1$ and $\beta 2.2$ subunits, which are expressed in the embryo and adult but subject to both temporal and spatial regulation. Only selected transcript variants of each $\beta 2$ subunit gene are expressed in the embryonic heart, and these differ from adult transcript variants. Thus, the heterogeneity of β subunits and their transcript variants in teleost species is extensive.

Gene structure of teleost calcium channel $\beta 2$ subunits

Alternative splicing occurs within the N-terminus as well as internally in the $\beta 2.1$ and $\beta 2.2$ subunit genes. Notably, each zebrafish $\beta 2$ subunit gene encodes at least two mutually exclusive 5' exons, each with a separate translation initiation site. Each gene has one 5' exon conserved with mammals, and one 5' exon unique to fish. Similarly, some of the internal alternatively spliced exons are similar to those in mammals, while others are unique to teleosts. This variety of transcript variants potentially encodes an array of different $\beta 2$ subunit proteins. Previous studies have demonstrated that 5' variation due to alternative splicing can be functionally significant in β subunits. When expressed in HEK 293 cells, human 5' $\beta 2$ subunit variants $\beta 2a$ and $\beta 2e$ (starting with human exons 4 or 6, respectively) showed differential sub-cellular localization compared to transcript variants $\beta 2b$, $\beta 2c$ and $\beta 2d$ (starting with human exons 4, 1 and 2, respectively) [57]. Moreover, the five human 5' $\beta 2$ subunit variants differentially affected open probability, peak current and availability of L-type channels [57,58]. Likewise, in *Xenopus* oocyte

expression studies, human $\beta 4$ subunits with alternatively spliced N-termini showed functionally distinct electrophysiological properties [59,60]. Domain swapping experiments further indicate the functional significance of the N-terminal sequences in β subunit proteins. Replacing the $\beta 1b$ N-terminus with $\beta 2a$ N-terminus created a chimeric β protein with slow inactivation kinetics [61]. Conversely, replacing $\beta 2a$ N-terminus with the non-palmitoylated $\beta 3$ N-terminus created a chimeric β protein with $\beta 3$ -like inactivation kinetics [62]. Post-translational modifications such as phosphorylation or palmitoylation, present on exons near the N-terminus, may account for some of the functional diversity, although this functional relationship has been experimentally demonstrated for only a few genes to date [63,64]. Intriguingly, a recent report indicates that the length of the $\beta 2$ N-terminus is, independent of sequence, an important factor in mediating the magnitude of channel modulation [58]. Further functional studies using the zebrafish model system will be necessary to determine whether the array of zebrafish $\beta 2$ subunit proteins indeed have different biochemical functions within the cell.

A curious feature of at least five species of teleost $\beta 2$ subunit genes, as well as the human $\beta 2$ subunit gene, is that some of the largest introns of the genome separate the several short, alternatively spliced 5' coding exons. Do these large introns have any adaptive significance for $\beta 2$ subunit biology? In several instances, introns have been shown to incorporate enhancer or repressor elements that influence transcription [65]. Expanding this idea, we propose that

the unusual $\beta 2$ gene structure may provide a mechanism for independent cis-regulation of each transcript variant; that is, sequences within individual introns might independently direct the tissue and temporally-specific expression of transcript variants utilizing the associated ATG-containing 5' exon. This hypothesis is supported by the observation that transcript variant-specific patterns of expression do in fact occur in both embryo and adult. However, it is increasingly becoming appreciated that, beyond regulatory elements, introns can impact mRNA metabolism in a number of other ways. Potential effects include modulating transcription rates by regulating DNA accessibility, modulating editing and polyadenylation of the pre-mRNA, and affecting nuclear export, translation and mRNA decay rates [65,66].

The inclusion of different subsets of 5' exons (and their associated introns) among each of the pufferfish or zebrafish $\beta 2$ subunit genes is interesting from an evolutionary standpoint. A canonical view is that variations in protein structure and function form the basis of evolutionary innovation and phenotypic divergence. However, as more genomes are sequenced and annotated, it is becoming evident that alternative splicing substantially increases the proteome in many species. (For example, about 40–50% of human and mouse genes contain alternative promoters [67]). A corollary to this hypothesis is that the regulatory circuitry regulating alternative splicing or the expression of particular transcript variants is also an important source of evolutionary innovation [67,68]. The mechanisms by which new introns arise during evolution, and the impact of large intron size per se on cis-regulation or other adaptive phenomena, remain actively debated areas of research [69-71].

The evolutionary origin of the duplicated calcium channel $\beta 2$ subunit genes

Zebrafish $\beta 2$ duplication may be the consequence of a genome-wide duplication thought to have occurred in teleosts approximately 300 million years ago [72-74]. Alternatively, duplication could be the result of an ancient region-specific duplication. The $\beta 2$ subunit gene does not appear to be duplicated in either the *Fugu* or *Tetraodon* genomes. The existence of single $\beta 2$ subunit homologs was of itself not particularly surprising since phylogeny and synteny data suggest that the common ancestor of zebrafish and pufferfish underwent a large-scale gene or complete genome duplication event [75]. Subsequently, pufferfish may have lost many duplicates that were retained in the zebrafish [75]. Data from the other β gene paralog groups ($\beta 1$ and $\beta 4$) are equivocal, as zebrafish have one described copy of $\beta 1$ and two copies of $\beta 4$ [41,76]. The fate of most duplicated genes is the accumulation of degenerative mutations in coding or regulatory regions that leads to gene loss or silencing [73]. Alterna-

tively, genes may acquire separable patterns of expression or separable functions that then require the maintenance of functional copies of both genes in the genome (subfunctionalization) [73]. Since the two zebrafish $\beta 2$ genes are both robustly but differentially expressed, our data are more consistent with subfunctionalization.

Developmental expression patterns of calcium channel $\beta 2$ subunit transcript variants

Calcium is an important signal in the embryo even prior to gastrulation [77]. Calcium gradients, waves and pulses have been described in the blastula and early gastrula that may represent key pattern forming events. The mechanism underlying these calcium signals, and whether they encompass voltage-gated calcium channels, is not clear. Surprisingly, both $\beta 2$ subunits show maternal as well as the early zygotic expression of several transcript variants. Our recent study of two zebrafish $\beta 4$ subunits, which are also expressed maternally and zygotically in the gastrulating embryo, indicates that these genes are essential for normal epiboly. Initially, we predicted that β subunits might be involved in the gastrulating embryo voltage-gated calcium channel-related functions. However, our study showed that the zebrafish $\beta 4$ subunits, at least, operate independently of voltage-gated calcium channels in mediating epiboly [76].

In addition to possible early roles, we find that expression of $\beta 2.1$ or $\beta 2.2$ is under strong temporal control in the post-gastrula embryo and in adult tissues. Of particular interest, given the cardiac lethality observed in mouse $\beta 2$ -deficient embryos, is the observation of differential subsets of $\beta 2.1$ and $\beta 2.2$ transcript variants expressed in the embryonic and adult heart. This observation supports the hypothesis, previously proposed by others, that the heterogeneity of β subunit transcript variant expression within the heart may provide a mechanism for fine-tuning the cardiac voltage-activated current as the organism progresses through embryonic, juvenile and adult stages [38,39]. In addition, the expression levels of particular $\beta 2$ transcript variants has also been linked to pathophysiology of heart failure in humans [78-80]. Clearly, the first step in interpreting the function of β subunits in any model system is simply to understand when and where the various transcript variants are expressed.

Conclusion

The primary novel findings of this study are as follows: 1) We have cloned two new $\beta 2$ subunit genes in zebrafish, one of which is phylogenetically quite divergent in amino acid sequence. They are classified as MAGUK proteins on the basis of high conservation with mammalian MAGUK core regions. 2) Alternative splicing occurs at the $\beta 2$ N-termini and internally. A comparative analysis showed that a subset of 5' exons present in several teleosts species is

shared with mammals, while a different subset appears to be unique to teleosts. 3) Some of the largest introns in the human or teleost genomes separate the small, alternatively spliced 5' exons of $\beta 2$ genes, leading us to hypothesize that β subunit transcript variant expression may be under independent cis-regulatory control. 4) The zebrafish $\beta 2$ genes are expressed maternally and zygotically in the gastrulating embryo and show strong evidence of temporal and spatial regulation in embryogenesis and in adults. 5) Only a subset of $\beta 2$ subunit transcript variants is expressed in the embryonic heart, and they differ from those expressed in the adult heart. In sum, this work provides the groundwork for a study of functional aspects of $\beta 2$ subunit biology. The high degree of temporal and spatial control of β subunits, combined with recently identified non-canonical functions for particular β subunits or transcript variants, suggests that a functional analysis of these genes would provide intriguing insight.

Methods

Zebrafish strains and care

This study used the WIK zebrafish strain. Zebrafish care was provided in accordance with animal care policies of Colorado State University. Embryos were staged as described [81].

Isolation of zebrafish $\beta 2$ cDNAs

Total RNA was prepared from embryos aged 24–72 hpf using Trizol (Invitrogen, Carlsbad, CA) as per manufacturer's protocol. Poly(A)⁺ mRNA was prepared using magnetic beads (μ MACS mRNA kit Miltenyi Biotec), and resuspended in RNase free dH₂O at a concentration of 1 mg/ml. RNA was stored in aliquots as an ethanol precipitate at -80 degrees Celsius. RACE Ready cDNA was created using the SMART RACE cDNA synthesis kit (Clontech, Palo Alto, CA). RACE reactions were carried out as per manufacturer's protocol using gene-specific primers designed to hybridize within the SH3 and GK domain encoding regions of β subunit genes identified by BLAST searches of the zebrafish genome (Fig. 1 and Additional File 2A). RACE PCR products were separated in ethidium bromide-stained agarose gels, excised and purified over a silica matrix column (Zymoclean Gel DNA Recovery Kit, Zymo Research). cDNA fragments were cloned into pCR 2.1-TOPO using the TOPO TA cloning method (Invitrogen, Carlsbad, CA).

Pufferfish sequence identification and amino acid alignments

We used human and zebrafish $\beta 2$ sequences to identify homologous *Fugu* and *Tetraodon* $\beta 2$ sequences by a multi-step process. First, we performed protein vs. translated DNA (TBLASTN) searches using human or zebrafish $\beta 2$ sequences against the NCBI, Sanger Centre, UCSC Genome Bioinformatics and pufferfish genome databases

[43–47]. This process identified the best matches among the *Fugu* or *Tetraodon* predicted proteins derived from contig assemblies and other sequences available for pufferfish. The homologous predicted *Fugu* or *Tetraodon* proteins were re-tested by TBLASTN against mammalian Refseq proteins to confirm they were more similar to $\beta 2$ subunit genes than other β subunit family members. Next, we used the homologous *Fugu* or *Tetraodon* sequences in BLAT searches of UCSC Genome Bioinformatics database sequences to identify the genomic location of *Fugu* or *Tetraodon* $\beta 2$ exons 3–14 and to determine conserved splice donor and splice acceptor sites [45]. To identify the short N terminal exons (exons 1 and 2) or search for alternative HOOK domain exons in pufferfish, we used zebrafish or human exons in 1) BLAT searches of UCSC Genome Bioinformatics pufferfish (*Fugu* or *Tetraodon*) sequences, 2) MEGABLAST, peptide motif and BLASTP searches of NCBI, and 3) BLAST or BLAT searches of the *Fugu* Genome Browser, the *Tetraodon* Genome Browser, or the Sanger Center Ensemble Genomic databases [43–47]. Exons were numbered according to their sequential location in the genome, and probable splice donor and splice acceptor sites were identified for each exon. The pufferfish sequences were submitted to the Third Party Annotation database of NCBI. Additional teleost sequences from trout, medaka, and stickleback were obtained by searching the NCBI EST or WGS databases, or from JGI [82].

Phylogenetic analysis

Amino acid sequences representing the β subunit "core" (i.e., from the amino acids "GSAD" just prior to the SH3 domain through the end of the GK domain) were aligned using ClustalW, varying gap opening and extension parameters; regions for which reliable homology could not be established because of indels were excluded from analysis [83]. Portions of the zebrafish $\beta 2.2$ gene were sufficiently divergent in sequence from all other genes (including those from *Drosophila* and mosquito) to be essentially random. To minimize the phylogenetic analytical problems associated with such extreme rate variation among lineages, we excluded regions of the alignment in which zebrafish $\beta 2.2$ contained autapomorphic amino acid substitutions for ≥ 3 adjacent amino acid positions. Such exclusions resulted in an alignment of 252 amino acids. Maximum likelihood analyses were implemented in TreePuzzle [84] using the Müller & Vingron (VT) model of amino acid substitution with a mixed model of rate heterogeneity; amino acid frequencies, the gamma distribution parameter alpha, and the proportion of invariant sites were estimated from the data [85]. Nodal support was assessed using quartet puzzling. Equally-weighted maximum parsimony analyses were carried out using PAUP* [55]. A heuristic search was performed with 25 random additional replicates and TBR branch-swapping.

Bootstrap proportions for clades were assessed with 1,000 pseudo-replicates. Resulting trees were rooted with *Drosophila* and mosquito. Teleost sequences other than pufferfish and zebrafish were not included because they only span a portion of the core domain.

Reverse transcription assays

As an RNA source, we used whole embryos, or individual organs or tissues dissected from adults aged approximately 1 year. Cardiac tissue from embryos was isolated on the basis of its GFP expression from the *cmlc2:GFP* transgenic line, in which the *cmlc2* promoter drives expression of GFP only in the heart [86]. RNA was extracted using the Trizol method and stored at -80°C. RT-PCR was performed for $\beta 2.1$ and $\beta 2.2$ genes using the Access RT-PCR System (Promega) using high-fidelity polymerases. The thermal cycling program was as follows: 48°C for 45 minutes (reverse transcription), followed by 94°C for 2 minutes (activation of PCR enzyme) and 40 cycles of 94°C for 30 seconds, 60°C for 1 minute, and 68°C for 2 minutes. A final step of 68°C for 7 minutes allowed for a final extension. Electrophoresis of the final product was performed on a 2% agarose gel containing ethidium bromide. Gels were imaged using a digital camera and imaging software (Scion Corporation, Frederick, MD).

Accession numbers

The following Genbank sequences were used for the phylogenetic analysis: CACNB1 genes: NP_000714.3, NP_660099.1, NP_059042.1, AAH76523.1; CACNB2 genes: EU301442 (mRNA), EU301434 (mRNA), NP_963864.1, NP_075605.1, XP_855731.1, NP_446303.1, XM_425981.1, BK006353, BK006349; CACNB3 genes: NP_000716.2, XP_543689.2, NP_031607.1, NP_036960.1, AAA75519.1; CACNB4 genes: ABU93250.1, ABU93253.1, XP_851697.1, NP_000717.2, NP_001032176.1, NP_001001733.1, A45982, CAG08264.1; other CACNB genes: NP_995685.1, XP_317433.2, NP_491193.1. In addition, if no appropriate sequence was available in Genbank, we used sequences from JGI database [82], including *X. tropicalis* genome assembly version 4.1, entitled FGGENESH1_PM.C_SCAFFOLD_1068000001 (*Xenopus* CACN β 2); C_scaffold_227000005 (*Xenopus* CACN β 4); and e_gw1.686.1.1 (*Xenopus* CACN β 1). From the *Fugu rubripes* genome assembly version 4.0, we used fgh5_pm.C_scaffold_38000036 (*Fugu* CACN β 4). The JGI sequences are available upon request from the authors.

Abbreviations

Reverse-transcriptase polymerase chain reaction (RT-PCR), membrane-associated guanylate-kinase protein (MAGUK), voltage-dependent L-type Ca²⁺ channel (V-LTCC), rapid amplification of cDNA ends (RACE), base-

pairs (bp), cardiac myosin light chain (cmlc), green fluorescent protein (GFP), days post-fertilization (dpf), hours post-fertilization (hpf), maximum parsimony bootstrap proportion (MPBP), maximum likelihood quartet puzzling (MLQP).

Authors' contributions

AS generated the cDNA used for RACE experiments; AME, CAM and DMG carried out the RACE and RT-PCR cloning experiments; AME carried out the RT-PCR expression experiments; DBG and DMG did the *Fugu*, *Tetraodon* and other teleost bioinformatics and alignments; RLM contributed the phylogenetic analysis; DMG wrote the manuscript with comments from all authors. All authors read and approved the final manuscript.

Additional material

Additional file 1

Amino acid sequences of 5' exons. The alternatively spliced 5' exons listed here occur as the first or second exon in the transcript variant. They occur in the N-terminus of the protein, prior to the beginning of the SH3 domain. For convenient comparison, human exons are co-listed, using data from [57]. See Fig. 1D for description of the transcript variants.

Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-2199-9-38-S1.ppt]

Additional file 2

RT-PCR analysis and list of primers. (A) A list of primers used for RT-PCR analysis, and their locations (see also Figure 1 for primer locations). (B) A representative RT-PCR analysis of *Danio rerio* $\beta 2.1$ and $\beta 2.2$ transcripts. In lanes 1–3, a single band of the expected sizes amplified for $\beta 2.2$, consistent with a lack of alternative splicing in the HOOK domain (spanning exons 5–8 in $\beta 2.2$). In lanes 6 and 7, products consistent with $\beta 2.1_{tw1}$ (697 bp) or $\beta 2.1_{tw6}$ (580 bp) tended to amplify robustly. Potentially, larger amplicons from less abundant transcript variants may be out-competed in these reactions. We therefore tested primers closely flanking the HOOK domain (spanning exons 5–10 in $\beta 2.1$). In lane 8, products consistent with all HOOK domain transcript variants were observed (a 647 bp band, a 415/394 bp doublet band, and a 280 bp band). Cloning and sequencing of these products of Lane 8 confirmed that the bands represented are the transcripts indicated. Lanes 4 and 9 are positive controls using primers to the house-keeping gene *EF1 α* (220 bp). Lanes 5 and 10 are negative controls. Reactions were run on two gels, as shown (some lanes have been removed). (C) For each primer pair, the expected sizes of PCR amplicons and the exons that comprise them are listed.

Click here for file

[http://www.biomedcentral.com/content/supplementary/1471-2199-9-38-S2.ppt]

Additional file 3

Amino acid alignments of zebrafish $\beta 2.1$ subunit "core" regions with other vertebrate genes. The light grey bars indicate the SH3 domain, black bars the HOOK domain, and dark grey bars the GK domain. ▼ denotes an exon border (exons numbered according to Danio rerio $\beta 2.1_{tw1}$, which lacks the alternatively spliced exons 8 and 9). The plus signs (+) indicate the alpha binding pocket residues suggested by crystallographic studies to contribute side-chain contacts with the AID domain [48-50]. Note that the zebrafish isoleucine at residue 325 differs from the conserved mammalian valine, but isoleucine occurs in this position in human CACNB3 and other CACNB genes [49]. Likewise the zebrafish threonine at residue 329 is either serine or threonine in the mammalian CACNB genes [50]. The crystal structure predicts the sequences PYDVVP (just 5' to the GK domain), conserved in all genes shown, forms a beta strand that folds back to interact with the SH3 domain [50]. See Fig. 1D for description of exons present in each transcript variant.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2199-9-38-S3.ppt>]

Additional file 4

Amino acid alignments of the more divergent zebrafish $\beta 2.2$ subunit "core" regions with a potential homolog in three-spined stickleback. The light grey bars indicate the SH3 domain, black bars the HOOK domain, and dark grey bars the GK domain. ▼ denotes an exon border (exons numbered according to Danio rerio $\beta 2.2_{tw1}$). The plus signs (+) indicate the alpha binding pocket residues that contribute side-chain contacts with the AID domain of the α subunit, whereas the minus sign (-) marks a valine residue that was not conserved in either species. Grey plus signs represent mammalian residues conserved in zebrafish but not stickleback [48-50]. Note that DW608729 EST is only a partial mRNA sequence which does not span the complete core region. See Fig. 1D for description of exons present in each transcript variant.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2199-9-38-S4.ppt>]

Additional file 5

Amino acid alignments of the C-terminal regions of several teleost $\beta 2$ clones with human $\beta 2$. The C terminal residues aligned here reside 3' to the GK domain, and are typically encoded by the final two exons of the $\beta 2$ gene. For stickleback, translations of available EST sequences mapping to the C terminal portion of the $\beta 2$ gene were included, although they may not represent the full sequences encoded by the final two exons.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2199-9-38-S5.ppt>]

Acknowledgements

The authors thank Dr. Geoff Burns for the *cmlc2:GFP* transgenic fish, Dr. William Horne for reading the manuscript, and several talented undergraduates in the Garrity lab for their support. Supported by NSF grant IOS 0719083 to DMG.

References

- Hidalgo P, Neely A: **Multiplicity of protein interactions and functions of the voltage-gated calcium channel beta-subunit.** *Cell Calcium* 2007, **42(4-5)**:389-396.
- Weissgerber P, Held B, Bloch W, Kaestner L, Chien KR, Fleischmann BK, Lipp P, Flockerzi V, Freichel M: **Reduced Cardiac L-Type Ca^{2+} Current in Cav $\beta 2$ - Embryos Impairs Cardiac**

Development and Contraction With Secondary Defects in Vascular Maturation. *Circ Res* 2006, **99(7)**:749-757.

- Frey N, McKinsey TA, Olson EN: **Decoding calcium signals involved in cardiac growth and function.** *Nat Med* 2000, **6(11)**:1221-1227.
- Mesaali N, Nakamura K, Zvaritch E, Dickie P, Dziak E, Krause KH, Opas M, MacLennan DH, Michalak M: **Calreticulin is essential for cardiac development.** *J Cell Biol* 1999, **144(5)**:857-868.
- Linask KK, Han MD, Artman M, Ludwig CA: **Sodium-calcium exchanger (NCX-1) and calcium modulation: NCX protein expression patterns and regulation of early heart development.** *Dev Dyn* 2001, **221**:249-264.
- Ebert AM, Hume GL, Warren KS, Cook NP, Burns CG, Moohideen MA, Siegal G, Yelon D, Fishman M, Garrity DM: **Calcium extrusion is critical for cardiac morphogenesis and rhythm in embryonic zebrafish hearts.** *Proc Natl Acad Sci U S A* 2005, **102(49)**:17705-17710.
- Rottbauer W, Baker K, Wo ZG, Moohideen MA, Cantiello HF, Fishman MC: **Growth and function of the embryonic heart depend upon the cardiac-specific L-type calcium channel alpha subunit.** *Dev Cell* 2001, **1**:265-275.
- Reamon-Buettner SM, Spanel-Borowski K, Borlak J: **Bridging the gap between anatomy and molecular genetics for an improved understanding of congenital heart disease.** *Ann Anat* 2006, **188(3)**:213-220.
- Feske S, Giltman J, Dolmetsch R, Staudt LM, Rao A: **Gene regulation mediated by calcium signals in T lymphocytes.** *Nat Immunol* 2001, **2(4)**:316-324.
- Porter GA Jr., Makuck RF, Rivkees SA: **Intracellular calcium plays an essential role in cardiac development.** *Dev Dyn* 2003, **227(2)**:280-290.
- Dalton S, Takahashi SX, Miriyala J, Colecraft HM: **A single Cav β can reconstitute both trafficking and macroscopic conductance of voltage-dependent calcium channels.** *J Physiol* 2005, **567(Pt 3)**:757-769.
- Arikkath J, Campbell KP: **Auxiliary subunits: essential components of the voltage-gated calcium channel complex.** *Curr Opin Neurobiol* 2003, **13(3)**:298-307.
- Brette F, Leroy J, Le Guennec JY, Salle L: **Ca^{2+} currents in cardiac myocytes: Old story, new insights.** *Prog Biophys Mol Biol* 2006, **91(1-2)**:1-82.
- Ludwig A, Flockerzi V, Hofmann F: **Regional expression and cellular localization of the alpha 1 and beta subunit of high voltage-activated calcium channels in rat brain.** *J Neurosci* 1997, **17(4)**:1339-1349.
- Murakami M, Fleischmann B, De Felipe C, Freichel M, Trost C, Ludwig A, Wissenbach U, Schwegler H, Hofmann F, Hescheler J, Flockerzi V, Cavalie A: **Pain perception in mice lacking the beta3 subunit of voltage-activated calcium channels.** *J Biol Chem* 2002, **277(43)**:40342-40351.
- Reimer D, Huber IG, Garcia ML, Haase H, Striessnig J: **beta subunit heterogeneity of L-type Ca^{2+} channels in smooth muscle tissues.** *FEBS Lett* 2000, **467(1)**:65-69.
- Mikami A, Imoto K, Tanabe T, Niidome T, Mori Y, Takeshima H, Narumiya S, Numa S: **Primary structure and functional expression of the cardiac dihydropyridine-sensitive calcium channel.** *Nature* 1989, **340(6230)**:230-233.
- Welling A, Ludwig A, Zimmer S, Klugbauer N, Flockerzi V, Hofmann F: **Alternatively spliced IS6 segments of the alpha 1C gene determine the tissue-specific dihydropyridine sensitivity of cardiac and vascular smooth muscle L-type Ca^{2+} channels.** *Circ Res* 1997, **81(4)**:526-532.
- Seisenberger C, Specht V, Welling A, Platzer J, Pfeifer A, Kuhbandner S, Striessnig J, Klugbauer N, Feil R, Hofmann F: **Functional embryonic cardiomyocytes after disruption of the L-type alpha 1C (Cav1.2) calcium channel gene in the mouse.** *J Biol Chem* 2000, **275(50)**:39193-39199.
- Bohn G, Moosmang S, Conrad H, Ludwig A, Hofmann F, Klugbauer N: **Expression of T- and L-type calcium channel mRNA in murine sinoatrial node.** *FEBS Lett* 2000, **481(1)**:73-76.
- Lipscombe D, Helton TD, Xu W: **L-type calcium channels: the low down.** *J Neurophysiol* 2004, **92(5)**:2633-2641.
- Takimoto K, Li D, Nerbonne JM, Levitan ES: **Distribution, splicing and glucocorticoid-induced expression of cardiac alpha 1C and alpha 1D voltage-gated Ca^{2+} channel mRNAs.** *J Mol Cell Cardiol* 1997, **29(11)**:3035-3042.

23. Gregg RG, Messing A, Strube C, Beurg M, Moss R, Behan M, Sukhareva M, Haynes S, Powell JA, Coronado R, Powers PA: **Absence of the beta subunit (cchb1) of the skeletal muscle dihydropyridine receptor alters expression of the alpha 1 subunit and eliminates excitation-contraction coupling.** *Proc Natl Acad Sci U S A* 1996, **93(24)**:13961-13966.
24. Ball SL, Powers PA, Shin HS, Morgans CW, Peachey NS, Gregg RG: **Role of the beta(2) subunit of voltage-dependent calcium channels in the retinal outer plexiform layer.** *Invest Ophthalmol Vis Sci* 2002, **43(5)**:1595-1603.
25. Namkung Y, Smith SM, Lee SB, Skrypnik NV, Kim HL, Chin H, Scheller RH, Tsien RW, Shin HS: **Targeted disruption of the Ca2+ channel beta3 subunit reduces N- and L-type Ca2+ channel activity and alters the voltage-dependent activation of P/Q-type Ca2+ channels in neurons.** *Proc Natl Acad Sci U S A* 1998, **95(20)**:12010-12015.
26. Murakami M, Aoyama M, Suzuki T, Sasano H, Nakayama S, Iijima T: **Genetic characterization of a new splice variant of the beta2 subunit of the voltage-dependent calcium channel.** *Mol Cell Biochem* 2003, **254(1-2)**:217-225.
27. Cork RJ, Namkung Y, Shin HS, Mize RR: **Development of the visual pathway is disrupted in mice with a targeted disruption of the calcium channel beta(3)-subunit gene.** *J Comp Neurol* 2001, **440(2)**:177-191.
28. Burgess DL, Jones JM, Meisler MH, Noebels JL: **Mutation of the Ca2+ channel beta subunit gene Cchb4 is associated with ataxia and seizures in the lethargic (lh) mouse.** *Cell* 1997, **88(3)**:385-392.
29. Walker D, Bichet D, Campbell KP, De Waard M: **A beta 4 isoform-specific interaction site in the carboxyl-terminal region of the voltage-dependent Ca2+ channel alpha 1A subunit.** *J Biol Chem* 1998, **273(4)**:2361-2367.
30. Herlitze S, Xie M, Han J, Hummer A, Melnik-Martinez KV, Moreno RL, Mark MD: **Targeting mechanisms of high voltage-activated Ca2+ channels.** *J Bioenerg Biomembr* 2003, **35(6)**:621-637.
31. Dolphin AC: **Beta subunits of voltage-gated calcium channels.** *J Bioenerg Biomembr* 2003, **35(6)**:599-620.
32. Hanlon MR, Berrow NS, Dolphin AC, Wallace BA: **Modelling of a voltage-dependent Ca2+ channel beta subunit as a basis for understanding its functional properties.** *FEBS Lett* 1999, **445(2-3)**:366-370.
33. te Velthuis AJ, Admiraal JF, Bagowski CP: **Molecular evolution of the MAGUK family in metazoan genomes.** *BMC Evol Biol* 2007, **7**:129.
34. Foell JD, Balijepalli RC, Delisle BP, Yunker AM, Robia SL, Walker JW, McEnery MW, January CT, Kamp TJ: **Molecular heterogeneity of calcium channel beta-subunits in canine and human heart: evidence for differential subcellular localization.** *Physiol Genomics* 2004, **17(2)**:183-200.
35. Harry JB, Kobrinisky E, Abernethy DR, Soldatov NM: **New short splice variants of the human cardiac Cavbeta2 subunit: redefining the major functional motifs implemented in modulation of the Cav1.2 channel.** *J Biol Chem* 2004, **279(45)**:46367-46372.
36. Hibino H, Pironkova R, Onwumere O, Rousset M, Charnet P, Hudspeith AJ, Lesage F: **Direct interaction with a nuclear protein and regulation of gene silencing by a variant of the Ca2+-channel beta 4 subunit.** *Proc Natl Acad Sci U S A* 2003, **100(1)**:307-312.
37. Haase H, Pfitzmaier B, McEnery MW, Morano I: **Expression of Ca(2+) channel subunits during cardiac ontogeny in mice and rats: identification of fetal alpha(1C) and beta subunit isoforms.** *J Cell Biochem* 2000, **76(4)**:695-703.
38. Acosta L, Haase H, Morano I, Moorman AF, Franco D: **Regional expression of L-type calcium channel subunits during cardiac development.** *Dev Dyn* 2004, **230(1)**:131-136.
39. Chu PJ, Larsen JK, Chen CC, Best PM: **Distribution and relative expression levels of calcium channel beta subunits within the chambers of the rat heart.** *J Mol Cell Cardiol* 2004, **36(3)**:423-434.
40. Vance CL, Begg CM, Lee WL, Haase H, Copeland TD, McEnery MW: **Differential expression and association of calcium channel alpha1B and beta subunits during rat brain ontogeny.** *J Biol Chem* 1998, **273(23)**:14495-14502.
41. Zhou W, Saint-Amant L, Hirata H, Cui WW, Sprague SM, Kuwada JY: **Non-sense mutations in the dihydropyridine receptor beta1 gene, CACNB1, paralyze zebrafish relaxed mutants.** *Cell Calcium* 2006, **39(3)**:227-236.
42. Pelster B, Burggren WW: **Disruption of hemoglobin oxygen transport does not impact oxygen-dependent physiological processes in developing embryos of zebra fish (Danio rerio).** *Circ Res* 1996, **79(2)**:358-362.
43. **The National Center for Biotechnology Information Entrez Database** [<http://www.ncbi.nlm.nih.gov/sites/gquery>]
44. **The Sanger Center Ensembl Databases** [<http://www.ensembl.org/index.html>]
45. **The UCSC Genome Bioinformatics Databases** [<http://genome.ucsc.edu/>]
46. **The Fugu Genomce Project Database** [<http://www.fugu-sg.org/>]
47. **The Tetraodon Genome Browser** [<http://www.genoscope.cns.fr/externe/tetranew/>]
48. Van Petegem F, Clark KA, Chatelain FC, Minor DL Jr.: **Structure of a complex between a voltage-gated calcium channel beta-subunit and an alpha-subunit domain.** *Nature* 2004, **429(6992)**:671-675.
49. Opatowsky Y, Chen CC, Campbell KP, Hirsch JA: **Structural analysis of the voltage-dependent calcium channel beta subunit functional core and its complex with the alpha 1 interaction domain.** *Neuron* 2004, **42(3)**:387-399.
50. Chen YH, Li MH, Zhang Y, He LL, Yamada Y, Fitzmaurice A, Shen Y, Zhang H, Tong L, Yang J: **Structural basis of the alpha 1-beta subunit interaction of voltage-gated Ca2+ channels.** *Nature* 2004, **429(6992)**:675-680.
51. Aparicio S, Chapman J, Stupka E, Putnam N, Chia JM, Dehal P, Christoffels A, Rash S, Hoon S, Smit A, Gelpke MD, Roach J, Oh T, Ho IY, Wong M, Detter C, Verhoeve F, Predki P, Tay A, Lucas S, Richardson P, Smith SF, Clark MS, Edwards YJ, Doggett N, Zharkikh A, Tavtigian SV, Pruss D, Barnstead M, Evans C, Baden H, Powell J, Glusman G, Rowen L, Hood L, Tan YH, Elgar G, Hawkins T, Venkatesh B, Rokhsar D, Brenner S: **Whole-genome shotgun assembly and analysis of the genome of Fugu rubripes.** *Science* 2002, **297(5585)**:1301-10. Epub 2002 Jul 25..
52. Venkatesh B, Yap WH: **Comparative genomics using fugu: a tool for the identification of conserved vertebrate cis-regulatory elements.** *Bioessays* 2005, **27(1)**:100-107.
53. Sakharkar MK, Chow VT, Kanguane P: **Distributions of exons and introns in the human genome.** *In Silico Biol* 2004, **4(4)**:387-393.
54. Hong X, Scofield DG, Lynch M: **Intron size, abundance, and distribution within untranslated regions of genes.** *Mol Biol Evol* 2006, **23(12)**:2392-404. Epub 2006 Sep 15..
55. Swofford DL: **PAUP*: phylogenetic analysis using parsimony (* and other methods). Ver. 4.0.** Sinauer Associates, Sunderland, MA; 1998.
56. Kane DA, Kimmel CB: **The zebrafish midblastula transition.** *Development* 1993, **119(2)**:447-456.
57. Takahashi SX, Mittman S, Colecraft HM: **Distinctive modulatory effects of five human auxiliary beta2 subunit splice variants on L-type calcium channel gating.** *Biophys J* 2003, **84(5)**:3007-3021.
58. Herzig S, Khan IF, Grundemann D, Matthes J, Ludwig A, Michels G, Hoppe UC, Chaudhuri D, Schwartz A, Yue DT, Hüllin R: **Mechanism of Ca(v)1.2 channel modulation by the amino terminus of cardiac beta2-subunits.** *Faseb J* 2007, **21(7)**:1527-38. Epub 2007 Feb 8..
59. Helton TD, Horne WA: **Alternative splicing of the beta 4 subunit has alpha 1 subunit subtype-specific effects on Ca2+ channel gating.** *J Neurosci* 2002, **22(5)**:1573-1582.
60. Helton TD, Kojetin DJ, Cavanagh J, Horne WA: **Alternative splicing of a beta4 subunit proline-rich motif regulates voltage-dependent gating and toxin block of Cav2.1 Ca2+ channels.** *J Neurosci* 2002, **22(21)**:9331-9339.
61. Olcese R, Neely A, Qin N, Wei X, Birnbaumer L, Stefani E: **Coupling between charge movement and pore opening in vertebrate neuronal alpha 1E calcium channels.** *J Physiol* 1996, **497(Pt 3)**:675-686.
62. Qin N, Olcese R, Zhou J, Cabello OA, Birnbaumer L, Stefani E: **Identification of a second region of the beta-subunit involved in regulation of calcium channel inactivation.** *Am J Physiol* 1996, **271(5 Pt 1)**:C1539-45.
63. Chien AJ, Carr KM, Shirokov RE, Rios E, Hosey MM: **Identification of palmitoylation sites within the L-type calcium channel**

- beta2a subunit and effects on channel function.** *J Biol Chem* 1996, **271(43)**:26465-26468.
64. Gao T, Yatani A, Dell'Acqua ML, Sako H, Green SA, Dascal N, Scott JD, Hosey MM: **cAMP-dependent regulation of cardiac L-type Ca²⁺ channels requires membrane targeting of PKA and phosphorylation of channel subunits.** *Neuron* 1997, **19(1)**:185-196.
 65. Le Hir H, Nott A, Moore MJ: **How introns influence and enhance eukaryotic gene expression.** *Trends Biochem Sci* 2003, **28(4)**:215-220.
 66. Nott A, Meislin SH, Moore MJ: **A quantitative analysis of intron effects on mammalian gene expression.** *Rna* 2003, **9(5)**:607-617.
 67. Baek D, Davis C, Ewing B, Gordon D, Green P: **Characterization and predictive discovery of evolutionarily conserved mammalian alternative promoters.** *Genome Res* 2007, **17(2)**:145-55. Epub 2007 Jan 8.
 68. Mattick JS: **A new paradigm for developmental biology.** *J Exp Biol* 2007, **210(Pt 9)**:1526-1547.
 69. Pozzoli U, Menozzi G, Comi GP, Cagliani R, Bresolin N, Sironi M: **Intron size in mammals: complexity comes to terms with economy.** *Trends Genet* 2007, **23(1)**:20-4. Epub 2006 Oct 30.
 70. Vinogradov AE: **"Genome design" model: evidence from conserved intronic sequence in human-mouse comparison.** *Genome Res* 2006, **16(3)**:347-54. Epub 2006 Feb 3.
 71. Lynch M, Conery JS: **The origins of genome complexity.** *Science* 2003, **302(5649)**:1401-1404.
 72. Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, Ho RK, Langeland J, Prince V, Wang YL, Westerfield M, Ekker M, Postlethwait JH: **Zebrafish hox clusters and vertebrate genome evolution.** *Science* 1998, **282(5394)**:1711-1714.
 73. Taylor JS, Van de Peer Y, Meyer A: **Genome duplication, divergent resolution and speciation.** *Trends Genet* 2001, **17(6)**:299-301.
 74. Hurley IA, Mueller RL, Dunn KA, Schmidt EJ, Friedman M, Ho RK, Prince VE, Yang Z, Thomas MG, Coates MI: **A new time-scale for ray-finned fish evolution.** *Proc Biol Sci* 2007, **274(1609)**:489-498.
 75. Taylor JS, Braasch I, Frickey T, Meyer A, Van de Peer Y: **Genome duplication, a trait shared by 22000 species of ray-finned fish.** *Genome Res* 2003, **13(3)**:382-390.
 76. Ebert AM, McAnelly CA, Srinivasan A, Linker JL, Horne WA, Garrity DM: **Ca²⁺ channel-independent requirement for MAGUK family CACNB4 genes in initiation of zebrafish epiboly.** *Proc Natl Acad Sci U S A* 2008, **105(1)**:198-203. Epub 2008 Jan 2.
 77. Webb SE, Miller AL: **Ca²⁺ signaling and early embryonic patterning during the blastula and gastrula periods of zebrafish and Xenopus development.** *Biochim Biophys Acta* 2006, **1763(11)**:1192-1208.
 78. Hullin R, Matthes J, von Vietinghoff S, Bodi I, Rubio M, D'Souza K, Friedrich Khan I, Rottlander D, Hoppe UC, Mohacsi P, Schmitteckert E, Gilsbach R, Bunemann M, Hein L, Schwartz A, Herzog S: **Increased expression of the auxiliary beta2-subunit of ventricular L-type Ca²⁺ channels leads to single-channel activity characteristic of heart failure.** *PLoS ONE* 2007, **2(3)**:e292.
 79. Hullin R, Khan IF, Wirtz S, Mohacsi P, Varadi G, Schwartz A, Herzog S: **Cardiac L-type calcium channel beta-subunits expressed in human heart have differential effects on single channel characteristics.** *J Biol Chem* 2003, **278(24)**:21623-30. Epub 2003 Feb 26.
 80. Haase H, Kresse A, Hohaus A, Schulte HD, Maier M, Osterziel KJ, Lange PE, Morano I: **Expression of calcium channel subunits in the normal and diseased human myocardium.** *J Mol Med* 1996, **74(2)**:99-104.
 81. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF: **Stages of embryonic development of the zebrafish.** *Dev Dyn* 1995, **203**:253-310.
 82. **The JGI Eukaryotic Genomes Databases** [<http://genome.jgi-psf.org/>]
 83. **The EMBL-EBI Tools Page for ClustlW** [<http://www.ebi.ac.uk/Tools/clustalw/index.html>]
 84. **Tree Puzzle Analysis Page** [<http://www.tree-puzzle.de/>]
 85. Muller T, Vingron M: **Modeling amino acid replacement.** *J Comput Biol* 2000, **7(6)**:761-776.
 86. Burns CG, Milan DJ, Grande EJ, Rottbauer W, MacRae CA, Fishman MC: **High-throughput assay for small molecules that modulate zebrafish embryonic heart rate.** *Nat Chem Biol* 2005, **1(5)**:263-264.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

