
Properties of *Bacillus cereus* hemolysin II: A heptameric transmembrane pore

GEORGE MILES,¹ HAGAN BAYLEY,^{1,2} AND STEPHEN CHELEY¹

¹Department of Medical Biochemistry & Genetics, The Texas A&M University System Health Science Center, College Station, Texas 77843-1114, USA

²Department of Chemistry, Texas A&M University, College Station, Texas 77843-3255, USA

(RECEIVED February 5, 2002; FINAL REVISION April 24, 2002; ACCEPTED April 24, 2002)

Abstract

The gene encoding hemolysin II (HlyII) was amplified from *Bacillus cereus* genomic DNA and a truncated mutant, HlyII(Δ CT), was constructed lacking the 94 amino acid extension at the C terminus. The proteins were produced in an *E. coli* cell-free in vitro transcription and translation system, and were shown to assemble into SDS-stable oligomers on rabbit erythrocyte membranes and liposomes. The hemolytic activity of HlyII was measured with rabbit erythrocytes yielding an HC_{50} value of 1.64 ng mL^{-1} , which is over 15 times more potent than staphylococcal α -hemolysin. HlyII(Δ CT) was about eight times less potent than HlyII in this assay. Limited proteolysis of the oligomers formed by HlyII and HlyII(Δ CT) on red cell membranes showed that the C-terminal extension is sensitive to digestion, while HlyII(Δ CT) is protease resistant and migrates with an electrophoretic mobility similar to that of digested HlyII. HlyII forms moderately anion selective, rectifying pores ($I_{+80}/I_{-80} = 0.57$, 1 M KCl, pH 7.4) in planar lipid bilayers of diphtanoylphosphatidylcholine with a unitary conductance of 637 pS (1 M KCl, 5 mM HEPES, pH 7.4) and exhibits no gating over a wide range of applied potentials (–160 to +160 mV). In addition, it was demonstrated that HlyII forms a homoheptameric pore by using gel shift electrophoresis aided by a genetically encoded oligoaspartate tag. Although they share limited primary sequence identity (30%), these data confirm that HlyII is a structural and functional homolog of staphylococcal α -hemolysin.

Keywords: β -Barrel; hemolysin; membrane protein; pore-forming toxin; staphylococcal α -hemolysin; subunit stoichiometry

Reprint requests to: Hagan Bayley, Department Medical Biochemistry & Genetics, Texas A&M Health Science Center, 440 Reynolds Medical Building, College Station, TX 77843-1114, USA; e-mail: bayley@tamu.edu; fax: (979) 847-9481.

Abbreviations: α HL, α -hemolysin of *Staphylococcus aureus*; α HL-D8, α -hemolysin with a C-terminal extension of eight aspartate residues; α HL-TL, α -hemolysin fusion protein with a C-terminal extension comprising the C-terminal 94 residues of HlyII; β -PFT, β -barrel pore forming toxin; CytK, cytotoxin K of *Bacillus cereus*; HEPES, N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]; MBSA, 10 mM Na MOPS, 150 mM NaCl, pH 7.4, containing 1 mg mL^{-1} bovine serum albumin; MOPS, 3-[N-morpholino]propanesulfonic acid; IVTT, in vitro transcription and translation; PMSF, phenylmethylsulfonyl fluoride; rRBC, rabbit erythrocyte; rRBCM, rabbit erythrocyte membranes; SDS, sodium dodecyl sulfate; HlyII, hemolysin II of *Bacillus cereus*; HlyII-D8, hemolysin II with a C-terminal extension of eight aspartate residues; HlyII(Δ CT), a truncation mutant of HlyII lacking 94 amino acid residues at the C terminus; HlyII(Δ CT)-D8, HlyII(Δ CT) with a C-terminal extension of eight aspartate residues; TL, a polypeptide comprising the C-terminal 94 amino acids of HlyII.

Article and publication are at <http://www.proteinscience.org/cgi/doi/10.1110/ps.0204002>.

In this work, we examine the properties of hemolysin II (HlyII), a β -barrel pore forming toxin (β -PFT) from *Bacillus cereus*. The β -PFTs consist of several subfamilies of bacterial exotoxins that are related by sequence (Gouaux et al. 1997; Menestrina et al. 2001) and structure (Song et al. 1996; Gouaux et al. 1997; Olson et al. 1999; Pédelacq et al. 1999). These polypeptides are secreted as water-soluble molecules that bind to the surfaces of susceptible cells and assemble into oligomeric transmembrane pores leading to cell permeation and lysis (Bhakdi et al. 2000; Menestrina et al. 2001). The crystal structure of a heptameric staphylococcal α -hemolysin (α HL) pore has been determined in detergent at 1.9-Å resolution (Song et al. 1996). α HL has also been shown to form heptamers on red cell membranes (Gouaux et al. 1994), in planar lipid bilayers (Krasilnikov et al. 2000), supported bilayers (Fang et al. 1997), and after spontaneous assembly in solution (Cheley et al. 1997). In a

working scheme for the assembly of α HL, the 293-residue polypeptide first binds to the membrane as a monomer, associates to form a heptameric prepore, and finally inserts into the bilayer to form the transmembrane pore. This scheme is supported by numerous biophysical and biochemical experiments (Cheley et al. 1997), and has been refined to accommodate recent structural data (Olson et al. 1999; Pédelacq et al. 1999).

Bacillus cereus is an opportunistic pathogen. It is associated with a wide range of clinical symptoms, but encountered primarily in cases of severe food poisoning (Drobeniowski 1993; Lund et al. 2000). Over 20 exotoxins are produced and secreted by *B. cereus*, including several nonhemolytic enterotoxins (Alouf and Freer 1999). Two distinct hemolytic proteins produced by this bacterium, hemolysin II (HlyII) (Baida et al. 1999) and cytotoxin K (CytK) (Lund et al. 2000), have recently been cloned and shown to be homologous with the β -PFTs. HlyII has the longest polypeptide chain in the β -PFT family, with 412 residues, and contains a C-terminal 94 amino acid extension that has no homology with any other known β -PFT. The

remainder of HlyII shares 30% sequence identity with α HL, and the level of identity with other members of the β -PFT family does not exceed this value. Although regions of similarity are dispersed throughout the aligned sequences (Fig. 1A), the majority are concentrated in the strands that comprise the “cap” domain (Song et al. 1996), and are probably necessary to preserve the fold of the cap (Fig. 1B).

Members of the β -PFT family other than α HL have not been thoroughly investigated. In particular, limited data exist on the molecular architecture of the pores formed by them. The bicomponent leukocidin pore has a relatively large unitary conductance (2.5 nS in 1 M KCl) compared to other members of the β -PFT family (Miles et al. 2001), and it has been recently demonstrated to form an octameric transmembrane pore (Miles et al. 2002). The clostridial β -toxin has been shown to oligomerize into an SDS-stable multimer of unknown composition on human endothelial cells (Steinthorsdottir et al. 2000), and forms cation-selective channels with two conductance states of 60 and 110 pS (100 mM NaCl, 10 mM HEPES, pH 7.4, +60 mV) (Shatursky et al. 2000). Recently, it was shown that CytK purified




Fig. 1. Comparison of *Bacillus* hemolysin II and staphylococcal α -hemolysin. (A) Primary sequence alignment of *Bacillus cereus* hemolysin II (HlyII) and staphylococcal α -hemolysin (α HL). Residues highlighted with a blue background are identical, while similar residues (Blosom62 similarity scoring matrix; Henikoff and Henikoff 1992) are shown as blue characters. The 94 residue C-terminal extension of HlyII is shown in maroon characters; it is the portion that was deleted to form the truncation mutant HlyII(Δ CT). The figure was generated using ClustalW 1.81 (Thompson et al. 1994) and rendered using ESPript 2.0 (Gouet et al. 1999). (B) Structure of one protomer taken from the crystal structure of the α HL heptamer (7aHL.pdb). Areas in dark blue and light blue correspond to identical and similar residues, respectively, as shown in (A). The image was created with SPOCK 6.3 (Christopher 1998) and rendered with Molscript (Kraulis 1991).

from *B. cereus* supernatants forms pores in planar lipid bilayers, and is cytotoxic towards intestinal epithelial cells (Hardy et al. 2001). The subunit stoichiometries of the *Bacillus* toxins, HlyII and CytK, have not been determined.

In this study, we demonstrate that HlyII, produced by *in vitro* transcription and translation, forms a heptameric transmembrane pore in red cell membranes, which is resistant to SDS. In planar lipid bilayers, the pores are rectifying and lack voltage-induced gating. HlyII with the C-terminal extension removed, HlyII(Δ CT), has similar properties. Knowledge of the subunit stoichiometry and channel properties of this relative of α HL adds to the understanding of the β -PFTs family, and will be helpful in protein engineering aimed at the construction of pore-forming proteins with new properties (Bayley 1999; Bayley and Cremer 2001).

Results and Discussion

HlyII and *HlyII*(Δ CT) form SDS-resistant oligomers on red cell membranes and liposomes

HlyII includes a 94-amino acid C-terminal extension, which is not homologous with any known β -PFT. The two most

similar candidates from a BLAST search (Altshul et al. 1997) were a 46-amino acid segment of the pX01–124 gene product from *Bacillus anthracis* (39% identity, 67% similarity, one gap) and a 78-amino acid segment of orf16 from *Streptococcus* phage Cp-1 (34% identity, 49% similarity, three gaps). The genes for these proteins are both associated with genetic material linked with virulence, but the roles of the proteins have not been fully defined (Welkos 1991; Martín et al. 1996). We constructed a truncation mutant, HlyII(Δ CT), which lacks the extension (residues 289–382) (Fig. 1A). 35 S-labeled HlyII and HlyII(Δ CT) were produced in an *E. coli* S30 transcription and translation system. When HlyII was translated in the presence of rabbit red cell membranes (rRBCM) or incubated with small unilamellar vesicles (SUVs) composed of egg yolk phosphatidylcholine, cholesterol, and phosphatidic acid at a molar ratio of 55:25:20, a single high-molecular mass band above the 220-kD marker appeared upon SDS-polyacrylamide gel electrophoresis of unheated samples (Fig. 2). The extent of oligomerization on rRBCM is comparable to that of α HL, with 74% of the total membrane-bound protein in the oligomeric form (compared to 83% seen with α HL). In




Fig. 2. 35 S-Labeled HlyII and HlyII(Δ CT) synthesized by coupled *in vitro* transcription and translation (IVTT) with an S30 extract from *E. coli*. An autoradiogram of a 10% SDS-polyacrylamide gel is shown. Lane 1, HlyII translated in the absence of rRBCM; lane 2, HlyII(Δ CT) translated in the absence of rRBCM; lane 3, HlyII translated in the presence of rRBCM; lane 4, HlyII(Δ CT) translated in the presence of rRBCM; lane 5, HlyII incubated in the presence of freshly prepared liposomes for 1 h at room temperature; lane 6, HlyII(Δ CT) incubated in the presence of liposomes. In the samples used in lanes 3 and 4, the rRBCM were washed before electrophoresis of the bound protein. In the samples used in lanes 5 and 6, the entire sample was loaded onto the gel. The high molecular weight bands (arrows) represent oligomerized HlyII and HlyII(Δ CT).

the case of HlyII(Δ CT), the extent of oligomerization is reduced to 49%. Oligomers formed by HlyII and HlyII(Δ CT) are stable in 2.3% SDS (1 \times Laemmli sample buffer) at room temperature and dissociate at 82°C and 78°C, respectively. α HL dissociates at 70°C in the sample buffer (data not shown).

A deletion variant lacking the extension (and two additional amino acids) was previously shown to retain some hemolytic activity towards human red blood cells (Baida et al. 1999). The removal of the two additional amino acids is expected to lead to a reduction in activity. We found earlier that the removal of three amino acids from the C terminus of α HL causes a dramatic reduction in hemolytic activity that is associated with an almost complete loss of ability to form SDS-stable oligomers (Walker et al. 1992a).

Examination of the HlyII oligomer by limited proteolysis suggests structural similarity to α HL

The susceptibility of HlyII and HlyII(Δ CT) to cleavage by proteinase K was compared with that of α HL (Fig. 3). The HlyII oligomer shows marked susceptibility to proteinase K, yielding multiple proteolytic fragments at low protease concentrations (5 μ g mL⁻¹; Fig. 3, lane 2). In contrast, the oligomeric form of HlyII(Δ CT), like the α HL heptamer, was protease resistant. At the highest concentration of proteinase K (500 μ g mL⁻¹), a protease-resistant, SDS-stable form of oligomeric HlyII was formed, which mi-

grated in SDS gels with a similar mobility to oligomers formed from the genetically truncated HlyII(Δ CT) (Fig. 3). These results suggest that the C-terminal extension either has a structure with pronounced sensitivity to proteolytic cleavage, or that the extension is connected to the rest of the protein through a readily accessible linker. The former is favored because the C terminus (TL, see below) expressed as a separate domain is protease sensitive (data not shown). In any case, the stability of the proteolyzed pore implies that it is unlikely that the extension contributes significantly to the stability of the cap domain, the formation of which initiates heptamerization at the prepore stage (Cheley et al. 1997). In addition, when the sequences of HlyII from several different *B. cereus* and *B. thuringiensis* strains, BGSC 4A1, 4A4, 4A7, 6A3, and 6E2, were compared to the HlyII sequence from *Bacillus cereus* strain 6A5, the nonredundant sequence differences were concentrated in the C-terminal extension (14 differences in the 94 residues, compared with 9 differences in the 288 residues of the remainder of the polypeptide). This relative lack of conservation again suggests that the extension does not play a major role in oligomerization.

Hemolysin II is a potent hemolytic toxin

The hemolytic activities of HlyII and HlyII(Δ CT), produced by IVTT, were compared with α HL in a quantitative assay. The concentration of HlyII at which 50% of the rabbit erythrocytes were lysed in 1 h at 20°C (HC₅₀) was 1.64 ng mL⁻¹ (Fig. 4A). By contrast, α HL gives an HC₅₀ value of 25 ng mL⁻¹ at 37°C (Walker et al. 1992b) and a similar value at 20°C (unpublished results). The HC₅₀ value for CytK is similar to that of α HL (Lund et al. 2000). Therefore, the specific activity of HlyII is more than 15 times greater than that of α HL or CytK. The HC₅₀ value amounts to about 1000 monomers per cell. HlyII(Δ CT) had a lower specific activity of 4.8 ng mL⁻¹ (Fig. 4B), in keeping with the somewhat less efficient oligomerization seen with this mutant (Fig. 3). The C-terminal extension by itself (TL) showed no hemolytic activity (Fig. 4B), and did not affect the activity of HlyII(Δ CT) or HlyII. We have found that other β -PFTs can accommodate large extensions at their C termini. For example, the mutant α HL-TL was constructed in which the *Bacillus* tail was spliced genetically onto the C terminus of α HL. The HC₅₀ of α HL-TL was 64 ng mL⁻¹, i.e. α HL-TL was 2.5 times less active than wild type α HL. In the case of the F and S subunits of leukocidin, the hemolytic activities were unchanged when TL was fused to the C termini (Miles et al. 2002). The cytotoxic effects of HlyII against other cell lines has not been examined. It has been shown recently that CytK has potent activity against human intestinal epithelia, and is believed to have caused necrotic enteritis in a food poisoning outbreak resulting in several deaths (Lund et al. 2000; Hardy et al. 2001).




Fig. 3. Conformational states of HlyII, HlyII(Δ CT) and α HL examined by limited proteolysis. HlyII, HlyII(Δ CT) and α HL were translated in the presence of rRBCM. The membranes were washed, treated with proteinase K, solubilized and subjected to electrophoresis in a 10% SDS-polyacrylamide gel prior to autoradiography. Lanes 1–4, HlyII; lanes 5–8, HlyII(Δ CT); lanes 9–12, α HL. The final proteinase K concentrations were: lanes 1, 5, and 9, 0 μ g mL⁻¹; lanes 2, 6, and 10, 5 μ g mL⁻¹; lanes 3, 7, and 11, 50 μ g mL⁻¹; lanes 4, 8, and 12, 500 μ g mL⁻¹.




Fig. 4. Quantitative hemolysis assays with rabbit erythrocytes. (A) One-hour activity assays on HlyII and α HL synthesized by IVTT. The first well in each panel contained IVTT mix diluted to a final volume of 100 μ L in MBSA. The concentration of α HL in an IVTT mix was determined as previously described (Walker et al. 1992b; Miles et al. 2001) and used to calculate the concentration of HlyII produced by IVTT (see Materials and Methods). On this basis, the final concentrations of HlyII or α HL were made equal in the first well (0.21 μ g/mL). Twofold serial dilutions from left to right were then made. The final concentration of rRBC in all wells was 0.5%. The arrows indicate \sim 50% lysis. (B) One-hour activity assays on various constructs synthesized by IVTT. IVTT mixes containing HlyII, HlyII(Δ CT), α HL, and α HL-TL were diluted 10-fold with MBSA before addition (1 μ L) to the first well. TL, the C-terminal extension of HlyII was translated as a separate polypeptide, and added where indicated in 32-molar excess as determined by phosphorimager quantitation (5 μ L of undiluted IVTT mix in the first well). Twofold serial dilutions were made from left to right.

By comparison with α HL, there is a shorter initial lag period associated with HlyII hemolysis (Fig. 4): HlyII, 5 min to 5% lysis at HC_{50} ; α HL, 18 min to 5% lysis at HC_{50} . The short lag period for HlyII suggests that the overall rate of pore formation (membrane binding, oligomerization, and membrane penetration) is greater than that of α HL. By contrast, the rates of hemolysis after the lag period were similar at comparable effective concentrations. Further investigation will be necessary to better define the origin of the difference in lag time.

HlyII forms ionic channels in planar lipid bilayers

Upon insertion into planar lipid bilayers, the gel-purified oligomeric form of HlyII obtained by IVTT (Fig. 2) pro-

duced discrete conductance steps of 25.5 pA in 1 M KCl at pH 7.4 at a transmembrane potential of -40 mV (Fig. 5A). If each step is presumed to correspond to a single ionic channel, the unitary conductance of the HlyII pore is 637 ± 51 pS ($n = 180$), under the prevailing conditions (Fig. 5A). A similar result was obtained with gel-purified HlyII(Δ CT) oligomers, 636 ± 73 pS (Table 1). By comparison, the conductance of the α HL pore was 775 ± 38 pS, under the same conditions (Miles et al. 2001) and that of CytK 627 ± 14 pS (1 M NaCl, 5 mM HEPES, pH 7.2) (Hardy et al. 2001). Therefore, the functional diameters of the HlyII, CytK and α HL pores are similar.

The HlyII pores show rectification when examined at applied potentials from -160 to $+160$ mV ($I_{+80}/I_{-80} = 0.566$, 1 M KCl, pH 7.4) (Fig. 5B, Table 1), which is the opposite of that observed with the clostridial β -toxin (Shatursky et al. 2000) (note that these authors use the opposite polarity sign convention in their bilayer recordings). The charge selectivity of the HlyII pore was determined by measuring the reversal potential of single-channel currents in asymmetrical KCl solutions. HlyII forms anion selective pores, with a permeability ratio (P_{K^+}/P_{Cl^-}) of 0.29 ± 0.03 (1000 mM KCl (*cis*), 200 mM KCl (*trans*), pH 7.5) (Fig. 5C). The HlyII pore is significantly more anion selective than the α HL pore ($P_{K^+}/P_{Cl^-} = 0.79$, under the same conditions) (Gu et al. 2000), whereas the leukocidin pore is cation selective ($P_{K^+}/P_{Cl^-} = 1.64$) (Miles et al. 2001). Of the known β -PFTs, HlyII and CytK share the greatest sequence identity within the putative transmembrane domain (48%). In HlyII, each subunit contributes seven charged residues that are predicted on the basis of their hydrophilicity and a sequence alignment with α HL to face the lumen of the transmembrane barrel (K106, E110, K120, D125, K126, D131, K141). However, only two charged side chains (K128, E139) are predicted to project into the lumen of the CytK pore, which is almost nonselective (Hardy et al. 2001). The overall positive charge in the HlyII pore is likely to contribute to the observed anion selectivity.

In addition, the HlyII pore remains predominantly open at the applied potentials tested (-160 to $+160$ mV) (Fig. 6). This is similar to what is observed with α HL (Korchev et al. 1995), but in contrast with the properties of most porins and many other β -PFTs, which exhibit voltage-induced gating. For example, the leukocidin pore gates at positive potentials above $+60$ mV, but remains fully open at negative potentials up to -160 mV (Miles et al. 2001). Two rings of charged residues facing the lumen of the HlyII pore form the boundaries of the putative transmembrane region, E110/K141 and K120/D131, based on the alignment of residues in the stem domain (unpublished results) with those in the α HL β -barrel (Song et al. 1996). We speculate that these side chains form salt bridges that strengthen the β -barrel, holding it in a rigid conformation even under high applied potentials. This argument is reinforced by the ruggedness of the HlyII




Fig. 5. Single-channel recordings from HlyII pores. (A) Individual insertions of HlyII pores into planar lipid bilayers. HlyII oligomer was prepared by IVTT in the presence of rRCBM, purified by SDS-polyacrylamide gel electrophoresis, and added to the *cis* chamber of a bilayer apparatus. Both chambers contained 1 M KCl, 5 mM HEPES, pH 7.4, and the applied potential was -40 mV. Stepwise changes in current are shown as a function of time. A histogram of the current steps is displayed and represents a compilation from 15 independent recordings. The histogram was fitted to a Gaussian function to obtain the single-channel conductance. (B) Single-channel current-voltage (I-V) relationship. A representative plot is shown. (Inset) Plot of I^+/I^- versus voltage representing data (\pm SD) from four independent recordings. The conditions were as described in (A). (C) HlyII is anion selective at neutral pH. The reversal potential (V_r), the applied voltage that gave zero current (arrow), was determined from single channel I-V plots with asymmetrical KCl solutions (1000 mM KCl *cis*, 200 mM KCl *trans*, both in 5 mM HEPES, pH 7.4). A representative plot is shown. The permeability ratio (P_{K^+}/P_{Cl^-}) was derived from the GHK equation.

oligomer, which dissociates in SDS at 82°C , as it is known that it is barrel formation that stabilizes the αHL heptamer to heat denaturation in SDS (Valeva et al. 1996; Cheley et al. 1999).

Hemolysin II assembles as a heptamer on membranes

Finally, we determined the subunit composition of the HlyII pore by gel shift electrophoresis. Whereas the original gel-

Table 1. Single-channel properties of wild-type and engineered HlyII pores

Protein	g (pS) ^a	n	I_{+80}/I_{-80} ^b	V_r (mV); P_{K^+}/P_{Cl^-} ^c
HlyII(WT)	637 ± 51	180	0.566 ± 0.004	-18.0 ± 1.3 ; 0.29 ($n = 4$)
HlyII(ΔCT)	636 ± 73	142	0.583 ± 0.001	n.d.
HlyII(ΔCT)-D8	610 ± 54	273	0.591 ± 0.032	-19.3 ± 1.2 ; 0.26 ($n = 3$)
[HlyII(ΔCT)] ₆ / [HlyII(ΔCT)-D8] ₁	642 ± 62	144	0.691 ± 0.077	n.d.

^a Recordings were made at -40 mV in 1 M KCl, 5 mM HEPES, pH 7.4. Single-channel conductances were determined by fitting the peaks in amplitude histograms to Gaussian functions (e.g., Figs. 5, 8).

^b Rectification ratios were determined from three or more experiments.

^c The reversal potentials (V_r) and permeability ratios (P_{K^+}/P_{Cl^-}) are mean values \pm SD determined with the following buffers: *cis*, 1 M KCl, 5 mM HEPES, pH 7.4; *trans*, 0.2 M KCl, 5 mM HEPES, pH 7.4. n.d., not determined.




Fig. 6. The HlyII pore exhibits minimal gating over a wide range of applied potentials. Each trace is representative of a single channel from HlyII oligomer prepared by IVTT in the presence of rRBCM and purified by SDS-polyacrylamide gel electrophoresis. The fully open state of the channel is shown at both polarities of the indicated voltage.

shift method used to evaluate the stoichiometry of the α HL pore was based on a mobility change brought about by site-specific chemical modification (Gouaux et al. 1994; Braha et al. 1997), gel-shift experiments based on genetically engineered truncations or extensions have proved successful in other cases (Heginbotham et al. 1997; Zitzer et al. 1999; Miyata et al. 2001; Miles et al. 2002).

In a similar approach to the one we used here, transthyretin heterotetramers were separated by using an N-terminal charged extension (Flag tag) (Schneider et al. 2001). Similarly, hybrid trimers of the catalytic subunits of aspartate transcarbamoylase were resolved with a six-residue aspartate tail (Sakash and Kantrowitz 2000). The mutant HlyII-D8 was constructed in which an extension encoding eight aspartate residues was incorporated at the C terminus of HlyII (Fig. 7A). Oligomers containing various ratios of HlyII to HlyII-D8 were produced by cotranslation in the presence of rRBCM. Analysis of the oligomers by SDS-polyacrylamide gel electrophoresis and autoradiography lacked sufficient resolution to enable a count of individual bands (Fig. 7B). It is unclear why the D8 tail on full-length HlyII fails to yield sharp bands; it is possible that the vari-

ous permutations of each combination of subunits (Braha et al. 1997) are spread out on the gel. By contrast, eight distinct bands were obtained by using various mixtures of α HL and α HL-D8 (Fig. 7C). Each downward shift in electrophoretic mobility corresponds to the incorporation of one α HL-D8 subunit into the α HL heptamer. Therefore, we engineered the D8 tail onto HlyII(Δ CT). When assembled with HlyII(Δ CT), eight species were resolved by gel electrophoresis (Fig. 7D). This result indicates that the HlyII(Δ CT) oligomer contains seven subunits.

Because the properties of the pores formed by HlyII and HlyII(Δ CT) are similar, HlyII is also likely to form a heptamer. HlyII and HlyII(Δ CT) produced pores with similar conductance values (Fig. 8A,B; Table 1). Together with the proteolysis experiments and measurements of hemolytic activity, these results support the idea that the 94-amino acid C-terminal extension is neither essential for pore formation nor for determining functional properties. Oligomers composed entirely of HlyII(Δ CT)-D8 subunits and a heteromer containing one D8 subunit [HlyII(Δ CT)]₆/[HlyII(Δ CT)-D8]₁ also exhibited similar conductance values, rectification properties and charge selectivities when compared with the pores formed by HlyII and HlyII(Δ CT) (Fig. 8A,B; Table 1). Therefore, despite the additional 56 negative charges in [HlyII(Δ CT)-D8]₇, the conductive pathway remains unaffected, in keeping with the location of the C terminus distant from the channel entrance.

Several different N- and C-terminal oligoaspartate extensions have been evaluated in our laboratory for their ability to produce electrophoretic shifts in β -PFTs. For instance, a D8 tail provides about 1.5 times the separation between oligomeric species in SDS gels compared with a D4 extension (G. Miles and S. Cheley, unpubl.). Besides its utility in evaluating the subunit composition of proteins, this approach provides a convenient way to separate functional heteromers (Sakash and Kantrowitz 2000; Howorka et al. 2001a, 2001b).

General implications and future prospects

Here, we have shown that the properties of the pore-forming protein HlyII from *Bacillus cereus* conforms, in general, with those of other members of the class of β -PFTs. HlyII has the most potent hemolytic activity yet found in the β -PFTs; 50% lysis of rabbit red cells occurs at a ratio of \sim 1000 HlyII monomers per cell. HlyII possesses additional properties that might be improved for applications in biotechnology (Eroglu et al. 2000; Bayley and Cremer 2001). For example, the heptameric pore is more stable than that formed by α HL, dissociating in SDS at 82°C. Remarkably, the pore formed by clostridial β toxin is yet more stable, resisting "boiling" in SDS (Steinthorsdottir et al. 2000). The pore formed by HlyII remains open at high transmembrane potentials, which is a useful property for applications in




Fig. 7. Subunit stoichiometry of the HlyII pore. (A) Constructs used in the study. The maroon block represents the 94-residue C-terminal extension. The oligoaspartate (D8) tail is shown in red. (B) SDS-polyacrylamide gel electrophoresis of heteromers generated from WT HlyII and/or HlyII-D8. (C) Heteromers generated from αHL and/or αHL-D8. (D) Heteromers generated from HlyII(ΔCT) and/or HlyII(ΔCT)-D8. The DNA constructs were mixed at various molar ratios, as indicated in (B–D), and cotranslated in the presence of rRBCM. The membranes were washed, solubilized in sample buffer without heating, and subjected to electrophoresis in a 5% gel. Autoradiograms of dried gels are shown. The results in B are from a long gel (36 cm), which was run for 72 h at 50 V.

sensor technology (Bayley et al. 2000; Bayley and Cremer 2001). The results provided here also demonstrate that the tail carried by HlyII has little influence on the properties of the pore. This suggests that the tail might be replaced by polypeptide sequences that provide additional functionality such as the ability to form lattices or bind to surfaces, or a linked catalytic activity. Finally, the use of an oligoaspartate extension provides a convenient means to purify heterooligomeric pores. Combined with new findings about the subunit–subunit interface, revealed by the ability of the related binary pore-forming toxin, leukocidin, to form octam-

ers (Miles et al. 2002), the present work suggests that it should be possible to extend the range of heteromeric pores that can be formed and purified.

Materials and methods

Isolation of genomic DNA from *Bacillus cereus*

Bacillus cereus strain 6A5 (Bacillus Genetic Stock Center, Ohio State University) (equivalent to ATCC #14579) was grown to saturation from log-phase inocula and stored frozen at -80°C in




Fig. 8. Representative channel insertion events and single channel conductance histograms for [HlyII]₇, [HlyII(ΔCT)]₇, [HlyII(ΔCT)-D8]₇, and the heteroheptamer [HlyII(ΔCT)]₆[HlyII(ΔCT)-D8]₁. (A) HlyII homoheptamer. (B) HlyII(ΔCT) homoheptamer. (C) HlyII(ΔCT)-D8 homoheptamer. (D) [HlyII(ΔCT)]₆[HlyII(ΔCT)-D8]₁ heteroheptamer. After a preparative IVTT reaction in the presence of rRCM and SDS-polyacrylamide gel electrophoresis, the desired band was excised from the gel and processed as described in Materials and Methods. Current traces were recorded in 1 M KCl, 5 mM HEPES (pH 7.4). The scale for the current trace in (D), applies to the entire figure. The histograms of the current steps represent compilations from at least 15 independent recordings. Conductance values, rectification properties, and ion selectivities are summarized in Table 1.

50% glycerol. To isolate genomic DNA, a portion of a culture (5 mL) grown overnight at 30°C was centrifuged at 3000g. The pellet was resuspended in buffer B1 (Qiagen #19060; 1 mL, 50 mM Tris.HCl, pH 8.0, 0.5% Tween-20, 0.5% Triton X-100), supplemented with 125 U of mutanolysin (Sigma, #M4782; 25 μL of a stock made at 5000 U mL⁻¹ in deionized water and stored in frozen aliquots), 0.4 mg proteinase K (GibcoBRL; 20 μL of a stock made at 20 mg mL⁻¹ in 1× TE [10 mM Tris.HCl, 1 mM EDTA, pH 8.0, supplemented with 10 mM NaCl]) and 0.2 mg RNase A (Qiagen; 2 μL of a 100 mg mL⁻¹ stock), and incubated at 37°C overnight. Genomic DNA was purified on an anion-exchange resin using a Qiagen Genomic tip 20/G (#10223) and re-dissolved in 1× TE. The DNA was then sheared by two passes through an 18-gauge needle.

Amplification of *Bacillus hemolysin-II* genes

The coding sequences for HlyII were amplified from the genomic DNA by using the following primers: HlyII (sense) 5'-ACATATGGCAGATTCTAAAGGAAGTGTAGAAAATC-3';

HlyII (antisense) 5'-CAAGCTTATCAGATTTTTTTAATCTCAATATAAGG-3'. The sense primer generated an *NdeI* site (CATATG), containing a new initiation codon, immediately before the first codon of the mature polypeptide, predicted by sequence alignment (Fig. 1A). The antisense primer encodes two stop codons and a *HindIII* site immediately following the last codon of the gene. PCR was carried out in a 50-μL mixture containing sheared genomic DNA (50 ng), primers (50 pmole of each), 200 μM dNTPs (Stratagene), and 1.5 U of Taq/Pwo DNA polymerase mixture in PCR buffer 1 (Expand Long Template PCR System) with the following program: 95°C for 2 min, 25 cycles of 94°C (60 sec), 45 to 56°C gradient (60 sec), 72°C (80 sec), followed by a final extension at 72°C for 7 min. The PCR product was then cloned into the TOPO-TA plasmid (#K4500-01, Invitrogen) to yield pHlyII-TA.

pHlyII-TA was digested with *NdeI* and *HindIII*, and the liberated DNA insert was ligated into the pT7 expression vector pT7-SC1 (Miles et al. 2001). Upon sequencing, two independently amplified clones from *Bacillus cereus* strain 6A5; the following differences were noted from the published sequence (*B. cereus* VKM-B771) (Baida et al. 1999): **G66S** (GGG→AGC), **S236T**

(TCT→ACT), **N276H** (AAC→CAC), P294L (CCT→CTT), I299N (ATT→AAT), G300S (GGT→AGT), N303S (AAC→AGC), N306D (AAC→GAT), Q307K (CAG→AAA), F317L (TTT→CTT), T358S (ACA→TCA). With the exception of three residues (in bold), the variations noted occurred in the C-terminal extension. In addition, there were 35 silent changes throughout the sequence. The DNA sequence from strain 6A5 has been deposited in GenBank with the accession number AF448485.

Hemolysin-II C-terminal truncation

Unless otherwise noted, the constructs used in this study were created by PCR and cloned into the TOPO-TA plasmid prior to being subcloned into pT7-SC1 by ligation. Each construct was verified by DNA sequencing.

HlyII(Δ CT), a mutant of HlyII truncated at the C terminus, was constructed by removing the sequence encoding the last 94 amino acids of HlyII (residues 289–382). PCR was carried out on linearized pT7-HlyII(6A5) using the forward primer SC001 5'-CAC TATAGGGAGACCACAACGG-3' and the reverse primer BAC6 A5TRN1 5'-TAAGCTTCATTAAGAGTAACTTGATG-3'. The latter encodes two stop codons and a *Hind*III site immediately after the Leu-288 codon in the HlyII gene.

The C-terminal extension (TL) was cloned separately into the pT7 vector by using PCR. The sense primer BAC6A5TAILBEGIN 5'-ACATATGGATAACCAAAAAGCCCTT-3' generated an *Nde*I site (CATATG), containing a new initiation codon, immediately before the first codon of the C-terminal extension (Asp-289). The antisense primer was SC011 5'-CCCCTCAAGACCCGTTTAG AGGC-3', which hybridized at a site in the vector downstream of the stop codons.

Construction of mutants with C-terminal (oligo)-aspartic acid extensions

We sought to develop a genetic alternative to the original gel shift method for counting subunits, which was based on a mobility change brought about by site-specific chemical modification (Gouaux et al. 1994). The mutants HlyII-D8, HlyII(Δ CT)-D8, and α HL-D8 were constructed with an extension encoding eight C-terminal aspartate residues (Fig. 7A). The "D8 tail" was expected to change the electrophoretic mobility of the assembled pore based on charge. PCR was carried out with SC001 as the sense primer, using each of the following reverse primers: HlyII-D8, 5'-AAGCT TATCAATCGTCATCGTCATCGTCATCGTCGATTTTTTTAA TCTCAA-3'; HlyII(Δ CT)-D8, 5'-AAGCTTATCAATCGTCATC GTCATCGTCGTCGTCAAGAGT ACAAGATGGTT-3'; α HL-D8, 5'-AAGCTTATCAATCATCGTCATCATCGTCATC ATTTGTCATTTCTTCTTTTCCCC-3'. These electrophoretically purified antisense oligonucleotides incorporated two stop codons (bold) after the D8 tail codons (underscored) followed by a *Hind*III site (italics).

Construction of the fusion protein, α HL-TL

In vivo recombination (Howorka and Bayley 1998) was used to fuse a 3' extension directly to the last codon of the α HL gene (Jones 1995). The extension encoded the 94 amino acids of the *Bacillus* hemolysin II C-terminal tail (residues 289–382). The fused gene (α HL-TL) was generated in pT7-SC1 by cotransforming *Escherichia coli* XL-10 Gold cells with two PCR products encompassing the *Bacillus* tail and the α HL gene. The primer sets

for the two amplification reactions were: (1) nonmutagenic primer (F-NM) (sense), 5'-GTATTCAACATTTCCGTGTCGCCCTTAT TC-3'; α HL-TAIL- α HL, (antisense), 5'-AAGGGCTTTAAGGTTA TCATTTGTCATTTCTTCTTT-3'; and (2) α HL-TAIL-TL (sense), 5'-AAAGAAGAAATGACAAATGATAACCAAAAAGCCCTT-3'; nonmutagenic primer (R-NM) (antisense) 5'-GAATAAGG CGACACGGAAATGTTGAATAC-3'. The underlined 18-nt sequences form the overlap for recombination between the two PCR products.

In vitro transcription and translation (IVTT)

Polypeptides were synthesized in a cell-free in vitro transcription and translation (IVTT) system by using an S30 extract from *E. coli* (T7 S30 No. L114A, Promega) supplemented with rifampicin (20 μ g mL⁻¹) (Walker et al. 1992b). Radiolabeled polypeptides were synthesized by IVTT with the complete amino acid mix supplemented with [³⁵S]methionine (ICN Biomedicals, Inc., 1175 Ci mmole⁻¹, 10 μ Ci per 25 μ L translation). The concentrations of the translated polypeptides were determined by phosphorimager analysis, by comparison with α HL standards radiolabeled in parallel IVTT reactions (Miles et al. 2001). The specific radioactivity of the α HL, in phosphorimager units, was determined after deducing the concentration of the protein from its hemolytic activity, assuming a specific hemolytic activity of 25 ng mL⁻¹ (Walker et al. 1992b). The specific radioactivity was normalized to the number of Met residues in the polypeptide chain (assuming that the N-terminal Met is intact). The concentration of HlyII, and its variants, in a translation mix could then be determined from the strengths of the phosphorimager signals and the number of Met residues in the polypeptide chains.

Quantitative hemolysis assay

HlyII proteins, synthesized by IVTT with the complete amino acid mix, were diluted into MBSA (10 mM 3-[*N*-morpholino]propane sulfonic acid; MOPS, Cat. No. AB1270, American Bioanalytical, 150 mM NaCl, pH 7.4, containing 1 mg mL⁻¹ bovine serum albumin, Cat. No. 4503, Sigma) and subjected to 12 twofold serial dilutions in the same buffer in microtiter wells (final volume 50 μ L). An equal volume of 1% washed rabbit erythrocytes (rRBC) in MBSA was quickly added to each well, beginning with the most diluted sample. Hemolysis was recorded for 1 h at 20°C by monitoring the decrease in light scattering at 595 nm with a Bio-Rad microplate reader (Model 3550-UV) and using the Microplate Manager 4.0 software.

HlyII oligomer formation on rabbit erythrocyte membranes

Radiolabeled HlyII oligomers were prepared by IVTT in the presence of [³⁵S]methionine and rRBCM (5 μ L, 3.0 mg protein mL⁻¹) (Cheley et al. 1999) in a total reaction volume of 25 μ L. After 1 h at 30°C, the mixture was centrifuged and the supernatant discarded. The membrane pellet was washed and resuspended in MBSA (80 μ L) prior to solubilization by the addition of 5 \times Laemmli sample buffer (20 μ L; Laemmli 1970). A portion (20 μ L) was subjected to electrophoresis in a 10% SDS-polyacrylamide gel. The gel was fixed for 1 h prior to drying and autoradiography.

Oligomerization of HlyII on liposomes

Egg yolk phosphatidylcholine, cholesterol, and phosphatidic acid in chloroform were mixed in the desired molar ratio of 55:25:20. After drying under vacuum, the lipid film was resuspended in buffer (10 mM Tris, pH 8.0, 150 mM NaCl, 1 mM EDTA) to a total lipid concentration of 5 mg mL⁻¹. Liposomes were prepared by ultrasonication for 30 min on ice using a probe sonicator (Dyna-tech Sonic Dismembrator Model 150) (relative output set to 50% power), followed by a brief centrifugation (30 sec, 16,000g) to remove titanium particles. [³⁵S]Methionine-labeled HlyII or HlyII(Δ CT) (2 μ L of an IVTT reaction mix) was incubated with freshly prepared liposomes (8 μ L) for 1 h at room temperature. After solubilization in Laemmli sample buffer, the mixture was subjected to SDS polyacrylamide gel electrophoresis.

Proteinase K treatment of HlyII, HlyII(Δ CT) and α HL polypeptides on membranes

Proteinase K (Sigma, #P-0390) solutions (5.0, 0.5, and 0.05 mg mL⁻¹ in water) were prepared by dilution of an enzyme stock (10 mg mL⁻¹ in water) and used immediately. Limited proteolysis was performed on HlyII, HlyII(Δ CT) and α HL bound to rRBCM. The membranes were resuspended in MBSA at 0.19 mg membrane protein mL⁻¹ and divided into four tubes (18 μ L in each). Proteinase K or water (2 μ L) was added to each tube. After 5 min at room temperature, the reactions were stopped by treatment with PMSF (9 mM final, added in 2 μ L of isopropanol, 5 min, room temperature), followed by the addition of 2 \times Laemmli loading buffer. The samples were subjected to electrophoresis in 10% SDS-polyacrylamide gels (unheated samples) or 12% gels (heated samples: 95°C, 5 min).

Gel purification of HlyII oligomers

HlyII, HlyII(Δ CT) and HlyII(Δ CT)-D8, and were prepared by translation in the presence of rRBCM as described above, but in preparative amounts (75 μ L IVTT reaction). To obtain, HlyII(Δ CT)₆(Δ CT)-D8₁, HlyII(Δ CT) and HlyII(Δ CT)-D8 were translated using the corresponding plasmids at a ratio of 5:1. The oligomers were purified by SDS-polyacrylamide gel electrophoresis in an 8% gel in the presence of 0.1 mM sodium thioglycolate (Movileanu et al. 2001; Miles et al. 2001), stored at -80°C in 10 mM Tris.HCl, pH 7.5, and used for bilayer recordings without further treatment.

Hetero-oligomer formation for the determination of stoichiometry

Hetero-oligomers of hemolysin II subunits containing HlyII and/or HlyII-D8, and HlyII(Δ CT) and/or HlyII(Δ CT)-D8 were prepared by mixing the corresponding plasmids in the desired molar ratios (see Fig. 7 legend) prior to IVTT in the presence of rRBCM. To obtain α HL heteromers, α HL and/or α HL-D8 were used. The washed membrane pellets were solubilized with Laemmli sample buffer and subjected, without heating, to electrophoresis in 5% SDS-polyacrylamide gels. Autoradiographs were made of the dried gels.

Planar lipid bilayer recordings

All measurements were performed at 25°C. Numerical values are given as the mean \pm SD (σ_{n-1}). Planar lipid bilayer membranes

were formed with 1,2-diphytanoyl-*sn*-glycero-3-phosphocholine (Avanti Polar Lipids) on a 150–160 μ m-diameter aperture in a 25 μ m-thick Teflon film (Goodfellow Corporation) separating the *cis* and *trans* compartments (2 mL each) of a bilayer apparatus (Montal and Mueller 1972). Prior to forming the lipid bilayer, the orifice was pretreated with 10% (v/v) hexadecane (#29,631-7; Aldrich) in *n*-pentane (Burdick & Jackson) and allowed to dry thoroughly. For most measurements, the *cis* and *trans* chambers contained 1 M KCl, 5 mM HEPES, pH 7.4. Various concentrations of KCl in 5 mM HEPES, pH 7.4 were used for ion selectivity measurements. Protein samples were added to the *cis* chamber, which was at ground.

Currents were recorded by using a Dagan 3900A patch clamp amplifier (Dagan Corporation) with a 3910 expander and a built-in low-pass four-pole Bessel filter set at 5 kHz. Data were stored on digital audio tape with a DAS-75 data recorder (Dagan Corporation). Prior to analysis, the signal was low-pass filtered at 1 kHz with an eight-pole Bessel filter (Model 902, Frequency Devices) and acquired with a Digidata 1200A A/D board with a sampling time interval of 200 μ sec. Data were acquired and analyzed with pClamp 8.0 software (Axon Instruments). Single-channel conductance values were determined by fitting the peaks in amplitude histograms to Gaussian functions. Current-voltage (I-V) relationships for single channels were determined by recording the currents obtained after stepwise changes in applied potential. The permeability ratios (P_{K^+}/P_{Cl^-}) were calculated from experimentally determined reversal potentials (V_p) by using the Goldman-Hodgkin-Katz (GHK) equation (Hille 2001) and the appropriate activity coefficients for KCl solutions (Zemaitis et al. 1986). In these measurements, the *cis* compartment contained 1000 mM KCl, while the other chamber contained 200 mM KCl. Any electrode DC offset was balanced prior to the addition of protein to the *cis* chamber. The applied voltage that gave zero current was noted. In addition, the reversal potential was more accurately determined by polynomial fits to current-voltage (I-V) data. Symmetrical solutions were then reestablished to evaluate whether or not any DC offset had built up during the course of the experiment. In all cases, the offset was less than 1 mV.

Acknowledgments

This work was supported by grants from the DOE and NIH. G.M. holds an MD-PhD training fellowship at The Texas A&M University System Health Science Center, College of Medicine, and was the recipient of an ASSERT (ARO) award. The authors thank Daniel Zeigler, BGSC Director, Ohio State University, for graciously supplying *Bacillus* strains; Orit Braha and Li-Qun Gu for their advice on ion selectivity; Brian Lauman for technical help; Michael Palmer for advice on liposomes; and Sean Conlan for guidance on using Spock.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Alouf, J.E. and Freer, J.H. 1999. *The comprehensive sourcebook of bacterial protein toxins*. Academic Press, New York.
- Altshul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* **25**: 3389–3402.
- Baida, G., Budarina, Z.I., Kuzmin, N.P., and Solonin, A.S. 1999. Complete

- nucleotide sequence and molecular characterization of hemolysin II gene from *Bacillus cereus*. *FEMS Microbiol. Lett.* **180**: 7–14.
- Bayley, H. 1999. Designed membrane channels and pores. *Curr. Opin. Biotechnol.* **10**: 94–103.
- Bayley, H. and Cremer, P.S. 2001. Stochastic sensors inspired by biology. *Nature* **413**: 226–230.
- Bayley, H., Braha, O., and Gu, L.-Q. 2000. Stochastic sensing with protein pores. *Adv. Mater.* **12**: 139–142.
- Bhakdi, S., Walev, I., Palmer, M., and Valeva, A. 2000. Staphylococcal α toxin. In *Bacterial protein toxins* (eds. K. Aktories and I. Just), pp. 509–527. Springer, Berlin.
- Braha, O., Walker, B., Cheley, S., Kasianowicz, J.J., Song, L., Gouaux, J.E., and Bayley, H. 1997. Designed protein pores as components for biosensors. *Chem. Biol.* **4**: 497–505.
- Cheley, S., Braha, O., Lu, X., Conlan, S., and Bayley, H. 1999. A functional protein pore with a “retro” transmembrane domain. *Protein Sci.* **8**: 1257–1267.
- Cheley, S., Malghani, M.S., Song, L., Hobaugh, M., Gouaux, J.E., Yang, J., and Bayley, H. 1997. Spontaneous oligomerization of a staphylococcal α -hemolysin conformationally constrained by removal of residues that form the transmembrane β barrel. *Protein Eng.* **10**: 1433–1443.
- Christopher, J.A. 1998. SPOCK: The structural properties observation and calculation kit (program manual). Center for Macromolecular Design, Texas A&M University, College Station, TX.
- Drobeniowski, F.A. 1993. *Bacillus cereus* and related species. *Clin. Microbiol. Rev.* **6**: 324–338.
- Eroglu, A., Russo, M.J., Bieganski, R., Fowler, A., Cheley, S., Bayley, H., and Toner, M. 2000. Intracellular trehalose improves the survival of cryopreserved mammalian cells. *Nat Biotechnol* **18**: 163–167.
- Fang, Y., Cheley, S., Bayley, H., and Yang, J. 1997. The heptameric prepore of a staphylococcal α -hemolysin mutant in lipid bilayers imaged by atomic force microscopy. *Biochemistry* **36**: 9518–9522.
- Gouaux, J.E., Braha, O., Hobaugh, M.R., Song, L., Cheley, S., Shustak, C., and Bayley, H. 1994. Subunit stoichiometry of staphylococcal α -hemolysin in crystals and on membranes: A heptameric transmembrane pore. *Proc. Natl. Acad. Sci.* **91**: 12828–12831.
- Gouaux, E., Hobaugh, M., and Song, L. 1997. α -Hemolysin, γ -hemolysin and leukocidin from *Staphylococcus aureus*: Distant in sequence but similar in structure. *Protein Sci.* **6**: 2631–2635.
- Gouet, P., Courcelle, E., Stuart, D.I., and Metz, F. 1999. ESPript: Analysis of multiple sequence alignments in PostScript. *Bioinformatics* **15**: 305–308.
- Gu, L.-Q., Dalla Serra, M., Vincent, J.B., Vigh, G., Cheley, S., Braha, O., and Bayley, H. 2000. Reversal of charge selectivity in transmembrane protein pores by using non-covalent molecular adapters. *Proc. Natl. Acad. Sci.* **97**: 3959–3964.
- Hardy, S.P., Lund, T., and Granum, P.E. 2001. CytK toxin of *Bacillus cereus* forms pores in planar lipid bilayers and is cytotoxic to intestinal epithelia. *FEMS Microbiol. Lett.* **197**: 47–51.
- Heginbotham, L., Odessey, E., and Miller, C. 1997. Tetrameric structure of a prokaryotic K^+ channel. *Biochemistry* **36**: 10335–10342.
- Henikoff, S. and Henikoff, J.G. 1992. Amino acid substitution matrices from protein blocks. *Proc. Natl. Acad. Sci.* **89**: 10915–10919.
- Hille, B. 2001. *Ion channels of excitable membranes*. Sinauer, Sunderland, MA.
- Howorka, S. and Bayley, H. 1998. Improved protocol for high-throughput cysteine scanning mutagenesis. *Biotechniques* **25**: 766–772.
- Howorka, S., Cheley, S., and Bayley, H. 2001a. Sequence-specific detection of individual DNA strands using engineered nanopores. *Nat Biotechnol* **19**: 636–639.
- Howorka, S., Movileanu, L., Braha, O., and Bayley, H. 2001b. Kinetics of duplex formation for individual DNA strands within a single protein nanopore. *Proc. Natl. Acad. Sci.* **98**: 12996–13001.
- Jones, D.H. 1995. PCR mutagenesis and recombination in vivo. In *PCR primer: A laboratory manual* (eds. C.W. Dieffenbach and G.S. Dveksler), pp. 591–601. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Korchev, Y.E., Alder, G.M., Bakhranov, A., Bashford, C.L., Joomun, B.S., Sviderskaya, E.V., Usherwood, P.N.R., and Pasternak, C.A. 1995. *Staphylococcus aureus* alpha-toxin-induced pores: Channel-like behavior in lipid bilayers and patch clamped cells. *J. Membr. Biol.* **143**: 143–151.
- Krasilnikov, O.V., Merzlyak, P.G., Yuldashева, L.N., Rodrigues, C.G., Bhakdi, S., and Valeva, A. 2000. Electrophysiological evidence for heptameric stoichiometry of ion channels formed by *Staphylococcus aureus* alpha-toxin in planar lipid bilayers. *Mol. Microbiol.* **37**: 1372–1378.
- Kraulis, P.J. 1991. MOLSCRIPT: A program to produce both detailed and schematic plots of protein structure. *J. Appl. Crystallogr.* **24**: 946–949.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680–685.
- Lund, T., De Buysse, M.L., and Granum, P.E. 2000. A new cytotoxin from *Bacillus cereus* that may cause necrotic enteritis. *Mol. Microbiol.* **38**: 254–261.
- Martín, A.C., López, R., and García, P. 1996. Analysis of the complete nucleotide sequence and functional organization of the genome of the *Streptococcus pneumoniae* bacteriophage Cp-1. *J. Virol.* **70**: 3678–3687.
- Menestrina, G., Dalla Serra, M., and Prévost, G. 2001. Mode of action of β barrel pore-forming toxins of the staphylococcal α -hemolysin family. *Toxicon* **39**: 1661–1672.
- Miles, G., Cheley, S., Braha, O., and Bayley, H. 2001. The staphylococcal leukocidin bicomponent toxin forms large ionic channels. *Biochemistry* **40**: 8514–8522.
- Miles, G., Movileanu, L., and Bayley, H. 2002. Subunit composition of a bicomponent toxin: Staphylococcal leukocidin forms an octameric transmembrane pore. *Protein Sci.* **11**: 894–902.
- Miyata, S., Matsushita, O., Minami, J., Katayama, S., Shimamoto, S., and Okabe, A. 2001. Cleavage of a C-terminal peptide is essential for heptamerization of *Clostridium perfringens* ϵ -toxin in the synaptosomal membrane. *J. Biol. Chem.* **276**: 13778–13783.
- Montal, M. and Mueller, P. 1972. Formation of bimolecular membranes from lipid monolayers and study of their electrical properties. *Proc. Natl. Acad. Sci.* **69**: 3561–3566.
- Movileanu, L., Cheley, S., Howorka, S., Braha, O., and Bayley, H. 2001. Location of a constriction in the lumen of a transmembrane pore by targeted covalent attachment of polymer molecules. *J. Gen. Physiol.* **117**: 239–251.
- Olson, R., Nariya, H., Yokota, K., Kamio, Y., and Gouaux, E. 1999. Crystal structure of staphylococcal LukF delineates conformational changes accompanying formation of a transmembrane channel. *Nat. Struct. Biol.* **6**: 134–140.
- Pédrelacq, J.-D., Maveyraud, L., Prévost, G., Baba-Moussa, L., González, A., Courcelle, E., Shepard, W., Monteil, H., Samama, J.-P., and Mourey, L. 1999. The structure of *Staphylococcus aureus* leukocidin component (LukF-PV) reveals the fold of the water-soluble species of a family of transmembrane pore-forming toxins. *Structure* **7**: 277–288.
- Sakash, J.B. and Kantrowitz, E.R. 2000. The contribution of individual inter-chain interactions to the stabilization of the T and R states of *Escherichia coli* aspartate transcarbamoylase. *J. Biol. Chem.* **275**: 28701–28707.
- Schneider, F., Hammarström, P., and Kelly, J.W. 2001. Transhyretin slowly exchanges subunits under physiological conditions: A convenient chromatographic method to study subunit exchange in oligomeric proteins. *Protein Sci.* **10**: 1606–1613.
- Shatursky, O., Bayles, R., Rogers, M., Jost, B.H., Songer, J.G., and Tweten, R.K. 2000. *Clostridium perfringens* beta-toxin forms potential-dependent, cation-selective channels in lipid bilayers. *Infect. Immun.* **68**: 5546–5551.
- Song, L., Hobaugh, M.R., Shustak, C., Cheley, S., Bayley, H., and Gouaux, J.E. 1996. Structure of staphylococcal α -hemolysin, a heptameric transmembrane pore. *Science* **274**: 1859–1865.
- Steinthorsdottir, V., Halldorsson, H., and Andresson, O.S. 2000. *Clostridium perfringens* beta-toxin forms multimeric transmembrane pores in human endothelial cells. *Microb. Pathog.* **28**: 45–50.
- Thompson, J.D., Higgins, D.G., and Gibson, T.J. 1994. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**: 4673–4680.
- Valeva, A., Weisser, A., Walker, B., Kehoe, M., Bayley, H., Bhakdi, S., and Palmer, M. 1996. Molecular architecture of a toxin pore: A 15-residue sequence lines the transmembrane channel of staphylococcal alpha-toxin. *EMBO J.* **15**: 1857–1864.
- Walker, B.J., Krishnaswamy, M., Zorn, L., and Bayley, H. 1992a. Assembly of the oligomeric membrane pore formed by staphylococcal α -hemolysin examined by truncation mutagenesis. *J. Biol. Chem.* **267**: 21782–21786.
- Walker, B.J., Krishnaswamy, M., Zorn, L., Kasianowicz, J.J., and Bayley, H. 1992b. Functional expression of the α -hemolysin of *Staphylococcus aureus* in intact *Escherichia coli* and in cell lysates. *J. Biol. Chem.* **267**: 10902–10909.
- Welkos, S.L. 1991. Plasmid-associated virulence factors of non-toxicogenic (pX01-) *Bacillus anthracis*. *Microb. Pathog.* **10**: 183–198.
- Zemaitis, J.F., D.M. Clark, M. Rafal, and N. Scriven. 1986. Handbook of aqueous electrolyte thermodynamics: Theory and application. American Institute of Chemical Engineers, New York.
- Zitzer, A., Zitzer, O., Bhakdi, S., and Palmer, M. 1999. Oligomerization of *Vibrio cholerae* cytotoxin yields a pentameric pore and has a dual specificity for cholesterol and sphingolipids in the target membrane. *J. Biol. Chem.* **274**: 1375–1380.