
Identification of four proteins from the small subunit of the mammalian mitochondrial ribosome using a proteomics approach

EMINE CAVDAR KOC,¹ WILLIAM BURKHART,² KEVIN BLACKBURN,² HASAN KOC,³ ARTHUR MOSELEY,² AND LINDA L. SPREMULLI¹

¹Department of Chemistry, University of North Carolina, Chapel Hill, North Carolina 27599, USA

²Glaxo Wellcome Research and Development Department of Analytical Chemistry, Research Triangle Park, North Carolina 27709, USA

³School of Public Health, Environmental Sciences and Engineering, University of North Carolina, Chapel Hill, North Carolina 27599–3290, USA

(RECEIVED August 15, 2000; FINAL REVISION November 16, 2000; ACCEPTED November 28, 2000)

Abstract

Proteins in the small subunit of the mammalian mitochondrial ribosome were separated by two-dimensional polyacrylamide gel electrophoresis. Four individual proteins were subjected to in-gel Endoprotease Lys-C digestion. The sequences of selected proteolytic peptides were obtained by electrospray tandem mass spectrometry. Peptide sequences obtained from in-gel digestion of individual spots were used to screen human, mouse, and rat expressed sequence tag databases, and complete consensus cDNAs for these species were deduced in silico. The corresponding protein sequences were characterized by comparison to known ribosomal proteins in protein databases. Four different classes of mammalian mitochondrial small subunit ribosomal proteins were identified. Only two of these proteins have significant sequence similarities to ribosomal proteins from prokaryotes. These proteins are homologs to *Escherichia coli* S9 and S5 proteins. The presence of these newly identified mitochondrial ribosomal proteins are also investigated in the *Drosophila melanogaster*, *Caenorhabditis elegans*, and in the genomes of several fungi.

Keywords: Mitochondria; protein synthesis; ribosome; proteomics; mass spectrometry; ribosomal protein

Mammalian mitochondria carry out the synthesis of 13 polypeptides that are essential for oxidative phosphorylation and, hence, for the synthesis of the majority of the adenosine triphosphate (ATP) used by eukaryotic organisms. The ribosomes present in mammalian mitochondria are 55–60 S particles and are composed of small (28 S) and large (39 S) subunits. They are characterized by a low percentage of rRNA and a compensating increase in the number of ribosomal proteins (De Vries et al. 1973; O'Brien et al. 1976). The small subunit of the ribosome contains a 12 S rRNA and about 30 proteins, whereas the large subunit consists of a 16 S rRNA and about 50 proteins (Matthews et al. 1982). This total of about 80 proteins is significantly higher than that observed in bacterial ribosomes and may exceed the number of proteins present in eukaryotic cytoplasmic ribosomes (Wittmann-Liebold et al. 1980).

Considerable progress has been made on the identification of the mitochondrial ribosomal proteins (MRPs) in yeast (*Saccharomyces cerevisiae*). About 50 different mitochondrial ribosomal proteins have been identified in this organism (Graack et al. 1998). Additional protein components in yeast mitochondrial ribosomes remain to be determined. Surprisingly, less than half of the MRPs in yeast show significant sequence identities to the ribosomal proteins of other systems (Graack and Wittmann-Liebold 1998).

Although the importance of the mitochondrial translation products is well known, limited information is available on the identities of the proteins that are parts of the translational machinery in mitochondria or on their relationships to bacterial ribosomal proteins. Recently, 18 proteins of the large subunit and 11 proteins of the small subunit of the mammalian mitochondrial ribosome have been characterized primarily by peptide sequencing coupled to the extensive use of the expressed sequence tag (EST) databases to deduce the full-length cDNAs and the corresponding amino acid sequences (Goldschmidt-Reisin et al. 1998; Graack et al. 1999; Koc et al. 1999, 2000; Mariottini et al. 1999;

Reprint request to: Linda L. Spremulli, Department of Chemistry, Campus Box 3290, University of North Carolina, North Carolina 27599-3290, USA; e-mail: Linda_Spremulli@unc.edu; fax: (999) 966-3675.

Article and publication are at www.proteinscience.org/cgi/doi/10.1110/ps.35301.

O'Brien et al. 1999). Of these proteins, 12 from the large subunit and 4 from the small subunit are homologs of bacterial ribosomal proteins. The remainder fall into new classes of ribosomal proteins. In the current work, peptide sequence information has been obtained for four new mitochondrial small subunit ribosomal proteins. The cDNAs and amino acid sequences have been assembled using EST databases, and the genomes of *S. cerevisiae*, *Drosophila melanogaster*, and *Caenorhabditis elegans* have been searched for homologs. Only two out of the four new small subunit proteins found in mammalian mitochondrial ribosomes are similar to prokaryotic ribosomal proteins. The remaining two small subunit proteins fall into new classes of ribosomal proteins.

Results

Characterization of bovine MRPs by tandem mass spectrometry

As a first step toward understanding the protein components of mammalian mitochondrial ribosomes, small subunit proteins from bovine mitochondrial ribosomes were separated by two-dimensional polyacrylamide gel electrophoresis using the system developed by Cahill et al. with rat mitochondrial ribosomes (1995). Four individual protein spots that were well resolved and that had not been identified previously were selected for analysis (Fig. 1). These spots were excised, and the proteins present were subjected to in-gel digestion using endoprotease Lys-C. After digestion, the resulting peptide mixtures were analyzed by nanoscale capillary LC/MS/MS using a quadrupole time-of-flight mass spectrometer (Q-TOF) with data-dependent MS to MS/MS switching. As peptides are eluted from the nanoscale LC column and detected by the mass spectrometer, the mass spectrometer switches from MS to MS/MS mode in which product ion (fragmentation) spectra are obtained from each peptide to be used for determining the primary sequence of the peptide.

Each uninterpreted peptide product ion spectrum was initially searched against the nonredundant protein and EST databases for exact matches using the Mascot search program (Perkins et al. 1999). In some cases, high-quality product ion spectra that had no exact matches in the databases were sequenced *de novo*. A representative spectrum obtained from the protein designated as MRP-S27 and sequenced *de novo* is shown in Figure 2. Peptide sequences obtained from Mascot database searches and *de novo* sequencing are shown in Table 1.

Nomenclature

Previous work tentatively identified 33 proteins in the small subunit of bovine mitochondrial ribosomes (Matthews et al. 1982). These proteins were designated S1 to S33 in order of decreasing molecular weight. However, this nomenclature




Fig. 1. Separation of bovine 28 S subunit proteins by two-dimensional gel electrophoresis using a NEPHGE-gel system developed previously (Cahill et al. 1995) for the separation of rat mitochondrial ribosomal proteins. The gel spots chosen for the in-gel tryptic digests are shown in ovals.

system does not provide a consistent set of designations for the MRPs from different mammals. For example, using this system, bovine MRP-S18 is the same protein as rat MRP-S13 (O'Brien et al. 2000). To simplify cross-species comparisons and to facilitate the correlation between proteins in mammalian mitochondrial ribosomes and in bacteria, we are using a nomenclature recently adopted for chloroplast ribosomal proteins (Yamaguchi and Subramanian 2000; Yamaguchi et al. 2000). In this system, proteins are designated by their prokaryotic homolog (e.g., S12 in bacteria is designated PRP-S12 in plastids and MRP-S12 in mammalian mitochondria). Proteins without bacterial homologs are given the next available number. Because there are 21 proteins in the bacterial ribosome, we began designating the new mammalian MRPs at MRP-S22 (Koc et al. 2000). The current manuscript describes two homologs of bacterial ribosomal proteins (MRP-S5 and MRP-S9) and two new small subunit ribosomal proteins (MRP-S27 and MRP-S28). The MRP-S28 that is described here is not related to yeast MRP-S28 (YDR337Wp) (Graack et al. 1998) or to the bovine MRP-S28 described previously (O'Brien et al. 2000).

Overall approach to the assembly of mammalian MRP sequences from cDNA clones

Sequences of the peptides obtained from bovine mitochondrial ribosomes (Table 1) were used to search the human EST database using the tBLASTN program (National Center for Biotechnology Information). A number of hits were obtained for some, but not all, of the peptides used as virtual screening probes. Overlapping clones for these hits were obtained using the initial hits as virtual probes to rescreen the human EST database. For all the peptide sequences obtained, consensus cDNAs were assembled by repetitive searching and comparison of EST sequences. The sequence




Fig. 2. Product ion spectrum of the tryptic peptide at m/z 831.8 from MRP-S27. Because there were no exact matches in the protein or EST databases, sequence was called de novo and is shown using the single letter amino acid symbols. Ions are labeled according to the nomenclature of Roepstorff and Fohlman (Roepstorff and Fohlman 1984), with the difference in m/z of adjacent singly charged y ions corresponding to the labeled amino acid residue mass. For the isobaric amino acids leucine and isoleucine, both possibilities are shown as L/I.

of the longest possible cDNA was then assembled in silico. Sequencing errors were corrected as much as possible by comparison of overlapping clones. Generally, the fully assembled human sequence was then used as a query against entries in the mouse and rat ESTs. Mouse and rat consensus cDNAs were assembled for each protein when possible.

MRP- S5

A protein spot with a molecular weight of about 35 kD (see Fig. 1) was excised, and peptide sequence information was obtained as described above. Only one peptide sequence was obtained using this approach (Table 1). However, a second peptide sequence was obtained from the proteolytic digestion of whole 28 S ribosomal subunits followed by direct analysis by nanoscale capillary LC/MS/MS (E.C. Koc, K. Blackburn, W. Burkhart, A. Moseley, and L.L. Sprelli, in prep.; Table 1). Database searches with these peptides hit a number of EST clones described as similar to ribosomal protein S5 or MRP S5. Assembly of overlapping EST clones allowed the deduction of the complete sequence of human MRP-S5 (Fig. 3). The chromosomal gene for mitochondrial S5 has been located on human chromosome 5 (contig AC011416). It covers about 1.3 kb of DNA and consists of only one exon. Analysis of the sequence near the N-terminus of MRP-S5 for a mitochondrial import signal was performed using PSort, which predicts cleavage following residue 86. MitoProt II gives this protein a 78% chance of being mitochondrially localized and predicts cleavage following residue 87. The predicted mature form of human mitochondrial MRP-S5 has a calculated molecu-

lar mass of 39 kD, which corresponds well to the predicted molecular weight from the two-dimensional gel.

The sequence of human MRP-S5 was used to probe the mouse EST databases and allowed the deduction of the full-length sequence from this species. Mouse MRP-S5 is 79% identical to the human MRP-S5 (Table 2). The *C. elegans* genome has a homolog of human MRP-S5 (Swiss Prot Q93425). The *D. melanogaster* genome also has a putative MRP-S5 homolog. However, the full-length *D. melanogaster* homolog could not be obtained from the given sequences (AE003072 and EST AA438342). Alignment of the putative partial fly and human sequences indicates that they are closely related (Table 2).

The yeast mitochondrial ribosome also contains a homolog to prokaryotic S5 (Fig. 3; Table 2). Interestingly, the fungal MRP-S5 is no more closely related to mammalian MRP-S5 than are the bacterial proteins (Table 2). This trend has been observed with a number of the MRPs in yeast and mammals (Koc et al. 1999). Human MRP-S5 is considerably longer than prokaryotic S5 (Fig. 3; Table 2). There is a long extension at the N-terminus (about 180 residues) that is longer than the bacterial factors even after removal of the mitochondrial import signal. There is also a long C-terminal extension of more than 50 amino acids. These additional sequences also appear to be present in worm MRP-S5 (Fig. 3). They may play a role in interacting with some of the MRPs not present in prokaryotic and yeast mitochondrial ribosomes. The longer form of MRP-S5 found in mammalian mitochondrial ribosomes may also reflect the small rRNA present in the 28 S subunit or reflect basic differences

Table 1. Amino acid sequences of mature MRPs of *Bos taurus* deduced from database searching of peptide product ion spectra of two-dimensional gel spots and whole 28S subunit^a

Protein name	Sequence	Swiss-Prot	Family
MRP-S5		P82675 (H)	S5
Peptide 1 ^a	LTADELWK	P82674 (B)	
Peptide 2	...EPEDEVDP(L/I)Q...*		
MRP-S27		Q92552 (H)	New
Peptide 1 ^a	EDCHLADLASLMDK	P82677 (B)	
Peptide 2 ^a	ALYTLVNK		
Peptide 3 ^a	VQYGIFPDNYTFNLLMDHFIKK		
Peptide 4 ^a	KTDFSWEERNNFGASLLLPGLK		
Peptide 5	...EQ(L/I)DVEETEQSK*		
Peptide 6	VESES(L/I)(L/I)T(L/I)TTQ(L/I)VK*		
MRP-S9		P82676 (B)	S9
Peptide 1 ^a	RHLANMMGEDPETFTQEDVDRAITYLFP SGLFEK		
Peptide 2 ^a	QRAVQWGEDGRPFHFLFYTGK		
Peptide 3	QSYYS(L/I)MHEAYGK*		
Peptide 4	EELEEMLVEK*		
Peptide 5	...IEP(L/I)QYDEQGM AFST...		
Peptide 6	QIEEFNIK		
MRP-S28		P82673 (H)	New
		P82672 (B)	
Peptide 1 ^a	EGNLELLK		
Peptide 2 ^a	IPNFLHLTPVAIK (IPNFLHLTPVAIKK)		
Peptide 3 ^a	LKNYCTEWPAAALDSDEK		
Peptide 4 ^a	LSSLNLDDHAK		
Peptide 5	(L/I)ET(L/I)FQ(L/I)K		

^a Peptides are obtained from the digestion of whole 28S subunit by Endoprotease Lys-C. Molecular masses and pIs of proteins were determined from the predicted mature protein sequences by PSort from fully assembled cDNAs. (*) Sequence for these peptides obtained de novo, with (...) representing internally called sequence; (H) SwissProt number for the complete human sequence; and (B) SwissProt number for the bovine peptides.

in the mechanism of chain initiation in animal mitochondria compared with other systems.

MRP-S27

Peptide sequence information from this protein allowed deduction of the full coding sequence from data in the EST databases and GenBank from both humans and mice. Human MRP-S27 is 414 amino acids in length (Fig. 4). The mature protein is predicted to be 378 residues in length (MitoProt II) with a molecular mass of 43.8. Homologs of human MRP-S27 are detectable in *C. elegans* (CE16258) and, interestingly, in *Fugu rupripes* (puffer fish; AAC60297; Fig. 4; Table 2). The fish protein is 50.7% identical to human MRP-S27. The *D. melanogaster* genome has a region with a sequence similar to human and *C. elegans* MRP-S27. However, this predicted protein is only 22.5% identical to human MRP-S27 and is not in a predicted reading frame. No protein corresponding to human MRP-S27 can be detected in the yeast genome.

Analysis of the amino acid sequence of the MRP-S27 using PROSITE does not indicate the presence of any common motifs, including RNA binding or ribosomal protein

motifs. However, MRP-S27 does possess one of three conserved blocks of sequence found in proteins that stimulate the dissociation of guanine nucleotides from G-proteins (Sone et al. 1997; Hoshino et al. 1999). Although this conservation is quite limited, the presence of a GTP-binding ribosomal protein in the small subunit of mammalian mitochondrial ribosomes (Denslow et al. 1991)¹ leaves open the possibility that MRP-S27 might be a functional partner of that GTP-binding ribosomal protein.

MRP-S9

Six peptides were obtained from a basic protein with a molecular mass of about 41 kD using mass spectrometry (Fig. 1; Table 1). Using the peptide sequences as virtual screening probes, the complete sequence of human MRP-S9 protein was assembled (Fig. 5). The MRP-S9 protein is 396 residues in length. The import signal of human MRP-S9

¹ Presence of a GTP-binding protein in the mammalian mitochondrial small subunit has been confirmed by the recent discovery of a putative GTP-binding protein (E.C. Koc, K. Blackburn, W. Burkhart, A. Moseley, and L.L. Spemulli, in prep.)

MRP-S5

Human	1	MATAVRAVGCLPVLCSGTAGHLLGRQCSLNTLPAASILAWKSVLGNHGLSSLGTRDTHPY
<i>C.elegans</i>	1	-----
<i>E.coli</i>	1	-----
Yeast	1	-----
		↓
Human	61	ASLSRALQTQCCISSP SH MSQ Q YRPY SFF EL LA DELWKGALAE GAGAKKGRGAR IK AK
<i>C.elegans</i>	1	-----MRRSGPELWKT TSVSK Q Q KKGR--RNT R
<i>E.coli</i>	1	-----
Yeast	1	-----MFKRQLS S RYL CH YDESLL RYYP SL LK K KL AQQ I PEDT FRV IN
Human	121	KKR DLNR GQI IG EGR GFL PGLN VPI ---MKN GA Q IA QR SK E Q E K V AD Q QR
<i>C.elegans</i>	29	QPV PLNR FYR IG SSE MKIE AGLN AP IRMRE TEN ON MS IA EQ EE E IRDSM GG T K IL
<i>E.coli</i>	1	-----
Yeast	52	VEFA P PYLDDFT KI HP WDYK PG N PHLHAQ--E EN N N F S I F RWDQ VQ PL P GE GN PPG
Human	177	EEWDR KKK M V KR---ER G SC N SWG G IS L G P PD F GP Q ET ED F T LE V N V FT M
<i>C.elegans</i>	89	EEEDT GKK R N RE K LHP M ERG S GT Q L Y Q K L G AP P LD G V N ED F TY C LE V K R IS N M
<i>E.coli</i>	1	-----A H TE K O A G---
Yeast	110	VSLPND GG K S K S -----AD V A A GL H K Q T G V D P---DY I TR K LT M K P L V K R VS N Q
Human	232	T K E G R K S I R V L V AV G NG K GA G ES G K A T L ---M D A F R K A K N A V E L L Y E R Y E D E T
<i>C.elegans</i>	148	T N V F G R M M A L V V T G N C R G A G A G K A P I R T T A I N G M G M A S R K L F H E L H E G R T
<i>E.coli</i>	23	T V K G G R I F S F I A L T V V G D G N G R V G F G Y G K A R E V P --A A I O K A M E K A R R N I N V A L N N G T
Yeast	158	T K -G A I A S F Y A L V V G D K N G V L G E G K S R E E -M S K A I F K A H W D A V R N L E P R Y E N R T
Human	290	I H D I S I R T R T K M K Q P G GL R CH A I T TC R I G I K D V Y A K V S G S I N-M L S L T O
<i>C.elegans</i>	208	I Y Q D F Y E C N T R V F A O N R P G F G L T C H P R I K I C E A I G I K D V Y V K V E G S T K N Y L A L T
<i>E.coli</i>	80	L Q H P K V H T G S R V F M Q P A S E T C I A G G A R A L E A G N N V L A K A Y G S T N -P I N V R A
Yeast	216	I Y G D I D F R V G V K T H R S A K P G F G L R V N V I F E I C E C A G I K D L S C K V Y K S R N-D I N A N G
Human	349	L F R G L R -Q E T H Q A A K G L H V V E I R E E C G P L I V A S P R G P L R E D P E P E V P D K L D
<i>C.elegans</i>	268	F I G L N-Q E T H Q A A K G L H V V E I S P S R H E L Q V A S P I S T E L K T E T L A D R N L D
<i>E.coli</i>	139	T I G L N-M N P E M A A K G K S V E I L G -
Yeast	275	T I L F K A Q K T L D V A L G E G K V V R K V Y Y S -----
Human	408	WEDV K T A Q G ---S V S N L K R A A T -----
<i>C.elegans</i>	327	D F Y G E G R Y P P K S L P F S N L E G H L D A R W R K H P F R N Q E S T M I R L I A D N M V P R W T R D A R A
<i>E.coli</i>		-----
Yeast		-----
Human		-----
<i>C.elegans</i>	387	AWAD Q R N E R M T T G V E P M P L G I G L S H V V P K K D
<i>E.coli</i>		-----
Yeast		-----

Fig. 3. Sequence of human MRP-S5 (P82675) and its alignment with homologs from *C. elegans* (Q93425), *E. coli* (P02356), and yeast (P33759, YBR251W or YBR1704). (↓) Shows the signal peptide cleavage site predicted by MitoProtII and, additionally, represents the predicted peptide signal cleavage sites in Figures 4, 5, and 6.

protein is predicted to be 52 residues in length with cleavage occurring at an R-2 motif as predicted by PSort and MitoProtII programs. Overall mouse and human MRP-S9 proteins are 78.8% identical (Table 2). However, the N termi-

nus of MRP-S9 was not found in the mouse EST database. This sequence indicates that only a few residues of the import signal of mouse MRP-S9 are missing from the assembled sequence.

Table 2. Percentage identity of human mitochondrial ribosomal proteins that are also found in other species

Human protein	Percentage identity to				
	Mouse	<i>Drosophila</i>	<i>Caenorhabditis elegans</i>	Yeast	<i>Escherichia coli</i>
MRP-S5	79.0	46.5 ^a	35.3	23.3	24.7
MRP-S27	77.1	22.5 ^b	22.3	n.d.	n.d.
MRP-S9	78.8	33.4	36.5	33.0	38.6
MRP-S28	76.8	44.3	37.9	27.1	n.d.

(n.d.) Homologs of human ribosomal proteins are not detected in database searches.

^a Only partial sequences could be found in the databases.

^b Percentage identity is calculated by using a putative *Drosophila* homolog that is not found in the same frame as the predicted reading frame of AAF47311.

Proteins related to human and mouse MRP-S9 are present in *C. elegans*, *D. melanogaster*, and yeast (Fig. 5; Table 2). Analysis of the sequences obtained indicates that this protein falls into the S9 family of ribosomal proteins. However, in agreement with the size of MRP-S9 on the two-dimensional gel, the mammalian mitochondrial protein is considerably longer than its bacterial counterparts. Mammalian MRP-S9 begins to align with the bacterial proteins around residue 255 (Fig. 5). The alignment continues through to the C terminus. Interestingly, the *C. elegans* MRP-S9 appears to be even longer than the mammalian protein. The N-terminal half of MRP-S9 may be interacting with one or more of the ribosomal proteins in mitochondria that do not have prokaryotic homologs.

Yeast mitochondrial MRP-S9 (YBR146W or YBR1123) is 30.1% identical to the mouse protein (Table 2). It is intermediate in length between the mammalian and prokaryotic proteins (Fig. 5). Again, the C-terminal region of the yeast protein corresponds to bacterial S9 with additional residues located at the N terminus. The mitochondrial import sequence of yeast MRP-S9 is predicted to be only 11 residues in length. Hence, yeast MRP-S9 is probably about 100 residues longer than the bacterial proteins. The human cytoplasmic homolog of S9 (Rsp 16, P17008) also has considerable sequence identity (37.1%) to human MRP-S9 especially near the C-terminal end.

MRP-S28

In-gel digestion of a spot on the two-dimensional gel at a molecular weight of about 33.5 kD generated a single peptide. However, additional peptides from this protein were obtained from proteolytic digests of whole 28 S subunits (Table 1). The complete cDNA for this protein was found in the TIGR Tentative Human Consensus (THC) sequences and encodes a protein of 323 amino acid residues (Fig. 6). MitoProtII assigns a 95% probability that this protein is

localized in mitochondria and predicts a cleavage position right after residue 30. Partial sequences for this protein were also found in the mouse and rat EST databases. Searches of the *C. elegans* and *D. melanogaster* databases also indicate homologs in these species (Table 2). A homolog of this protein was detected in the yeast genome using the *C. elegans* and *D. melanogaster* sequences as the query in advanced BLAST searches. Yeast MRP-S28 (NP_010460, YDR175Cp) is 319 aa residues in length and 27.1% identical to the human protein. The yeast MRP-S28 (YDR175Cp) described here was not previously identified as a ribosomal protein. MRP-S28 is the third representative of a MRP that is shared between mammals and yeast that is not represented in prokaryotic ribosomes.

Database searches did not reveal any proteins with known functions that correspond to MRP-S28, and no ribosomal proteins from prokaryotic organisms show significant similarities to it. Therefore, MRP-S28 represents a new class of ribosomal protein.

Discussion

The identification and characterization of the proteins present in mitochondrial ribosomes has been difficult because of their low abundance. During the past several years, significant progress has been made in identifying some of the proteins present by using a combination of protein sequencing and database searching. In the present report, two proteomics-based approaches consisting of either two-dimensional PAGE coupled with high-sensitivity peptide sequencing by mass spectrometry or direct peptide sequencing by mass spectrometry have been used to identify four different classes of mammalian mitochondrial ribosomal small subunit proteins that had not been characterized previously. This work increases the number of proteins identified in the small subunit to 15.

The small subunit of the mammalian mitochondrial ribosome is thought to have 33 different proteins based on two-dimensional PAGE. This number is considerably higher than the 21 proteins found in the 30S subunit of the prokaryotic ribosome (Wittmann 1986). Therefore, a number of the mammalian MRPs are not expected to be homologs of prokaryotic ribosomal proteins. Two of the four ribosomal proteins described in this paper and 4 of the previously characterized ribosomal proteins show significant sequence similarities to bacterial ribosomal proteins (Koc et al. 1999, 2000). These are S5, S7, S9, S10, S12, and S14.

Four of the six MRPs that have been identified having prokaryotic homologs (S7, S9, S10, and S14) are located in the head of the small subunit (Samaha et al. 1994). S9, the prokaryotic homolog of MRP-S9 identified in the current work, is classified as a tertiary rRNA-binding protein because its assembly into the small subunit is strongly dependent on the presence of other proteins rather than arising

MRP-S9

Mouse	1	-----WFTPAVR-----ASL
Drosophila	1	-----MALRVFGNIVKNNRNLL-QNACK
E.coli	1	-----
Yeast	1	-----
C.elegans	181	TLSGIFPAIMITAGCLSFADSRVLITYFCNQVLVGI GLVCAVHG VYMPTLLAIFGSDFYQ
		↓
Mouse	12	CQRPGYWTASAVGWQ GTRFQLSKLIHTTVVTT KKNVQASQESYTED KKQIEEFNIG
Drosophila	24	SRAPMPCYMAAPFA DVTVQAAPAAVQQRVS KAMRAYLKRATFDESKTQHL EEFQIG
E.coli	1	-----
Yeast	1	-----MFSRLSLFRRAPAPAPM MPTTIKQKTEDELPRRI
C.elegans	241	NMLRIVLARSSRAMS ASPASASDSDTSVRKIG KAEETYLKSSQIVAM EKHRA EEFETG
Mouse	72	RRLANMMGDEPETFAEEDTDRATAYLFPSGLF KRAPMMPHPHHPKQRATQ GEDG
Drosophila	84	RRLANMMGADAETFTQEDIDEAISYLFPSGLY QKARPAMSPVLPARKAAE DETG
E.coli	1	-----
Yeast	38	VPKLATFYSAIPN--HEERTYRLERL LRKYIKLPSQNNNEAQQTAP ISFDEYALIGGG
C.elegans	301	RRLAKMMSIDTIELDQEAIDRAILYLFPSGLT PNARPVMPPEELPKFQRFTEDEEG
Mouse	132	PFHFLFYTGKQSYSLHHDVYGY---KMQLEK---HRGPLSASAESRD IGSRWLIKQ
Drosophila	144	PFHSMFYTGKPNFQLLHDI VEETNKADLEERML RRGNKPDENQKLEACFQILPKQ
E.coli	1	-----
Yeast	96	TKLPTQYTLQLYMLNKLHNTDP-----CLTN-
C.elegans	361	PEGSRFETLSPKIGLISDITGVKTHS MFKFYDEHVGSRSVNRSDLEPANSGSQWITAE
Mouse	185	EIEEMLVEKLSDEDYAFIRLLKLLTLP CGPAEEEFYQRERRS YTIQSKKQIEPVQYD
Drosophila	203	QIELLLVESIADIEYSFTNSMDIL IASBYAYKSKAFERYMPLMDQSKQLEVPKPRID
E.coli	1	-----AENQYYG-----
Yeast	123	-----DEITSELSQYYKSSMLSNN--KIKT-LD
C.elegans	421	KLKKKLSKFSKELYGVIIAFGLASL EGSIAEQFMEPEPTASGSKFGPAIP
Mouse	245	EQMAFSTSEGLEIERHGTGGRKRSATQORVVYEHGSGKIHGXGVDYIYFPITDREQL
Drosophila	263	EQRQYITTY-----ECLRKARADVTVRLPCGKISINGEDSYFEDENCREQL
E.coli	8	-----TGR-----RKSAAARVFKP-GNGKIVINQSLQYFGETARMV
Yeast	150	EQRSIAVG-----RKSAAKVFVVR-GEGLINGQLNDYFLMKDRES
C.elegans	481	VHCAVTNR-----YAEVTRCKDRATVKVTDAGKGFDTGLQ-HDFRHLAREIL
Mouse	305	MPFHFLRLEHHDVTCVTS-----GGGRSQAQAIRLAAALCSFV
Drosophila	313	LPPIQFSELLGKVDVEANVE-----GGGPSGQAGAIRWGIAIRSEV
E.coli	48	QPLEMMEKLDYITVK-----GGGISGQAGAIRHGITAEIMED
Yeast	197	PLOYSKGYIFATTS-----CCCPGQAEIMHIAALVVEN
C.elegans	535	APMIVSQSLGRFDVTATTS CISNTLPEAPNKAPLMRS GGMSLPRAVRHGTALCAAALQ
Mouse	348	TEEEVEWRQAGLLTDPRI RERKKPGQEGARRK TWKKR
Drosophila	356	DQKTESMRLAGLLTRDYRRERKKFGQEGARRK TWKKR
E.coli	91	-ESRSELRKAGEVTRDARQVERKKVGLRKRARRPQSKR
Yeast	240	-PLKSRLLKACGLTRDYRVERKKKPKKKARMPITWVKR
C.elegans	595	-PAAEPLRLSGLLTLDPRI RERSKVNCPGARAK TWKKR

Fig. 5. Sequence of the human MRP-S9 and its alignment with homologs from *C. elegans* (P34388), *D. melanogaster* (CG2957), yeast (P38120, YBR146W or YBR1123), and *E. coli* (P02363).

1997). This region has been conserved in the 12 S rRNA of mammalian mitochondria. The structure of *Bacillus stearothermophilus* S5 has been determined (Ramakrishnan et al. 1992; Davies et al. 1998). This protein folds into two do-

main that interact strongly with each other through a hydrophobic core formed by the packing of one strand of the β -sheet in domain 1 against two α -helices in domain 2. As expected, the overall fold of MRP-S5 is predicted to be

MRP-S28

Human	1	-----AAALPAWLSLQSRART-----RAFSTAVYSATP-----PTP
Drosophila	1	---SATRIILRLPLKQTAQSG-----RVFSNQEQRCIEEDDETFVLSIR
C.elegans	1	-MRLHVSFRNRSSLAEKLRNAVGGQDFGKAKQIWAAGSIAVKDEHGEDFRRELYVMPKR
Yeast	1	MKVPLGHWKVSRLGNLWSTQKRVLTMSRCLN-----SDAGNEAKTVRFEGPAFSADLYMHPE
		↓
Human	36	SLPERTPGNRP-----PRKALPPRTEMAVEQDWPVVPVAAPTPSA
Drosophila	48	TKQQMQKROQV-----RRDPITPPRTGMAVDQDWTAVWPPRFPPAS
C.elegans	60	KLGGOTQLEQATGRTEYKTRSRFDLMTKTRPPRFEMDPAQRWCDVWPARFASS
Yeast	56	KWKGLPPQRIILELY-----WRRALGSEYKPNKDLALLTSEYNVPVNDL
Human	81	VPLPVRMGTP--VKKGVPAKEGNLELKIPNFLHLTPVAIKHCEA--KDFCTEWPAA
Drosophila	93	VPLPVRMGTP--ERGFAPSKFNAELKIPNFLHLTPPAIQCEA--IKKFTCPWPK
C.elegans	120	VPLPVRMGTRPNVEKRAFKKEGNLELKIPNFLHLTPPAIQQHCA--IKKFTCPPE
Yeast	105	KKLYHRGEQGAIKGNVNSLRPFMFDELPSQAQELAQHREORFYNLAAVELPL
Human	137	LSI-EKCFHFPIETTDYSGPSRNPRARVVLRVKLSLNLDAKAKKILVLG
Drosophila	149	LSI-AKQRHFPEETDYCQSLPIRNPEARVTTIKLSLKFDAHARDKELRLVG
C.elegans	178	LSNPSATCOHPIPISTYTYHCGTSIRDIRSRVVTKIELKLSNQMBKLRILA
Yeast	165	LAQYRQEKRPSPESHVYRYSYVGEHFNRSKVVLSVKTKEGLLELHKFRILAR
Human	196	RYCKTTDTIKTDRCPLRQNYDYAVYLLTVLYHESNTEWEKAKTEADMEYFEN
Drosophila	208	RYKDDTDTFTDRCPLRQNYDYAVYLLTACYHESVTEPWEAKKTEADMEYFEN
C.elegans	238	NRYDEKTGKTLTDRCHTRQONLDYAHYLLTVLYHEQKVEKDELKNRTDALKVEFDG
Yeast	225	SRYDHTTDFKSSDKFEHASQNRARYHDILQRLLAE-----S-KDLTEDDFSIVPLT
Human	256	SSERNILETLLQMKAAEKVINKHEELLGTKEIEEKKSQVSLNNEEENENSQYKES
Drosophila	268	NSKVSACGLWNAKGAQKVPSS-----K--S--FECVEITINEGENEYNGKYKEE
C.elegans	298	SNKTKLIDLLEKAKLTPSPSAAGCGDQKSTIEEEMWKAYNSEEETVEKTREYGRQ
Yeast	278	RH-----AKSLRKKRQ-----YE--FPEHWKPEAPKPK--FDI----
Human	316	VKLLNVT-
Drosophila	319	VKKLNIA-
C.elegans	357	VKLLGLQQ
Yeast	312	VDQLLSTL-

Fig. 6. Sequence of the human MRP-S28 (P82673) and its alignment with homologs from *D. melanogaster* (CG2101), *C. elegans* (CAB61063), and yeast (NP_010460 or YDR175Cp).

similar to that observed with prokaryotic S5 (Swiss Model). The residues forming the hydrophobic core are generally conserved or are conservative replacements. A series of Gly residues creating an overall pattern of turns in S5 are also generally conserved in MRP-S5. A surface loop rich in basic amino acids in the bacterial S5 is thought to interact with the 16 S rRNA. Mutations in this region are known to confer resistance to spectinomycin and are thought to be important for accurate translation. This region in MRP-S5 is modeled as a loop and is also rich in basic residues. Presumably, this region in MRP-S5 interacts with 12 S rRNA and may be important in maintaining ribosomal accuracy.

The two mammalian MRPs classified in new families identified in this study are not functionally characterized yet. Careful investigations of available databases identified homologs of these proteins in the yeast genome, but not in

prokaryotic genomes. The presence of homologs of these two proteins in yeast brings yeast and mammalian mitochondrial translational systems closer to each other. Determination of the location and function of small subunit ribosomal proteins that are not homologs of known prokaryotic ribosomal proteins will require an extensive investigation of the structure of the mammalian mitochondrial ribosome.

Materials and methods

Preparation of bovine MRPs for two-dimensional gel electrophoresis

Bovine mitochondria and 28 S subunits were prepared as described previously by Matthews et al. (1982), and the 28 S subunits were collected by centrifugation at 48,000 rpm for 6 h in a Beckman

Type-50 rotor. The pellet containing about 5 A₂₆₀ (~420 pmole; Hamilton et al. 1974) was resuspended in isoelectric focusing gel buffer containing 9.5 M urea, 2% Triton X-100, 2% ampholytes (consisting of 1.6% [v/v] pH 5–7 and 0.4% [v/v] pH 3–10), and 0.24 M 2-mercaptoethanol. The sample was prepared as described previously before loading on nonequilibrium pH gradient tube gels (Cahill et al. 1995; Koc et al. 1999). Following electrophoresis in the first dimension, gels were equilibrated in buffer (10% glycerol, 2% sodium dodecyl sulfate, 1% dithiothreitol, 62.5 mM Tris-HCl pH 6.8) and subjected to electrophoresis in the second dimension on 12% sodium dodecyl sulfate polyacrylamide gels (SDS-PAGE; Laemmli 1970). Gels were stained with Coomassie Brilliant Blue G-250 to locate protein bands.

Peptide sequencing by mass spectrometry

Four randomly picked spots from the two-dimensional PAGE of the mitochondrial ribosomal small subunit (28 S) were excised and digested in-gel with Endoprotease Lys-C (Boehringer Mannheim) in 10 mM Tris-HCl pH 8.0 according to the procedure of Shevchenko et al. (1996), except that sulfhydryl groups were alkylated using 4-vinylpyridine. Nanoscale capillary liquid chromatography-tandem mass spectrometric (LC/MS/MS) analyses of in-gel digests were performed using an Ultimate capillary LC system (LC Packings) coupled to a quadrupole time-of-flight (Q-TOF) mass spectrometer (Micromass) fitted with a Z-spray ion source as described previously (Koc et al. 1999).

Additionally, 3 pmole of the whole 28 S subunit prepared by the traditional method (Matthews et al. 1982) were subjected to in situ endoprotease Lys C digestion to identify the mitochondrial small subunit ribosomal proteins in the intact subunit (Burkhart 1992). The peptides that were obtained from the whole 28 S subunit digests were also analyzed by LC/MS/MS (E.C. Koc, W. Burkhart, K. Blackburn, A. Moseley, and L.L. Spemulli, in prep.; a preliminary report was presented at the American Society of Mass Spectrometry in June 2000). Uninterpreted peptide product ion spectra generated by LC/MS/MS were searched against the nonredundant protein database and human and rat EST databases for exact matches using the Mascot search program (Perkins et al. 1999). High quality spectra that had no exact matches in either the protein or EST databases were sequenced de novo either manually or with the aid of the PepSeq program (Micromass).

Computational analysis

Peptide sequences obtained from Mascot searches of the protein and EST databases and those obtained by de novo sequencing from peptide product ion spectra were searched against the nonredundant protein database using the FASTA algorithm (Pearson et al. 1988). For peptides with no exact matches in the databases, sequences obtained by de novo sequencing were used for FASTA searches. Because mass spectrometry cannot distinguish between the isobaric (same nominal residue molecular weight) amino acids Leu and Ile, initial database searches were performed using Leu in the peptide sequences. Hits with an Ile at these positions were considered exact matches. If no hits were obtained when Leu was present in the search sequence, the search was reperformed with Ile. The isobaric amino acids Phe and oxidized Met (a common artifact of PAGE) were distinguished by diagnostic loss of methanesulfenic acid (64 Da) from oxidized Met (Jiang et al. 1996). Because the protease endoprotease Lys C, which cleaves on the C-terminal side of Lys residues, was used for in-gel and in situ digestions, Lys residues could be distinguished with a fairly high

certainty from isobaric Gln residues. EST database and genomic DNA searches of the peptide sequences were performed using the BLAST search program (Altschul et al. 1997). Sequence analysis and homology comparisons were performed using the GCG DNA analysis software package (Wisconsin Package Version 10, Genetics Computer Group [GCG]), Vector NTI (Informax Inc.), and Biology WorkBench 3.2. The results were displayed using BOXSHADE (version 3.21, written by K. Hofmann and M. Baron). Prediction of the cleavage sites for the mitochondrial signal sequence was performed using PSort and MitoProt II (Nakai et al. 1992; Claros et al. 1996).

Acknowledgments

The authors thank Mary Moyer (Glaxo Wellcome) for her outstanding technical contributions to this work. This work has been supported by funds provided by the National Institutes of Health (grant GM32734).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Abdurashidova, G.G., Tsvetkova, E.A., and Budowsky, E.I. 1991. Direct tRNA-protein interactions in ribosomal complexes. *Nucleic Acids Res.* **19**: 1909–1915.
- Altschul, S.F., Madden, T.L., Schaffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D.J. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nuc. Acids Res.* **25**: 3389–3402.
- Burkhart, W. 1992. In situ proteolytic digestion performed on proteins bound to the Hewlett Packard hydrophobic sequencing column. In: *Techniques in Protein Chemistry IV* (ed. R. Angeletti), pp. 399–406. Academic Press, New York.
- Cahill, A., Baio, D., and Cunningham, C. 1995. Isolation and characterization of rat liver mitochondrial ribosomes. *Anal. Biochem.* **232**: 47–55.
- Claros, M.G. and Vincens, P. 1996. Computational method to predict mitochondrially imported proteins and their targeting sequences. *Eur. J. Biochem.* **241**: 770–786.
- Clemons, W.M.J., May, J.L., Wimberly, B.T., McCutcheon, J.P., Capel, M.S., and Ramakrishnan, V. 1999. Structure of a bacterial 30S ribosomal subunit at 5.5 Å resolution. *Nature* **400**: 833–840.
- Davies, C., Bussiere, D., Golden, B., Porter, S.J., Ramakrishnan, V., and White, S.W. 1998. Ribosomal proteins S5 and L6: High resolution crystal structures and roles in antibiotic resistance. *J. Mol. Biol.* **279**: 873–888.
- Denslow, N., Anders, J., and O'Brien, T.W. 1991. Bovine mitochondrial ribosomes possess a high affinity binding site for guanine nucleotides. *J. Biol. Chem.* **266**: 9586–9590.
- De Vries, H. and van der Koogh-Schuuring, R. 1973. Physicochemical properties of isolated 55-S mitochondrial ribosomes from rat-liver. *Biochem. Biophys. Res. Comm.* **54**: 308–314.
- Goldschmidt-Reisin, S., Kitakawa, M., Herfurth, E., Wittmann-Liebold, B., Grohmann, L., and Graack, H-R. 1998. Mammalian mitochondrial ribosomal proteins: N-terminal amino acid sequencing, characterization, and identification of corresponding gene sequences. *J. Biol. Chem.* **273**: 34828–34836.
- Graack, H-R., Bryant, M., and O'Brien, T.W. 1999. Identification of mammalian mitochondrial ribosomal proteins (MRPs) by N-terminal sequencing of purified bovine MRPs and comparison to data bank sequences: The large subribosomal particle. *Biochem. J.* **38**: 16569–16577.
- Graack, H-R. and Wittmann-Liebold, B. 1998. Mitochondrial ribosomal proteins (MRPs) of yeast. *Biochem. J.* **329**: 433–448.
- Hamilton, M.G. and O'Brien, T.W. 1974. Ultracentrifugal characterization of the mitochondrial ribosome and subribosomal particles of bovine liver: Molecular size and composition. *Biochem. J.* **13**: 5400–5403.
- Heilek, G.M. and Noller, H.F. 1996. Site-directed hydroxyl radical probing of the rRNA neighborhood of ribosomal protein S5. *Science* **272**: 1659–1662.
- Hoshino, M., Sone, M., Fukata, M., Kuroda, S., Kaibuchi, K., Nabeshima, Y., and Hama, C. 1999. Identification of the stef gene that encodes a novel

- guanine nucleotide exchange factor specific for Rac1. *J. Biol. Chem.* **274**: 17837–17844.
- Jiang, X., Smith, J.B., and Abraham, E.C. 1996. Identification of a MS-MS fragment diagnostic for methionine sulfoxide. *J. Mass. Spect.* **31**: 1309–1310.
- Koc, E.C., Blackburn, K., Burkhart, W., and Spremulli, L.L. 1999. Identification of a mammalian mitochondrial homolog of ribosomal protein S7. *Biochem. Biophys. Res. Comm.* **266**: 141–146.
- Koc, E.C., Blackburn, K., Burkhart, W., Moseley, A., Koc, H., and Spremulli, L.L. 2000. A proteomics approach to the identification of mammalian mitochondrial small subunit ribosomal proteins. *J. Biol. Chem.* **275**: 32585–32591.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680–685.
- Lodmell, J. and Dahlberg, A. 1997. A conformational switch in *Escherichia coli* 16S ribosomal RNA during decoding of messenger RNA. *Science* **277**: 1262–1267.
- Mariottini, P., Shah, Z.H., Toivonen, J., Bagni, C., Spelbrink, J., Amaldi, F., and Jacobs, H. 1999. Expression of the gene for mitoribosomal protein S12 is controlled in human cells at the levels of transcription, RNA splicing, and translation. *J. Biol. Chem.* **274**: 31853–31862.
- Mathews, D.E., Hessler, R.A., Denslow, N.D., Edwards, J.S., and O'Brien, T.W. 1982. Protein composition of the bovine mitochondrial ribosome. *J. Biol. Chem.* **257**: 8788–8794.
- Mueller, F. and Brimacombe, M. 1997. A new model for the three-dimensional folding of *Escherichia coli* 16 S ribosomal RNA. II. The RNA-protein interaction data. *J. Mol. Biol.* **271**: 545–565.
- Nakai, K. and Kanehisa, M. 1992. A knowledge base for predicting protein localization sites in eukaryotic cells. *Genomics* **14**: 897–911.
- Noller, H., Moazed, D., Stern, S., Powers, T., Allen, P., Robertson, J., Weiser, B., and Triman, K. Structure of rRNA and its functional interactions in translation. In *The Ribosome: Structure, function and evolution* (eds. W. Hill, et al.), pp. 73–92. ASM Press, Washington D.C.
- O'Brien, T.W., Fiesler, S., Denslow, N.D., Thiede, B., Wittmann-Liebold, B., Mouge, E., Sylvester, J.E., and Graack, H-R. 1999. Mammalian mitochondrial ribosomal proteins: Amino acid sequences, characterization and identification of corresponding gene sequences. *J. Biol. Chem.* **274**: 36043–36051.
- O'Brien, T.W., Liu, J., Sylvester, J., Mourgey, E.B., Fischel-Ghodsian, N., Thiede, B., Wittmann-Liebold, B., and Graack, H-R. 2000. Mammalian mitochondrial ribosomal proteins (4): Amino acid sequencing, characterization and identification of corresponding gene sequences. *J. Biol. Chem.* **275**: 18153–18159.
- O'Brien, T.W., Mathews, D.E., and Denslow, N.D., 1976. In *Genetics and Biogenesis of Chloroplasts and Mitochondria* (eds. Bucher, T.L., et al.), pp 741–748. Elsevier, Amsterdam
- Pearson, W.R. and Lipman, D.J. 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**: 2444–2448.
- Perkins, D.N., Pappin, D.J., Creasy, D.M., and Cotrell, J.S. 1999. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* **20**: 3551–3567.
- Ramakrishnan, V. and White, S.W. 1992. The structure of ribosomal protein S5 reveals sites of interaction with 16 S rRNA. *Nature* **358**: 768–771.
- Roepstorff, P. and Fohlman, J. 1984. Proposal for a common nomenclature for sequence ions in mass spectra of peptides. *Biomedical Mass Spectrometry* **11**: 601.
- Samaha, R.R., O'Brien, B., O'Brien, T.W., and Noller, H.F. 1994. Independent in vitro assembly of a ribonucleoprotein particle containing the 3' domain of 16S rRNA. *Proc. Natl. Acad. Sci.* **91**: 7884–7888.
- Sayle, R.A. and Milner-White, E.J. 1995. RASMOL: Biomolecular graphics for all. *Trends Biochem. Sci.* **20**: 374–376.
- Shevchenko, A., Wilm, M., Vorm, O., and Mann, M. 1996. Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels. *Anal. Chem.* **68**: 850–858.
- Schlutzen, F., Tocilj, A., Zarivach, R., Harms, J., Gluehmann, M., Janell, D., Bashan, A., Bartels, H., Agmon, I., Franceschi, F., et al. 2000. Structure of functionally activated small ribosomal subunit at 3.3 angstroms resolution. *Cell* **102**: 615–623.
- Sone, M., Hoshino, M., Suzuki, E., Kuroda, S., Kaibuchi, K., Nakagoshi, H., Saigo, K., Nabeshima, Y., and Hama, C. 1997. Still life, a protein in synaptic terminals of *Drosophila* homologous to GDP-GTP exchangers. *Science* **275**: 543–547.
- Wittmann-Liebold, B. 1985. Ribosomal proteins. In *Structure, function and genetics of ribosomes*. (eds. B. Hardesty and G. Kramer) pp. 326–361. Springer-Verlag, New York.
- Wittmann-Liebold, B. and Littlechild, J. 1980. Structure of ribosomal proteins. In *Ribosomes: Structure, function and genetics*. (eds. G. Chambliss et al.), pp 51–88. University Park Press, Baltimore.
- Wittmann, H. 1985. Structure of ribosomes. In *Structure, function and genetics of ribosomes* (eds. B. Hardesty and G. Kramer), pp. 1–27. Springer-Verlag, New York.
- Yamaguchi, K. and Subramanian, A.R. 2000. The plastid ribosomal proteins (2): Identification of all the proteins in the 50S subunit of an organelle ribosome. *J. Biol. Chem.* **275**: 28466–28482.
- Yamaguchi, K., von Knoblauch, K., and Subramanian, A.R. 2000. The plastid ribosomal proteins (1): Identification of all the proteins in the 30S subunit of an organelle ribosome (chloroplast). *J. Biol. Chem.* **275**: 28455–28465.