

Published in final edited form as:

*Tetrahedron Lett.* 2008 March 10; 49(11): 1782–1785.

## Synthesis of deacetyl-1,10-didehydrosalvinorin G

Zhongze Ma and David Y. W. Lee\*

Bio-Organic and Natural Products Laboratory, McLean Hospital, Harvard Medical School, 115 Mill Street, Belmont, MA 02478

### Abstract

To unambiguously confirm the actual product in autoxidation of salvinorin A under basic conditions, deacetyl-1,10-didehydrosalvinorin G was synthesized from salvinorin C via intermediate salvinorin H. Furthermore, oxidation of salvinorin D with manganese dioxide gave salvinorin G in good yield.

### Keywords

Salvinorin A; Oxidation; Manganese dioxide; Synthesis

*Salvia divinorum*, a Mexican medicinal plant, has been used traditionally for its psychoactive (hallucinogenic) effects in divination rites.<sup>1</sup> Previous phytochemical studies have resulted in the isolation of 34 compounds, including salvinorins A (**1a**), C (**2a**), D (**2b**), G (**3**) and H (**2c**).<sup>2–9</sup> Of those compounds, **1a** was identified as a potent and selective kappa opioid receptor (KOR) agonist.<sup>10,11</sup> Because of the unique non-nitrogenous structure and potent binding activities to KOR, much effort has been directed toward a better understanding of structure-activity relationships (SAR) of **1a**.<sup>12–26</sup> Salvinorin derivatives readily underwent epimerization at C-8 under basic conditions.<sup>3,4,13–17</sup> Surprisingly, treatment of **1a** and its derivative with strong bases, such as Ba(OH)<sub>2</sub>,<sup>15</sup> KOH<sup>18,26</sup> and NaOH,<sup>25</sup> yielded corresponding natural salvinorin analogs, and no epimerization at C-8 was observed. It was reported that treatment of **1a** with KOH in methanol produced deacetyl-1,10-didehydrosalvinorin G (**4a**).<sup>18</sup> Recently, we revised the structure of **4a** to its 8-epimer (**4b**) based on comparison of <sup>1</sup>H and <sup>13</sup>C NMR data with those of **1a** and **1b**, NOESY data and chemical conversion.<sup>27</sup> In order to unambiguously confirm the actual product in autoxidation of **1a** under harsh basic conditions,<sup>18</sup> it is necessary to synthesize the natural salvinorin derivative **4a** and to complete its NMR data. In this Letter, we report the synthesis of **4a** from **2a** via intermediate **2c**, and chemical conversion of **3** from **2b**.

Following the published procedure,<sup>18</sup> the diol **2c** was prepared by deacetylation of **2a**. Subsequent oxidation of **2c** with manganese dioxide (Scheme 1) yielded **4a** and deacetylsalvinorin G (**5**).<sup>28</sup> Using 2D NMR techniques, including COSY, NOESY, HMQC and HMBC, permitted the full assignment of all <sup>1</sup>H and <sup>13</sup>C NMR chemical shifts of **4a** (Tables 1 and 2), and the key <sup>13</sup>C-<sup>1</sup>H correlations in the HMBC spectrum of **4a** were shown in Figure 1. In the <sup>1</sup>H NMR spectrum of **4a**, the C-8-H shifted much upper field to δ 2.42, and the coupling constants (dd, *J* = 12.6 and 3.3 Hz) of H-8 are more suitable for axial orientation than those (δ 2.99, dd, *J* = 9.6 and 5.1 Hz) of **4b** (Table 1). The C-12-H of **4a** shifted low-field slightly compared with that of **4b** (Table 1), and the H-12 of **4a** showed the *J* values (10.5 and 6.6 Hz)

Corresponding author. Tel: +1-617-855-2038; Fax: +1-617-855-2040; e-mail: dlee@mclean.harvard.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

for axial orientation. In addition, the H-11 $\alpha$  ( $\delta$  3.76) of **4a** shifted much lower field compared with that of **4b** ( $\delta$  3.11), and the chemical shift changes are consistent with those of **2a** ( $\delta$  2.49)<sup>4</sup> and its 8-epimer ( $\delta$  2.14).<sup>29</sup> Comparison of the <sup>13</sup>C resonances of C-6, C-8, C-12, C-13, C-17, C-19 and C-20 (Table 2) of **4a** and **4b** also confirms that H-8 in **4a** is the  $\beta$  configuration. In the NOESY spectrum of **4a**, H-12 ( $\delta$  5.62) showed cross peaks to H-11 $\alpha$  ( $\delta$  3.76) and H-20 ( $\delta$  1.54), while H-19 ( $\delta$  1.76) related to H-6 $\alpha$  ( $\delta$  2.51), H-7 $\alpha$  (1.94) and H-20 ( $\delta$  1.54). It should be noted that there is no crossed peak between H-8 and H-12/H-19/H-20. Based on these data, the structure of **4a** was confirmed as deacetyl-1,10-didehydrosalvinorin G.

The coupling constants of H-6 and H-7 (Table 1) in **4a** are significantly different from those of **4b**. The *J* values of H-6 $\beta$  (td, 13.5, 3.9) and H-7 $\alpha$  (dtd, 14.4, 3.3, 13.5) of **4a** indicate that both protons are axial. It has been reported that the protons in anti-periplanar relationships show stronger correlations in the COSY spectrum.<sup>30</sup> This was also evidenced by the protons (H-7 $\alpha$  and H-6 $\beta$ , H-7 $\alpha$  and H-8) of **4a** in the COSY spectrum. On the other hand, only H-7 $\beta$  and H-8 of **4b** exhibited stronger correlations in the COSY spectrum. These findings suggest that the B-ring in **4a** should be an identical chair conformation (Figure 1), which is different from that of **4b**. Obviously, the double bond between C-1 and C-10 in **4a** and **4b** distorts the B-ring conformation to a different extent.

Compounds **4a** and **5** were screened for binding affinity at opioid receptors in vitro, as reported previously.<sup>7</sup> Both compounds were inactive at mu, delta and kappa opioid receptors at 3  $\mu$ M.

Salvinorin G (**3**) presents in *S. divinorum* in much lower level than **1a** and **2a**, and it showed a moderate binding affinity at KOR.<sup>7</sup> **3** can be prepared by oxidation of the natural occurring salvinorin D (**2b**)<sup>7</sup> with manganese dioxide in an excellent yield (Scheme 2).<sup>31</sup> This reaction not only confirms the structure of **3** but also demonstrates that no epimerization occurs under the mild oxidation conditions as shown in Scheme 1.

Numerous salvinorin derivatives have been prepared in recent years for SAR study and improvement of KOR binding affinity.<sup>12–26</sup> Compounds **6a**, **6b**, **7a**, **7b**, **8a** and **8b** have served as key intermediates for C-2 and C-18 SAR studies. Among these compounds, only **6a** and **8a** were reported with full NMR assignments.<sup>30,13</sup> However, **6a** was measured in acetone-*d*<sub>6</sub> at higher temperature (40 °C).<sup>30</sup> Furthermore, **1b** is the only compound with full <sup>1</sup>H and <sup>13</sup>C NMR assignments in numerous 8-*epi*-salvinorin derivatives.<sup>13</sup> Therefore, we assigned all <sup>13</sup>C NMR chemical shifts of **6a**, **6b**, **7a**, **7b**, **8a** and **8b** in comparison with those of **1a** and **1b** (Table 2).<sup>27</sup> On the other hand, both aldehydes **9a** and **9b** were synthesized in our laboratory, and the incorrect <sup>1</sup>H and <sup>13</sup>C NMR data of **9a** were presented in our previous paper.<sup>21</sup> The <sup>13</sup>C NMR data of **9a** were revised and shown in Table 2.

In conclusion, deacetyl-1,10-didehydrosalvinorin G (**4a**) was readily synthesized from salvinorin H (**2c**). The product obtained by the treatment of **1a** with hydroxides in MeOH<sup>18</sup> has been unambiguously identified as 8-*epi*-deacetyl-1,10-didehydrosalvinorin G (**4b**). Finally, conversion of salvinorin G (**3**) from salvinorin D (**2b**) provides an authentic sample with intact stereochemistry at C-8 for further confirmation of **4a**.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements


The authors wish to thank Ms. Jane F. McNeil for EI-MS determination and Drs. Wei Xu and Lee-Yuan Liu-Chen for the binding assay at opioid receptors. This work was supported by a research grant (R01-DA019688) from NIDA to D.Y.L.

## References and notes


1. Siebert DJ. *J Ethnopharmacol* 1994;43:53–56. [PubMed: 7526076]
2. Ortega A, Blount JF, Manchand PS. *J Chem Soc, Perkin Trans 1* 1982:2505–2508.
3. Valdes LJ III, Butler WM, Hatfield GM, Paul AG, Koreeda M. *J Org Chem* 1984;49:4716–4720.
4. Valdes LJ III, Chang H-M, Visger DC, Koreeda M. *Org Lett* 2001;3:3935–3937. [PubMed: 11720573]
5. Munro TA, Rizzacasa MA. *J Nat Prod* 2003;66:703–705. [PubMed: 12762813]
6. Bigham AK, Munro TA, Rizzacasa MA, Robins-Browne RM. *J Nat Prod* 2003;66:1242–1244. [PubMed: 14510607]
7. Lee DYW, Ma Z, Liu-Chen LY, Wang Y, Chen Y, Carlezon WA, Cohen B. *Bioorg Med Chem* 2005;13:5635–5639. [PubMed: 16084728]
8. Harding WW, Tidgewell K, Schmidt M, Shah K, Dersch CM, Snyder J, Parrish D, Deschamps JR, Rothman RB, Prisinzano TE. *Org Lett* 2005;7:3017–3020. [PubMed: 15987194]
9. Shirota O, Nagamatsu K, Sekita S. *J Nat Prod* 2006;69:1782–1786. [PubMed: 17190459]
10. Roth BL, Baner K, Westkaemper R, Siebert DJ, Rice KC, Steinberg S, Ernsberger P, Rothman RB. *Proc Natl Acad Sci* 2002;99:11934–11939. [PubMed: 12192085]
11. Sheffler DJ, Roth BL. *Trends Pharmacol Sci* 2003;24:107–109. [PubMed: 12628350]
12. Chavkin C, Sub S, Jin W, Stewart J, Zjawiony JK, Siebert DJ, Toth BA, Hufeisen SJ, Roth BL. *J Pharmacol Exp Ther* 2004;308:1197–1203. [PubMed: 14718611]
13. Munro TA, Rizzacasa MA, Roth BL, Toth BA, Yan F. *J Med Chem* 2005;48:345–348. [PubMed: 15658846]
14. Beguin C, Richards MR, Wang Y, Chen Y, Liu-Chen LY, Ma Z, Lee DYW, Carlezon WA, Cohen BM. *Bioorg Med Chem Lett* 2005;15:2761–2765. [PubMed: 15869877]
15. Lee DYW, Karnati VVR, He M, Liu-Chen LY, Kondaveti L, Ma Z, Wang Y, Chen Y, Beguin C, Carlezon WA, Cohen B. *Bioorg Med Chem Lett* 2005;15:3744–3747. [PubMed: 15993589]
16. Lee DYW, He M, Kondaveti L, Liu-Chen LY, Ma Z, Wang Y, Chen Y, Li JG, Beguin C, Carlezon WA, Cohen B. *Bioorg Med Chem Lett* 2005;15:4169–4173. [PubMed: 16051487]
17. Harding WW, Tidgewell K, Byrd N, Cobb H, Dersch CM, Butelman ER, Rothman RB, Prisinzano TE. *J Med Chem* 2005;48:4765–4771. [PubMed: 16033256]
18. Munro TA, Goetchius GW, Roth BL, Vortherms TA, Rizzacasa MA. *J Org Chem* 2005;70:10057–10061. [PubMed: 16292839]
19. Harding WW, Schmidt M, Tidgewell K, Kannan P, Holden KG, Dersch CM, Rothman RB, Prisinzano TE. *Bioorg Med Chem Lett* 2006;16:3170–3174. [PubMed: 16621556]
20. Beguin C, Richards MR, Li JG, Wang Y, Xu W, Liu-Chen LY, Carlezon WA, Cohen BM. *Bioorg Med Chem Lett* 2006;16:4679–4685. [PubMed: 16777411]
21. Lee DYW, He M, Liu-Chen LY, Wang Y, Li JG, Xu W, Ma Z, Carlezon WA, Cohen B. *Bioorg Med Chem Lett* 2006;16:5498–5502. [PubMed: 16945525]
22. Tidgewell K, Harding WW, Lozama A, Cobb H, Shah K, Kannan P, Dersch CM, Parrish D, Deschamps JR, Rothman RB, Prisinzano TE. *J Nat Prod* 2006;69:914–918. [PubMed: 16792410]
23. Munro, TA.; Duncan, KK.; Staples, RJ.; Xu, W.; Liu-Chen, L-Y.; Beguin, C.; Carlezon, WA.; Cohen, BM. *Beilstein J Org Chem*. 2007. [http://bjoc.beilstein-journals.org/content/pdf/1860-5397-3-1.pdf]
24. Bikbulatov RV, Yan F, Roth BL, Zjawiony JK. *Bioorg Med Chem Lett* 2007;17:2229–2232. [PubMed: 17303418]
25. Holden KG, Kidgewell K, Marquam A, Rothman RB, Navarro H, Prisinzano TE. *Bioorg Med Chem Lett* 2007;17:6111–6115. [PubMed: 17904842]
26. Bikbulatov RV, Stewart J, Jin W, Yan F, Roth BL, Ferreira D, Zjawiony JK. *Tetrahedron Lett* 2008;49:10.1016/j.tetlet.2007.12.041 in press
27. Ma Z, Lee DYW. *Tetrahedron Lett* 2007;48:5461–5464.
28. Synthesis of 4a. To a solution of 2c (12 mg, 31  $\mu\text{mol}$ ) in  $\text{CH}_2\text{Cl}_2$  (3 ml) was added manganese dioxide (50 mg, 575  $\mu\text{mol}$ ), and the suspension was stirred at room temperature for 3 h. The solution was filtered and evaporated in vacuo. The residue was purified by silica gel column [ $\text{CH}_2\text{Cl}_2$ :AcOEt


(10:1)] to give 4a (6.2 mg, yield 52%) and 5 (2.5 mg, yield 21%). Data for 4a:  $^1\text{H}$  and  $^{13}\text{C}$  NMR data, see Tables 1 and 2; EI-MS  $m/z$  386 ( $\text{M}^+$ ). Data for 5:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ , 300 MHz)  $\delta$  7.46 (1H, br s, H-16), 7.43 (1H, br s, H-15), 6.41 (1H, br s, H-14), 6.38 (1H, s, H-3), 5.61 (1H, dd,  $J = 11.7$  and 5.4 Hz, H-12), 4.33 (1H, d,  $J = 3.3$  Hz, H-1), 3.84 (3H, s,  $\text{COOCH}_3$ ), 2.54 (1H, brs, OH), 2.52 (1H, dd,  $J = 12.6$  and 5.1 Hz, H-11a), 2.12–2.32 (3H, m, H-6a, H-7a, H-8), 1.70–1.94 (3H, m, H-7b, H-10, H-11b), 1.73 (3H, s, H-19), 1.53 (3H, s, H-20), 1.34 (1H, m, H-6b);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ , 75 MHz)  $\delta$  198.9 (C-2), 171.4 (C-17), 166.4 (C-18), 162.3 (C-4), 143.9 (C-15), 139.4 (C-16), 127.1 (C-3), 125.4 (C-13), 108.3 (C-14), 71.8 (C-12), 70.1 (C-1), 54.0 (C-10), 52.5 (C-8), 52.0 ( $\text{COOCH}_3$ ), 43.2 (C-11), 38.0 (C-5), 37.3 (C-9), 35.1 (C-6), 23.3 (C-19), 18.2 (C-7), 16.8 (C-20); EI-MS  $m/z$  388 ( $\text{M}^+$ ).

29. Munro, TA. Ph.D. Thesis. University of Melbourne; Victoria, Australia: 2006. The Chemistry of *Salvia divinorum*; p. 212[<http://eprints.infodiv.unimelb.edu.au/archive/00002327>]
30. Giner JL, Kiemle DJ, Kutrzeba L, Zjawiony J. Magn Reson Chem 2007;45:351–354. [PubMed: 17351976]
31. The quaternary carbon at C-17 and the carbonyl carbon of the acetyl group in 3 should be reassigned as  $\delta$  170.9 and  $\delta$  169.7, respectively.








**Figure 1.**  
Key HMBC correlations of **4a** and **4b**



**Scheme 1.**




Scheme 2.



**3**







**Table 1**<sup>1</sup>H NMR Data (300 MHz, CDCl<sub>3</sub>) for **4a** and **4b** [ $\delta$  (ppm), m, *J* (Hz)]

Proton	4a	4b
3	6.88 s	7.00 s
6 $\alpha$	2.51 dt (13.2, 3.3)	2.54 dt (13.8, 7.5)
6 $\beta$	1.46 td (13.5, 3.9)	1.67–1.77 m
7 $\alpha$	1.94 ddt (14.4, 3.3, 13.5)	2.24 ddd (14.4, 7.8, 5.1)
7 $\beta$	2.27 dq (14.7, 3.6)	1.98 ddd (14.1, 9.6, 6.6)
8	2.42 dd (12.6, 3.3)	2.99 dd (9.6, 5.1)
11 $\alpha$	3.76 dd (14.4, 6.3)	3.11 dd (14.7, 3.0)
11 $\beta$	2.09 dd (14.4, 10.5)	2.02 dd (14.4, 12.3)
12	5.62 dd (10.5, 6.6)	5.44 dd (12.3, 2.4)
14	6.42 dd (1.2, 0.9)	6.42 dd (1.8, 0.9)
15	7.41 t (1.8)	7.41 t (1.8)
16	7.44 d (0.9)	7.49 d (0.9)
19	1.76 s	1.72 s
20	1.54 s	1.67 s
OH	7.07 s	6.92 s
CO <sub>2</sub> CH <sub>3</sub>	3.86 s	3.85 s

**Table 2**  
 $^{13}\text{C}$  NMR Data (75 MHz,  $\text{CDCl}_3$ ) for **4a**, **4b**, **6a**, **6b**, **7a**, **7b**, **8a**, **8b**, **9a** and **9b** [ $\delta$  (ppm)]

Carbon	4a	4b	6a <sup>a</sup>	6b <sup>b</sup>	7a <sup>b</sup>	7b <sup>b</sup>	8a <sup>b</sup>	8b <sup>b</sup>	9a <sup>c</sup>	9b <sup>c</sup>
1	145.3	145.1	208.9	209.2	201.9	202.3	203.7	204.1	202.2	202.6
2	180.9	180.7	74.4	74.4	74.9	75.1	75.9	76.2	75.2	75.5
3	127.6	128.2	34.5	33.8	30.6	30.5	31.8	31.5	28.3	28.1
4	159.0	157.5	53.1	52.3	53.4	52.6	50.7	50.2	59.7	59.1
5	43.3	42.3	42.6	42.6	42.0	42.1	41.9	42.0	42.5	42.6
6	32.0	28.3	38.1	34.2	38.1	33.8	38.0	33.8	38.0	33.8
7	18.1	21.9	18.1	17.5	18.1	17.5	18.1	17.6	17.9	17.4
8	50.6	44.8	51.3	45.2	51.3	45.2	51.4	45.3	51.2	45.1
9	39.4	37.6	35.3	34.5	35.4	34.7	35.2	34.6	35.4	34.6
10	138.5	140.0	63.8	63.6	63.9	63.9	64.4	64.5	63.6	63.5
11	42.4	36.8	43.5	48.2	43.3	47.9	43.3	48.0	43.2	47.9
12	71.5	70.8	71.9	70.0	72.1	72.1	72.1	70.1	72.0	70.1
13	126.0	124.4	125.3	123.5	125.1	123.3	125.2	123.3	125.1	123.2
14	108.6	108.4	108.3	108.4	108.4	108.5	108.4	108.5	108.3	108.5
15	143.8	143.7	143.8	143.6	143.7	143.6	143.7	143.6	143.8	143.6
16	139.4	139.6	139.3	139.6	139.4	139.7	139.4	139.7	139.4	139.7
17	171.6	173.2	171.0	173.4	171.3	173.7	171.4	173.7	170.9	173.5
18	166.0	165.4	171.8	172.1	175.8	176.2	171.8	176.2	200.6	200.9
19	31.3	30.3	16.5	15.3	16.4	15.3	16.6	15.5	17.9	16.7
20	18.8	24.4	15.2	24.6	15.2	24.6	15.3	24.8	15.1	24.4
$\text{CO}_2\text{CH}_3$	52.7	52.6	51.9	51.6	-	-	-	-	-	-
$-\text{COCH}_3$	-	-	-	-	170.0	169.9	170.1	169.9	170.0	169.8
$-\text{COCH}_3$	-	-	-	-	20.5	20.5	20.6	20.6	20.6	20.6

<sup>a</sup> **6a** was isolated from the leaves of *Salvia divinorum*.<sup>7</sup>

<sup>b</sup> **6b**, **7a**, **7b**, **8a** and **8b** were synthesized and reported by our group.<sup>15,16,21</sup>

<sup>c</sup> The chemical shift assignments were based on the comparison with those in **1a** and **1b**.<sup>27</sup>