

Proceedings

Open Access

Signaling role of iron in NF-kappa B activation in hepatic macrophages

Shigang Xiong*¹, Hongyun She¹ and Hidekazu Tsukamoto^{1,2}

Address: ¹Department of Pathology, Keck School of Medicine, University of Southern California, USA and ²Greater Los Angeles VA Health Care Systems, Los Angeles, California, USA

Email: Shigang Xiong* - shigangx@usc.edu; Hongyun She - hshe@usc.edu; Hidekazu Tsukamoto - htsukamo@usc.edu

* Corresponding author

from 11th International Symposium on the Cells of the Hepatic Sinusoid and their Relation to Other Cells
Tucson, Arizona, USA, 25–29 August, 2002

Published: 14 January 2004

Comparative Hepatology 2004, **3**(Suppl 1):S36

This article is available from: <http://www.comparative-hepatology.com/content/3/S1/S36>

Abstract

Iron is both essential and toxic for cells and impaired iron homeostasis has been shown to cause or potentiate various forms of liver injury. Research in our laboratory suggests that iron also plays a pivotal role in intracellular signaling for NF-kappa B activation in hepatic macrophages (HM). Our results showed: 1) HM from alcohol-fed rats had a increase in the nonheme iron content accompanied by NF-kappa B activation; 2) iron chelation normalized nonheme iron concentration and blocked enhanced NF-kappa B activation and TNF-alpha expression in these cells; 3) LPS-induced NF-kappa B activation was also blocked by iron chelator; 4) iron directly induced TNF-alpha expression via IKK and NF-kappa B activation in normal HM. We propose that iron acts as an independent proinflammatory molecule via induction of the intracellular signaling for NF-kappa B activation in HM and primes the liver for chronic inflammation and injury.

Iron and HM NF-kappa B activation in alcohol model

Our earlier study showed that hepatic macrophages (HM) from rats fed ethanol and high fat diet had a significant 70% increase in the nonheme iron content as compared to controls [1]. This study also suggested enhanced heme turnover as a cause of the increased iron storage in HM. To test this notion, an increase in HM iron content was recapitulated in vitro by phagocytosis of heat-treated autologous red blood cells. To extend this observation to the whole animal situation, the effects of splenectomy on alcohol-fed animals were also examined. The most intriguing and critical finding from these cellular or animal model experimentations, was that activation of NF-kappa B was tightly correlated with the increased nonheme iron content in HM, suggesting the priming role of iron in NF-kappa B activation and proinflammatory cytokine expression by HM in alcoholic liver disease.

Direct iron induction of TNF-alpha in cultured HM

Direct addition of ferrous but not ferric iron in cultured HM increased TNF-alpha release 8 fold at 10 and 50 micromolar during a 4 hr treatment period without cell toxicity. Cuprous (Cu¹⁺) but not cupric (Cu²⁺) copper also stimulated TNF-alpha release at 50 micromolar to less extent. Thus, these results demonstrate direct stimulation of HM TNF-alpha release by iron and copper in a redox status dependent manner. We then tested whether Fe²⁺ stimulates TNF-alpha promoter in cultured HM. The promoter activity was indeed increased 2~3 fold with 10~50 micromolar Fe²⁺. Cu¹⁺ (50 micromolar) also slightly increased TNF-alpha promoter activity but not Cu²⁺ or Fe³⁺ (Figure 1). Co-transfection of a super repressor I-kappa B-alpha vector completely abrogated the stimulation with 50 micromolar Fe²⁺ (Figure 1). The enhanced promoter activity with 50 micromolar Fe²⁺ was about half




Figure 1

(A.) Cultured HM were transfected with a TNF- α promoter-luciferase construct followed by the treatment with Fe²⁺, Fe³⁺, Cu¹⁺ or Cu²⁺. The data were normalized by co-transfected Renilla luciferase activity. Note the Fe²⁺ induces the promoter activity by 2–3 fold at 10 and 50 micromolar. Cu¹⁺ slightly induces but oxidized metals (Fe³⁺ and Cu²⁺) do not. (B.) HM were co-transfected with the promoter-luciferase construct with a vector of super-repressor I- κ B- α (DN-I- κ B), followed by addition of Fe²⁺. Note DN-I- κ B completely blocks iron induced promoter activity.




Figure 2

I-kappa B kinase (IKK) and c-Jun NH₂-terminal kinase (JNK) activity were determined on HM lysate collected at different time points after FeSO₄ treatment. Note that IKK activity as assessed by phosphorylation of GST-I-kappa B-alpha (P-I-kappa B-alpha), is increased at 15 min after addition of iron while no activation of JNK is evident.

of the maximal response achieved with LPS (500 ng/ml) in a serum-free condition. These results establish that Fe²⁺ activates TNF-alpha promoter in a NF-kappa B dependent manner.

Iron-induced activation of IKK and NF-kappa B and ROS generation in cultured HM

As shown in the top panel of Figure 2, IKK activity, as assessed by phosphorylation of GST-I-kappa B-alpha, was increased at 15 min after addition of ferrous iron. The timing of IKK activation preceded a disappearance of cytosolic I-kappa B-alpha and an increase in the nuclear level of p65 at 30~45 min. Iron did not induce JNK and p38 activities (unpublished results). These results correlated well with induced NF-kappa B binding in iron-treated HM at 30 min and no effects on AP-1 binding in these cells (data not shown). Thus, these results unequiv-

ocally demonstrate that ferrous iron can directly and selectively stimulates the signaling leading to IKK and NF-kappa B activation in cultured HM. Addition of ferrous iron to these cells resulted in an enhancement of the hydroxyl and methyl-POBN adduct signals. Both signals increased with incubation time up to a maximum at 15–20 min, which coincided with IKK activation at 15~30 min and preceded activation of NF-kappa B at 30 min, suggesting the possible signaling role of the former in the latter events. In fact, this notion was made by previous studies which demonstrated activation of NF-kappa B by hydroxyl radical generating systems and a reversal of this effect by hydroxyl radical scavengers or metal chelators in Jurkat cells [2].

Discussion and Conclusion

Our studies to date strongly suggest the causal link between iron and activation of NF-kappa B in HM in both normal and alcohol-fed rats. These findings raise a question as to how iron signals to activate NF-kappa B. Since NF-kappa B is a redox sensitive transcription factor and ROS are implicated in its activation [3,4], it is reasonable to speculate that iron stimulates ROS production in HM and in turn ROS activates NF-kappa B. Conversely, stimulation of HM with an agonist such as LPS, induces ROS generation and ROS may initiate intracellular signaling that is dependent on a chelatable pool of iron. Nitric oxide (NO) is known to cause mobilization of intracellular iron [5] and to inhibit enzymes with catalytically active iron-sulfur groups [6]. Superoxide anion can also release iron from ferritin [7]. Indeed, our preliminary results demonstrate that a selective inhibitor of iNOS and Cu/Zn SOD overexpression abolish a LPS-mediated transient rise in the intracellular level of chelatable iron and NF-kappa B activation (unpublished observation). We propose that iron acts as a proinflammatory effector molecule via selective induction of the intracellular signaling for NF-kappa B activation and that dysregulation of this signaling mechanism may prime HM for chronic liver inflammation and injury.

References

1. Tsukamoto H, Lin M, Ohata M, Giulivi C, French SW, Brittenham G: **Iron primes hepatic macrophages for NF-kappa B activation in alcoholic liver injury.** *Am J Physiol* 1999, **277**:G1240-G1250.
2. Shi X, Dong Z, Huang C, Ma W, Liu K, Ye J, Chen F, Leonard SS, Ding M, Castranova V, Vallyathan V: **The role of hydroxyl radical as a messenger in the activation of nuclear transcription factor NF-kappa B.** *Mol Cell Biochem* 1999, **194**:63-70.
3. Anderson MT, Staal FJ, Gitler C, Herzenberg LA, Herzenberg LA: **Separation of oxidant-initiated and redox-regulated steps in the NF-kappa B signal transduction pathway.** *Proc Natl Acad Sci* 1994, **91**:11527-11531.
4. Schreck R, Baeuerle PA: **Assessing oxygen radicals as mediators in activation of inducible eukaryotic transcription factor NF-kappa B.** *Methods Enzymol* 1994, **234**:151-163.
5. Drapier JC, Hirling H, Wietzerbin J, Kaldy P, Kuhn LC: **Biosynthesis of nitric oxide activates iron regulatory factor in macrophages.** *EMBO J* 1993, **12**:3643-3649.

6. Drapier JC, Hibbs JB Jr: **Differentiation of murine macrophages to express nonspecific cytotoxicity for tumor cells results in L-arginine-dependent inhibition of mitochondrial iron-sulfur enzymes in the macrophage effector cells.** *J Immunol* 1988, **140**:2829-2838.
7. Cairo G, Tacchini L, Pogliaghi G, Anzon E, Tomasi A, Bernelli-Zazzera A: **Induction of ferritin synthesis by oxidative stress. Transcriptional and post-transcriptional regulation by expansion of the "free" iron pool.** *J Biol Chem* 1995, **270**:700-703.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

