

A computational screen for mouse signaling pathways targeted by microRNA clusters

JIANZHEN XU and CHIWAI WONG

Center of Integrative Biology, Guangzhou Institute of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China 510663

ABSTRACT

MicroRNAs (miRNAs) are one class of short, endogenous RNAs which can regulate gene expression at the post-transcriptional level. Previous analysis revealed that mammalian miRNAs tend to cluster on chromosomes. However, the functional consequences of this clustering and conservation property are largely unknown. In this study we present a method to identify signaling pathways targeted by clustered miRNAs. We performed a computational screen for mouse signaling pathways targeted by miRNA clusters. Here, we report that the target genes of 3 miRNA clusters are overrepresented in 15 signaling pathways. We provided experimental evidence that one miRNA cluster, *mmu-mir-183-96-182* targets *Irs1*, *Rasa1*, and *Grb2*, all of which are located in the insulin signaling pathway. These results suggest that by targeting components with different roles along a signaling pathway, different members of one miRNA cluster can act as a whole to coordinately control the signal transduction process.

Keywords: clustered microRNAs; signaling pathways; insulin pathway

INTRODUCTION

MicroRNAs (miRNAs) are a class of ~22-nucleotide (nt) endogenous single-stranded RNA which can cleave mRNA or repress the initiation of protein translation by imperfect base pairing with the targeting sequences on 3'-UTR in animals (Bartel 2004). miRNAs are transcribed by RNA polymerase II or III as long primary miRNAs (pri-miRNAs) which range from hundreds to thousands of nucleotides in length. The pri-miRNAs are then processed into mature miRNAs by Drosha and Dicer enzymes sequentially. Studies have shown that miRNAs play crucial roles in a broad range of biological processes including development, differentiation, tissue morphogenesis, and also in several types of diseases such as carcinogenesis and viral infections (Bartel 2004; Kloosterman and Plasterk 2006).

Previous analysis has revealed conserved patterns as clustering of mammalian miRNAs on chromosomes (Altuvia et al. 2005). Megraw et al. (2007) reported that in human, mouse, rat, and chicken, even at a very conservative maximum inter-miRNA distance (MID) of 1 kb, over 30% of all miRNAs are organized into clusters. If the

MID extends to 50 kb, over half of all miRNAs are clustered on chromosomes. Expression profiling analysis showed that proximal pairs of miRNAs are generally coexpressed, and an abrupt transition in the correlation between pairs of expressed miRNAs occurs at a distance of 50 kb (Baskerville and Bartel 2005). According to their chromosomal positions, several miRNA clusters have also been experimentally confirmed by RT-PCR or Northern blotting (Cai et al. 2004; Lee et al. 2004; Weston et al. 2006). These evidences strongly suggested that members of a particular miRNA cluster, which are defined by their chromosomal positions, are highly likely to be processed as cotranscribed units. However, the functional consequences of this clustering and conserved property remain unclear.

Analogous to transcriptional regulation (between transcriptional factors and *cis*-regulatory elements), most miRNAs function to fine tune the expression of hundreds of genes in a combinatorial manner (Bartel 2004). Specifically, different miRNA species have been proposed to regulate a single mRNA target through binding to the same mRNA transcript. Several such cases have also been experimentally verified. For example, Krek et al. (2005) experimentally validated that myotrophin (*Mtpn*) as the first mammalian gene regulated by *miR-375*, *miR-124*, and *let-7b* coordinately. Other studies also showed that miRNAs *17-5p*, *20a*, and *106a* control monocytopenia via binding to the 3'-UTR of transcription factor acute myeloid leukaemia-1 (*AML1*) mRNA (Fontana et al. 2007).

Reprint requests to: Chiwai Wong, Center of Integrative Biology, Guangzhou Institute of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, China 510663; e-mail: wong_chiwai@gibh.ac.cn; fax: (80) 20-32290606.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.997708>.

Dews et al. (2006) recently reported the first evidence of a cluster of miRNAs that could regulate functionally related proteins in cancer. They found that in *Myc* overexpressing tumors, two anti-angiogenesis genes *Tsp1* and *Ctgf* are downregulated by one *Myc* activated miRNA cluster *miR-17-miR-92*. Noticeably, they confirmed that within the cluster, *miR-19* is primarily responsible for *Tsp1* downregulation and *miR-18* for *Ctgf* downregulation in response to *Myc* activation. Inspired by their findings, in this paper, we combined the combinatorial regulation concept with the clustering property to screen mouse signaling pathways which are targeted by clustered miRNAs. Especially, we explored another layer of combinatorial regulation, i.e., we hypothesize that each member of a miRNA cluster could regulate one component within the same signaling pathway; thus, the cluster acts coordinately as a whole to control a biological process.

RESULTS

A computational screen framework for mouse signaling pathways targeted by miRNA clusters

As indicated in Figure 1, to identify mouse signaling pathways targeted by miRNA clusters, we initially extracted mouse miRNA clusters according to mouse chromosomal positions. Although previous studies had indicated that defining clustered miRNA memberships according to chromosomal location is highly consistent with experimental evidences as RT-PCR or Northern blotting, we further filtered miRNA clusters through (1) an expression profiling filter and (2) a regulatory element filter.




FIGURE 1. Flowchart of the computational screen framework for mouse signaling pathways targeted by the miRNA cluster. The algorithm is divided into two stages. The initial stage is to preprocess both the mouse miRNA clusters and the Biocarta signaling pathways. The second stage involves selecting the significant pathways that are enriched with target genes of one miRNA cluster.

For any particular miRNA cluster that shares the same transcriptional unit, the expression profiles of miRNAs within this cluster should be similar, and thus positively, correlated. We filtered out those miRNA clusters with the mean correlation measures that are less than those of the background correlation in at least two of the three analyzed expression profiles. In addition, we examined for the presence of regulatory elements such as transcription start site (TSS), CpG window, or predicted first exon in the inter-miRNA region of the same cluster. All of these possibly indicated a different transcription initiation. We extensively searched for the elements in the inter-miRNA region of the same cluster, and we filtered out the miRNA clusters in which regulatory elements are found. After filtering signaling pathways, we selected the significant signaling pathways that are enriched with those predicted target genes according to the optimized target prediction data.

Filtering miRNA clusters

Forty-five miRNA clusters which contained 169 miRNAs, were extracted when MID was set to 5 kb. The largest miRNA cluster *mmu-mir-669a-3* is located on chromosome 2 with 25 miRNA members (Supplemental Table S1). Among these 45 miRNA clusters, 40 clusters have a correlation value higher than 0.35, the background correlation value of data set published by Thomson et al. (2004). Nineteen out of the 45 clusters have correlation measures significantly higher than the background (nominal $P < 0.05$) (Supplemental Table S1). Previously, the miRNA members of the *mir-17* cluster show consistent upregulation patterns in a *Myc*-overexpressing *Ras*-transformed colonocytes (Dews et al. 2006). However, the expressions of these miRNA members are not correlated in two out of three analyzed profiles (Supplemental Table S1), perhaps suggesting that this cluster is only regulated conditionally. In addition, three other miRNA clusters *mmu-mir-212*, *mmu-mir-363*, and *mmu-mir-382* also do not coexpress in various cellular conditions, indicating that the members within these clusters have independent regulation mechanisms (Supplemental Table S1). Nonetheless, most miRNA clusters defined by chromosomal locations have coexpressed patterns under different cellular conditions, consistent with previous observations (Baskerville and Bartel 2005).

We also filtered the clusters according to regulatory element presence using both FirstEF and Eponine, and found regulatory elements within *mmu-mir-212-132* cluster (Supplemental Table S1). Interestingly, the expression profiling filter also shows that the expressions of these two miRNAs are not correlated in all three analyzed expression profiles. Collectively, these evidences strongly suggested that these two miRNAs have independent transcription mechanisms, and they may not function coordinately to regulate signaling pathways.

After these preprocessing phases, we finally obtained 41 miRNA clusters, which show coregulation patterns. Similar analysis were also conducted when MID was set to 50 kb. In all, 55 miRNA clusters were initially selected, and after the preprocessing phases, 45 clusters were used in the following analysis (Supplemental Table S2).

Selection of significant pathways that are enriched with target genes of one miRNA cluster

Next, we selected significant pathways that are enriched with target genes of one miRNA cluster among the filtered 97 signaling pathways. As indicated in Table 1, totally 18 cluster–pathway interactions were found by false discovery rate analysis (false discovery rate [FDR] <0.3; nominal *P* < 0.03), which involved three miRNA clusters and 15 signaling pathways.

Interestingly, one miRNA cluster *mmu-mir-183-96-182* controls 12 signaling pathways. Detailed analysis of the target genes by this cluster in these 12 significant pathways suggested that *miR-96* is the major regulator. Other than a few exceptions, *miR-96* targets most of the member genes in these 12 signaling pathways (Table 2). Previously, the pri-miRNA polycistronic organization of this cluster has been identified by experimental methods and the co-expression patterns of this cluster were also confirmed (Weston et al. 2006; Xu et al. 2007).

Consistent with our results, this cluster seems to be involved in at least three biological functions according to the current literature. First, it was reported that one member of this cluster, *mir-96*, targets protein arginine methyltransferase (*PRMT5*). Aberrant expression of *PRMT5*

leads to altered epigenetic modification of chromatin, and consequently impacts on the growth of transformed lymphoid cells (Pal et al. 2007). Second, in colorectal cancer cell lines, all three members of this cluster show consistent upregulations. Forkhead family of transcription factors including *CHES 1*, *FOXF2*, *FOXK2*, *FOXO1A*, *FOXO3A*, and *FOXQ1*, were proposed as putative targets (Bandres et al. 2006). Finally, Xu et al. (2007) reported that *Mitf*, a transcription factor required for the establishment and maintenance of retinal pigmented epithelium, is a direct target of *miR-96* and *miR-182*, indicating that this miRNA cluster is also important for retinal development and function. We experimentally proved that this cluster is also involved in insulin signaling pathway (see below). Therefore, these evidences suggested that this cluster plays an important role in multiple signaling pathways.

We also generated a set of experimentally testable hypotheses. For example, the epigenetic regulation of chromatin domains by miRNAs has recently been demonstrated through the finding that *miR-26a* targets the histone methyltransferase enhancer of zeste homolog 2 during myogenesis (Wong and Tellam 2008). Our prediction found that both *mmu-mir-206* and *mmu-mir-133a-1* clusters target chromatin remodeling hSWI/SNF ATP-dependent complexes, which is a major group of chromatin-modifying complexes. It would be interesting to test whether these miRNA clusters can regulate the chromatin structure via a mechanism involving hSWI/SNF ATP-dependent chromatin remodeling complexes. The same set of cluster–pathway interactions were obtained when the MID set to 50 kb (data not shown), indicating robustness of our prediction results.

TABLE 1. Significant pathways that are enriched with target genes of miRNA clusters (MID <5 kb)

miRNA cluster	Signaling pathway	Target gene
<i>mmu-mir-206</i>	Chromatin remodeling by hSWI/SNF ATP-dependent complexes	<i>Smarca4, Smarcb1, Tbp, Smarcd1</i>
<i>mmu-mir-206</i>	FOSB gene expression and drug abuse	<i>Fosb Junb</i>
<i>mmu-mir-206</i>	Sumoylation by RanBP2 regulates transcriptional repression	<i>Ranbp2, Sumo1, Hdac4</i>
<i>mmu-mir-183</i>	Integrin signaling pathway	<i>Cav1, Crkl, Fyn, Grb2, Itgb1, Shc1</i>
<i>mmu-mir-183</i>	Bioactive peptide-induced signaling pathway	<i>Fyn, Gnb1, Grb2, Prkca, Plcg1, Shc1</i>
<i>mmu-mir-183</i>	Activation of cAMP-dependent protein kinase, PKA	<i>Gnb1, Prkar1a</i>
<i>mmu-mir-183</i>	G-Protein signaling through tubby proteins	<i>Gnb1, Htr2c, Tub</i>
<i>mmu-mir-183</i>	PTEN-dependent cell cycle arrest and apoptosis	<i>Grb2, Itgb1, Shc1, Foxo3a</i>
<i>mmu-mir-183</i>	Ion channel and phorbol esters signaling pathway	<i>Prkca, Plcg1</i>
<i>mmu-mir-183</i>	Eph Kinases and ephrins support platelet aggregation	<i>Epha4, Fyn, Itgb1, L1cam</i>
<i>mmu-mir-183</i>	Insulin signaling pathway	<i>Grb2, Irs1, Shc1, Rasa1</i>
<i>mmu-mir-183</i>	IGF-1 signaling pathway	<i>Grb2, Irs1, Shc1, Rasa1</i>
<i>mmu-mir-183</i>	Role of Erk5 in neuronal survival	<i>Creb1, Grb2, Plcg1, Shc1</i>
<i>mmu-mir-183</i>	Links between Pyk2 and map kinases	<i>Crkl, Grb2, Pak1, Plcg1, Rac1, Shc1, Map2k3</i>
<i>mmu-mir-183</i>	Rac1 cell motility signaling pathway	<i>Cfl1, Pak1, Pdgfra, Pld1, Rac1</i>
<i>mmu-mir-133a-1</i>	Chromatin remodeling by hSWI/SNF ATP-dependent complexes	<i>Smarca4, Smarcb1, Tbp</i>
<i>mmu-mir-133a-1</i>	FOSB gene expression and drug abuse	<i>Fosb Junb</i>
<i>mmu-mir-133a-1</i>	Sumoylation by RanBP2 regulates transcriptional repression	<i>Ranbp2, Sumo1, Hdac4</i>

A FDR <0.3(nominal *P* < 0.03) was considered significant.

TABLE 2. Target genes by *mmu-mir-183-96-182* in the 12 significant pathways

Target gene	<i>miR-96</i>	<i>miR-182</i>	<i>miR-183</i>
<i>Cav1</i>	X		
<i>Cfl1</i>	X	X	
<i>Creb1</i>	X	X	
<i>Crkl</i>	X		
<i>Epha4</i>			X
<i>Foxo3a</i>	X	X	
<i>Fyn</i>	X		
<i>Gnb1</i>			X
<i>Grb2</i>	X	X	
<i>Htr2c</i>	X	X	
<i>Irs1</i>	X		X
<i>Itgb1</i>			X
<i>L1cam</i>	X	X	
<i>Map2k3</i>	X	X	
<i>Pak1</i>	X		
<i>Pdgfra</i>	X	X	
<i>Plcg1</i>	X	X	
<i>Pld1</i>	X	X	
<i>Prkar1a</i>	X	X	
<i>Prkca</i>			X
<i>Rac1</i>	X	X	
<i>Rasa1</i>	X	X	
<i>Shc1</i>	X		
<i>Tub</i>			X

Target genes set include all genes in the 12 selected significant pathways targeted by at least one member of the *mir-183-96-182* cluster. "X" means a hit by the intersection of PicTar and TargetScanS predictions.

Experimental verification of one miRNA cluster targeting the insulin signaling pathway

Insulin signaling pathway is involved in both metabolic and mitogenic aspects of cellular function. In our prediction, one miRNA cluster *mmu-mir-183-96-182* targets four members of the insulin signaling pathway. In the intersection of PicTar and TargetScanS prediction, both *mir-96* and *mir-183* target insulin receptor substrate 1 (*irs1*), whereas *mir-96* and *mir-182* target both ras p21 protein activator 1 (*rasa1*) and growth factor receptor bound protein 2 (*grb2*).

We set up luciferase reporter assays to validate these predicted targets. Fragments of the 3'-UTR of *irs1*, *rasa1*, and *grb2*, including the miRNA target sites, were individually cloned into a firefly luciferase expression vector pGL3. Cells were respectively transfected with these reporters together with a renilla luciferase vector pRL-TK as an internal control. While using mock transfected cells as a control (no oligos), we examined the effects of over-expressing different miRNA oligos on the expressions of these reporters. First, we found that a nonspecific oligo (NC), even at a relatively high concentration, did not alter the expression of all three reporters (Fig. 2A–C). Second, a *mir-153* oligo which is predicted to target *irs2*, did not significantly alter the gene expression of a reporter containing




FIGURE 2. Repression of *irs1*, *rasa1*, and *grb2* 3'-UTR by *mmu-mir-183-96-182* cluster. 3'-UTR of target genes were inserted into the 3' end of the luciferase gene in the pGL3 control vector. Cells were transfected with or without 10 nM miRNA oligos indicated, pGL3 vectors containing the 3'-UTR of target genes, and an internal control Renilla luciferase vector. The luciferase activity levels were determined after 24 h. Data were generated from triplicate in at least three independent experiments. The ratios of Firefly luciferase expression to Renilla luciferase expression were normalized to no oligos transfection. ** $P < 0.01$ for no oligos versus various oligos treatments, NS: no significant difference. (A) pGL-3-Irs1 plasmid cotransfected with *miR-153*, *miR-183*, *miR-96*, *miR-182*, and nonspecific oligos (NC). NC, *miR-153*, and *miR-182* were employed as negative controls as predicted by both PicTar and TargetScanS algorithms. (B) pGL-3-Rasa1 plasmid cotransfected with *miR-183*, *miR-96*, *miR-182*, and nonspecific oligos (NC). Both NC and *miR-183* were employed as negative controls as predicted by both PicTar and TargetScanS algorithms. (C) pGL-3-Grb2 plasmid cotransfected with *miR-183*, *miR-96*, *miR-182*, and nonspecific oligos (NC). Both NC and *miR-183* were employed as negative controls as predicted by both PicTar and TargetScanS algorithms. A dotted rectangle indicates the miRNA cluster *mmu-mir-183-96-182*.

the 3'-UTR of *irs1* (Fig. 2A). Third, we experimentally confirmed that both *mir-96* and *mir-183* repressed the gene expression of this reporter about 40% via targeting the 3'-UTR of *irs1* (Fig. 2A). The 3'-UTR of *irs1* does not contain a targeting sequence for another member of this cluster *mir-182*. We did not observe any significant effects of *mir-182* oligo on the expression of this reporter, demonstrating the specificity and validity of our reporter system (Fig. 2A).

We also observed strong repression of a luciferase reporter through presumed binding of *mir-182* to the 3'-UTR of *rasa1* (Fig. 2B). Moderate but significant repression of a luciferase reporter was also observed in pGL-3-*grb2* plasmid cotransfected with *mir-96* or *mir-182* oligos (Fig. 2C). Although *rasa1* was also a predicted target of *mir-96* by optimized prediction method, there was no significant difference between cotransfections with or without *mir-96* oligos, which possibly indicated that this was a false positive prediction.

DISCUSSION

One principle of gene regulation by miRNA is coordinate gene expression. In this model, individual miRNA may reduce the expression of target genes moderately while a collection of miRNAs may act in a combinatorial fashion to exert significant effects (Bartel 2004; Stark et al. 2005). Bioinformatics analysis agrees with this principle as well, since it is common to find multiple and independent binding sites on most, if not all, target genes. Different groups had also reported in experimental instances that miRNAs can act cooperatively through multiple target sites on the same gene. Recent studies have widened the scope of this combinatorial regulation such that a few miRNAs can regulate a group of functionally related genes instead of a single gene (Hua et al. 2006). These are convenient and efficient strategies to regulate cellular processes, but the key issue is that these regulatory miRNAs should themselves be regulated coordinately. Therefore, certain mechanisms must exist to guarantee the coordinate expression of these miRNAs.

Combining the coregulation concept with the observation that mammalian miRNAs tend to cluster on chromosomes, we proposed that the clustering and conserved patterns provide a natural mechanism to explain the principle of coregulation by miRNAs. Since those clustered miRNAs are largely regulated under the same control, upon stimulation, members within the same cluster could express and function in a cooperative manner. Thus, the downstream effect, for example, to completely shut down a biological process, is ensured. However, the above regulation model does not rule out the possibility that a single miRNA could greatly alter a cellular process. For instance, a miRNA may target an essential regulator of a signaling pathway which leads to alterations in multiple downstream effectors.

From the discovery of miRNAs, plenty of evidences have demonstrated the increasingly important role of these tiny molecules in developmental processes and carcinogenesis. However, relatively few are known for their regulation on cellular metabolic processes. Here, we provided the first evidence of an interplay between one miRNAs cluster *mmu-mir-96-182-183* and the insulin signaling pathway.

The IRS proteins are major players for the metabolic aspect of the insulin signaling cascade because they mediate the activations of intracellular signaling molecules, such as PI 3-kinase (*PI3k*), *Akt*, and molecular target of rapamycin (*mTOR*). On the other hand, the proliferative and mitogenic effects of insulin are mediated largely via the activation of *Ras* signaling pathway (Draznin 2006). *Grb2* is a classical adapter molecule that controls *Ras* and *MAPK* activations, which are important for growth factor signaling. Therefore, *Grb2* is a potential target of antitumor therapy (Dharmawardana et al. 2006). *Rasa1*, which transforms active GTP-bound *Ras* into inactive GDP-bound form, links the insulin signaling with *Ras* signaling pathway. Loss of *Rasa1* could lead to increased activation of *Ras*, which has been linked to various tumors (Downward 2003).

Importantly, we showed that three members of *mmu-mir-96-182-183* have distinct but coordinate roles in controlling insulin signaling. *Mir-96* and *mir-183* are mainly responsible for regulating *Irs1*, which is the key mediator of metabolic aspect of insulin pathway, whereas *mir-182* targets *Rasa1* and *Grb2*, both of which play important roles in linking insulin signaling with the *Ras* pathway. Thus we proved that by targeting components with different roles along the insulin signaling pathway, *mmu-mir-183-96-182* can coordinately control the insulin signal transduction. This regulation model is illustrated in Figure 3. Due to the potential bias of a reporter assay, further *in vivo* experiments are required to confirm the regulation and to establish the physiological consequences.

Recently, several groups reported about the global analysis of predicted miRNA targets with sets of genes (i.e., pathway or gene ontology [GO] term). Gusev et al. (2007) reported GO terms as well as biological functions targeted by five coexpressed miRNAs sets. Cui et al. (2006) analyzed distribution of the predicted miRNA targets in the cellular signaling network. But no one explicitly explored the functional consequence of clustered miRNAs with signaling pathways. Our approach can be extended forward to GO and other pathway system (i.e., KEGG) analysis as well as human pathway analysis. It is also important to identify signaling pathways that are conditionally and tissue specifically regulated by miRNA clusters. This would require further fine tuning the mining methods and obtaining additional miRNA expression profiles under specific conditions. However, this direction of research is beyond the scope of our current study. Such systems biological approaches will provide insight into the principles of the




FIGURE 3. Model of the regulation of the insulin signaling pathway by miRNA cluster *mmu-mir-183-96-182*. *Mir-96* and *mir-183* are mainly responsible for regulating *Irs1*, which is the key mediator of the metabolic aspect of the insulin pathway, whereas *mir-182* targets *Rasa1* and *Grb2*, both of which play important roles in linking insulin signaling with the *Ras* pathway. A dotted arrow indicates the experimentally verified miRNA and target gene interactions via luciferase activity assay.

regulation of signaling networks by miRNAs and facilitate experimentally identifying interactions between miRNAs and target mRNAs.

MATERIALS AND METHODS

Data sources

Mouse miRNA clusters data

According to chromosome positional relationships between miRNAs, any two miRNAs with maximum distance less than defined distance were considered to be in the same cluster. We derived mouse miRNAs clusters from miRGen database (Megraw et al. 2007). Forty-five miRNA clusters which contains 169 miRNAs were extracted when MID was set to 5 kb. Fifty-five miRNA clusters which contains 202 miRNAs were found when MID was set to 50 kb.

Optimized targets prediction data

Sethupathy et al. (2006) compared various miRNA target-prediction programs both individually and in combination with other programs. They suggested that the intersection of PicTar and TargetScanS prediction could achieve both high sensitivity and specificity compared to other programs or program combinations. This optimized target prediction data was used in this study. It includes 21,225 miRNA–target gene pairs, which totally involved 112 miRNAs and 3586 target genes.

Mouse Biocarta pathway data

We use the compiled pathway data by Tian et al. (2005). It is consisted of 277 Mouse Biocarta pathway data, and could be

downloaded from <http://compbio.med.harvard.edu/Supplements/PNAS05.html>.

miRNA expression data

We used three mouse miRNA expression profile data sets. Thomson data: the investigators used a dual-channel microarray to monitor expression levels of 124 mouse miRNAs in seven mouse tissues and four different embryonic stages, embryonic stem cells, and embryoid bodies (Thomson et al. 2004). Beuvink data: chemically modified 2'-O-(2-methoxyethyl)-(MOE) oligo-ribonucleotide probes were arrayed onto Evanescent Resonance microchips to detect 200 miRNAs expression in eight mouse tissues (Beuvink et al. 2007). Betel data: miRNA expression profiles derived from a comprehensive cloning and sequencing effort of 64 mouse small RNA libraries extracted from major organs and cell types. Expression values were reported as normalized clone counts for 400 miRNAs (Betel et al. 2008).

Computational framework

miRNA profiling filter

We computed the Pearson coefficient for each pair of miRNAs within the same cluster. The mean correlation coefficients were used as a correlation measure for a miRNA cluster. For each miRNA profile, we also randomly sampled 1000 pairs of miRNAs and computed the Pearson coefficient. The average value of positive correlation measures was used as a background correlation. Finally, we filtered out such clusters that the mean correlation measure is less than those of the background correlation in at least two of the three miRNA profiles. If there are <2 miRNAs within the same cluster measured in the three expression profiles, the miRNA cluster passes the filter automatically as currently there are not enough evidences to show this cluster does not coexpressed. To assess the statistical significance of the correlation measure for each miRNA cluster, the fraction of 1000 random sampling pairs of miRNAs having equal to or larger than the correlation measure was reported as nominal *P* value.

Regulatory elements filter

We used FirstEF and Eponin programs to search for possible regulatory elements among the inter-miRNA region within the same cluster. FirstEF is a decision tree based promoter detection programs to predict exon, promoter, and CpG window of sequences (Davuluri et al. 2001). The Eponine program provides a probabilistic method for detecting transcription start sites (TSS) in the mammalian genomic sequence (Down and Hubbard 2002). For the analysis, we used the UCSC human multiz28way conservation track to get the corresponding mouse regions. All the sequence analyses were done at Galaxy platform (Giardine et al. 2005; Karolchik et al. 2008). We filtered out such clusters that the regulatory elements were found by any one of the two detection programs. miRNA cluster that passes both miRNA profiling filter and regulatory elements filter were used in the following analysis.

Filtering pathways

For the 277 Biocarta signaling pathways, there are some pathways that only a few or even none of its component genes are predicted

as miRNA targets in the optimized target prediction data. Only those signaling pathways with at least 50% of their component genes listed in the optimized target prediction data as target genes were selected for further analysis. After this selection, there remained 97 signaling pathways.

Finding significant interactions between clusters and pathways

For each cluster and each signaling pathway, we first found predicted target genes of all the miRNA members within this cluster. Then, we calculated the number of irredundant overlapping target genes with the component genes on the signaling pathway. One thousand simulated miRNA clusters of the same size were also randomly sampled from the 112 miRNAs with target prediction and then the corresponding number of overlapping targets was recorded. We set the *P* value of the cluster-pathway pair as the fraction of 1000 random sampling having equal to or larger than the number of overlapping targets of the real miRNA cluster. The *P* value based on such randomizations was used to assess whether the observed number of overlapping genes for a cluster-pathway pair was achieved by chance. Multiple statistical tests were controlled by false discovery rate (FDR) (Benjamini and Hochberg 1995). After applying the same procedure to all the cluster-pathway pairs, we selected the cluster-pathway pair at FDR <0.3 (nominal *P* < 0.03).

Experimental verification

The siRNA duplexes *mir-96*, *mir-153*, *mir-182*, and *mir-183* were purchased from Shanghai Gene-Pharma Co. siRNA duplexes with nonspecific sequences were used as a negative control (NC). Mouse genes 3'-UTR sequences were downloaded from Pictar website (<http://pictar.bio.nyu.edu/cgi-bin/PicTar Vertebrate.cgi>). The fragments of *irs1* (mm_NM_005544.0) 3'-UTR (nucleotides 33–1001), *grb2* (mm_NM_203506.0) 3'-UTR (nucleotides 82–598), and *rasa1* (NM_002890.0) 3'-UTR (nucleotides 9–936) were PCR-amplified and subcloned into the XbaI site in pGL3-Control vector (Promega Corp.). Constructs were all sequence verified (Guangzhou Invitrogen Biotechnology Co., Ltd.). In addition to nonspecific sequences as a negative control, specific negative controls were employed (both *mir-153* and *mir-182* for *irs1*, and *mir-183* for *rasa1*, and *grb2*) because no target sites were predicted for the corresponding miRNA oligos by both Pictar and TargetScanS algorithms.

For luciferase activity assay, HeLa cells were seeded 24 h prior to transfection in 96-well tissue culture plates. The next day, cells were cotransfected with various constructs and internal control pRL-TK and 10 nM of each miRNA duplex using Lipofectamine 2000 in accordance with the manufacturer's instructions. Cell lysate was collected 24 h after transfection. Firefly and Renilla luciferase activities were measured using a Dual Luciferase Reporter Assay System (Promega Corp.). All of the assays were performed in triplicate in at least three independent experiments. We assessed statistical significance using two-tailed Student's *t*-test.

SUPPLEMENTAL DATA

Supplemental material can be found at <http://www.rnajournal.org>.

ACKNOWLEDGMENTS

We thank the UCSC Genome Bioinformatics Group for helpful discussion on genome analysis for this work, and Molly Megraw at the University of Pennsylvania for kindly providing us with the optimized target prediction data. We also thank Zhanfang Kang for his assistance on graphic preparation. The research is supported by grants from the National Science Foundation of China #30672463, the Chinese National Science and Technology Department #2006CB50390, the Chinese Academy of Sciences #KSCX2-YW-R-085, and a start-up grant from GIBH.

Received January 10, 2008; accepted April 10, 2008.

REFERENCES

- Altuvia, Y., Landgraf, P., Lithwick, G., Elefant, N., Pfeffer, S., Aravin, A., Brownstein, M.J., Tuschl, T., and Margalit, H. 2005. Clustering and conservation patterns of human microRNAs. *Nucleic Acids Res.* **33**: 2697–2706.
- Bandres, E., Cubedo, E., Agirre, X., Malumbres, R., Zarate, R., Ramirez, N., Abajo, A., Navarro, A., Moreno, I., Monzo, M., et al. 2006. Identification by real-time PCR of 13 mature microRNAs differentially expressed in colorectal cancer and non-tumoral tissues. *Mol. Cancer* **5**: 29. doi: 10.1186/1476-4598-5-29.
- Bartel, D.P. 2004. MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Baskerville, S. and Bartel, D.P. 2005. Microarray profiling of microRNAs reveals frequent coexpression with neighboring miRNAs and host genes. *RNA* **11**: 241–247.
- Benjamini, Y. and Hochberg, Y. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* **57**: 289–300.
- Betel, D., Wilson, M., Gabow, A., Marks, D.S., and Sander, C. 2008. The microRNA.org resource: Targets and expression. *Nucleic Acids Res.* **36**: D149–D153.
- Beuvink, I., Kolb, F.A., Budach, W., Garnier, A., Lange, J., Natt, F., Dengler, U., Hall, J., Filipowicz, W., and Weiler, J. 2007. A novel microarray approach reveals new tissue-specific signatures of known and predicted mammalian microRNAs. *Nucleic Acids.* **35**: e52. doi: 10.1093/nar/gkl1118.
- Cai, X., Hagedorn, C.H., and Cullen, B.R. 2004. Human microRNAs are processed from capped, polyadenylated transcripts that can also function as mRNAs. *RNA* **10**: 1957–1966.
- Cui, Q., Yu, Z., Purisima, E.O., and Wang, E. 2006. Principles of microRNA regulation of a human cellular signaling network. *Mol. Syst. Biol.* **2**: 46. doi: 10.1038/msb4100089.
- Davuluri, R.V., Grosse, I., and Zhang, M.Q. 2001. Computational identification of promoters and first exons in the human genome. *Nat. Genet.* **29**: 412–417.
- Dews, M., Homayouni, A., Yu, D., Murphy, D., Seignani, C., Wentzel, E., Furth, E.E., Lee, W.M., Enders, G.H., Mendell, J.T., et al. 2006. Augmentation of tumor angiogenesis by a Myc-activated microRNA cluster. *Nat. Genet.* **38**: 1060–1065.
- Dharmawardana, P.G., Peruzzi, B., Giubellino, A., Burke Jr., T.R., and Bottaro, D.P. 2006. Molecular targeting of growth factor receptor-bound 2 (Grb2) as an anti-cancer strategy. *Anticancer Drugs* **17**: 13–20.
- Down, T.A. and Hubbard, T.J. 2002. Computational detection and location of transcription start sites in mammalian genomic DNA. *Genome Res.* **12**: 458–461.
- Downward, J. 2003. Targeting RAS signalling pathways in cancer therapy. *Nat. Rev. Cancer* **3**: 11–22.
- Draznin, B. 2006. Molecular mechanisms of insulin resistance: Serine phosphorylation of insulin receptor substrate-1 and increased expression of p85alpha: The two sides of a coin. *Diabetes* **55**: 2392–2397.

- Fontana, L., Pelosi, E., Greco, P., Racanicchi, S., Testa, U., Liuzzi, F., Croce, C.M., Brunetti, E., Grignani, F., and Peschle, C. 2007. MicroRNAs 17-5p-20a-106a control monocytopenesis through AML1 targeting and M-CSF receptor upregulation. *Nat. Cell Biol.* **9**: 775–787.
- Giardine, B., Riemer, C., Hardison, R.C., Burhans, R., Elnitski, L., Shah, P., Zhang, Y., Blankenberg, D., Albert, I., Taylor, J., et al. 2005. Galaxy: A platform for interactive large-scale genome analysis. *Genome Res.* **15**: 1451–1455.
- Gusev, Y., Schmittgen, T.D., Lerner, M., Postier, R., and Brackett, D. 2007. Computational analysis of biological functions and pathways collectively targeted by co-expressed microRNAs in cancer. *BMC Bioinformatics (Suppl. 7)* **8**: S16. doi: 10.1186/1471-2105-8-S7-S16.
- Hua, Z., Lv, Q., Ye, W., Wong, C.K., Cai, G., Gu, D., Ji, Y., Zhao, C., Wang, J., Yang, B.B., et al. 2006. MiRNA-directed regulation of VEGF and other angiogenic factors under hypoxia. *PLoS ONE* **1**: e116. doi: 10.1371/journal.pone.0000116.
- Karolchik, D., Kuhn, R.M., Baertsch, R., Barber, G.P., Clawson, H., Diekhans, M., Giardine, B., Harte, R.A., Hinrichs, A.S., Hsu, F., et al. 2008. The UCSC Genome Browser Database: 2008 update. *Nucleic Acids Res.* **36**: D773–D779.
- Kloosterman, W.P. and Plasterk, R.H. 2006. The diverse functions of microRNAs in animal development and disease. *Dev. Cell* **11**: 441–450.
- Krek, A., Grun, D., Poy, M.N., Wolf, R., Rosenberg, L., Epstein, E.J., MacMenamin, P., da Piedade, I., Gunsalus, K.C., Stoffel, M., et al. 2005. Combinatorial microRNA target predictions. *Nat. Genet.* **37**: 495–500.
- Lee, Y., Kim, M., Han, J., Yeom, K.H., Lee, S., Baek, S.H., and Kim, V.N. 2004. MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.* **23**: 4051–4060.
- Megraw, M., Sethupathy, P., Corda, B., and Hatzigeorgiou, A.G. 2007. miRGen: A database for the study of animal microRNA genomic organization and function. *Nucleic Acids Res.* **35**: D149–D155.
- Pal, S., Baiocchi, R.A., Byrd, J.C., Grever, M.R., Jacob, S.T., and Sif, S. 2007. Low levels of miR-92b/96 induce PRMT5 translation and H3R8/H4R3 methylation in mantle cell lymphoma. *EMBO J.* **26**: 3558–3569.
- Sethupathy, P., Megraw, M., and Hatzigeorgiou, A.G. 2006. A guide through present computational approaches for the identification of mammalian microRNA targets. *Nat. Methods* **3**: 881–886.
- Stark, A., Brennecke, J., Bushati, N., Russell, R.B., and Cohen, S.M. 2005. Animal MicroRNAs confer robustness to gene expression and have a significant impact on 3'-UTR evolution. *Cell* **123**: 1133–1146.
- Thomson, J.M., Parker, J., Perou, C.M., and Hammond, S.M. 2004. A custom microarray platform for analysis of microRNA gene expression. *Nat. Methods* **1**: 47–53.
- Tian, L., Greenberg, S.A., Kong, S.W., Altschuler, J., Kohane, I.S., and Park, P.J. 2005. Discovering statistically significant pathways in expression profiling studies. *Proc. Natl. Acad. Sci.* **102**: 13544–13549.
- Weston, M.D., Pierce, M.L., Rocha-Sanchez, S., Beisel, K.W., and Soukup, G.A. 2006. MicroRNA gene expression in the mouse inner ear. *Brain Res.* **1111**: 95–104.
- Wong, C.F. and Tellam, R.L. 2008. microRNA-26a targets the histone methyltransferase Enhancer of Zeste homolog 2 during myogenesis. *J. Biol. Chem.* **283**: 9836–9843.
- Xu, S., Witmer, P.D., Lumayag, S., Kovacs, B., and Valle, D. 2007. MicroRNA (miRNA) transcriptome of mouse retina and identification of a sensory organ-specific miRNA cluster. *J. Biol. Chem.* **282**: 25053–25066.