



Published in final edited form as:

Mol Ecol. 2008 May ; 17(9): 2276–2286.

Multigene analysis suggests ecological speciation in the fungal pathogen *Claviceps purpurea*

G. W. DOUHAN¹, M. E. SMITH², K. L. HUYNH¹, A. WESTBROOK², P. Beerli³, and A. J. FISHER⁴

¹ Department of Plant Pathology and Microbiology, University of California, Riverside, CA (USA) 92521

² Department of Plant Pathology, One Shields Avenue, University of California, Davis, CA (USA) 956161

³ School of Computational Science and Biological Sciences, Department Florida State University Tallahassee, FL, (USA) 32306-4120

⁴ USDA, ARS, Exotic and Invasive Weeds Research Unit, Albany, CA (USA) 94710

Abstract

Claviceps purpurea is an important pathogen of grasses and source of novel chemical compounds. Three groups within this species (G1, G2, and G3) have been recognized based on habitat association, sclerotia and conidia morphology, and alkaloid production. These groups have further been supported by RAPD and AFLP markers, suggesting this species may be more accurately described as a species complex. However, all divergent ecotypes can coexist in sympatric populations with no obvious physical barriers to prevent gene flow. In this study, we used both phylogenetic and population genetic analyses to test for speciation within *C. purpurea* using DNA sequences from ITS, a RAS-like locus, and a portion of beta-tubulin. The G1 types are significantly divergent from the G2/G3 types based on each of the three loci and the combined dataset, whereas the G2/G3 types are more integrated with one another. Although the G2 and G3 lineages have not diverged as much as the G1 lineage based on DNA sequence data, the use of three DNA loci does reliably separate the G2 and G3 lineages. However, the population genetic analyses strongly suggest little to no gene flow occurring between the different ecotypes and we argue that this process is driven by adaptations to ecological habitats; G1 isolates are associated with terrestrial grasses, G2 isolates are found in wet and shady environments, and G3 isolates are found in salt marsh habitats.

Keywords

Claviceps; fungi; genealogy; genetic structure; speciation

Introduction

Species concepts have been intensely debated in the literature and no one concept is generally accepted among biologists. Traditional species concepts for the classification of fungi are based on morphology and reproductive biology, but phylogenetic approaches have become increasingly popular in recent years to resolve species identification (Taylor *et al.* 2000; Harrington & Rizzo 1999). Moreover, phylogenetic analyses have challenged morphological species concepts and have been especially helpful in delineating fungal species with few morphological characters. Phylogenetic approaches have also been sought for the many fungi in which sexual reproduction is not known to occur, making biological species concepts

impossible to implement. Phylogenetic studies have routinely identified cryptic species within morphological species in various fungal genera including *Fusarium* (Skovgaard *et al.* 2002; O'Donnell *et al.*, 2000), *Stachybotrys* (Cruse *et al.* 2002), *Tricholoma* (Horton 2002), *Coccidioides* and close relatives (Koufopanou *et al.* 2001), *Cenococcum* (Douhan & Rizzo 2005), *Neurospora* (Dettman *et al.* 2003), and lichenized genera such as *Physcia* (Myllys *et al.* 2001) and *Letharia* (Kroken & Taylor 2001).

What has not been determined in most of these studies is the mechanism or driving force behind the speciation process. In some instances, geographic isolation has been suggested, as in *Coccidioides* (Koufopanou *et al.* 2001) and *Fusarium* (O'Donnell *et al.* 2000), whereas in other cases these putative cryptic species may be found in the same geographic location and in some instances even isolated from the same soil core (Douhan & Rizzo 2005; Moyersoen *et al.* 2003; Skovgaard *et al.* 2002). Ecological theory predicts that the stable coexistence of identical competitors will not occur (Hardin 1960), suggesting that cryptic species occupying the same apparent niche may play different ecological roles. Therefore, ecological factors likely play a significant role in the speciation process of co-occurring organisms. This 'ecological speciation' process has received little attention in the fungi. We will adopt the terminology of Rundle and Nosil (2005) that defines 'ecological speciation' as the "process by which barriers to gene flow evolve between populations as a result of ecologically-based divergent selection."

One group of fungi where ecological aspects may have influenced speciation is within the *Claviceps purpurea* complex. These are plant pathogenic fungi that infect the flowers of grasses and cause a disease known as 'ergot.' The infection results in a replacement of the grass floral tissue with fungal mycelium that grows to form a darkly pigmented survival structure called a sclerotium (Alexopolous *et al.* 1996). Sclerotia of *C. purpurea* contain potent alkaloids that are toxic to humans and other animals (Hudler, 1998). This has been important during human history because sclerotia have been unwittingly mixed with grain and the fungal toxins have contaminated bread flour. The consumption of bread tainted with *C. purpurea* causes the debilitating disease known as 'Holy Fire' or 'St. Anothony's Fire,' which includes a range of symptoms such as itching, headache, hallucinations, gangrene, seizures, and even death (Hudler, 1998). Despite the negative impacts of *C. purpurea* alkaloids, interesting and beneficial chemical compounds have also been derived from these fungi, including the infamous lysergic acid diethylamide (LSD) and pharmaceuticals used to treat both postpartum bleeding and migraine headaches (Hart 1999, Rehacek & Sajdl 1993)

Pazoutová *et al.* (2000) synthesized previous research on *C. purpurea* morphology, alkaloid chemistry and genetics, and identified three distinct groups within the species, describing them as 'chemoraces.' Rather than species delimitation based on host range, which was historically common in *C. purpurea* taxonomy, intraspecific groups were defined based on habitat specialization. The largest group, G1, is associated with land grasses. G2 isolates are associated with grasses in 'wet and shady' environments, whereas G3 isolates are found only on grasses in salt marsh habitats. G3 isolates are synonymous with *C. purpurea* var. *spartinae* (Duncan *et al.* 2002) and are referred to herein as G3. Sclerotia of both G2 and G3 *C. purpurea* float in water while sclerotia from terrestrial *C. purpurea* (G1) sink, clarifying the significance of earlier reports that sclerotia of some isolates float, and the association of this trait with host habitat (Stager 1922).

Our objective in this study was to investigate speciation within the *C. purpurea* complex using phylogenetic and population genetic approaches based on representative samples. Isolates included in this study were selected as representative of the variation within *C. purpurea sensu lato* and chosen to cover a large geographic area, a wide host range within the Poaceae, and all three habitat types. The group identity of each isolate used in this study was determined previously by RAPD and AFLP (Fisher *et al.* 2005b; 2005c; Pazoutová *et al.* 2000). RAPD

and AFLP analyses of these, and additional isolates, supports the recognition of three discrete groups within *C. purpurea* and revealed high genetic variability between groups, with less than 2% of polymorphic markers shared across all isolates. Similarly, analysis of molecular variation (AMOVA) revealed that genetic variability was mainly due to variations between groups rather than within groups or populations, revealing a high probability of phylogenetic substructure (Fisher *et al.* 2005b; 2005c). Determining if the *C. purpurea* complex is made up of one or multiple species is important because it is common practice in wide-ranging studies to make no distinction between the three different *C. purpurea* lineages (Tooley *et al.* 2001; Stensrud *et al.* 2005; Scheffer & Tudzynski 2006; Dabkeviccius & Mikaliunaite 2006; Scheffer *et al.* 2005).

Materials and methods

Molecular analyses

Fungal isolates & DNA extraction—*Claviceps purpurea* isolates used in this study and their origins are listed in Table 1. For field-collected isolates, sclerotia were collected during the fall of 2000, 2001 and 2002 and brought to University of California, Davis for culturing. Sclerotia were surface sterilized, cultured, dried and refrigerated as described in Fisher *et al.* (2005c). Numbered isolates were obtained as pure cultures from S. Paloutová (Institute of Microbiology, Czech Republic) including: G1 isolates 165, 204, 428 and G2 isolates 236 and 434 (Fisher *et al.* 2005c; Paloutová *et al.*, 2000).

Tissue for DNA extraction was obtained by growing isolates on cellophane overlaid on PDA as described by Fisher *et al.* (2005c) or by scraping fungal mycelium directly from PDA plates. Total DNA was extracted using the methods described by Daehler *et al.* (1999) or slightly modified from Gardes and Bruns (1993). After DNA was recovered in solution, 75 μ l of supernatant was withdrawn and mixed with 200 μ l Tris-EDTA buffer (10 mM Tris, 1 mM EDTA (pH 7.8). DNA concentration was quantified by spectrophotometry.

PCR, Sequencing and phylogeny construction—PCR was performed in 40 μ l reaction mixtures containing 2 μ l of a 1:10 to 1:25 dilution of template DNA, 1X PCR buffer (Invitrogen, Carlsbad, CA), 2.5 mM MgCl₂, 0.2 mM each dNTP (Invitrogen), 7.5 μ M of each primer, and 0.5 U of Taq polymerase (Invitrogen). Three loci were amplified and sequenced; ITS rDNA using the primers ITS1F and ITS 4 (Gardes & Bruns 1993; White *et al.* 1990), a portion of a beta-tubulin gene using the primers described in Annis and Panaccione (1998), and a putative RAS-like protein using primers described by Carbone and Kohn (1999). Thermocycling conditions consisted of an initial hold at 94°C (4 min), followed by 30 cycles with a denaturing step of 94°C (30 sec), annealing temperature of 50° to 55°C (30 sec), and an extension temperature of 72°C (1 min). All amplifications were performed in a PE-9700 thermocycler (Perkin Elmer Corp., Norwalk, CT) or a MyCycler (Bio-Rad Laboratories Inc., Hercules, CA, USA).

Each locus was sequenced in both directions using Big Dye® Terminator v3.1 chemistry (Applied Biosystems, Foster City, CA, USA). The PCR products were cleaned using ExoSapit (USB Amersham, Uppsala, Sweden) following the manufactures instructions and sequenced at the Core Instrumentation Facility (CIF) of the University of California's Institute of Integrative Genome Biology at UC Riverside. The sequences were edited using Sequencher (version 4.1.2, Gene Codes Corporation, Ann Arbor, MI), aligned using Clustal X (version 1.81) (Chenna *et al.* 2003) and manually edited in MacClade version 4 (Maddison & Maddison 2001). Maximum Parsimony (MP) analysis using the heuristic search procedure with 1000 random-addition-sequence replicates and tree-bisection-reconnection branch swapping were conducted using PAUP* version 4.0 beta 10 (Swofford 2002). Confidence in tree topology

was examined using bootstrap with 1,000 replicates under the heuristic option. All trees were midway rooted.

Population genetic analyses—We investigated the genetic separation of the three ecotypes (G1, G2, & G3) using the program MIGRATE-N (Version 2.3.3: Beerli and Felsenstein 1999, 2001, Beerli 2006) that is based on coalescence theory (cf Kingman 2000). The three ecotypes were treated as independent genetic units where only migrants and mutation could import new alleles into a unit. We compared two different models: one single population versus 3 populations with 6 different migration rates between all populations. This allowed us to test whether the 3 loci were powerful enough to reject if these three ecotypes were generated by a single random mating population.

MIGRATE-N does not take into account potential splitting of the ancestral population. Under such a model, the different ecotypes should show low migration rates if there was no recent divergence or no ongoing gene flow. We tested whether high gene flow between the ecotypes could be rejected using a Bayes Factor approach (Kass and Raftery, 1995) by running the Bayesian module of MIGRATE-N for both models and compared the marginal likelihoods.

We used the default settings for MIGRATE-N, except for the following run-options: we used the Bayesian inference module; one single long run using heating with temperatures of 1.0, 1.5, 3.0, and 10,000, totaling 50,100,000 visited parameter and genealogy changes in the cold (1.0) chain; sampling 10 replicates of each 50,000 in intervals of 100; after discarding the first 100,000 visits. We used uniform prior distributions with ranges from 0 to 0.1 for the mutation scaled population size Θ , that is the effective population size N_e times the mutation rate μ per site and generation, and ranges from 0 to 5000 for M , that is the mutation scaled immigration rate m/μ . The mutation model used was the F84 model (Felsenstein 2004) with a transition-transversion ratio set to 2. We report the mode and median, and the 95% credibility set of the posterior distribution for all estimated parameters of the two models.

Estimates of the average number of nucleotide differences, and shared, fixed and unique number of mutations between the G1, G2, and G3 ecotypes were calculated in DNAsp 4.0 (Rozas *et al.* 2003). *Fst* values were also calculated in DNAsp 4.0, with permutation tests used to test significance using 1000 randomizations.

Results

Phylogenetic inference

For beta-tubulin, 426 characters were analyzed with 5 parsimony un-informative and 13 parsimony informative characters. Figure 1 shows one of the 29 most parsimonious trees that were found with a length of 19. Consistency index, retention index, and the rescaled consistency index were 0.947, 0.987, and 0.935, respectively. All G2 isolates except 236 had two copies of the repeat AACTG starting at position 113 in the alignment whereas all G1 and G3 isolates had a single copy of the AACTG repeat. All G1 isolates clustered in a clade with a bootstrap support of 100. The G2 and G3 isolates clustered together into one weakly supported clade. Although all G3 isolates clustered together with weak bootstrap support, G2 isolates were more variable with two isolates, 434 and 236, notably different from the rest (Fig. 1).

For ITS, 586 characters were analyzed with 4 parsimony un-informative and 8 parsimony informative characters. Figure 1 shows the single most parsimonious tree with a length of 14. Consistency index, retention index, and the rescaled consistency index were 0.929, 0.977, and 0.907, respectively. The G1 clade was supported with a bootstrap score of 83. The G2 and G3 isolates clustered together with a bootstrap value of 88. Although G2 and G3 isolates clustered separately into their own subclades, these subclades had weak bootstrap support.

For RAS, 375 characters were analyzed with 5 parsimony un-informative and 13 parsimony informative characters. Figure 1 shows one of a total of 354 most parsimonious trees that were found with a length of 15. Consistency index, retention index, and the rescaled consistency index were 0.800, 0.955, and 0.764, respectively. The G1 isolates clustered together with a bootstrap support of 99. The G2 and G3 isolates clustered together with poor bootstrap support for any sub-groupings.

In the combined dataset including all three genes, 1294 characters were analyzed with 8 parsimony un-informative and 32 parsimony informative characters. The consistency index, retention index, and the rescaled consistency index were 0.840, 0.956, and 0.803, respectively. The G1, G2, and G3 clades were supported by bootstrap values of 83, 68, and 82, respectively. The G2/3 clade was supported with a bootstrap value of 100. No obvious substructuring based on geography was found for any of the three loci or the combined dataset.

Population Genetic analyses

3-population model (Each ecotype is treated as a population): Results on migration rates are all low between all populations with large credibility intervals, that for individual loci always include zero; the combined estimates exclude zero migration rates. The posterior distribution for the scaled migration rates are all heavily skewed to the right, and peak all near zero compared to the whole distribution. We are confident that the prior distribution, a uniform distribution, had little influence on our results. The population sizes Θ reveal that the ecotype G3 has more variability and a larger population than the other types, G1 and G2. Table 2 gives credibility intervals and mode and medians. Typical values for the number of immigrants Nm ($\Theta M/4$) are 1 immigrant every four generations.

1-population model (All ecotypes are pooled into one population): the population is much larger than the combined population size of the 3-population model suggesting that, because we ignored migration, the length of the genealogies are lengthened which results in an overestimate of the population size (Excoffier, 2004). We compared our results from the 3-population model with 1-population model using a Log-Bayes factor (BF). Using the reported marginal likelihoods from the two MIGRATE-N analyses: $\log L(3\text{-population model}) = -2073.887689$; $\log L(1\text{-population model}) = -2204.2369$. Then the $\text{Log}(\text{BF})$ is $\log L(3) - \log L(1) = 130.34$. Values higher than 10 suggest that we should strongly prefer the first model, in our case the three-population model (Kass and Raftery, 1995).

Table 3 shows the summary statistics in the comparison between the G1, G2, and G3 isolates of *C. purpurea*. In the G2–G3 comparisons, the average number of nucleotide differences and nucleotide diversity were consistently lower than in the comparisons of G1–G3 and G1–G2. This is also consistent with lower F_{st} values for G2–G3 comparisons as compared with G1–G3 and G1–G2 comparisons, except for the ITS locus in which the lowest F_{st} value was for the G1–G3 comparison. All F_{st} comparisons were significant ($P < 0.001$), suggesting limited gene flow between the three *C. purpurea* lineages from different habitats, which was consistent with the coalescence analyses.

Discussion

In this study, phylogenetic and population genetic analyses showed marked genetic differences among the different ecotypes and suggest little or no gene flow among the different ecotypes. We can also definitely reject models that assume random mating between the different ecotypes based on the analyses. The G1 types are significantly divergent from the G2/G3 habitat types based on each of the three loci and the combined dataset, whereas the G2/G3 types are more integrated with one another. However, although the G2 and G3 lineages have not diverged as much as the G1 lineage based on DNA sequence data, the use of three DNA loci does reliably

separate the G2 and G3 lineages. The fact that the G2 and G3 lineages are more closely related to each other than to the G1 lineage is strongly supported by the fact that sclerotia from G2 and G3 isolates float in water while those of G1 isolates sink (Pazoutová *et al.* 2000). Results from this study are in agreement with previous conclusions based on AFLP and RAPD data, in which only 2% of genetic markers were shared among the three lineages of *C. purpurea*. Perhaps because of their high rate of polymorphism or because of their assessment of variation across the entire genome, AFLP and RAPD data were more informative than DNA sequences for separating the different ecotypes. A similar situation was recently shown in two species complexes of mycoparasites, *Hypomyces microspermus* and *H. chrysospermus* (Douhan & Rizzo 2003). In this case, AFLP clearly differentiated cryptic lineages within both *H. microspermus* and *H. chrysospermus*, whereas ITS rDNA sequence data recovered the same cryptic lineages, but in some cases with weak bootstrap support (Douhan & Rizzo 2003).

Should three distinct species within the *C. purpurea* complex be recognized taxonomically? As previously mentioned, traditional species concepts for the classification of fungi are based on morphology and reproductive biology, but phylogenetic approaches may be more powerful to accurately resolve fungal lineages (Taylor *et al.* 2000; Harrington & Rizzo 1999). During the speciation process, reproductive barriers arise between groups of individuals. The two groups at first share allelic polymorphisms until one of the two groups become fixed for certain alleles whereas the other group remains polymorphic (Avice 1994; Geiser *et al.* 1998). Therefore, based on previous AFLP and RAPD data, it would appear that the justification for species recognition would be warranted. However, Taylor *et al.* (2000) advocate identifying phylogenetic species of fungi by analyzing evolutionary relationships among multiple genes, which they call the Genealogical Concordance Phylogenetic Species Recognition (GCPSR). They suggest the use of multiple genes to determine the transition from concordance to conflict among taxa, which can be used to determine species boundaries. The conflict is thought to be due to recombination occurring between individuals. Based on the GCPSR criteria we would reject that there are three species within the *C. purpurea* complex and conclude that two species exist, representing the G1 and G2/G3 lineages. However, for more recently derived taxa, a marker system such as AFLP may be more powerful than DNA sequence loci to determine species boundaries because methods such as AFLP screen many loci across a genome. Despite the strengths of markers such as AFLP, inferring phylogenies based on randomly amplified fragments can be problematic, and there are arguments both for (Buntjer *et al.* 2002; Després *et al.* 2002; Kardolus *et al.* 1998) and against this approach (Seberg & Peterson 1998).

The overall evidence from our phylogenetic and population genetic analyses and previous studies suggest that the three lineages (G1, G2, and G3) should be recognized as unique species, or at least as varieties. Duncan *et al.* (2002) proposed renaming the G3 lineage as *C. purpurea* var. *spartinae* based on a combination of host identity (*Spartina* spp.), sclerotia ecology (floatation) and conidial morphology. However, since the variety status was proposed, G3 isolates were found infecting a non-*Spartina* host in nature (*Distichlis spicata* – Fisher *et al.* 2005b). Therefore, the named variety based on host association does not seem warranted. Regardless of what these fungi are called, it is important that some distinction be made since *C. purpurea sensu lato* is widely studied as a model plant pathogen of economic importance (Tudzynski & Scheffer 2004). Several lines of evidence suggest three divergent lineages within *C. purpurea* and these lineages likely have important biological differences relevant for studies in the wider scientific community. Currently, it is common practice in wide-ranging studies of everything from pathogenicity of *C. purpurea* on various hosts to molecular studies of virulence genes to phylogenetic assessments of the genus *Claviceps* to make no distinction between the three different *C. purpurea* lineages (Stensrud *et al.* 2005; Scheffer & Tudzynski 2006; Dabkevicius & Mikaliunaite 2006; Scheffer *et al.* 2005).

Some authors have argued that species concepts should have an ecological basis as well as a genetic one (Harrington & Rizzo 1999), and we suggest that ecological factors have driven the speciation process in *C. purpurea*. In estuarine habitats, the transition from salt-water tolerant species such as *Spartina* spp. and *Distichlis spicata*, to riparian and terrestrial grasses, is often gradual, with no physical barrier between habitat types (Mitsch & Gosselink 1993). The boundaries of the high tide and the slope of the terrain delimit the change from one habitat to the next, and this will differ by site. For example, in Willapa Bay, Washington (USA) the G1, G2 and G3 lineages coexist within a distance of less than 100 meters, with no physical barriers to spore dispersal by rain splash or insect vectors (Fisher *et al.* 2005b). It is not known if G1, G2 and G3 *C. purpurea* diverged sympatrically in a habitat such as a coastal estuary, or allopatrically with geographic barriers to gene flow. However, since there was no phylogeographic structure found in this diverse sampling of isolates and host grass species, it seems likely that ecological factors were more important in the speciation process of these fungi. The genus *Claviceps* is thought to have a Gondwanan origin and most species in the genus are tropical or subtropical. It has been hypothesized that species close to *C. purpurea* migrated from South America to North America after the formation of the Panama land bridge, and they later spread to Europe and Africa (Pazoutová 2003). It is only these species that developed the ability to deal with cold winters and semi-arid conditions, which is potentially more evidence suggesting that ecological habitat helped shaped this species complex.

Although it remains to be determined if the three habitat-associated lineages are reproductively isolated, results from our phylogenetic analysis suggest that reproduction among G2/3 isolates would be more likely than between either G1 and G2 or G1 and G3 isolates. However, the population genetic analyses strongly suggest little to no gene flow occurring between the different ecotypes. In order for reproduction to occur, these different ecotypes must be able to infect the same hosts. The results of a host range study showed that G3 isolates can infect both riparian and terrestrial grasses after artificial inoculation (Pazoutová *et al.* 2002), though its range in nature is so far limited to the C4 grasses *Spartina* spp. and *Distichlis spicata*. Similarly, Pazoutová *et al.* (2000; 2002) reported that G2 isolates can infect both riparian and terrestrial grasses in the greenhouse but this has not been documented under natural conditions. Currently we have not identified the barriers to gene flow in sympatric G1, G2 and G3 *C. purpurea* populations like in the Willapa Bay's upper marsh, but possibilities include differences in flowering time or flowering duration among hosts, or differences in plant biochemistry which might preclude infection of C3 grasses by a pathogen adapted to C4 grass hosts. Thus, as in other fungi, there may be multiple habitat-related factors that drive the speciation process. For example, it has been suggested that the behavior of insect vectors or physiological barriers to mating or infection may prevent gene flow among host races of the anther smut fungus *Microbotryum violaceum* in sympatric populations of three host plant species (Shykoff *et al.* 1999).

Regardless of whether the divergence among *C. purpurea* lineages occurred in allopatric or sympatric populations, selection pressures in maritime habitats are distinct from those necessary for dispersal and survival in terrestrial habitats and *C. purpurea* from *Spartina* and *D. spicata* (G3) exhibit characteristics uniquely suited for maritime environments. For example, sclerotia from G2 and G3 *C. purpurea* float on water (Pazoutová *et al.* 2000), presumably due to large intercellular spaces (Duncan *et al.* 2002), and flotation presumably aids both survival and dispersal in areas with flooding or tides. In addition, G3 sclerotia do not require a cold stratification prior to germination, a requirement common in *C. purpurea* from terrestrial habitats (G1) not likely to be met in coastal environments (Duncan *et al.* 2002). Although data is lacking, tolerance for highly salinated leaves, water, and soil is likely involved in directional selection leading to the isolation of the G3 lineage.

Host range within *C. purpurea* also suggests that ecology is more important than host in the evolution of these fungi, which is different from many plant-associated fungi. For example, within the plant pathogenic *Magnaporthe grisea* species complex there is strong evidence that speciation and genetic divergence are highly coupled with virulence on particular species or even varieties of host grasses. Although host switches have been documented, radiation events in this group of fungal pathogens appear to primarily follow the evolutionary history of hosts (Kohn 2005; Couch *et al.* 2005; Couch & Kohn 2002). A similar pattern of host-associated speciation has also been suggested for some symbiotic fungi, such as in the *Pinus*-associated ectomycorrhizal mushroom genus *Suillus*. In a study of disjunct *Suillus* species complexes from East Asia and Eastern North America, Wu *et al.* (2000) suggested that high host-fidelity for species or subgenera of *Pinus* has led to speciation through comigration.

“Ecological speciation” has previously been suggested as the mode of speciation among many diverse animal groups (Via 2001) including fishes (Rocha *et al.* 2005; Hatfield & Schluter 1999), lizards (Richmond & Reeder 2002; Ogden & Thorpe 2002; Rosenblum 2006), toads (Kruuk & Gilchrist 1997), brittle stars (Muths *et al.* 2006), snails (Cruz *et al.* 2004) and various phytophagous and non-phytophagous insects (Via *et al.* 2000; Runle & Nosil 2005). Similarly, at least six examples of putative ecological speciation have been inferred among angiosperm plants from diverse habitats on several continents (Wang *et al.* 1997; Rieseberg 2000; Lamont *et al.* 2003; Nagy 1997; Fine *et al.* 2005; Hall & Willis 2006). Ecological speciation in fungi has not received much attention and speciation is usually attributed to hosts for pathogenic fungi or due to geographic vicariance, both of which may or may not have ecological similarities. However, as far as we know, *Claviceps purpurea* is only the second specific example of a fungus where ecological speciation is the main proposed mode of speciation. The other example of possible ecological speciation in fungi is in the insect-pathogenic fungus *Metarhizium anisopliae*. In a study of *M. anisopliae* in Ontario (Canada), Bidochka *et al.* (2001) showed that this fungus was clearly segregated into two genetic groups, one common in agricultural areas and the other in forested habitats. The authors did not find consistent differences in host preference between the cryptic species, but they detected distinct differences in tolerance to UV light and temperature sensitivity. Although these are the only examples that we know of where ecological speciation has been indicated in fungi specifically, we expect that this is a product of the lack of study and we expect that many more cases of ecological speciation will be found among fungi in the future.

Acknowledgements

Financial support of the Agricultural Experiment Station, University of California Riverside, for GD is gratefully acknowledged. PB is supported by the joint NSF/NIGMS Mathematical Biology program under NIH grant R01 GM 078985. We also thank Sylvia Pazoutová for cultures of *C. purpurea*.

References

- Alexopolous, C.J.; Mims, C.; Blackwell, M. *Introductory Mycology*. John Wiley and Sons, Inc; New York, USA: 1996.
- Annis SL, Panaccione DG. Accumulation of peptide synthetase gene transcripts and ergopeptines in cultures of ergopeptine-producing fungi. *Canadian Journal of Microbiology* 1998;44:80–86. [PubMed: 9546869]
- Avise, J.C. *Molecular Markers, Natural History and Evolution*. Chapman & Hall; New York, NY: 1994.
- Beerli P, Felsenstein J. Maximum-likelihood estimation of migration rates and effective population numbers in two populations using a coalescent approach. *Genetics* 1999;152:763–73. [PubMed: 10353916]
- Beerli P, Felsenstein J. Maximum likelihood estimation of a migration matrix and effective population sizes in n subpopulations by using a coalescent approach. *Proceedings of the National Academy of Sciences of the United States of America* 2001;98:4563–4568. [PubMed: 11287657]

- Beerli P. Effect of unsampled populations on the estimation of population sizes and migration rates between sampled populations. *Molecular Ecology* 2004;13:827–836. [PubMed: 15012758]
- Beerli P. Comparison of Bayesian and maximum-likelihood inference of population genetic parameters. *Bioinformatics* 2006;22:341–345. [PubMed: 16317072]
- Bidochka MJ, Kamp AM, Lavender TM, Dekoning J, De Croos JNM. Habitat association in two genetic groups of the insect-pathogenic fungus *Metarhizium anisopliae*: Uncovering cryptic species? *Applied and Environmental Microbiology* 2001;67:1335–1342. [PubMed: 11229929]
- Buntjer JB, Otsen M, Nijman IJ, Kuiper MTR, Lenstra JA. Phylogeny of bovine species based on AFLP fingerprinting. *Heredity* 2002;88:46–51. [PubMed: 11813106]
- Carbone I, Kohn LM. A method for designing primer sets for speciation studies in filamentous ascomycetes. *Mycologia* 1999;91:553–556.
- Chenna R, Sugawara H, Koide T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Research* 2003;31:3497–3500. [PubMed: 12824352]
- Couch BC, Fudal I, Lebrun M-H, Tharreau D, Valent B, van Kim P, Nottéghem J-L, Kohn LM. Origins of host-specific populations of the blast pathogen, *Magnaporthe oryzae*, in crop domestication with subsequent expansion of pandemic clones on rice and weeds of rice. *Genetics* 2005;170:613–630. [PubMed: 15802503]
- Couch BC, Kohn LM. A multilocus gene genealogy concordant with host preference indicates segregation of a new species, *Magnaporthe oryzae*, from *M. grisea*. *Mycologia* 2002;94:683–93.
- Cruse M, Telerant R, Gallagher T, Lee T, Taylor JW. Cryptic species in *Stachybotrys chartarum*. *Mycologia* 2002;94:814–822.
- Cruz R, Carballo M, Conde-Padin P, Rolan-Alvarez E. Testing alternative models for sexual isolation in natural populations of *Littorina saxatilis*: indirect support for by-product ecological speciation? *Journal of Evolutionary Biology* 2004;17:288–293. [PubMed: 15009262]
- Dabkevicius Z, Mikaliunaite R. The effect of fungicidal seed treatments on germination of rye ergot (*Claviceps purpurea* (Fr.) Tul.) sclerotia and on ascocarp formation. *Crop Protection* 2006;25:677–683.
- Daehler CC, Anttila CK, Ayres DR, Strong DR, Bailey JP. Evolution of a new ecotype of *Spartina alterniflora* (Poaceae) in San Francisco Bay, California, USA. *American Journal of Botany* 1999;86:543–546. [PubMed: 10205074]
- Després L, Gielly L, Redoutet B, Tabarlet P. Using AFLP to resolve phylogenetic relationships in a morphologically diversified plant species complex when nuclear and chloroplast sequences fail to reveal variability. *Molecular Phylogenetics and Evolution* 2002;27:185–196.
- Dettman JR, Jacobson DJ, Turner E, Pringle A, Taylor JW. Reproductive isolation and phylogenetic divergence in *Neurospora*: comparing methods of species recognition in a model eukaryote. *Evolution* 2003;57:2721–2741. [PubMed: 14761052]
- Douhan GW, Rizzo DM. Host-parasite relationships among bolete infecting *Hypomyces* species. *Mycological Research* 2003;107:1342–1349. [PubMed: 15000236]
- Douhan GW, Rizzo DM. Phylogenetic divergence in a local population of the ectomycorrhizal fungus *Cenococcum geophilum*. *New Phytologist* 2005;166:263–271. [PubMed: 15760369]
- Duncan RA, Sullivan R, Alderman SC, Spatafora JW, White JF. *Claviceps purpurea* var. *spartinae* var. nov.: an ergot adapted to the aquatic environment. *Mycotaxon* 2002;81:11–25.
- Excoffier L. Patterns of DNA sequence diversity and genetic structure after a range expansion: lessons from the infinite-island model. *Molecular Ecology* 2004;13(4):853–864. [PubMed: 15012760]
- Felsenstein, J. *Inferring Phylogenies*. Sinauer Associates; Sunderland MA: 2004. p. 664
- Fine PVA, Daly DC, Munoz GV, Mesones I, Cameron KM. The contribution of edaphic heterogeneity to the evolution and diversity of Burseraceae trees in the western Amazon. *Evolution* 2005;59:1464–1478. [PubMed: 16153032]
- Fisher AJ, DiTomaso JM, Gordon TR. Conidial morphology and ecological characteristics as diagnostic tools for identifying *Claviceps purpurea* from salt-marsh habitats. *Canadian Journal of Plant Pathology* 2005a;27:389–395.

- Fisher AJ, DiTomaso JM, Gordon TR. Sub-specific groups of *Claviceps purpurea* associated with grass species in Willapa Bay, Washington, and the prospects for biological control of invasive *Spartina alterniflora*. *Biological Control* 2005b;34:170–179.
- Fisher AJ, Gordon TR, DiTomaso JM. Geographic distribution and diversity in *Claviceps purpurea* (Fr.) Tul from salt marsh habitats and characterization of Pacific Coast populations. *Mycological Research* 2005c;109:439–446. [PubMed: 15912931]
- Gardes M, Bruns TD. ITS primers with enhanced specificity for Basidiomycetes -application to the identification of mycorrhizae and rusts. *Molecular Ecology* 1993;2:113–118. [PubMed: 8180733]
- Geiser DM, Pitt JI, Taylor JW. Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:388–393. [PubMed: 9419385]
- Hall MC, Willis JH. Divergent selection on flowering time contributes to local adaptation in *Mimulus guttatus* populations. *Evolution* 2006;60:2466–2477. [PubMed: 17263109]
- Hardin G. The competitive exclusion principle. *Science* 1960;131:1292–1297. [PubMed: 14399717]
- Harrington, TC.; Rizzo, DM. *Defining species in the fungi*. Academic Publishers; London: 1999.
- Hart C. Forged in St. Anthony's Fire: drugs for migraine. *Modern drug discovery* 1999;2:20–31.
- Hatfield T, Schluter D. Ecological speciation in sticklebacks: environment dependent hybrid fitness. *Evolution* 1999;53:866–873.
- Horton TR. Molecular approaches to ectomycorrhizal diversity studies: variation in ITS at a local scale. *Plant and Soil* 2002;244:29–39.
- Hudler, GW. *Magical Mushrooms, Mischievous Molds*. Princeton University Press; 1998.
- Kass RE, Raftery AE. Bayes factors. *Journal of the American Statistical Association* 1995;90(430):773–795.
- Kardolus JP, van Eck HJ, van den Berg RG. The potential of AFLPs in biosystematics: a first application in *Solanum* taxonomy (*Solanaceae*). *Plant Systematics and Evolution* 1998;210:87–103.
- Kingman JF. Origins of the coalescent. 1974–1982. *Genetics* 2000;156(4):1461–1463. [PubMed: 11102348]
- Kohn LM. Mechanisms of fungal speciation. *Annual Review of Phytopathology* 2005;43:279–308.
- Koufopanou V, Burt A, Szaro T, Taylor JW. Gene genealogies, cryptic species, and molecular evolution in the human pathogen *Coccidioides immitis* and relatives (Ascomycota, Onygenales). *Molecular Biology and Evolution* 2001;18:1246–1258. [PubMed: 11420364]
- Kroken S, Taylor JW. A gene genealogical approach to recognize phylogenetic species boundaries in the lichenized fungus *Letharia*. *Mycologia* 2001;93:38–53.
- Kruuk LEB, Gilchrist JS. Mechanisms maintaining species differentiation: predator mediated selection in a *Bombina* hybrid zone. *Proceedings of the Royal Society of London B* 1997;264:105–110.
- Lamont BB, He T, Enright NJ, Krauss SL, Miller BP. Anthropogenic disturbance promotes hybridization between *Banksia* species by altering their biology. *Journal of Evolutionary Biology* 2003;16:551–557. [PubMed: 14632219]
- Maddison, DR.; Maddison, WP. *MacClade 4: Analysis of phylogeny and character evolution*. Version 4.03. Sinauer Associates; Sunderland, Massachusetts: 2001.
- Mitsch, WJ.; Gosselink, JG. *Wetlands*. John Wiley & Sons, Inc; New York: 1993.
- Moyersoen B, Beever RE, Martin F. Genetic diversity of *Pisolithus* in New Zealand indicates multiple long-distance dispersal from Australia. *New Phytologist* 2003;160:569–579.
- Muths D, Davoult D, Gentil F, Jollivet D. Incomplete cryptic speciation between intertidal and subtidal morphs of *Acrocnida brachiata* (Echinodermata: Ophiuroidea) in the Northeast Atlantic. *Molecular Ecology* 2006;15:3303–3318. [PubMed: 16968272]
- Myllys L, Lohtander K, Tehler A. Beta-tubulin, ITS and group I intron sequences challenge the species pair concept in *Physcia aipolia* and *P. caesia*. *Mycologia* 2001;93:335–343.
- Nagy ES. Selection for native characters in hybrids between two locally adapted plant subspecies. *Evolution* 1997;51:1469–1480.
- O'Donnell K, Kistler HC, Tack BK, Casper HH. Gene genealogies reveal global phylogeographic structure and reproductive isolation among lineages of *Fusarium graminearum*, the fungus causing wheat scab. *Proceedings of the National Academy of Sciences* 2000;97:7905–7910.

- Ogden R, Thorpe RS. Molecular evidence for ecological speciation in tropical habitats. *Proceedings of the National Academy of Sciences* 2002;99:13612–13615.
- Pazoutová S, Olsovska J, Linka M, Kolínská R, Flieger M. Chemoraces and habitat specialization of *Claviceps purpurea* populations. *Applied and Environmental Microbiology* 2000;66:5419–5425. [PubMed: 11097923]
- Pazoutová S, Raybould AF, Honzátko A, Kolínská R. Specialized populations of *Claviceps purpurea* from salt marsh *Spartina* species. *Mycological Research* 2002;106:210–214.
- Pazoutová, S. Evolutionary strategy of *Claviceps*. In: White, JF.; Bacon, CW.; Hywel-Jones, NL.; Spatafora, JW., editors. *Clavicipitalean Fungi: Evolutionary Biology, Chemistry, Biocontrol and Cultural Impacts*. Marcel Dekker; New York, Basel: 2003. p. 329-354.
- Rehacek, Z.; Sajdl, P. *Ergot alkaloids: Chemistry, Biological Effects, Biotechnology*. New York: Elsevier Science publishing; 1993.
- Richmond JQ, Reeder TW. Evidence for parallel ecological speciation in scincid lizards of the *Eumeces skiltonianus* species group (Squamata: Scincidae). *Evolution* 2002;56:1498–1513. [PubMed: 12206249]
- Rieseberg LH. Crossing relationships among ancient and experimental sunflower hybrid lineages. *Evolution* 2000;54:859–865. [PubMed: 10937259]
- Rocha LA, Robertson DR, Roman J, Bowen BW. Ecological speciation in tropical reef fishes. *Proceedings of the Royal Society of London B* 2005;272:573–579.
- Rosenblum EB. Convergent evolution and divergent selection: Lizards at the White Sands ecotone. *American Naturalist* 2006;167:1–15.
- Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 2003;19:2496–2. [PubMed: 14668244]
- Rundle HD, Nosil P. Ecological speciation. *Ecology Letters* 2005;8:336–352.
- Scheffer J, Tudzynski P. In vitro pathogenicity assay for the ergot fungus *Claviceps purpurea*. *Mycological Research* 2006;110:465–470. [PubMed: 16483754]
- Scheffer J, Chen CB, Heidrich P, Dickman MB, Tudzynski P. A CDC42 Homologue in *Claviceps purpurea* is involved in vegetative differentiation and is essential for pathogenicity. *Eukaryotic Cell* 2005;4:1228–1238. [PubMed: 16002649]
- Seberg, OG.; Peterson, G. Constructing phylogenies from discrete data-parsimony methods. In: Karp, A.; Isaac, PG.; Ingram, DS., editors. *Molecular Tools for Screening Biodiversity*. Chapman & Hall; London: 1998. p. 344-355.
- Shykoff JA, Meyhöfer A, Bucheli E. Genetic isolation among host races of the anther smut fungus *Microbotryum violaceum* on three host plant species. *International Journal of Plant Science* 1999;160:907–916.
- Skovgaard K, Bodker L, Rosendahl S. Population structure and pathogenicity of members of the *Fusarium oxysporum* complex isolated from soil and root necrosis of pea (*Pisum sativum* L.). *FEMS Microbiology and Ecology* 2002;42:367–374.
- Stager R. Beitrag zur Verbreitungsbiologie der *Claviceps*-Sklerotien. *Zentralblatt Fur Bakteriologie Parasitenkunde Infefektionskrankheiten Und Hygiene* 1922;56:329–339.
- Stensrud O, Hywel-Jones NL, Schumacher T. Towards a phylogenetic classification of *Cordyceps*: ITS rDNA sequence data confirm divergent lineages and parafyly. *Mycological Research* 2005;109:41–56. [PubMed: 15736862]
- Swofford, DL. *PAUP* Phylogenetic Analysis Using Parsimony (* and other methods)*. Version 4.0b10. Sunderland, Massachusetts (USA): Sinauer Associates, Inc; 2002.
- Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fisher MC. Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* 2000;31:21–32. [PubMed: 11118132]
- Tooley PW, Goley ED, Carras MM, Frederick RD, Weber EL, Kuldauc GA. Characterization of *Claviceps* species pathogenic on sorghum by sequence analysis of the β -tubulin gene intron 3 region and EF-1a gene intron 4. *Mycologia* 2001;93:541–551.
- Tudzynski P, Scheffer J. *Claviceps purpurea*: molecular aspects of a unique pathogenic lifestyle. *Molecular Plant Pathology* 2004;5:377–388.

- Via S. Sympatric speciation in animals: the ugly duckling grows up. *Trends in Ecology and Evolution* 2001;16:381–390. [PubMed: 11403871]
- Via S, Bouck AC, Skillman S. Reproductive isolation between divergent races of pea aphids on two hosts. II. Selection against migrants and hybrids in the parental environments. *Evolution* 2000;54:1626–1637. [PubMed: 11108590]
- Wang H, McArthur ED, Sanderson SC, Graham JH, Freeman DC. Narrow hybrid zone between two subspecies of big sagebrush (*Artemisia tridentata*: Asteraceae). IV. Reciprocal transplant experiments. *Evolution* 1997;51:95–102.
- White, TJ.; Bruns, TD.; Lee, S.; Taylor, J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, MA.; Gelfand, DH.; Sninsky, JJ.; White, TJ., editors. *PCR Protocols: A guide to methods and applications*. San Diego, USA: Academic Press; 1990.
- Wu QX, Mueller GM, Lutzoni FM, Huang YQ, Guo SY. Phylogenetic and biogeographic relationships of eastern Asian and eastern North American disjunct *Suillus* species (Fungi) as inferred from ITS sequences of nuclear ribosomal RNA ITS sequences. *Molecular Phylogenetics and Evolution* 2000;17:37–47. [PubMed: 11020303]




Fig. 1. Maximum parsimony analyses of three individual loci (ITS, Beta-tubulin, Ras) and the combined data set from G1, G2, G3, isolates of *Claviceps purpurea*. Bootstrap support of 50% and above are indicated above nodes based on 1,000 replicates. Sequences have been deposited in GenBank under accession numbers (XXXXX-XXXXXX).

Table 1*Claviceps purpurea* isolates used in this study.

Code	Group	Location	Year collected	Host
165	G1	Zubri, Czech Republic	1994	<i>Poa pratensis</i>
204	G1	Lauderdale, Alabama	1996	<i>Festuca arundinaceae</i>
428	G1	Hohenheim, Germany	?	<i>Secale cereale</i>
NGE1	G1	Newfoundland, Canada	2001	<i>Leymus mollis</i>
WFA	G1	Nahcotta, Washington	2002	<i>Festuca arundinaceae</i>
WLS	G1	Nahcotta, Washington	2002	<i>Lolium spp.</i>
236	G2	Vlei Pole u Bousova, Czech Republic	?	<i>Molinia coerulea</i>
434	G2	Phillipsreuth, Germany	1998	<i>Dactylis spp</i>
WAB-1	G2	Long Beach, Washington	2002	<i>Ammophila breviligulata</i>
WCN2	G2	Willapa Bay, Washington	2002	<i>Calamagrostis nutkaensis</i>
WDG	G2	Leadbetter State Park, Washington	2002	<i>Dactylis glomerata</i>
WDS	G2	Willapa River, Washington	2002	<i>Deschampsia caespitosa</i>
WHS	G2	Long Beach, Washington	2002	<i>Holcus lanatus</i>
ARG1	G3	Argentina Celpa Marsh, Argentina	2002	<i>Spartina densiflora</i>
CDE1	G3	Point Reyes, CA	2002	<i>Spartina foliosa</i>
CMD1	G3	MacDoel, CA	2002	<i>Secale cereale</i>
CPE10	G3	Palo Alto, CA	2001	<i>Spartina foliosa</i>
FSA1	G3	St. Augustine, Florida	2000	<i>Spartina alterniflora</i>
IRE12	G3	Dublin, Ireland	2001	<i>Spartina anglica</i>
RH2	G3	Rhode Island	2001	<i>Spartina alterniflora</i>
WDI-1	G3	Willapa River, Washington	2002	<i>Distichlous spicata</i>

Table 2

Combined estimates over three loci of mutation scaled population size Θ and mutation scaled migration rates M . The estimates were found using the Bayesian inference module in MIGRATE-N.

Model	Parameter	Mode	Median	95% Credibility interval
3-population	Θ_{G1}	0.00062	0.001	0.00008 – 0.00356
	Θ_{G2}	0.00049	0.00087	0.00008 – 0.00302
	Θ_{G3}	0.00353	0.00478	0.00137 – 0.00923
	$M_{G2 \text{ to } G1}$	543	868	25 – 2370
	$M_{G3 \text{ to } G1}$	508	848	5 – 2405
	$M_{G1 \text{ to } G2}$	663	948	30 – 2530
	$M_{G3 \text{ to } G2}$	688	982	10 – 2695
	$M_{G1 \text{ to } G3}$	333	528	0 – 1490
	$M_{G2 \text{ to } G3}$	283	548	5 – 1560
	1-population	$\Theta_{G1+G2+G3}$	0.01125	0.01185

DNA divergence between G1, G2, and G3 *Claviceps purpurea* isolates in the three analyzed loci and the combined data set measured as number of segregating sites, fixed and shared polymorphic sites, average number of nucleotide differences (k), average number of nucleotide substitutions per site between habitat types (π), and population differentiation values between habitat types (Fst).

Table 3

	No. of polymorphic sites	Fixed differences	No. of polymorphic compared to monomorphic*	No. of Polymorphic compared to monomorphic [†]	Shared differences	k	π	Fst
Beta-tubulin								
G1-G2	15	0	10	3	2	6.385	0.0152	0.8216
G2-G3	9	1	2	6	0	3.121	0.0083	0.6875
G1-G3	13	7	4	2	0	5.747	0.0137	0.9034
ITS								
G1-G2	11	5	1	6	0	4.121	0.0072	0.8369
G2-G3	5	3	1	1	0	2.121	0.0037	0.8800
G1-G3	9	3	1	6	0	3.495	0.0061	0.7719
Ras								
G1-G2	7	4	1	2	0	3.648	0.0134	0.9091
G2-G3	4	0	2	3	0	1.802	0.0065	0.5172
G1-G3	7	5	0	2	1	3.692	0.0135	0.8600
Combined								
G1-G2	33	14	11	8	0	14.08	0.0115	0.8504
G2-G3	17	4	6	8	0	6.516	0.0053	0.7042
G1-G3	29	15	10	5	1	12.93	0.0102	0.8576

* Number of sites that are polymorphic in the first species compared and monomorphic in the second species

[†] Number of sites that are polymorphic in the second species compared and monomorphic in the first species