

ERRATA

cis-Dominant Defect in Activation of Adenovirus Type 5 E1b Early RNA Synthesis

CHRISTOPHER L. PARKS AND DAVID J. SPECTOR

*Department of Microbiology and Immunology, Pennsylvania State University, College of Medicine,
Hershey, Pennsylvania 17033*

Volume 64, no. 6, p. 2783: Figure 2 should appear as shown below.




FIG. 2

A Herpes Simplex Virus Type 1 Latency-Associated Transcript Mutant Reactivates with Normal Kinetics from Latent Infection

TIMOTHY M. BLOCK, JORDAN G. SPIVACK, ISRAEL STEINER, SATISH DESHMANE,
MICHAEL T. McINTOSH, RONALD P. LIRETTE, AND NIGEL W. FRASER

*Thomas Jefferson University Medical School, Philadelphia, Pennsylvania 19107, and
The Wistar Institute, Philadelphia, Pennsylvania 19104*

Volume 64, no. 7, p. 3423: The first three sentences in column 1 should read: "Ho and Mocarski (7) have also described an HSV-1 mutant, RH142, that has a β -galactosidase gene insertion at nucleotide +136 of the LAT RNA that reactivates normally. They were unable to detect transcripts from this region during latency. Since β -galactosidase activity was present, RNA must have been. . . ."