

Deletion of the mouse *RegIIIβ* (*Reg2*) gene disrupts ciliary neurotrophic factor signaling and delays myelination of mouse cranial motor neurons

L. A. Tebar*, S. M. Géranton†, C. Parsons-Perez†, A. S. Fisher†, R. Bayne‡, A. J. H. Smith‡, M. Turmaine†, S. Perez-Luz§, A. Sheasby†, C. De Felipe¶, C. Ruff||, G. Raivich||, and S. P. Hunt†***

*National Centre for Cancer Research, Melchor Fernández Almagro, 3. E-28029 Madrid, Spain; †Department of Anatomy and Developmental Biology, University College London, Medawar Building, Gower Street, WC1E 6BT London, United Kingdom; ‡Gene Targeting Laboratory, Institute for Stem Cell Research, University of Edinburgh, The King's Buildings, West Mains Road, Edinburgh EH9 3JQ, Scotland; §Department of Molecular Biology, Universidad Autónoma de Madrid Cantoblanco, 28049 Madrid, Spain; ¶Consejo Superior de Investigaciones Científicas, Universidad Miguel Hernández, Campus de San Juan, Sant Joan d'Alacanti. Nacional 33203550, Alicante, Spain; and ||Perinatal Brain Repair Group, Departments of Obstetrics and Gynaecology and Anatomy, University College London, 86-96 Chenies Mews, London WC1E 6HX, United Kingdom

Edited by Tomas Hökfelt, Karolinska Institutet, Stockholm, Sweden, and approved June 16, 2008 (received for review January 2, 2008)

A large number of cytokines and growth factors support the development and subsequent maintenance of postnatal motor neurons. *RegIIIβ*, also known as *Reg2* in rat and *HIP/PAP1* in humans, is a member of a family of growth factors found in many areas of the body and previously shown to play an important role in both the development and regeneration of subsets of motor neurons. It has been suggested that *RegIIIβ* expressed by motor neurons is both an obligatory intermediate in the downstream signaling of the leukemia inhibitory factor/ciliary neurotrophic factor (CNTF) family of cytokines, maintaining the integrity of motor neurons during development, as well as a powerful influence on Schwann cell growth during regeneration of the peripheral nerve. Here we report that in mice with a deletion of the *RegIIIβ* gene, motor neuron survival was unaffected up to 28 weeks after birth. However, there was no CNTF-mediated rescue of neonatal facial motor neurons after axotomy in KO animals when compared with wild-type. In mice, *RegIIIβ* positive motor neurons are concentrated in cranial motor nuclei that are involved in the patterning of swallowing and suckling. We found that suckling was impaired in *RegIIIβ* KO mice and correlated this with a significant delay in myelination of the hypoglossal nerve. In summary, we propose that *RegIIIβ* has an important role to play in the developmental fine-tuning of neonatal motor behaviors mediating the response to peripherally derived cytokines and growth factors and regulating the myelination of motor axons.

Schwann cells | suckling | hypoglossal nerve | LIF

A large number of neurotrophic factors and cytokines have been shown to sustain developing motor neurons and to encourage the survival of postnatal motor neurons after damage (1–6). Rat *Reg2* [also known as *RegIIIβ* in mouse and *HIP/PAP1* in humans (7)] has been reported to play an important role in both the support of motor neurons during development and in the process of axon regeneration through axon–Schwann cell signaling in the adult peripheral nervous system (8, 9). *Reg2* is a member of a large family of over 17 related genes divided into four subtypes (types I, II, III, and IV) based on the primary structures of the encoded proteins of the genes (10–13). *Reg2* is the equivalent of mouse *RegIIIβ* gene, which share 90% homology at both the nucleotide and protein level. First identified as a transcript up-regulated in pancreatitis, *Reg2*, a secreted protein (relative M_r 16,000) found in many sites throughout the body, was shown to have an anti-apoptotic action on pancreatic cell lines (14). *Reg2* also promotes the growth of epithelial intestinal cells, whereas loss of *Reg2/RegIIIβ* delayed liver regeneration (10, 15–18).

Two roles have been proposed for *Reg2* in the nervous system. First, *in vivo* studies have suggested that *Reg2* is released from

damaged motor and sensory neurons and has a proregenerative function (9, 19, 20). Secondly, *in vitro* data have cast *Reg2* as a neurotrophic factor for motor neurons acting as an obligatory intermediate for the ciliary neurotrophic factor (CNTF)/leukemia inhibitory factor (LIF) family of cytokines (8). In the adult rat, *Reg2* is massively up-regulated in motor neurons and some sensory neurons after nerve crush (19) and is rapidly transported to the lesion site, where it is thought to be secreted and act on Schwann cells. *In vitro*, *Reg2* has a mitogenic effect on Schwann cells and inhibition of *Reg2* with a neutralizing antibody retards the progress of regeneration. Taken together, it seemed likely that the Schwann cell response at the point of axotomy was potentiated by release of *Reg2* from damaged axons (9).

Here, we describe the effects of targeted ablation of the *RegIIIβ* gene in mice and show a developmental role of *RegIIIβ* in axon–Schwann cell signaling. We also report that *RegIIIβ* is required to promote the response of subsets of motor neurons to CNTF but not for motor neuron survival.

Results

Generation of *RegIIIβ*-Deficient Mice. KO mice were homozygous for a targeted disruption of the *RegIIIβ* gene locus created in ES cells, in which a region from exons 2 to 5 was deleted and replaced by an IRES-*Tau-LacZ-loxP/MC1neopA/loxP* reporter/selection cassette, thus resulting in a null allele. Mice homozygous for the *RegIIIβ*-null allele were phenotypically indistinguishable from wild-type or heterozygous littermates. No embryonic lethality or significant developmental defects were observed in *RegIIIβ*^{-/-} animals. Adult *RegIIIβ*^{-/-} mice of both sexes were fertile, and litter size was normal. We compared the expression of *RegIIIβ* between wild-type and KO animals by using quantitative RT-PCR (RT-qPCR) and found that, as expected, the expression of *RegIIIβ* mRNA dropped from 100 ± 25.2% (wild type) to 0.08 ± 0.02% (KO) ($n = 6$ per group). Immunohistochemistry also showed a complete lack of *RegIIIβ* protein-like immunoreactivity in postnatal KO mice (Fig. 1). To check for potential compensatory up-regulation, we also mea-

Author contributions: A.J.H.S., C.D.F., and S.P.H. designed research; L.A.T., S.M.G., C.P.-P., A.S.F., R.B., A.J.H.S., M.T., S.P.-L., A.S., C.D.F., C.R., G.R., and S.P.H. performed research; L.A.T., S.M.G., C.P.-P., A.S.F., A.J.H.S., S.P.-L., and S.P.H. analyzed data; and S.M.G. and S.P.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

**To whom correspondence should be addressed. E-mail: hunt@ucl.ac.uk.

This article contains supporting information online at www.pnas.org/cgi/content/full/0711978105/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. RegIII β immunoreactivity is found in subsets of facial motor neurons in P5 wild-type mice but was not found in the postnatal KO mouse central nervous system. (A) Facial motor nucleus (P5) of a wild-type mouse. (B) Absence of staining in KO facial nucleus. (Scale bar, 120 μ m.)

sured the expression of *RegIII α* and *RegIII γ* : There was no difference in *RegIII γ* expression ($100 \pm 67\%$ in wild type vs. $62 \pm 43\%$ in KO), but we found a substantial elevation of *RegIII α* in the brainstem of KO mice ($100 \pm 26\%$ in wild type vs. $667 \pm 181\%$ in KO).

RegIII β Labeling of Neuronal Populations Is Restricted and Developmentally Regulated. In wild-type embryos, RegIII β proteins are first detected in subsets of motor neurons at embryonic day (E)13–E14 in cervical spinal cord, as previously described in rat (8), with expression peaking at E18.5 (2). Within the brainstem at postnatal day (P)1–P5, subsets of neurons along the medial half of the motor nucleus of the trigeminal and in the dorsal and medial regions of the facial nucleus were immunoreactive for RegIII β (Fig. 2). The heaviest concentrations of RegIII β positive neurons (up to 50% at P1) were found throughout the caudal half of the hypoglossal nucleus (XII) and nucleus ambiguus notably at day 1, but staining intensity reached maximal levels by




Fig. 2. RegIII β immunoreactivity is found in subsets of cranial and spinal motor nuclei up to P12. (A) Facial motor nucleus with labeled axons (arrow) streaming into the facial nerve. (B–D) RegIII β -positive neurons within the nucleus ambiguus (B), the hypoglossal nuclei (C), and the motor nucleus of the trigeminal (D). All specimens were from P5 mice. [Scale bars: 100 μ m (A, B, and D); 150 μ m (C).]




Fig. 3. β -Galactosidase activity is present postnatally in subsets of motor neurons in the RegIII β KO mouse. (A) Motor neuron in the facial motor nucleus at P3. (B) Sacral motor neurons at P1. (C) Punctate β -galactosidase staining in the facial motor nucleus at P5. [Scale bars: 20 μ m (A); 60 μ m (B); 40 μ m (C).]

P4. By P11, RegIII β staining had been lost from all areas of the brain and spinal cord.

Motor Neurons Are Not Lost After RegIII β Deletion. Two lines of evidence imply that motor neuron survival is not compromised by loss of RegIII β . First, in KO mice, β -galactosidase staining is still visible in motor neurons similar in position to those that show RegIII β expression in wild-type mice (Fig. 3). The tau-lac-Z strategy designed to transport β -galactosidase into axons met with limited success because the reaction product was only obviously visualized in motor neuron cell bodies and dendrites. These motor neurons included the spinal motor neurons and trigeminal, facial, hypoglossal, and nucleus ambiguus motor neurons up to P10. Postnatally, sacral motor neurons stained positively for β -galactosidase (Fig. 3B), but brainstem expression was restricted to small foci within the cytoplasm of motor neurons (Fig. 3C) or the occasional heavily stained motor neuron (Fig. 3A). In the spinal cord, β -galactosidase reaction product was found in only the medial column of neurons comparable in position to RegIII β immunoreactive neurons seen in wild-type mice (Fig. 3B). β -Galactosidase expression was not present beyond P10. Second, the total number of motor neurons in the facial motor nucleus of P5.5–P10.5 mice or mice 24 weeks old were not significantly different between wild-type and KO mice implying that no motor neuron cell death had resulted from the loss of RegIII β [see supporting information (SI) Fig. S1].

Lack of Effect of CNTF on Facial Motor Neuron Survival in RegIII β KO Mice. CNTF applied to the cut end of the facial nerve delays the axotomy provoked death of neonatal rat facial motor neurons (21). To examine the effects of CNTF in RegIII β KO mice, we cut the facial motor nerve unilaterally at P3.5 and then immediately applied CNTF or saline to the cut end of the nerve. Mice were then allowed to survive for 3 days. The loss of facial motor neurons after facial nerve lesion (FNL) was comparable between wild-type and KO mice (45% and 49% loss when compared with contralateral side in wild-type and KO, respectively) (Fig. 4). There was a significant effect of CNTF on wild-type mice when




Fig. 4. Application of CNTF to the cut facial nerve results in increased survival of facial motor neurons in wild-type but not KO mice counted 3 days after nerve section. The data show means \pm SEM ($n = 4-5$ in each group). **, $P < 0.001$.

compared with both CNTF-treated KO and saline-treated wild-type mice (24% and 32% savings respectively, $P < 0.001$ for both) (Fig. 4). When mice were allowed to survive for four days after FNL, cell loss was considerable but CNTF produced significant savings in the numbers of dying neurons in wild-type mice when compared with KO mice (Fig. S2).

We have previously shown (9) that section of the sciatic nerve results in down-regulation of Reg2/RegIII β in motor neurons in rat pups and up-regulation in adult rats. However, in mouse pups with facial nerve axotomy at P3.5 and perfused 2 days later, we did not detect any change in numbers of RegIII β -positive neurons in the facial motor nucleus. However, application of CNTF after facial nerve section at P3.5 in mouse pups significantly increased the number of RegIII β -positive neurons seen 2 days later to 148% compared with the contralateral side ($P < 0.0001$) (Fig. 5 and Fig. S3). Finally, whereas in adult rats, Reg2 is expressed by all facial motor neurons within 24h of axotomy, in adult mice, RegIII β is not reexpressed in facial motor neurons 1–7 days after axotomy (Fig. S4).

Loss of RegIII β Impairs Suckling Behavior. In newborn mouse pups, we noted that neurons immunoreactive for RegIII β were found




Fig. 5. CNTF increases the number of RegIII β positive neurons. (A) Application of CNTF to the cut facial nerve results in increased numbers of RegIII β -positive facial motor neurons in wild-type mice 2 days after nerve section. The data show means \pm SEM ($n = 7$ in each group). ***, $P < 0.0001$. (B and C) RegIII β -positive neurons in wild-type mice after facial nerve cut and CNTF (B) or saline (C) application. The arrow indicates the increased number of RegIII β -positive neurons induced by CNTF. [Scale bar, 150 μ m (B and C).]

to be concentrated in cranial motor neurons that have been associated with suckling and swallowing, such as the hypoglossal nucleus and nucleus ambiguus. We therefore examined the ability of RegIII β KO mice to ingest milk during a 1-h period of suckling after a 2-h period of isolation from the mother. Compared with wild-type pups, we found a significant reduction in milk/colostrum ingestion: percentage weight increase in wild-type mice, $3.1 \pm 0.2\%$; and in KO mice, 0.45 ± 0.2 ($P < 0.01$).

Loss of RegIII β Delays Myelination of the Hypoglossal Nerve. Because of the influence of Reg2 on Schwann cells (9) and the reduced efficiency of suckling described above, we used electron microscopy to analyze myelination in the hypoglossal nerve at P5 in wild-type and KO mice. Oculomotor neurons never express RegIII β , and we, therefore, used the P5 oculomotor nerve as a control. We found a significant increase in the numbers of unmyelinated nerve fibres in both the medial and lateral hypoglossal nerves in KO mice at P5 (Fig. 6A–C). One-way ANOVA revealed a significant effect of genotype on axon type in both the medial ($F_{1,14} = 24.8$; $P < 0.01$) and lateral hypoglossal nerves ($F_{1,13} = 21.5$; $P < 0.01$). There were no differences in the numbers of unmyelinated axons in the oculomotor nerves in P5 wild-type and KO mice. In the lateral hypoglossal nerve, the total numbers of axons in the RegIII β KO mice exceeded that in the wild-type animals (Fig. 6C). We therefore counted the total number of myelinated axons in the lateral and medial hypoglossal nerves at P21. We found that there were no differences between the two groups (wild type: lateral, 352.3 ± 25.8 ; medial, 980.3 ± 9.5 ; KO: lateral, 360.0 ± 16.5 ; medial, 950.3 ± 41.9 KO; $n = 3$ in each group). We concluded that excess collateralization of hypoglossal axons occurs because of delayed myelination during the early postnatal period in KO mice and is lost later by selective pruning of excess axonal branches (22, 23).

The impaired suckling phenotype, thus, may result from a disruption of myelination in mutant mice. We also examined the hypoglossal innervation of tongue musculature in wild-type and KO mice by using protein gene product (PGP) as a marker for nerve fibers and α -bungarotoxin for muscle end plates. We found no evidence of changes in innervation density or size of muscle end plates (data not shown).

Discussion

Previous research has suggested that *in vitro* rat Reg2 is a motor neuron survival factor essential for the actions of CNTF-like cytokines and a powerful Schwann cell mitogen that potentiates axonal repair and regeneration *in vivo* (8, 9). However, we show here that in mice with a genetic deletion of the RegIII β gene (the equivalent gene to Reg2 in rats), there is no increased motor neuron cell death during development. Nevertheless, we report that the efficacy of CNTF in reducing neuronal cell death was diminished in RegIII β KO mice indicating that RegIII β is important for the survival functions of CNTF. Finally, we found that myelination was delayed in subsets of hypoglossal motor neurons in KO animals and the efficiency of milk ingestion reduced.

RegIII β Is an Intermediate in the CNTF Survival Pathway. The suggestion that Reg2/RegIII β was a motor neuron neurotrophic factor and a signaling intermediary in the CNTF survival pathway stems from elegant *in vitro* studies demonstrating that purified Reg2 can act as a paracrine/autocrine neurotrophic factor for a subset of motor neurons (8). Furthermore, it has been shown that CNTF, as well as the related factors LIF, cardiotrophin 1, and oncostatin-M, rapidly induces Reg2 mRNA in some motor neurons and that released Reg2 acts in a paracrine/autocrine fashion to support motor neurons (8, 9). The CNTF/LIF family of cytokines signal through a receptor complex that includes the LIF receptor (LIFR) subunit, and it has been




Fig. 6. Increased number of unmyelinated axons in *RegIII β* KO mice. Electron micrograph of the medial hypoglossal nerve of *RegIII β* wild-type (A) or *RegIII β* KO (B) mice. There is an increase in the number of unmyelinated fibers in the lateral and medial hypoglossal motor nerve of *RegIII β* KO mice ($n = 8$ in each group). (C) No differences were observed in the oculomotor nerve ($n = 3$ in each group). The data show means \pm SEM. **, $P < 0.001$; ***, $P < 0.0001$. All nerves were studied at P5. [Scale bar, 2 μ m (A and B).]

shown previously that in *LIFR* KO mice, there is no expression of *Reg2/RegIII β* during development (9). Thus, *Reg2/RegIII β* expression in some developing motor (and sensory) neurons depends on cytokines of the LIF family acting through a receptor complex containing *LIFR*.

The absence of motor neuron cell death in KO mice, therefore, is puzzling, especially given that *RegIII β* is essential for the *in vitro* neuronal survival effects of CNTF to be manifest. The evidence for a lack of effect on motor neuron survival comes from the presence of β -galactosidase reaction product in postnatal neurons up to P10 and the normal numbers of facial motor neurons in KO mice. However, in *CNTF* KO mice, there is similarly no increased loss of motor neurons during the first weeks of life (24, 25). This, perhaps, is not surprising, given that levels of CNTF are low during development and, although Schwann cells are the richest source of CNTF in the adult peripheral nervous system, levels only rise during the first postnatal week (26–28). Knockout of the *CNTF* gene did result in motor neuron loss 28 weeks after birth (24); however, *RegIII β* mice did not show motor neuron loss in later adult stages.

Previous studies suggest that CNTF itself is not the key ligand acting at the *CNTF/LIF* receptor complex. Nishimune *et al.* (8) found that *RegIII β* expression was unimpaired or delayed in a range of KOs including *cntf*, *ct1*, and *cntf/lif* double mutants, leading them to suggest that a factor as yet unknown was key to driving the expression of *RegIII β* in motor neurons.

That CNTF-like factors are as important *in vivo* as *in vitro* and require *RegIII β* expression is implied by the observation that *RegIII β* expression increases in axotomized neurons only when CNTF is applied to the nerve stump and that CNTF has no survival effect on motor neurons in *RegIII β* KO mice. We also show that the number of *RegIII β* neurons increased in wild-type mice when CNTF was applied to the cut nerve. This implies that many motor neurons have the capacity to express *RegIII β* , but this number is restricted by availability of CNTF-like factors during development. Nevertheless, *RegIII β* may not be the only intermediary involved in CNTF-like factor signaling as only ≈ 15 – 20% of facial motor neurons express *RegIII β* , and CNTF has previously been shown to rescue $\approx 75\%$ of rat facial motor neurons from cell death when axotomized soon after birth (24). However, the substantially larger concentration of CNTF used in these experiments (5 μ g vs. 250 ng used in the present study) may well have driven the expression of *RegIII β* in many more axotomized facial motor neurons than seen here and resulted in greater levels of survival (24). Also the dynamics of *Reg2*

expression in rat are different from that of *RegIII β* in mice. *RegIII β* never reappeared in the adult mouse after section of the facial nerve (Fig. S4) or sciatic nerve (data not shown) and did not noticeably decrease 24 h after axotomy, as was observed after sciatic transection in neonatal rats. Presumably other factors may play a similar role to *RegIII β* in adult mouse motor neurons, although it is unlikely to be *RegIII α* , which although up-regulated in the KO mouse, did not potentiate CNTF-mediated survival.

Growth Factors and Myelination. We show here that myelination of a subset of hypoglossal motor neuron axons is delayed in *RegIII β* KO mice at P5.5 but normal by P21. This seems likely to be attributable to the loss of *RegIII β* because myelination of oculomotor axons in KO mice was normal and oculomotor motor neurons did not express *RegIII β* at any stages of development. This delayed myelination would be expected to disrupt the transmission of signals along the axon and may account for the reduced efficiency of suckling behavior. Delayed myelination did not, however, result in any reduction in the size or number of motor neuron end plates in target muscles of the hypoglossal such as the glossohyoid (unpublished data). However, how then does *RegIII β* promote myelination in discrete populations of motor neuron axons, and why is there such a restricted pattern of *RegIII β* expression?

Axons regulate Schwann cells during development of the peripheral nervous system (29–33). The axon signals for regulating Schwann cell differentiation are known to include both proteins encoded by the neuregulin gene (*NRG*) signaling through the *erbB* family of receptors, adhesion molecules such as L1 and N-cadherin and neurotrophic factors such as brain-derived neurotrophic factor (*BDNF*) (34–37). It has been suggested that cell adhesion molecules on the axon surface as well as signals from the extracellular matrix and neurotrophins are also required for myelination to proceed efficiently and accurately. For example, *BDNF* supports postnatal facial motor neurons after axotomy and acts through p75 NTR to inhibit Schwann cell migration and promote myelination (35, 36, 38). It seems likely that *RegIII β* released from subsets of motor neuron axons is playing much the same role as *BDNF* in driving the process of myelination. As with *NRG1* type III, it has been shown that *RegIII β* signals through the PI3-kinase pathway (8, 17), and this may represent a common pathway for axonally released *RegIII β* to influence Schwann cells. Why *RegIII β* expression is restricted to motor neurons concerned with the suckling and swallowing is unclear, but we suggest that the patterning of this

critical function in neonatal mice may be plastic and regulated by release of factors from target musculature concerned with suckling in an activity-dependent fashion (8).

Materials and Methods

All procedures complied with the United Kingdom Animals (Scientific Procedures) Act 1986.

Generation of the RegIII β -Deficient Mice by Gene Targeting. The generation of these mice has been described elsewhere (18). Complete knockout of the gene was confirmed in nervous tissue with immunocytochemistry by using antibodies generated against rat Reg2 protein, and absence of RegIII β mRNA was confirmed by using real-time PCR in brain, pancreas, and regenerating liver (16, 18).

Real-Time RT-qPCR Assay. Tissue preparation and RNA extraction were as described previously (39). RNA samples were treated with DNase I (Qiagen). Equal amounts (3 μ g) of total RNA were reverse transcribed by using random nonamers (Sigma) and SuperScript TM III RT (Invitrogen) for 1 h at 50°C in a total reaction volume of 20 μ l. cDNAs were immediately quantified by real-time PCR or kept at -20°C until further experiments. Real-time PCRs were performed with a DNA Engine (Bio-Rad) by using SYBR Green Jump Start RT-PCR master mix (Sigma) with each gene-specific primer (RegIII β forward, 5'-AAGAATATACCTCCGCACGC-3'; RegIII β reverse, 5'-CAGACAT-AGGGCAACTTACC-3'; RegIII α forward, 5'-GAAGTGCCTCTCCACGTACC-3'; RegIII α reverse, 5'-ACAAATGGTAATGTCCCATCG-3'; RegIII γ forward, 5'-GCTCCTATTGCTATGCTTGTGGT-3'; RegIII γ reverse 5'-CATGGAGGACAG-GAAGGAAGC-3'; β -actin forward, 5'-CAACGAGCGGTTCCGATG-3'; β -actin reverse, 5'-GCCACAGGATTCCATACCA-3'). One microliter of cDNA was amplified in a three-step cycling program in a final reaction volume of 25 μ l. Control cDNA samples (obtained without transcriptase) were always included, as well as samples without any cDNA template. Reactions were performed in triplicate for five to six biological replicates, and threshold cycle values were normalized to β -actin gene expression. The specificity of the products was determined by melting curve analysis. The ratio of the relative expression of target genes to β -actin was calculated by using the $2^{-\Delta\Delta C_T}$ formula.

Facial Nerve Section. Pups were cooled on wet ice (4°C), and the extent of anesthesia was determined by assessing reflex responses to tail pinch. Adult mice were anesthetized with Fluothane. The right facial nerve was transected at the stylomastoid foramen. One to 10 days later, mice were deeply anesthetized with Euthatal i.p. and transcardially perfused with 4% paraformaldehyde (PFA) in 0.1 M sodium phosphate buffer (PB) (pH 7.4) preceded by a brief wash with heparinized saline. After a 2 h postfix in PFA, tissue was moved to sucrose (30%) PB solution in preparation for freeze sectioning. In some pups, 5 μ l of CNTF (250 ng) applied to a piece of Gelfoam in 5 μ l saline, or saline alone, was applied to the cut end of the facial nerve at the time of section. Animals received only one treatment: either CNTF or saline.

Detection of β -Galactosidase Staining by X-Gal Staining. PFA-fixed tissue was washed in PBT (PBS/0.1% Tween-20) several times. After a rinse in X-Gal staining solution [5 mM K₃Fe(CN)₆, 5 mM K₄Fe(CN)₆, 1 mM MgCl₂, 0.01% sodium deoxycholate, 0.02% Triton plus X-Gal (4-chloro,5-bromo,3-indolyl-

β -galactosidase) at 1 mg/ml, suspended in PBS (pH 7.2)], tissue was transferred to fresh staining solution. Tissue was incubated at 37°C in the dark overnight. Sections were washed several times with PBT.

Cell Counting. Sections (40 μ m) were cut through the facial nucleus and all sections (the facial nucleus is \approx 700 μ m in length in the anterior-posterior direction) were mounted and stained with neutral red. All neuronal profiles in every section were counted and total counts corrected by using the Abercrombie correction (40).

Immunohistochemistry. Mice were deeply anesthetized with pentobarbitone (60 mg/kg, i.p.) and transcardially perfused with 4% PFA in 0.1 M sodium phosphate buffer (PB) (pH 7.4) preceded by a brief wash with heparinized saline. Tissue was dissected and postfixed for 2 h at 4°C and then transferred to 30% (wt/vol) sucrose in 0.1 M PB containing 0.02% (wt/vol) sodium azide. 40 μ m free-floating tissue sections were processed as previously described but for detection of RegIII β (9, 39). Hypoglossal innervation of tongue musculature in wild-type and KO mice was assessed with PGP, a marker for nerve fibres, and α -bungarotoxin, a marker for muscle end plates.

Electron Microscopy. Mouse pups or adult mice (P21) were deeply anesthetized with Nembutal and perfused with 10 ml heparinized saline followed by 20 ml 4% PFA plus 0.5% glutaraldehyde. Tissue was fixed overnight before dissection of the caudal tongue and attached hypoglossal nerves and the oculomotor nerve still attached to the optic nerve and eye cup. After being osmicated (30 min in 1% OsO₄ in 0.1 M PB), the sections were stained for 15 min in 0.1% uranyl acetate in sodium acetate buffer at 4°C, dehydrated in ethanol, cleared in propylene oxide, and embedded in Araldite. Semithin sections were cut with glass knives and stained with toluidine blue adjacent to thin sections cut with a diamond knife on an Ultracut ultramicrotome (Reichert). The sections were collected on mesh grids coated with a thin Formavar film, counterstained with lead citrate, and viewed in a JEOL 1010 electron microscope. Counts were made from four photographs of the lateral and medial nerves for each animal taken at \times 4,000 magnification. Total counts of myelinated fibers at P21 were made from photomicrographs of the lateral and medial hypoglossal nerves at \times 1,000 magnification.

Measurement of Milk/Colostrum Intake. The procedure described by Fujita and colleagues (41, 42) was followed. Briefly, pups were separated from the dam for 2 h and kept in a warm, dark chamber. The pups were then weighed and placed back with the mothers for 1 h before reweighing. Thirty to 40 pups of each genotype were used for the measurement of milk intake.

Statistical Analysis. The data are expressed as means \pm SEM. The data were analyzed by general linear model univariate or multivariate test, as appropriate, followed by Bonferroni post hoc tests or Student's *t* test, as appropriate. For all statistical analysis, statistical significance was set at $P < 0.05$.

ACKNOWLEDGMENTS. We thank Austin Smith for support and advice during the early stages of this project and Rhona Mirsky for discussion. This work was supported by the Motor Neuron Disease Society (United Kingdom) and the Medical Research Council (United Kingdom).

- Brujin LI, Miller TM, Cleveland DW (2004) Unraveling the mechanisms involved in motor neuron degeneration in ALS. *Annu Rev Neurosci* 27:723-749.
- Henderson CE, et al. (1998) Role of neurotrophic factors in motoneuron development. *J Physiol Paris* 92:279-281.
- Oppenheim RW (1996) Neurotrophic survival molecules for motoneurons: An embarrassment of riches. *Neuron* 17:195-197.
- Sendtner M, Holtmann B, Hughes RA (1996) The response of motoneurons to neurotrophins. *Neurochem Res* 21:831-841.
- Thoenen H, Hughes RA, Sendtner M (1993) Trophic support of motoneurons: Physiological, pathophysiological, and therapeutic implications. *Exp Neurol* 124:47-55.
- Wiese S, Beck M, Karch C, Sendtner M (2004) Signalling mechanisms for survival of lesioned motoneurons. *Acta Neurochir Suppl* 89:21-35.
- Narushima Y, et al. (1997) Structure, chromosomal localization and expression of mouse genes encoding type III Reg, RegIII alpha, RegIII beta, RegIII gamma. *Gene* 185:159-168.
- Nishimune H, et al. (2000) Reg-2 is a motoneuron neurotrophic factor and a signalling intermediate in the CNTF survival pathway. *Nat Cell Biol* 2:906-914.
- Livesey FJ, et al. (1997) A Schwann cell mitogen accompanying regeneration of motor neurons. *Nature* 390:614-618.
- Iovanna JL, Dagorn JC (2005) The multifunctional family of secreted proteins containing a C-type lectin-like domain linked to a short N-terminal peptide. *Biochim Biophys Acta* 1723:8-18.
- Okamoto H (1999) The Reg gene family and Reg proteins: With special attention to the regeneration of pancreatic beta-cells. *J Hepatobiliary Pancreat Surg* 6:254-262.
- Okamoto H, Takasawa S (2002) Recent advances in the Okamoto model: The CD38-cyclic ADP-ribose signal system and the regenerating gene protein (Reg)-Reg receptor system in beta-cells. *Diabetes* 51 Suppl 3:S462-S473.
- Zhang YW, Ding LS, Lai MD (2003) Reg gene family and human diseases. *World J Gastroenterol* 9:2635-2641.
- Malka D, et al. (2000) Tumor necrosis factor alpha triggers antiapoptotic mechanisms in rat pancreatic cells through pancreatitis-associated protein I activation. *Gastroenterology* 119:816-828.
- Christa L, et al. (1999) Hepatocarcinoma-intestine-pancreas/pancreatic associated protein (HIP/PAP) is expressed and secreted by proliferating ductules as well as by hepatocarcinoma and cholangiocarcinoma cells. *Am J Pathol* 155:1525-1533.
- Gironella M, et al. (2007) Experimental acute pancreatitis in PAP/HIP knock-out mice. *Gut* 56:1091-1097.
- Lieu HT, et al. (2005) HIP/PAP accelerates liver regeneration and protects against acetaminophen injury in mice. *Hepatology* 42:618-626.
- Lieu HT, et al. (2006) Reg2 inactivation increases sensitivity to Fas hepatotoxicity and delays liver regeneration post-hepatectomy in mice. *Hepatology* 44:1452-1464.
- Averill S, Davis DR, Shortland PJ, Priestley JV, Hunt SP (2002) Dynamic pattern of reg-2 expression in rat sensory neurons after peripheral nerve injury. *J Neurosci* 22:7493-7501.
- Tam J, Rosenberg L, Maysinger D (2004) INGAP peptide improves nerve function and enhances regeneration in streptozotocin-induced diabetic C57BL/6 mice. *FASEB J* 18:1767-1769.

21. Sendtner M, et al. (1992) Ciliary neurotrophic factor prevents degeneration of motor neurons in mouse mutant progressive motor neuronopathy. *Nature* 358:502–504.
22. Bishop DL, Misgeld T, Walsh MK, Gan WB, Lichtman JW (2004) Axon branch removal at developing synapses by axosome shedding. *Neuron* 44:651–661.
23. Yamamoto M, Ueda R, Takahashi K, Saigo K, Uemura T (2006) Control of axonal sprouting and dendrite branching by the Nrg-Ank complex at the neuron-glia interface. *Curr Biol* 16:1678–1683.
24. Masu Y, et al. (1993) Disruption of the CNTF gene results in motor neuron degeneration. *Nature* 365:27–32.
25. Sendtner M, Arakawa Y, Stockli KA, Kreutzberg GW, Thoenen H (1991) Effect of ciliary neurotrophic factor (CNTF) on motoneuron survival. *J Cell Sci Suppl* 15:103–109.
26. Holtmann B, et al. (2005) Triple knock-out of CNTF, LIF, and CT-1 defines cooperative and distinct roles of these neurotrophic factors for motoneuron maintenance and function. *J Neurosci* 25:1778–1787.
27. Schweizer U, et al. (2002) Conditional gene ablation of Stat3 reveals differential signaling requirements for survival of motoneurons during development and after nerve injury in the adult. *J Cell Biol* 156:287–297.
28. Stockli KA, et al. (1991) Regional distribution, developmental changes, and cellular localization of CNTF-mRNA and protein in the rat brain. *J Cell Biol* 115:447–459.
29. French-Constant C, Colognato H, Franklin RJ (2004) Neuroscience. The mysteries of myelin unwrapped. *Science* 304:688–689.
30. Jessen KR, Mirsky R (2002) Signals that determine Schwann cell identity. *J Anat* 200:367–376.
31. Jessen KR, Mirsky R (2005) The origin and development of glial cells in peripheral nerves. *Nat Rev Neurosci* 6:671–682.
32. Taveggia C, et al. (2005) Neuregulin-1 type III determines the ensheathment fate of axons. *Neuron* 47:681–694.
33. Taveggia C, Salzer JL (2007) PARSing the events of myelination. *Nat Neurosci* 10:17–18.
34. Bunge RP, Bunge MB, Eldridge CF (1986) Linkage between axonal ensheathment and basal lamina production by Schwann cells. *Annu Rev Neurosci* 9:305–328.
35. Chan JR, Cosgaya JM, Wu YJ, Shooter EM (2001) Neurotrophins are key mediators of the myelination program in the peripheral nervous system. *Proc Natl Acad Sci USA* 98:14661–14668.
36. Tolwani RJ, et al. (2004) BDNF overexpression produces a long-term increase in myelin formation in the peripheral nervous system. *J Neurosci Res* 77:662–669.
37. Zhang JY, Luo XG, Xian CJ, Liu ZH, Zhou XF (2000) Endogenous BDNF is required for myelination and regeneration of injured sciatic nerve in rodents. *Eur J Neurosci* 12:4171–4180.
38. Yan Q, Elliott J, Snider WD (1992) Brain-derived neurotrophic factor rescues spinal motor neurons from axotomy-induced cell death. *Nature* 360:753–755.
39. Geranton SM, Morenilla-Palao C, Hunt SP (2007) A role for transcriptional repressor methyl-CpG-binding protein 2 and plasticity-related gene serum- and glucocorticoid-inducible kinase 1 in the induction of inflammatory pain states. *J Neurosci* 27:6163–6173.
40. Voegelzang MG, et al. (2001) Alpha4 integrin is expressed during peripheral nerve regeneration and enhances neurite outgrowth. *J Neurosci* 21:6732–6744.
41. Fujita K, et al. (2006) Effects of hypoglossal and facial nerve injuries on milk-suckling. *Int J Dev Neurosci* 24:29–34.
42. Fukuyama T, et al. (2006) Differential effects of hypoglossal and facial nerve injuries on survival and growth of rats at different developmental stages. *Int J Dev Neurosci* 24:307–317.