



Published in final edited form as:

*J Am Chem Soc.* 2006 December 20; 128(50): 16038–16039. doi:10.1021/ja066452e.

## Vinyldiazolactone as a Vinylcarbene Precursor: Highly Selective C-H Insertion and Cyclopropanation Reactions

Darren Bykowski, Kou-Hui Wu, and Michael P. Doyle

University of Maryland College Park, MD, Department of Chemistry and Biochemistry 20742, [mdoyle3@umd.edu](mailto:mdoyle3@umd.edu)

### Abstract

Vinylcarbenes are versatile synthetic intermediates, capable of asymmetric cyclopropanation and insertion into unactivated CH bonds. The vinyldiazolactone precursor to the metal vinylcarbene possesses superior stability in comparison to previously known vinyldiazoacetates and, for the first time, allows the use of *Z*-vinylcarbenes in asymmetric C-H insertion and cyclopropanation reactions. Dirhodium azetidinone ligated compounds are optimal catalysts. Cyclopropanation coupled with the Cope rearrangement provides access to *trans*-3,4-disubstituted hydroazulenes.

Rhodium(II)-stabilized vinylcarbenes are valuable synthetic intermediates capable of undergoing a spectrum of transformations in a highly selective fashion.<sup>1</sup> Their versatility is illustrated in applications directed to the construction of complex molecular architectures.<sup>2</sup> However, limitations of this catalytic methodology have become apparent. Vinyldiazoacetates undergo a spontaneous [1,5]-cyclization to yield pyrazoles,<sup>3</sup> and this transformation has limited their applications with catalysts that are less reactive towards diazo decomposition than dirhodium tetracarboxylates. In addition, although *E*-styrylcarbenes readily undergo intermolecular addition and insertion reactions, the *Z*-vinylcarbene isomer produces the intramolecular aromatic substitution product exclusively,<sup>4</sup> and asymmetric intermolecular reactions of *Z*-vinylcarbenes have been observed only when the *Z*-substituent is alkyl.<sup>5</sup> In the course of our research with vinyldiazoacetates,<sup>2c,d</sup> we became intrigued by the possibility of utilizing vinyldiazolactones as carbene precursors. Restriction of the vinyldiazo moiety within a ring was expected to prevent the undesirable intramolecular processes that occur with acyclic vinyldiazoacetates and vinylcarbene intermediates. We report here the first synthesis of a vinyldiazolactone and its unexpected suitability for asymmetric induction in intermolecular insertion and addition reactions with a chiral dirhodium carboxamidate.

The synthesis of vinyldiazolactone **1** was accomplished in one step from commercially available 5,6-dihydro-2*H*-pyran-2-one using direct diazo transfer conditions that are reported in the preparation of select vinyldiazoacetates.<sup>6</sup> However, unlike vinyldiazoacetates, lactone **1** exhibited a long shelf-life. Throughout this study, **1** was stored for several weeks at 4° C and used without further purification.

Perhaps the most remarkable property of rhodium(II)-stabilized vinylcarbenes is their applicability toward intermolecular C-H insertion reactions.<sup>7</sup> Carbenes generated from vinyldiazoacetates and aryldiazoacetates in the asymmetric environment of Rh<sub>2</sub>(DOSP)<sub>4</sub> (Figure 1) and related structures are unique in their ability to undergo insertion into unactivated C-H bonds with synthetically viable yields and enantioselectivities. To investigate the utility of **1** in intermolecular C-H insertion reactions, asymmetric dirhodium(II) carboxylate and carboxamidate catalysts were screened in the reaction of **1** with 1,4-cyclohexadiene (Table 1).


Surprisingly, Rh<sub>2</sub>(*S*-DOSP)<sub>4</sub>,<sup>8</sup> ordinarily the optimal catalyst for enantioselective C-H insertion reactions of vinyldiazoacetates,<sup>7</sup> was not suitable in this application. Not only was

the enantiomeric excess of C-H insertion product **2a** negligible, but the propensity of this catalyst for C-H insertion relative to cyclopropanation was the lowest of all catalysts evaluated (Table 1, entry 2). The carboxamidate catalyst  $\text{Rh}_2(\text{S-MEPY})_4$ ,<sup>9</sup> which has seen some use in C-H insertion reactions but was found to be unsuitable for reactions with vinyl diazoacetates, **3** provided a greater chemoselectivity for **2a** than  $\text{Rh}_2(\text{S-DOSP})_4$ , but enantiocontrol with this catalyst was low (Table 1, entry 3). With the azetidinone ligated catalyst  $\text{Rh}_2(\text{S-MEAZ})_4$  (Table 1, entry 4),<sup>10</sup> however, promising selectivities were obtained. Optimization of the azetidinone catalyst structure led to the use of diastereomeric catalysts  $\text{Rh}_2(\text{S,S-MenthAZ})$  and  $\text{Rh}_2(\text{S,R-MenthAZ})_4$ ,<sup>11</sup> the selectivities from which were high and comparable (Table 1, entries 5 and 6). Absolute stereochemistry of **2a** was determined by oxidative aromatization of the diene of **2a**, followed by reduction of the olefin to provide the known compound  $\alpha$ -phenyl- $\gamma$ -valerolactone. Comparison of the optical rotation to that reported for (*R*)- $\alpha$ -phenyl- $\gamma$ -valerolactone<sup>12</sup> allowed the assignment of absolute stereochemistry of **2a** as shown (Table 1). All of the dirhodium catalysts evaluated provided comparable isolated yields of **2a,b**.

Encouraged by the results of this C-H insertion study, we turned our attention to asymmetric cyclopropanation of **1** (Table 2). Reactions occurred cleanly and in high yield with notable diastereoselectivities. As with the previously described C-H insertion, azetidinone ligated dirhodium catalysts are the optimal catalysts for the cyclopropanation of styrene [ $\text{Rh}_2(\text{S-DOSP})_4$ , **3a** = 60% yield, 14% ee, dr > 20:1;  $\text{Rh}_2(\text{S-MEPY})_4$ , **3a** = 69% yield, 10% ee, dr > 20:1;  $\text{Rh}_2(\text{S,R-MenthAZ})_4$ , **3a** = 74% yield, 84% ee, dr > 20:1]. Other cyclopropanation reactions catalyzed by  $\text{Rh}_2(\text{S,R-MenthAZ})_4$  provided enantioselectivities ranging from 73–86% ee, with the *E*-cyclopropane as the dominant or exclusive diastereomer.

Extensive studies of vinyl diazoacetates by Davies have demonstrated that upon formation of *cis*-divinylcyclopropanes a Cope rearrangement occurs to provide hydroazulenes.<sup>13</sup> Construction of seven-membered carbocycles is important in natural product synthesis, and the development of stereoselective strategies toward these targets remains an ongoing challenge.<sup>14</sup> However, unlike the transient *cis*-divinylcyclopropanes formed from vinyl diazoacetates, conformational restrictions on spirocyclic cyclopropanes **3b-c** prevent subsequent Cope rearrangement and allow isolation of the cyclopropane products without rearrangement. Cleavage of the lactone overcomes this restriction and allows the dienes to access the necessary conformation for rearrangement.

Cope rearrangement product **4** was obtained upon reduction of lactone *E*-**3d** with  $\text{LiAlH}_4$  in refluxing THF (eq 1). Recrystallization provided a facile means of separating **4** from the cyclopropane-diol arising from *Z*-**3d** and, concomitantly, the enantiomeric excess of **4** was enriched to 92%.<sup>15</sup> Thus, in two steps the hydroazulene **4** is conveniently accessed in high enantiomeric excess from the diastereomeric mixture of **3d**. The relative stereochemistry of **4** was confirmed by X-ray diffraction.



(1)

Vinyl diazylactone **1** is an effective and stable vinyl carbene precursor. Over the course of this study comparisons were made of the reactivity of **1**, which had been stored for several weeks,

with that of freshly purified **1**; no significant diminution in yield or selectivity was observed, emphasizing the operational convenience of **1** as a vinylcarbene precursor. The striking differential in selectivity between dirhodium(II) carboxylates and carboxamides was unexpected but is probably a result of conformational influences in the reacting carbenes.<sup>8, 16</sup> The spirocyclic structure of **3b-d** allows the isolation of unrearranged divinylcyclopropanes, and reduction of divinylcyclopropane lactones (i.e., **3d**) provides a convenient route to hydroazulenes such as **4**. The stereochemistry for substitution at the 3,4-position of **4** is opposite that which would be obtained from reactions of *E*-substituted vinyl diazoacetates. Future research will be focused on evaluating the scope of cyclic vinyl diazo compounds as carbene precursors and providing a better understanding of the unique selectivities obtained with azetidinone ligated dirhodium catalysts.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.


## Acknowledgements

We are grateful to Prof. H. M. L. Davies for providing a sample of Rh<sub>2</sub>(S-DOSP)<sub>4</sub>. Peter Y. Zavalij of the University of Maryland X-ray Crystallographic Center obtained the X-ray crystal structure of **4**. Funding for this work was provided by the National Institutes of Health (GM 46503) and the National Science Foundation. DB would like to thank NSERC (Natural Sciences and Engineering Research Council of Canada) for a postgraduate scholarship.

## References


1. Doyle, MP.; McKervey, MA.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds: From Cyclopropanes to Ylides*. 1998. p. 652. Davies, HML.; Walji, AM. *Modern Rhodium-Catalyzed Organic Reactions*. Evans, PA., editor. Wiley-VCH; 2005. p. 301
2. (a) Davies HML, Dai X, Long MS. *J Am Chem Soc* 2006;128:2485. [PubMed: 16478205] (b) Nyong AM, Rainier JD. *J Org Chem* 2005;70:746. [PubMed: 15651836] (c) Yan M, Jacobsen N, Hu W, Gronenberg Luisa S, Doyle MP, Colyer John T, Bykowski D. *Angew Chem, Int Ed Engl* 2004;43:6713. [PubMed: 15593151] (d) Doyle MP, Yan M, Hu W, Gronenberg LS. *J Am Chem Soc* 2003;125:4692. [PubMed: 12696871] (e) Davies HML, Stafford DG, Hansen T. *Org Lett* 1999;1:233. [PubMed: 10905868] (f) Davies HML, Doan BD. *J Org Chem* 1998;63:657. [PubMed: 11672057]
3. Davies HML, Huby NJS, Cantrell WR Jr, Olive JL. *J Am Chem Soc* 1993;115:9468.
4. Davies HML, Hodges LM, Matasi JJ, Hansen T, Stafford DG. *Tetrahedron Lett* 1998;39:4417.
5. (a) Bulugahapitiya P, Landais Y, Parra-Rapado L, Planchenault D, Weber V. *J Org Chem* 1997;62:1630. (b) Landais Y, Planchenault D, Weber V. *Tetrahedron Lett* 1994;35:9549.
6. Baum JS, Shook DA, Davies HML, Smith HD. *Synthetic Communications* 1987;17:1709.
7. (a) Davies HML, Beckwith REJ. *Chem Rev* 2003;103:2861. [PubMed: 12914484] (b) Davies HML, Antoulinakis EG. *J Organomet Chem* 2001;617-618:47.
8. Davies HML. *Eur J Org Chem* 1999:2459.
9. Doyle MP, Van Oeveren A, Westrum LJ, Protopopova MN, Clayton TW Jr. *J Am Chem Soc* 1991;113:8982.
10. Doyle MP, Davies SB, Hu W. *Org Lett* 2000;2:1145. [PubMed: 10804575]
11. Hu W, Timmons DJ, Doyle Michael P. *Org Lett* 2002;4:901. [PubMed: 11893181]
12. Nakamura Y, Takeuchi S, Ohgo Y, Yamaoka M, Yoshida A, Mikami K. *Tetrahedron* 1999;55:4959.
13. Davies HML, Stafford DG, Doan BD, Houser JH. *J Am Chem Soc* 1998;120:3326.
14. For recent synthetic advances (a) Wender PA, Haustedt LO, Lim J, Love JA, Williams TJ, Yoon JY. *J Am Chem Soc* 2006;128:6302. [PubMed: 16683779] (b) Sperry JB, Wright DL. *J Am Chem Soc* 2005;127:8034. [PubMed: 15926826] For a recent review (c) Battiste MA, Pelphey PM, Wright DL. *Chem Eur J* 2006;12:3438.
15. Prior to recrystallization, measured enantiomeric excess of **5** matched that of **4d**.

16. Doyle MP, Winchester WR, Hoorn JAA, Lynch V, Simonsen SH, Ghosh R. *J Am Chem Soc* 1993;115:9968.



**Figure 1.**  
Asymmetric dirhodium catalysts.

Table 1

C-H insertion of **1** into 1,4-cyclohexadiene.<sup>a</sup>

Entry	Catalyst	2a:b <sup>b</sup>	Yield (%)	ee (%) <sup>c</sup>
1	Rh <sub>2</sub> (OAc) <sub>4</sub>	2:3	53	—
2 <sup>d</sup>	Rh <sub>2</sub> (S-DOSP) <sub>4</sub>	1:9	56	18, <i>R</i>
3	Rh <sub>2</sub> (S-MEPY) <sub>4</sub>	4:1	47	8, <i>R</i>
4	Rh <sub>2</sub> (S-MEAZ) <sub>4</sub>	9:1	43	60, <i>R</i>
5	Rh <sub>2</sub> (S,S-MenthAZ) <sub>4</sub>	9:1	42	79, <i>R</i>
6	Rh <sub>2</sub> (S,R-MenthAZ) <sub>4</sub>	9:1	50	80, <i>R</i>


<sup>a</sup>Unless otherwise noted, to a solution of 5 equivalents diene and 1.0 mol% of catalyst in refluxing CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under N<sub>2</sub> was added a solution of **1** (1.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) over 8h.

<sup>b</sup>Determined by <sup>1</sup>H NMR spectroscopy prior to chromatography.

<sup>c</sup>% ee of **2a** determined by GC.

<sup>d</sup>Reaction run at room temperature.

Table 2

Asymmetric cyclopropanation of selected alkenes with **1**.<sup>a</sup>

<sup>a</sup>To olefin (2.5 equiv) and catalyst (1.0 mol %) in refluxing  $\text{CH}_2\text{Cl}_2$  (20 mL) under  $\text{N}_2$  was added a solution of **2** (1.3 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) over 8h. E:Z ratio determined by  $^1\text{H}$  NMR prior to chromatography. Reported yields obtained upon isolation via silica gel chromatography.

<sup>b</sup>E-cyclopropane.