

Minireview

Role of *LKB1* in lung cancer development

L Makowski¹ and DN Hayes^{*,2}

¹Division of Endocrinology, Department of Medicine, Sarah W Stedman Nutrition and Metabolism Center, Duke University Medical Center, Metabolism, and Nutrition, Durham, NC, USA; ²Division of Medical Oncology, Department of Internal Medicine, Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC, USA

Three phenotypically related genetic syndromes and their lesions (*LKB1*, *PTEN*, and *TSC1/2*) are identified as frequently altered in lung cancer. *LKB1*, a kinase inactivated in 30% of lung cancers, is discussed in this review. Loss of *LKB1* regulation often coincident with *KRAS* activation allows for unchecked growth and the metabolic capacity to accommodate the proliferation.

British Journal of Cancer (2008) **99**, 683–688. doi:10.1038/sj.bjc.6604515 www.bjcancer.com

Published online 19 August 2008

© 2008 Cancer Research UK

Keywords: *LKB1*; *STK11*; Peutz-Jeghers syndrome; review; lung cancer; heritable tumour syndromes

HERITABLE TUMOUR SYNDROMES AND LUNG CANCER

Heritable tumour syndromes have provided invaluable insights into human disease. These focal perversions of normal growth and development shine as a beacon on particular genes and pathways in a manner, which, once deciphered, can impact far beyond the heritable syndrome itself. Just as a jigsaw puzzle assembled in parts suddenly comes together when a key piece is recognised, the genetic syndrome can bring together vastly disparate biologic knowledge to yield a bit of clarity where none existed before. Although the field of lung cancer has lacked such a fortuitous discovery to date, a series of recent events appears likely to open a new era in our understanding of the disease. Three genetic syndromes, Peutz–Jeghers syndrome (PJS), Cowden's disease (CD), and tuberous sclerosis (TS), all sharing parallel phenotypes primarily of non-malignant and non-pulmonary disease, have crystallised a common pathway, which is activated in a significant proportion of all human lung cancers. Key pieces have been identified and joined, such as nutrient metabolism, proliferative signalling, and anti-apoptosis pathways. The puzzle of lung tumour genesis, invasion, and metastasis is somewhat more complete.

LKB1 AND PJS

The central figure in the current discussion is the gene *LKB1*. Its discovery as well as an appreciation of its biology is dominated by its causative role in the autosomal-dominant inherited PJS. First described in the 1920s by Johannes Peutz, an association with cancer was made by Harold Jeghers in the 1940s (Peutz, 1921; Jeghers *et al*, 1949). The syndrome includes an overgrowth of differentiated tissues called hamartomas, primarily polyps in the gastrointestinal tract (GI), as well as abnormal pigmentation of the mucous membranes and skin. Validating Dr Jegher's observations, large cohorts have clarified the risk and spectrum of malignancies

in PJS, including a cumulative cancer risk of 60% by the age 60 years (Hearle *et al*, 2006). Of primary concern for PJS patients are tumours of GI origin, although breast, gynaecologic, pancreas, and lung have been reported.

The pattern of inheritance of PJS, as well as the high frequency of cancer in affected, indicated that a tumour suppressor gene was likely the causative agent. In 1997, Hemminki *et al* (1997) identified the chromosome region 19p13.3 as the relevant cytoband. A year later a single gene called serine/threonine kinase 11 (*STK11*) was identified as the culprit (Hemminki *et al*, 1998; Jenne *et al*, 1998). The transcript for *STK11* had previously been reported in an unmapped fashion by the more commonly used gene alias *LKB1*. The identification of a tumour suppressor and its genomic coordinates inspired a flurry of tumour resequencing surveys of *LKB1* in a range of tissues. Table 1 reviews a sample of the largest of these reports in the area of lung cancer, and includes associated epigenetic findings. Although infrequently mutated in spontaneous tumours, *LKB1* point mutations were seen in a pattern mirroring those seen in association with PJS, including cervix, gastrointestinal, and pancreas (Avizienyte *et al*, 1999; Sanchez-Cespedes, 2007). A range of atypical PJS tumours was also identified, including prostate and melanoma. Surprisingly, the highest mutation rate, hovering around 25% in multiple studies, was in lung adenocarcinomas, an established but uncommon PJS-associated tumour (Virmani *et al*, 1998; Sanchez-Cespedes *et al*, 2001) (Table 1).

LKB1 LOSS IN CANCER

The *LKB1* locus had been previously indicated as important in lung cancer, although admittedly without the laser focus of a known genomic syndrome and target gene. Indeed, 19p13.3 has been repeatedly indicated for its high frequency of loss of heterozygosity (LOH), often above 50%, in lung cancer by chromosomal analysis (Sobottka *et al*, 2000). Interestingly, the frequency of LOH for many tumours, including lung cancer, continues to be significantly higher than the reported rate of point mutations. The discrepancy between the reported rate of mutation and LOH raises a number of questions, including the accuracy of

*Correspondence: Dr DN Hayes; E-mail: hayes@med.unc.edu

Received 21 February 2008; revised 3 June 2008; accepted 25 June 2008; published online 19 August 2008

Table 1 Summary results of studies reporting *LKB1* mutation and other related events in lung cancer

Reference	Year	Finding	Method	Comments/observations
Virmani <i>et al</i> (1998)	1998	10/12 LOH in NSCLC cell lines and 3/9 in SmCC cell lines in 19p13.2	Allelotyping by PCR	Lung cancer cell lines used to define common, specific known, and novel deletions in NSCLC and SmCC cell lines
Sobottka <i>et al</i> (2000)	2000	7/12 (58%) patients with LOH at 19p in brain metastases of lung cancer patients, whereas glioma <i>LKB1</i> LOH was <20%	Microsatellite PCR analysis and genomic sequencing	None of brain metastases showed <i>LKB1</i> mutations by direct genomic sequencing
Sanchez-Cespedes <i>et al</i> (2002)	2002	5/20 of primary AC, 0/12 primary SCC; and 2/9 of lung cancer cell lines with mutation of <i>LKB1</i> 1/20 tumor with <i>LKB1</i> promoter hypermethylation	Microsatellite PCR, direct sequencing, long-range PCR Methylation-specific PCR	0/12 mutations in SCC
Ghaffar <i>et al</i> (2003)	2003	9/35 loss of <i>LKB1</i> protein in AC, 10/96 loss in AAH (7/33 loss in high-grade AAH; 3/63 loss in low grade; 9/35 invasive tumors)	Immunohistochemistry	<i>LKB1</i> protein loss from AAH correlated with severe dysplasia, suggesting loss as essential to transition from pre- to malignant tumor. Also showed strong positive association between biallelic gene inactivation and negative protein expression
Fernandez <i>et al</i> (2004)	2004	5/19 lung AD with point mutations of <i>LKB1</i> 0/19 samples with promoter methylation of <i>LKB1</i> 10/18 of samples with LOH at 19p	Direct sequencing, long-range PCR Methylation-specific PCR Fluorescent <i>in situ</i> hybridization	Frequent observation of concurrent <i>KRAS</i> (4/11) and <i>p53</i> (6/11) in <i>LKB1</i> mutant samples. Performed microarray analysis with several genes dependent on <i>LKB1</i> status
Carretero <i>et al</i> (2004)	2004	6/11 AD contained <i>LKB1</i> deletion or truncating mutant, 0/11 SCC	Allelotyping by PCR and sequencing	Several AD lines with <i>LKB1</i> mutant also have <i>KRAS</i> mutant
Matsumoto <i>et al</i> (2007)	2007	Lung cancer cell lines with mutations in <i>LKB1</i> : 13/31 AD, 1/2 ADSCC, 3/11 SCC, 3/7 LCC, and 1/19 SmCC 8/70 lung cancer cell lines demonstrated shortened <i>LKB1</i> transcripts, 9/70 demonstrated absent <i>LKB1</i> transcript Primary tumors: 1/106 stage I tumor with <i>LKB1</i> mutation, 3/24 stages I–III. 7/91 male smokers, 0/64 female non-smokers 3/25 brain metastases of lung cancer patients with <i>LKB1</i> mutation	RT-PCR, sequencing and genomic sequencing	Strong positive correlation with <i>KRAS</i> status, not correlation with <i>p53</i> status or <i>EGFR</i> mutation status Two cell lines with absent gene expression did not demonstrate mutation, suggesting alternate gene silencing method. The 9/70 with loss of <i>LKB1</i> had deletions in promoter or exon 1 Small numbers limit interpretation for <i>KRAS</i> and <i>EGFR</i> associations, appears to be associated with poorly differentiated tumors
Ji <i>et al</i> (2007)	2007	Primary tumors: 19/80 AD, 6/42 SCC, 1/7 LCC, and 1/4 ADSCC, respectively, with single copy alterations of <i>LKB1</i> ; 8/80 AD, 2/42 SCC with homozygous mutations	Direct sequencing, multiplex ligation-dependent probe amplification	SCC, AD, SCC, LCC, and AD: frequent loss of a single copy of <i>LKB1</i> , concomitant <i>p53</i> , and <i>KRAS</i> mutations found frequently
Onozato <i>et al</i> (2007)	2007	Cell lines with mutations in <i>LKB1</i> : 3/8 AD, 1/6 SCC, 1/3 LCC, and 0/5 SmCC 3/100 primary tumors with <i>LKB1</i> mutations in Japanese cohort 3/33 male smokers	Direct sequencing Direct sequencing	Lung cancer cell lines examined <i>LKB1</i> mutation rare in Japanese lung cancer population. All mutations were in male smokers with moderately or poorly differentiated tumors

ADSCC = adenocarcinoma; AD = adenocarcinoma; AAH = atypical adenomatous hyperplasia; LCC = large cell carcinoma; LOH = loss of heterozygosity; NSCLC = non-small-cell lung cancer; PCR = polymerase chain reaction; SmCC = small cell carcinoma; SCC = squamous cell carcinoma.

the reported mutation studies, the potential for other mechanisms of gene silencing (such as, promoter methylation or other epigenetic event), or the possibility that a second gene target lies in the region. In fact, careful attention to resequencing is crucial, and techniques that detect large homozygous deletions are required to fully characterize tumour *LKB1* loss (Ji *et al*, 2007). Yet, thorough sequencing alone does not fully account for the LOH mutation frequency discrepancy, prompting investigators to pursue other mechanisms of gene silencing. Promoter methylation accounts for a small fraction of repressed *LKB1* expression cases, but this is unlikely a major contributing mechanism (Esteller *et al*, 2000; Trojan *et al*, 2000). Other mechanisms of gene silencing, including microRNA, might contribute to some of the gap but these remain under investigation. Perhaps, the most interesting possibility is that *LKB1* haploinsufficiency itself could be tumori-

genic, as we have recently shown in animal models (Ji *et al*, 2007). At this time, however, we cannot exclude the possibility that there is another gene target in the 19p13.3 region that could account for additional cases of LOH, such as *BRG1* (Medina *et al*, 2008). In contrast to the challenges of detecting functional *LKB1* gene dosage described above, two authors have reported that immunohistochemical assessment of *LKB1* protein status as a straightforward proxy for both biallelic loss and promoter methylation (Ghaffar *et al*, 2003; Fenton *et al*, 2006). Taken together, information on genetic status of *LKB1* garnered through examining PJS patients and screening multiple tumour types has pointed to *LKB1* as an important mediator in the development of cancer.

Frequent LOH at 19p13.3 points only indiscriminately to the importance of the *LKB1* locus in lung cancer, as many regions have chromosomal alterations in this disease. Similarly, the dramatic

clinical histories of a minority of PJS patients include lung cancer, but not at such a high rate as to suggest the prevalence of *LKB1* alteration in this tumour (Hirano *et al*, 2002; Estrada Trigueros *et al*, 2005; von Herbay *et al*, 2005). Yet, cancers in a few young patients with PJS, indeed, suggest a role as a potential early carcinogenic event. Ghaffar *et al* (2003) have bolstered this possibility by detecting frequent *LKB1* loss in the progression of the premalignant lesion atypical adenomatous hyperplasia (AAH) towards frank invasive. Developmentally, *LKB1* is expressed in foetal lung and ubiquitously expressed in adult bronchial mucosa, underscoring its importance in normal lung physiology and development (Luukko *et al*, 1999). Step by step, pieces of a puzzle assemble to document both the importance of *LKB1* to the normal lung and its high frequency of aberrancy in cancer. Yet, the reader might still rightfully wonder why does a gene that primarily causes genetic disease of the gut appear to be mutated so early and often in sporadic cases of lung cancer?

LKB1 EXPRESSION, REGULATION, AND ACTION

Like individual pieces of a jigsaw puzzle, the various facets of *LKB1* need further consideration before they reveal a coherent picture. The functional domains of *LKB1*'s 10 exons (9 coding) include a central catalytic domain, nuclear localisation signal, putative cytoplasmic retention signal, and a prenylation motif (Alessi *et al*, 2006). Localised to the nucleus, it is transported to the cytosol and activated by binding two associated proteins STRAD-related adaptor (STRAD), a catalytically inactive kinase-like protein, and an armadillo repeat-containing protein named mouse protein 25 (MO25). The complex is 100-fold more active as a kinase than *LKB1* alone. *LKB1* also associates with a chaperone complex consisting of HSP90 and Cdc37, which likely aids in stabilisation of

LKB1 protein and assembly of the STRAD–MO25 complex. *LKB1* is phosphorylated on several sites, either activating or inhibiting, by several kinases, including p90 ribosomal S6 kinase (RSK), protein kinase A (PKA), or ataxia-telangiectasia-mutated (ATM) kinase. Despite the apparently rich regulatory aspects of *LKB1*, however, no mutation hot spots have emerged.

LKB1 functions as a tumour suppressor that regulates cell polarity, differentiation, and metastasis as well as responds to energy status to regulate cell metabolism (Figure 1) (see Alessi *et al*, 2006; and references within). One function of *LKB1* directly linked to lung cancer is the phosphorylation, ubiquitination, and the degradation of polyomavirus enhancer activator 3 (PEA3). Mutant *LKB1* increases the stability of PEA3, which, in turn, increases transcription of genes involved in metastasis, including *COX-2*, *MMP-9*, and *MMP-14* (Upadhyay *et al*, 2006; Cowden Dahl *et al*, 2007). Interestingly, *COX-2* is commonly elevated in many disease states, including PJS and lung cancer (Achiwa *et al*, 1999; Rossi *et al*, 2002; McGarrity *et al*, 2003). Indeed, *COX-2* inhibition decreased polyps in a small study of PJS patients and a large study of patients with familial adenomatous polyposis (Udd *et al*, 2004; Bertagnolli *et al*, 2006). Finally, increased *COX-2* is associated with decreased survival in patients with lung adenocarcinoma (Achiwa *et al*, 1999). Although *LKB1* clearly regulates proteins such as PEA3, current understanding is that its primary action is through the target 5'-AMP-activated protein kinase (AMPK) (Towler and Hardie, 2007). AMPK is a master regulator of metabolism that orchestrates efficient energy production with minimal waste in times of energy stress. It is a serine/threonine kinase activated by metabolic signals and stressors, including exercise, starvation, hypoxia, and ischaemia. In short, consumption of ATP, decreased ATP production, and/or increased concentration of AMP activate AMPK to alter numerous metabolic and proliferative pathways (Figure 1). Downstream of *LKB1*, AMPK tightly regulates the




Figure 1 Energy sentinels in proliferation: *LKB1* and AMPK. *LKB1* is activated by binding STRAD and MO25. AMPK is a master regulator of metabolism, which is activated by *LKB1* phosphorylation when the AMP/ATP ratio is high. AMPK drives catabolic pathways such as glucose uptake, glycolysis, mitochondrial proliferation, and fatty acid oxidation to generate ATP in times of stress. It also blunts ATP-consuming anabolic pathways including fatty acid, cholesterol, glycogen, and protein synthesis. The latter role of inhibiting protein synthesis through TSC1/2 and mTOR underlie AMPK's role in cell growth and proliferation. However, *LKB1*–AMPK regulation of fuel availability and the metabolic status is also central to proliferation. In fact, the *LKB1* signal that cells should halt division when sufficient nutrients are not present overrides growth factor signalling through EGFR, for example. In sum, *LKB1* and AMPK act together to sense energy stress to provide cells with nutrients for rapid energy production, while limiting growth. The sentinel activity of *LKB1* and AMPK is lost in cancer, which allows unlimited proliferation despite metabolic cues to the contrary.

metabolic status of cells. For example, when carbon sources are scarce, AMPK ensures ATP generation by activating catabolic pathways such as glycolysis and fatty acid oxidation, as well as upregulates mitochondrial biogenesis. In parallel, it switches off ATP-consuming anabolic pathways, including fatty acid, cholesterol, glycogen, and protein synthesis. Of relevance to this review, AMP binding to AMPK only causes a fivefold increase in activity, while the AMP-induced conformational change in AMPK primes it as a substrate for LKB1, allowing LKB1 phosphorylation to induce activation by 100-fold (Alessi *et al*, 2006). Therefore, AMPK action and LKB1 activity are tightly coordinated to sense energy stress and attempt to provide cells with fuel for rapid energy production, while limiting growth and energy loss when nutrients are scarce.

LKB1 PROLIFERATION, METASTASIS, AND METABOLISM

A central target for AMPK's control of proliferation is the mammalian target of rapamycin (mTOR) kinase, which regulates numerous downstream targets, such as amino acid transporters, VEGF, p70 ribosomal protein S6 kinase 1 (S6K), and eukaryotic initiation factor 4E-binding protein 1 (4E-BP1) pathways, among others, to increase the translation of proteins and cell growth (Figure 1). In fact, AMPK acts on mTOR through phosphorylating and activating tumour suppressor tuberous sclerosis complex-2 (TSC2), an upstream negative regulator of mTOR. AMPK activation also blocks cell cycle progression from G1 to S phase through phosphorylation and accumulation of the tumour suppressor p53 and the downstream cyclin-dependent kinase inhibitors p21^{WAF1/CIP1} and p27. Other cell cycle regulation occurs through the alteration of cytoplasmic/nuclear ratios of RNA-binding protein human antigen R (HuR), thereby reducing the ability to stabilise mRNAs encoding cyclins (Wang *et al*, 2002). Finally, hypoxia-induced AMPK activation of endothelial NOS (eNOS) drives angiogenesis. The relative importance of each mechanism discussed remains to be determined. In sum, LKB1 inactivation in cancer releases the AMPK-mediated breaks on energy waste and permits proliferation. The capacity to sense a nutrient deficit is a function of healthy cells that is lost in proliferating cells, which highlights a distinctly separate but parallel piece of the cancer puzzle.

HAMARTOMATOUS SYNDROMES, mTOR, AND LUNG CANCER

Decoding the regulatory networks of LKB1, particularly an interaction with the AMPK–mTOR axis, crystallises a link between a genetic disease with little pulmonary implications and a gene frequently altered in lung cancer. Clearly, mTOR is a lung cancer target; yet, the jumble of alternate pathways raises doubts as to whether this is the primary mechanism (Figure 1). Quite remarkably, upstream of the mTOR pathway, however, is the invocation of three genes and two autosomal-dominant inherited syndromes with phenotypes remarkably similar to PJS, including PTEN (CD) and TSC1/TSC2 (tuberous sclerosis, TS). Like PJS, both CD and TS manifest themselves as primary hamartomatous diseases. Patients with CD demonstrate hamartomas of the skin, mucous membranes, breast, and thyroid, with 85% of patients showing PTEN germline mutation. Like PJS, an increased risk of tumours is seen, although again lung cancer is not prominent. The PTEN phosphatase controls the levels of PIP₃, induced by growth factor activation of PI3 kinase, and acts to negatively regulate mTOR activity. Patients with TS do not have malignancy as their primary concern, but instead hamartomas of the skin, brain, kidney, skin, lungs, and other tissues (Crino *et al*, 2006). As in CD,

the TS complex acts as a negative regulator of mTOR. In summary, the related phenotypes of PJS, CD, and TS are joined by shared regulation of the mTOR signalling cascade.

Consideration of these syndromes begs the question, is there a broader lung cancer malignant phenotype equivalent to that shared by germline PJS, CD, and TS? In fact, mutations of PTEN have been reported in many tumours, including 4–8% of non-small-cell lung cancers (NSCLC) (Forbes *et al*, 2006). PTEN protein expression is lost in more than 25% of NSCLC tumours with evidence of epigenetic silencing at work as well (Luukko *et al*, 1999; Marsit *et al*, 2005). Additionally, PTEN is co-expressed with LKB1 in foetal lung development. Similarly, both the TSC1 locus (9q34) and the TSC2 locus (16p) are frequent targets of LOH in both lung adenocarcinoma and the pre-invasive lung lesion AAH (Takamochi *et al*, 2001, 2004). Although the frequency with which these events occur in concert remains unknown, the conclusion must be that targets coalescing on the mTOR pathway defined in large part by dissecting the disease of germline mutations of LKB1, PTEN, and TSC1/2 appear to be frequently altered in lung cancer, suggesting a broad and important contribution of this specific set of interacting proteins to the disease.

TURNING TO ANIMAL MODELS: LKB1 AND KRAS

Recent years have seen progress similar to that we document for LKB1 in a number of other lung cancer genes and cancer networks, such as EGFR, KRAS, p53, and CDKN2A (including both transcripts: p16^{Ink4a} and INK4a/ARF). As we suggest above, it is expected that the alteration of only one gene in a pathway may be sufficient for tumorigenesis. As such, an understanding of the relationships between aberrations may help clarify the minimal set of events necessary to cause cancer. To this end, we documented several interactions between LKB1 and other genes commonly altered in lung cancer in experimental animal models (Ji *et al*, 2007). As predicted from the natural human experiment of PJS, mice deficient in LKB1 do not get lung tumours. The alteration of other lung cancer genes in combination with LKB1, however, did produce tumours of a striking phenotype. First, we observed that homozygous loss of LKB1 in combination with KRAS resulted in an aggressive tumour phenotype with high tumour multiplicity, short survival, and frequent metastases well above those of KRAS alone. The KRAS/LKB1 double knockout was in fact the most aggressive phenotype of all tumours considered in the study, and had an additional feature that had not previously been reported in a mouse model: squamous cell carcinoma histology. Additionally, it was noted that even hemizygous loss of LKB1 in combination with KRAS resulted in a more aggressive phenotype than in KRAS mutation alone.

The clinical relevance of LKB1 alteration remains an area of active investigation, although preliminary results appear likely to confirm findings of the animal models, including the frequent co-mutation of LKB1 inactivation and activating mutations of KRAS (Table 1). As a mirror opposite to the established positive association of EGFR gene mutation with females and nonsmokers, we detect a positive association of LKB1 inactivation with both smoking and male gender. Other interesting reports, including prevalence of brain metastasis, associations with poorly differentiated tumours, the importance of the hemizygous LKB1 state, and apparent independence with p53 mutation, remain to be confirmed. Certainly, we hope for clinical associations of this type on which to base important patient management decisions. In broader terms, one has to note that many of the tumours associated with PJS, including GI and gynaecologic malignancies are also those where KRAS is a frequent target of mutation. The question remains, is the LKB1–KRAS connection part of a broad phenotype of interconnecting cancer pathways? We propose that perhaps KRAS activation and LKB1 loss permit liberation from cell

cycle and energy status checkpoints, respectively, which allow for a metabolic advantage.

mTOR AND KRAS IN THE CLINIC

Paramount in the *LKB1* story is the focus now thrust upon very specific cancer pathways, several of which have attractive therapeutic targets. At the top of the list is the mTOR pathway where three Food and Drug Administration (FDA)-approved inhibitors are currently available, such as rapamycin, everolimus, and temsirolimus, with more in development, including the recently described deforolimus (Cohen, 2008). Inhibitors of mTOR have already been investigated in lung cancer, with clear evidence of anticancer activity both as single agents and in combination with cytotoxic chemotherapy and radiation (Milton *et al*, 2007; Sarkaria *et al*, 2007; Mita *et al*, 2008). Interestingly, a recent report in the *New England Journal of Medicine* documented a striking therapeutic effect of mTOR-targeted therapy by sirolimus to angiomyolipomas, which are lesions attributable to TSC dysfunction in patients with TS (Bissler *et al*, 2008). Patients with angiomyolipomas, including the pulmonary form lymphangioleiomyomatosis, almost uniformly experienced reduction in the size of their tumours with this therapy. The efficacy of mTOR-targeted therapy in this benign tumour of the lung, marked by one aspect of mTOR activation, certainly is of interest due to its parallel biology in lung cancer.

Although activity in these early studies has been documented, there is little evidence that this will be of a broader spectrum than

for other classes of lung cancer therapeutics. Therefore, there is an incentive to define biomarkers of response to therapy, such as mutation status or, perhaps, more encompassing markers such as those indicating AMPK activity. There is increasing evidence that broad patterns of tumour behaviour can be captured as general phenotypes using profiling techniques such as gene expression arrays. Specifically, the squamoid gene expression subtype of lung adenocarcinoma is known to have higher rates of *KRAS* mutation and demonstrates gene expression correlated with *LKB1* inactivation (Hayes *et al*, 2006). It is also interesting to note that as *LKB1* mutation appears to pair with *KRAS* activation, specific combinations of therapy targeting parallel pathways might be appropriate. Indeed, initial work on this similar line has begun, including targeting downstream elements of RAS signalling (MEK) and mTOR simultaneously (Legrier *et al*, 2007).

CONCLUSION

Three phenotypically related genetic syndromes, including PJS, CD, and TS, are united by careful alignment of their mechanisms acting through the mTOR pathway. The lesions responsible for the syndromes, *LKB1*, *PTEN*, and *TSC1/2*, are identified as frequently altered in lung cancer, suggesting that they comprise elements of a common lung cancer phenotype. Taken together, when the phenotype of *LKB1* mutation is examined in the setting of known alterations of lung cancer, an interesting association of *KRAS* dependence appears, with specific clinical and treatment potential. Like all good puzzles, the pieces are starting to fall into place.

REFERENCES

- Achiwa H, Yatabe Y, Hida T, Kuroishi T, Kozaki K, Nakamura S, Ogawa M, Sugiura T, Mitsudomi T, Takahashi T (1999) Prognostic significance of elevated cyclooxygenase 2 expression in primary, resected lung adenocarcinomas. *Clin Cancer Res* 5: 1001–1005
- Alessi DR, Sakamoto K, Bayascas JR (2006) LKB1-dependent signaling pathways. *Annu Rev Biochem* 75: 137–163
- Avizienyte E, Loukola A, Roth S, Hemminki A, Tarkkanen M, Salovaara R, Arola J, Butzow R, Husgafvel-Pursiainen K, Kokkola A, Jarvinen H, Aaltonen LA (1999) LKB1 somatic mutations in sporadic tumors. *Am J Pathol* 154: 677–681
- Bertagnoli MM, Eagle CJ, Zauber AG, Redston M, Solomon SD, Kim K, Tang J, Rosenstein RB, Wittes J, Corle D, Hess TM, Woloj GM, Boisserie F, Anderson WF, Viner JL, Bagheri D, Burn J, Chung DC, Dewar T, Foley TR, Hoffman N, Macrae F, Pruitt RE, Saltzman JR, Salzberg B, Sylwestrowicz T, Gordon GB, Hawk ET (2006) Celecoxib for the prevention of sporadic colorectal adenomas. *N Engl J Med* 355: 873–884
- Bissler JJ, McCormack FX, Young LR, Elwing JM, Chuck G, Leonard JM, Schmithorst VJ, Laor T, Brody AS, Bean J, Salisbury S, Franz DN (2008) Sirolimus for angiomyolipoma in tuberous sclerosis complex or lymphangioleiomyomatosis. *N Engl J Med* 358: 140–151
- Carretero J, Medina PP, Pio R, Montuenga LM, Sanchez-Cespedes M (2004) Novel and natural knockout lung cancer cell lines for the LKB1/STK11 tumor suppressor gene. *Oncogene* 23: 4037–4040
- Cohen EE (2008) mTOR: the mammalian target of replication. *J Clin Oncol* 26: 348–349
- Cowden Dahl KD, Zeineldin R, Hudson LG (2007) PEA3 is necessary for optimal epidermal growth factor receptor-stimulated matrix metalloproteinase expression and invasion of ovarian tumor cells. *Mol Cancer Res* 5: 413–421
- Crino PB, Nathanson KL, Henske EP (2006) The tuberous sclerosis complex. *N Engl J Med* 355: 1345–1356
- Esteller M, Avizienyte E, Corn PG, Lothe RA, Baylin SB, Aaltonen LA, Herman JG (2000) Epigenetic inactivation of LKB1 in primary tumors associated with the Peutz–Jeghers syndrome. *Oncogene* 19: 164–168
- Estrada Trigueros G, Lopez-Encuentra A, Garcia Quero C (2005) Non-small cell bronchogenic carcinoma and Peutz–Jeghers syndrome. *Arch Bronconeumol* 41: 296
- Fenton H, Carlile B, Montgomery EA, Carraway H, Herman J, Sahin F, Su GH, Argani P (2006) LKB1 protein expression in human breast cancer. *Appl Immunohistochem Mol Morphol* 14: 146–153
- Fernandez P, Carretero J, Medina PP, Jimenez AI, Rodriguez-Perales S, Paz MF, Cigudosa JC, Esteller M, Lombardia L, Morente M, Sanchez-Verde L, Sotelo T, Sanchez-Cespedes M (2004) Distinctive gene expression of human lung adenocarcinomas carrying LKB1 mutations. *Oncogene* 23: 5084–5091
- Forbes S, Clements J, Dawson E, Bamford S, Webb T, Dogan A, Flanagan A, Teague J, Wooster R, Futreal PA, Stratton MR (2006) Cosmic 2005. *Br J Cancer* 94: 318–322
- Ghaffar H, Sahin F, Sanchez-Cepedes M, Su GH, Zahurak M, Sidransky D, Westra WH (2003) LKB1 protein expression in the evolution of glandular neoplasia of the lung. *Clin Cancer Res* 9: 2998–3003
- Hayes DN, Monti S, Parmigiani G, Gilks CB, Naoki K, Bhattacharjee A, Socinski MA, Perou C, Meyerson M (2006) Gene expression profiling reveals reproducible human lung adenocarcinoma subtypes in multiple independent patient cohorts. *J Clin Oncol* 24: 5079–5090
- Hearle N, Schumacher V, Menko FH, Olschwang S, Boardman LA, Gille JJ, Keller JJ, Westerman AM, Scott RJ, Lim W, Trimbath JD, Giardiello FM, Gruber SB, Offerhaus GJ, de Rooij FW, Wilson JH, Hansmann A, Moslein G, Royer-Pokora B, Vogel T, Phillips RK, Spigelman AD, Houlston RS, Mehenni H, Resta N, Park JG, Miyaki M, Guanti G, Costanza MC, Lim W, Olschwang S, Keller JJ, Westerman AM, Menko FH, Boardman LA, Scott RJ, Trimbath J, Giardiello FM, Gruber SB, Gille JJ, Offerhaus GJ, de Rooij FW, Wilson JH, Spigelman AD, Phillips RK, Houlston RS (2006) Frequency and spectrum of cancers in the Peutz–Jeghers syndrome. Cancer risks in LKB1 germline mutation carriers. Relative frequency and morphology of cancers in STK11 mutation carriers. *Clin Cancer Res* 12: 3209–3215
- Hemminki A, Markie D, Tomlinson I, Avizienyte E, Roth S, Loukola A, Bignell G, Warren W, Aminoff M, Hoglund P, Jarvinen H, Kristo P, Pelin K, Ridanpaa M, Salovaara R, Toro T, Bodmer W, Olschwang S, Olsen AS, Stratton MR, de la Chapelle A, Aaltonen LA (1998) A serine/threonine kinase gene defective in Peutz–Jeghers syndrome. *Nature* 391: 184–187
- Hemminki A, Tomlinson I, Markie D, Jarvinen H, Sistonen P, Bjorkqvist AM, Knuutila S, Salovaara R, Bodmer W, Shibata D, de la Chapelle A, Aaltonen LA (1997) Localization of a susceptibility locus for

- Peutz-Jeghers syndrome to 19p using comparative genomic hybridization and targeted linkage analysis. *Nat Genet* 15: 87–90
- Hirano S, Takiguchi Y, Igari H, Hiroshima K, Shingyoji M, Watanabe R, Moriya T, Tanabe N, Tatsumi K, Kuriyama T (2002) A case of pulmonary adenocarcinoma accompanied by superior vena caval thrombosis in a patient with Peutz-Jeghers syndrome. *Jpn J Clin Oncol* 32: 307–309
- Jeghers H, Mc KV, Katz KH (1949) Generalized intestinal polyposis and melanin spots of the oral mucosa, lips and digits; a syndrome of diagnostic significance. *N Engl J Med* 241: 993, illust; passim
- Jenne DE, Reimann H, Nezu J, Friedel W, Loff S, Jeschke R, Muller O, Back W, Zimmer M (1998) Peutz-Jeghers syndrome is caused by mutations in a novel serine threonine kinase. *Nat Genet* 18: 38–43
- Ji H, Ramsey MR, Hayes DN, Fan C, McNamara K, Kozlowski P, Torrice C, Wu MC, Shimamura T, Perera SA, Liang MC, Cai D, Naumov GN, Bao L, Contreras CM, Li D, Chen L, Krishnamurthy J, Koivunen J, Chirieac LR, Padera RF, Bronson RT, Lindeman NI, Christiani DC, Lin X, Shapiro GI, Janne PA, Johnson BE, Meyerson M, Kwiatkowski DJ, Castrillon DH, Bardeesy N, Sharpless NE, Wong KK (2007) LKB1 modulates lung cancer differentiation and metastasis. *Nature* 448: 807–810
- Legrier ME, Yang CP, Yan HG, Lopez-Barcons L, Keller SM, Perez-Soler R, Horwitz SB, McDaid HM (2007) Targeting protein translation in human non small cell lung cancer via combined MEK and mammalian target of rapamycin suppression. *Cancer Res* 67: 11300–11308
- Luukko K, Ylikorkala A, Tiainen M, Makela TP (1999) Expression of LKB1 and PTEN tumor suppressor genes during mouse embryonic development. *Mech Dev* 83: 187–190
- Marsit CJ, Zheng S, Aldape K, Hinds PW, Nelson HH, Wiencke JK, Kelsey KT (2005) PTEN expression in non-small-cell lung cancer: evaluating its relation to tumor characteristics, allelic loss, and epigenetic alteration. *Hum Pathol* 36: 768–776
- Matsumoto S, Iwakawa R, Takahashi K, Kohno T, Nakanishi Y, Matsuno Y, Suzuki K, Nakamoto M, Shimizu E, Minna JD, Yokota J (2007) Prevalence and specificity of LKB1 genetic alterations in lung cancers. *Oncogene* 26: 5911–5918
- McGarrity TJ, Peiffer LP, Amos CI, Frazier ML, Ward MG, Howett MK (2003) Overexpression of cyclooxygenase 2 in hamartomatous polyps of Peutz-Jeghers syndrome. *Am J Gastroenterol* 98: 671–678
- Medina PP, Romero OA, Kohno T, Montuenga LM, Pio R, Yokota J, Sanchez-Cespedes M (2008) Frequent BRG1/SMARCA4-inactivating mutations in human lung cancer cell lines. *Hum Mutat* 29: 617–622
- Milton DT, Riely GJ, Azzoli CG, Gomez JE, Heelan RT, Kris MG, Krug LM, Pao W, Pizzo B, Rizvi NA, Miller VA (2007) Phase 1 trial of everolimus and gefitinib in patients with advanced nonsmall-cell lung cancer. *Cancer* 110: 599–605
- Mita MM, Mita AC, Chu QS, Rowinsky EK, Fetterly GJ, Goldston M, Patnaik A, Mathews L, Ricart AD, Mays T, Knowles H, Rivera VM, Kreisberg J, Bedrosian CL, Tolcher AW (2008) Phase I trial of the novel mammalian target of rapamycin inhibitor deforolimus (AP23573; MK-8669) administered intravenously daily for 5 days every 2 weeks to patients with advanced malignancies. *J Clin Oncol* 26: 361–367
- Onozato R, Kosaka T, Achiwa H, Kuwano H, Takahashi T, Yatabe Y, Mitsudomi T (2007) LKB1 gene mutations in Japanese lung cancer patients. *Cancer Sci* 98: 1747–1751
- Peutz J (1921) Over een zeer merkwaardige, gecombineerde familiäre polyposis van de slijmvliesen van den tractus intestinalis met die van de neuskeelholte en gepaard met eigenaardige pigmentaties van huid-en slijmvliesen. *Ned Maandschr v Gen* 10: 134–146
- Rossi DJ, Ylikorkala A, Korsisaari N, Salovaara R, Luukko K, Launonen V, Henkemeyer M, Ristimaki A, Aaltonen LA, Makela TP (2002) Induction of cyclooxygenase-2 in a mouse model of Peutz-Jeghers polyposis. *Proc Natl Acad Sci USA* 99: 12327–12332
- Sanchez-Cespedes M (2007) A role for LKB1 gene in human cancer beyond the Peutz-Jeghers syndrome. *Oncogene* 26: 7825–7832
- Sanchez-Cespedes M, Ahrendt SA, Piantadosi S, Rosell R, Monzo M, Wu L, Westra WH, Yang SC, Jen J, Sidransky D (2001) Chromosomal alterations in lung adenocarcinoma from smokers and nonsmokers. *Cancer Res* 61: 1309–1313
- Sanchez-Cespedes M, Parrella P, Esteller M, Nomoto S, Trink B, Engles JM, Westra WH, Herman JG, Sidransky D (2002) Inactivation of LKB1/STK11 is a common event in adenocarcinomas of the lung. *Cancer Res* 62: 3659–3662
- Sarkaria JN, Schwingler P, Schild SE, Grogan PT, Mladek AC, Mandrekar SJ, Tan AD, Kobayashi T, Marks RS, Kita H, Miller RC, Limper AH, Leof EB (2007) Phase I trial of sirolimus combined with radiation and cisplatin in non-small cell lung cancer. *J Thorac Oncol* 2: 751–757
- Sobottka SB, Haase M, Fitze G, Hahn M, Schackert HK, Schackert G (2000) Frequent loss of heterozygosity at the 19p13.3 locus without LKB1/STK11 mutations in human carcinoma metastases to the brain. *J Neurooncol* 49: 187–195
- Takamochi K, Ogura T, Suzuki K, Kawasaki H, Kurashima Y, Yokose T, Ochiai A, Nagai K, Nishiwaki Y, Esumi H (2001) Loss of heterozygosity on chromosomes 9q and 16p in atypical adenomatous hyperplasia concomitant with adenocarcinoma of the lung. *Am J Pathol* 159: 1941–1948
- Takamochi K, Ogura T, Yokose T, Ochiai A, Nagai K, Nishiwaki Y, Suzuki K, Esumi H (2004) Molecular analysis of the TSC1 gene in adenocarcinoma of the lung. *Lung Cancer* 46: 271–281
- Towler MC, Hardie DG (2007) AMP-activated protein kinase in metabolic control and insulin signaling. *Circ Res* 100: 328–341
- Trojan J, Brieger A, Raedle J, Esteller M, Zeuzem S (2000) 5'-CpG island methylation of the LKB1/STK11 promoter and allelic loss at chromosome 19p13.3 in sporadic colorectal cancer. *Gut* 47: 272–276
- Udd L, Katajisto P, Rossi DJ, Lepisto A, Lahesmaa AM, Ylikorkala A, Jarvinen HJ, Ristimaki AP, Makela TP (2004) Suppression of Peutz-Jeghers polyposis by inhibition of cyclooxygenase-2. *Gastroenterology* 127: 1030–1037
- Upadhyay S, Liu C, Chatterjee A, Hoque MO, Kim MS, Engles J, Westra W, Trink B, Ratovitski E, Sidransky D (2006) LKB1/STK11 suppresses cyclooxygenase-2 induction and cellular invasion through PEA3 in lung cancer. *Cancer Res* 66: 7870–7879
- Virmani AK, Fong KM, Kodagoda D, McIntire D, Hung J, Tonk V, Minna JD, Gazdar AF (1998) Allelotyping demonstrates common and distinct patterns of chromosomal loss in human lung cancer types. *Genes Chromosomes Cancer* 21: 308–319
- von Herbay A, Arens N, Friedl W, Vogt-Moykopf I, Kayser K, Muller KM, Back W (2005) Bronchioloalveolar carcinoma: a new cancer in Peutz-Jeghers syndrome. *Lung Cancer* 47: 283–288
- Wang W, Fan J, Yang X, Furer-Galban S, Lopez de Silanes I, von Kobbe C, Guo J, Georas SN, Foufelle F, Hardie DG, Carling D, Gorospe M (2002) AMP-activated kinase regulates cytoplasmic HuR. *Mol Cell Biol* 22: 3425–3436