

at high risk of death. Thereafter the value of adjuvant chemotherapy should be tested in a randomised, prospective trial.

We thank all staff at the Birmingham and West Midlands Regional Cancer Registry and clinicians and pathologists throughout the region who contributed to these data. The research was supported in part by grants from the Cancer Research Campaign. CAM was supported by Boehringer Ingelheim and SW by the United Birmingham Hospitals Endowment Fund.

References

- Cook GA, Draper GJ. Trends in cervical cancer and carcinoma in situ in Great Britain. *Br J Cancer* 1984;50:367-75.
- Hall SW, Monaghan JM. Invasive carcinoma of the cervix in younger women. *Lancet* 1983;ii:731.
- Shepherd JH. Surgical management of early invasive cervical cancer. *Clin Obstet Gynecol* 1985;12:183-202.
- Ward BG, Shepherd JH, Monaghan JM. Occult advanced cervical cancer. *Br Med J* 1985; 290:1301-2.
- Waterhouse JAH. *Cancer handbook of epidemiology and prognosis*. Edinburgh: Churchill Livingstone, 1974.
- General Register Office. *Census 1961 England and Wales*. London: HMSO, 1963. (Table 6 in county reports for Herefordshire, Shropshire, Staffordshire, Warwickshire, and Worcestershire.)
- Office of Population Censuses and Surveys. *Census 1971 England and Wales*. London: HMSO, 1973. (Table 8 in part 1 of county reports for Herefordshire, Shropshire, Staffordshire, Warwickshire, and Worcestershire.)
- Office of Population Censuses and Surveys. *Census 1981 England and Wales*. London: HMSO, 1982. (Table 6 in part 1 of county reports for Hereford and Worcester, Shropshire, Staffordshire, Warwickshire, and West Midlands.)
- Dixon WJ, ed. *BMDP statistical software*. Berkeley: University of California Press, 1983.
- Peto R, Pike MC, Armitage P, et al. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. II. Analysis and examples. *Br J Cancer* 1977;35:1-39.
- Cox DR. Regression models and life tables. *Journal of the Royal Statistical Society [Series B]* 1972;34:187-220.
- Hopkins A. Survival analysis with covariates—Cox models. In: Dixon WJ, ed. *BMDP statistical software*. Berkeley: University of California Press, 1983:576-94.
- Zweifel E. Comments during discussion of paper by Graff-Wien E. (Über den Einfluss der Gravidität auf das Wachstum maligner Tumoren). In: *85. Versammlung zu Wien, September 1913. Verhandlungen der Gesellschaft deutscher Naturforscher und Ärzte*. Leipzig: F C W Vogel, 1914:477.
- Truelsen F. Cancer of the uterine cervix. Copenhagen: University of Copenhagen, 1949. (Dissertation.) Cited in: Lindell A. Carcinoma of the uterine cervix; incidence and influence of age; statistical study. *Acta Radiologica [Stockh]* 1952;92 (suppl):1-102.
- Kamnicker H. Das weibliche Genitalkarzinom bei Jugendlichen. *Wien Med Wochenschr* 1939;89:415-7.
- Decker DG, Fricke RE, Pratt JH. Invasive carcinoma of the cervix in young women. *JAMA* 1955;158:1417-20.
- Adcock LL, Julian TM, Okagaki T, et al. Carcinoma of the uterine cervix FIGO stage I-B. *Gynecol Oncol* 1982;14:199-208.
- Baltzer J, Koepcke W, Lohe KJ, Ober KG, Zander J. Age and 5-year survival rates in patients with operated carcinoma of the cervix. *Gynecol Oncol* 1982;14:220-4.
- Dodds JR, Latour JPA. Relationship of age to survival rate in carcinoma of the cervix. *Am J Obstet Gynecol* 1961;82:33-6.
- Kyriakos M, Kempson RL, Perez CA. Carcinoma of the cervix in young women. *Obstet Gynecol* 1971;38:930-44.
- Mann WJ, Levy D, Hatch KD, Shingleton HM, Soong S. Prognostic significance of age in stage I carcinoma of the cervix. *South Med J* 1980;73:1186-8.
- Carmichael JA, Clarke DH, Moher D, Ohlke ID, Karchmar EJ. Cervical carcinoma in women aged 34 and younger. *Am J Obstet Gynecol* 1986;154:264-9.
- Russell JM, Blair V, Hunter RD. Cervical carcinoma: prognosis in younger patients. *Br Med J* 1987;295:300-3.
- Gauwerky F. Ergebnisse und Erfahrungen bei der Behandlung des Uteruskarzinoms. *Strahlentherapie* 1948;77:325-48.
- Ashley DJB. Evidence for the existence of two forms of cervical carcinoma. *Journal of Obstetrics and Gynaecology of the British Commonwealth* 1966;73:382-9.
- Lindell A. Carcinoma of the uterine cervix; incidence and influence of age; statistical study. *Acta Radiologica [Stockh]* 1952;92 (suppl):1-102.
- Stanhope CR, Smith JP, Wharton JT, Rutledge FN, Fletcher GH, Gallager HS. Carcinoma of the cervix: the effect of age on survival. *Gynecol Oncol* 1980;10:188-93.
- Singer A. Cancer of the cervix; diagnosis and treatment. *Clin Obstet Gynecol* 1985;12:ix-x.
- Morgan LS, Nelson JH. Surgical treatment of early cervical cancer. *Semin Oncol* 1982;9:312-30.
- Campion MJ, Singer A, Mitchell HS. Complacency in diagnosis of cervical cancer. *Br Med J* 1987;294:1337-9.
- Piver MS, Chung WS. Prognostic significance of cervical lesion size and pelvic node metastases in cervical carcinoma. *Obstet Gynecol* 1975;46:507-10.
- Wentz WB, Reagan JW. Survival in cervical cancer with respect to cell type. *Cancer* 1959;12:384-8.
- Wentz WB, Lewis GC. Correlation of histologic morphology and survival in cervical cancer following radiation therapy. *Obstet Gynecol* 1965;26:228-32.
- Swan DS, Roddick JW. A clinical pathological correlation of cell type classification for cervical cancer. *Am J Obstet Gynecol* 1970;116:666-70.
- Van Nagel JR Jr, Donaldson ES, Parker JC, et al. The prognostic significance of cell type and lesion size in patients with cervical cancer treated by radical surgery. *Gynecol Oncol* 1977;5:142-51.
- Chung CF, Stryker JA, Ward SP, Nahhas WA, Mortel R. Histologic grade and prognosis of cancer of the cervix. *Obstet Gynecol* 1981;57:636-41.
- Beecham JB, Halvorsen T, Kolbenstvedt A. Histologic classification, lymph node metastases and patients survival in stage IB cervical cancer: an analysis of 245 uniformly treated cases. *Gynecol Oncol* 1978;6:95-105.
- Shingleton HM, Gore H, Bradley DH, et al. Adenocarcinoma of the cervix. I. Clinical evaluation and pathologic features. *Am J Obstet Gynecol* 1981;139:799-814.
- Rutledge FN, Galakatos AE, Wharton JT, et al. Adenocarcinoma of the uterine cervix. *Am J Obstet Gynecol* 1975;122:236-45.

(Accepted 13 October 1987)

Passive smoking and lung cancer: a publication bias?

JAN P VANDENBROUCKE

Abstract

To assess the likelihood of publication bias in a recent review of the effect of passive smoking on lung cancer the evidence from the reviewed papers was visualised on a "funnel" plot. In such a plot if the relative risks from various studies are plotted according to sample size they should scatter round some underlying true value, the scatter being greatest where the studies have the lowest statistical power—thus showing a "funnel" pattern. If there is publication bias and studies with non-significant results are not being published there should be a "gap" in the plot. The logarithm of the relative risks was plotted against the standard error of the logarithm of the relative risk (which was used instead of sample size as a measure of statistical uncertainty). The resulting plot was compatible with a publication bias but only in studies on men.

Further studies of passive smoking and lung cancer in men seem to be warranted.

Introduction

A recent review on passive smoking and lung cancer by Wald *et al* concluded, in line with other reviews, that passive smoking causes a 30% extra risk of lung cancer—that is, a relative risk of 1.30.¹ This conclusion was challenged by Mantel, who held, among others, that publication bias was responsible for this result and concluded, "Whether or not the risk is raised remains to be taken as a matter of faith according to one's choice."²

The objection of publication bias is interesting, since it is amenable to statistical analysis by the use of "funnel plotting."³

Methods and results

The principle is straightforward. When a diverse number of estimates of some value exist one expects some scatter around the underlying truth. The scatter will be largest, however, for the studies which contain the smallest number of subjects—that is, those which have lowest statistical power.

Department of Clinical Epidemiology, University Hospital, PO Box 9600, 2300 RC Leiden, The Netherlands

JAN P VANDENBROUCKE, MD, PHD, professor of clinical epidemiology

So, ideally, if some value X , which has the true underlying value X_T , has to be estimated and we plot the estimates from different studies according to sample size we expect to see a "funnel." Results will scatter around the truth and the more so in the regions of low statistical power. In figure 1 the borders of the imaginary scatter plot are given by the dashed lines.

When publication bias exists we also expect something else. A publication bias occurs when papers with non-significant results are either not submitted or not accepted for publication.³ Non-significant results are most likely when the statistical power is low—that is, when the numbers are small. So in the event of a positive true effect X_T and the existence of a publication bias we would expect to find few if any published papers within the lower left bottom of the funnel display—that is, around the null effect, indicated by the word "gap" in figure 1.




FIG 1—Idealised funnel plot of expected scatter of study results according to study size. X_T indicates a true positive effect; 0 indicates the null-effect; and "Gap" indicates the expected lack of published results in the event of publication bias.

Following this line of thinking, I set out to plot the studies reviewed by Wald *et al*¹ according to their degree of statistical uncertainty (see references 2-14 from that paper). Simply plotting the relative risk (RR) by study size would not do. Firstly, the plain relative risk does not have a symmetrical distribution and will thus never yield a funnel unless transformed logarithmically. Thus on the Y axis I plotted the logarithm of the relative risk ($\ln(RR)$). Secondly, ranking the reviewed studies by size was not feasible since they contained both cohort and case-control studies, and among the latter there were studies with different case to control ratios. Therefore, I sought some measure of statistical uncertainty independent of the type of study. I used the standard error of the logarithm of the relative risk, which could be calculated from the relative risk and its 95% confidence interval as given by Wald *et al*¹ (see table I of that paper). This was accomplished by logarithmic transformation of these three quantities, calculation of the average distance between upper and lower limits and $(\ln)RR$, and division by 1.96. After an initial inspection I judged it better to present the points for studies in men and women separately. The result is shown in figure 2.




FIG 2—Funnel plot of relative risk (on logarithmic scale) according to standard error of $\ln(RR)$ for men (●) and women (○) calculated on 13 studies reviewed by Wald *et al*.¹ Dashed line indicates proposed passive smoking effect of 1.30.

Discussion

An overall view of figure 2 is consistent with a pattern brought about by publication bias. The lower left of the funnel plot, the "non-significant relative risks around unity" is empty. On closer inspection, however, we see that studies with a larger standard error—that is, those with lower statistical power—are predominantly those in men and that it is among these studies that the possibility of publication bias seems most likely. The pattern of the studies in women is much more symmetrical around the value of 1.30, the average relative risk.

In summary, the funnel plot, used to verify the charge of publication bias by Mantel, shows us that for review purposes the published papers on passive smoking and lung cancer ought to be separated on the basis of sex—as so often in epidemiology—and that for studies in men the objection of publication bias seems reasonable.

The mechanism of such a bias can be imagined. Given the near unanimity in medical circles about the risk of active smoking, epidemiologists will have difficulty in exonerating the smoking habit from causing harm. Confronted with weak data on men alongside stronger data on women, authors or reviewers might be inclined to drop the former in favour of the latter. Theoretically two possible remedies for this unsatisfactory situation exist: either call for a new and large study on men only or invite researchers to submit their unpublished low power studies to accumulate their hidden information. In practice, this aim might be frustrated because the number of men who are passive smokers—for example, through being married to a smoker—is likely to be much lower than among women.

References

- 1 Wald NJ, Nanchackal K, Thompson SG, *et al*. Does breathing other people's tobacco smoke cause lung cancer? *Br Med J* 1986;293:1217-22.
- 2 Mantel N. Lung cancer and passive smoking. *Br Med J* 1987;294:440-1.
- 3 Light RS, Pillemer DB. *Summing up, the science of reviewing research*. Cambridge: Harvard University Press, 1984:65-9.

(Accepted 6 November 1987)