

Early origin of coronary heart disease

Maternal nutrition in early pregnancy may affect placental ratio

EDITOR,—In their editorial discussing the early origins of coronary heart disease Nigel Paneth and Mervyn Susser emphasise the importance of further investigation of the influences on the ratio of placental to fetal weight.¹ Godfrey *et al* have proposed that iron deficiency has a role since low maternal haemoglobin concentrations and a decrease in mean red cell volume during pregnancy are associated with an increase in the ratio.² In contrast, Perry *et al* found no relation between maternal haemoglobin concentrations in early pregnancy and placental ratio at delivery³ but confirmed the positive association between the ratio and maternal body mass index.²

Timing is another important consideration when the influence of the maternal environment on the developing fetus and placenta is studied. Many women start pregnancy with low iron stores, and we hypothesised that placental size would be increased by mid-pregnancy in those with low ferritin and haemoglobin concentrations. To investigate this we developed a method of measuring placental volume during the routine scan at around 18 weeks of pregnancy. In a study of 568 white women we found that low haemoglobin and ferritin concentrations at 14 weeks' gestation were associated with increased placental volume.⁴ Placental volume was also greater in taller women, in multiparous women, and in those who smoked heavily at the time of their last period. We found no relation between maternal weight or body mass index and placental volume.

We have since examined the factors affecting the ratio of placental to fetal size in mid-pregnancy by estimating fetal weight using Shepard *et al*'s algorithm.⁵ The relations between maternal characteristics and the ratio of placental volume to estimated fetal weight were examined by multiple regression analysis (table). The ratio was increased

Simultaneous effects of maternal characteristics on ratio of placental volume to estimated fetal weight (ml/g) at 18 weeks' gestation

Maternal factor	Regression coefficient (95% confidence interval)	P value
Gestation at which scan obtained (days)	-0.013 (-0.018 to -0.009)	<0.0001
Haemoglobin (g/l)	-0.002 (-0.0004 to 0.0)	0.05
Ferritin (square root $\mu\text{g/l}$)	-0.014 (-0.022 to -0.006)	0.001
Smoking at last menstrual period	0.027 (0.006 to 0.048)	0.012
Previous deliveries	0.007 (0.0004 to 0.014)	0.038

in scans obtained at earlier gestations, with lower maternal haemoglobin and ferritin concentrations, with increasing parity, and in mothers who smoked. None of these factors except gestation affected the estimated fetal weight, and the change in ratio was therefore principally due to changes in placental volume.

Perry *et al* concluded that their findings did not support a link between placental ratio and maternal nutrition at birth.³ In contrast, our study suggests that the ratio is affected early in pregnancy by the mother's nutrition as shown by the haemoglobin and ferritin concentrations. The evidence that maternal nutrition affects the placental ratio is further strengthened by experiments in sheep, which have shown that restricted maternal food intake results in an increase in placental size.⁶

DAVID HOWE
Senior registrar in fetal medicine

Birmingham Maternity Hospital,
Birmingham B15 2TG

TIMOTHY WHEELER
Reader in obstetrics and gynaecology

Princess Anne Hospital,
Southampton SO16 5YA

- Paneth N, Susser M. Early origin of coronary heart disease (the "Barker hypothesis"). *BMJ* 1995;310:411-2. (18 February.)
- Godfrey KM, Redman CWG, Barker DJP, Osmond C. The effect of maternal anaemia and iron deficiency on the ratio of fetal weight to placental weight. *Br J Obstet Gynaecol* 1991;98:886-91.
- Perry JJ, Beevers DG, Whincup PH, Bareford D. Predictors of placental weight to fetal weight in multiethnic community. *BMJ* 1995;310:436-9.
- Howe DT, Wheeler T, Osmond C. The influence of maternal haemoglobin and ferritin on mid-pregnancy placental volume. *Br J Obstet Gynaecol* 1995;102:213-9.
- Shepard MJ, Richards VA, Berkowitz RL, Warsof SL, Hobbins JC. An evaluation of two equations for predicting fetal weight by ultrasound. *Am J Obstet Gynaecol* 1982;142:47-54.
- Faichney GJ, White GA. Effects of maternal nutritional status on fetal and placental growth and on fetal urea synthesis in sheep. *Australian Journal of Biological Sciences* 1987;40:365-77.

The influence of parental somatic features on childhood weight has been extensively studied

EDITOR,—In their editorial on the early origin of coronary heart disease (the "Barker hypothesis") Nigel Paneth and Mervyn Susser overlooked my contribution.¹ In 1982 I suggested that some heavier children were relatively immune to non-insulin dependent diabetes and coronary heart disease in adult life and that the decline in rates of coronary heart disease that was occurring in some communities was linked to the increasing average weight of children in those communities.² I was also the first to publish this hypothesis in major North American journals.^{3,4} In 1989 Barker and his colleagues put forward similar ideas.

I have had 11 contributions to the discussion on this subject published, including constructive appraisals of Barker *et al*'s early papers. For example, I advised against generalised postnatal nutritional intervention to promote heavier infants, suggested that the importance of placental weight was as a marker for maternal obesity, and warned that the prenatal nutrition theory was not consistent with the existing data on twins or on prenatal exposure to famine. To date, studies have vindicated these concerns.

There were differences in approach, whereas Barker *et al* focused on maternal nutrient intake during pregnancy I focused on the influence of parental somatic features on weight in childhood. For example, birth weight reflects both maternal adiposity and maternal muscle mass.⁵ Adult somatic features, in turn, are determined by both hereditary and environmental factors.

Primary care doctors do not have the resources of major research teams. Nevertheless, our contributions should be acknowledged. Thus it seems fairer to describe the Barker hypothesis as the "Barker-Bradley hypothesis."

PATRICK J BRADLEY
Primary care doctor

Chatswood,
Sydney,
Australia 2057

- Paneth N, Susser M. Early origin of coronary heart disease (the "Barker hypothesis"). *BMJ* 1995;310:411-2. (18 February.)
- Bradley PJ. Is obesity an advantageous adaptation? *Int J Obes Relat Metab Dis* 1982;6:43-52.
- Bradley PJ. Obesity and risk of coronary heart disease in women. *N Engl J Med* 1990;323:1144.
- Bradley PJ. Decline in the incidence of epidemic glucose intolerance in Nauruans: implications for the "thrifty genotype." *Am J Epidemiol* 1992;136:499-500.
- Frisancho AR, Klayman JE, Matos J. Influence of maternal nutritional status on prenatal growth in a Peruvian urban population. *Am J Phys Anthropol* 1977;46:265-74.

Earlier published work supports the "Barker hypothesis"

EDITOR,—Nigel Paneth and Mervyn Susser discuss the fetal and early life origin of coronary heart disease, referring to the "Barker hypothesis."¹ We acknowledge the important contributions of Barker and his coworkers. More strongly than previous authors, they have pinpointed the rele-

vance of fetal nourishment to later coronary risk. Readers of the *BMJ* should also be aware, however, of the earlier publications by Anders Forsdahl of the University of Tromsø. In 1973, in a paper published in Norwegian with an English summary, he advanced "the theory that extremely poor social conditions during childhood and adolescence can be considered as risk factors for cardiovascular diseases."² This was more than 10 years before the paper by Barker and Osmond in 1986.³ Since 1977 Forsdahl has published several studies supporting his hypothesis in international journals.^{4,5}

Thus we believe that if one person alone is to be given the honour for the theory that experiences in early life are of major importance for the risk of cardiovascular diseases several decades later it should probably be Anders Forsdahl.

INGER TORHILD GRAM
Senior lecturer
BJØRN STRAUME
Senior lecturer
MAJA-LISA LØCHEN
Researcher
BJARNE K JACOBSEN
Professor
EGIL ARNESEN
Professor
EILIV LUND
Professor

Department of Epidemiology and Medical Statistics,
Institute of Community Medicine,
University of Tromsø,
N-9037 Tromsø,
Norway

- Paneth N, Susser M. Early origin of coronary heart disease (the "Barker hypothesis"). *BMJ* 1995;310:411-2. (18 February.)
- Forsdahl A. Momenter til belysning av den høye dødelighet i Finnmark fylke. *Tidsskr Nor Laegeforen* 1973;93:661-7. (Norwegian with English summary.)
- Barker DJP, Osmond C. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* 1986;i:1077-81.
- Forsdahl A. Are poor living conditions in childhood and adolescence an important risk factor for arteriosclerotic heart disease? *British Journal of Preventive and Social Medicine* 1977;31:91-5.
- Forsdahl A. Living conditions in childhood and subsequent development of risk factors for arteriosclerotic heart disease. *J Epidemiol Community Health* 1978;32:34-7.
- Arnesen E, Forsdahl A. The Tromsø heart study: coronary risk factors and their association with living conditions in childhood. *J Epidemiol Community Health* 1985;39:210-4.

A definition of sexual health

EDITOR,—No seminar or conference on the subject of sexual health seems complete without a presentation or debate rhetorically titled "What is sexual health?" Those who have discovered the subject since the launch of *Health of the Nation* face a daunting choice of definitions. Most definitions, coming from a political¹ or health promotion² background, extol in lengthy prose the positive personal benefits of sexual relationships for self esteem and fulfilment, apparently ignoring the medical perspective; the medical perspective appears as a more or less comprehensive list of serious health problems associated with sexual activity, giving an entirely negative impression.

A concise definition of sexual health that unites both benefits and deficits in one sentence of plain English is clearly needed. It must be simple enough to be understood and remembered easily; applicable to a wide range of social, religious, and sexual cultures; yet be robust enough to stand critical analysis and sufficiently unambiguous to be capable of expansion without significant distortion. Such a sentence is: Sexual health is the enjoyment of sexual activity of one's choice, without causing or suffering physical or mental harm.³

This definition allows the term "enjoyment," in its widest interpretation, to be expanded to the lengths of previous statements without prescribing the type of sexual relationship—be it heterosexual, gay or lesbian, monogamous, or otherwise—provided it is by mutual consent without coercion or damage. It can also promote two of the safest options, masturbation and abstinence, as positively healthy choices. It is consistent with the traditional humanist medical principle of first do