

prevalence may be biased. Furthermore, STI screening was offered to but not performed on all patients. Furthermore, repeat testing and the interval of such testing were, ultimately, decided by each individual patient.

Despite these limitations, there is value in assessing the local sex work environment and using this as the basis for public health interventions. Indeed, additional work must also be done to maximise the uptake of voluntary STI screening, particularly among HIV infected sex workers. This study suggests that STI prevention efforts in San Francisco must increasingly target sex workers who are African-American, male, and work independently of other sex workers. Moreover, anti-violence interventions should be an integral part of STI prevention. Our finding of decreased risk of STI among those individuals who have worked collectively with other sex workers is intriguing and warrants further study.

The St James Infirmary represents a novel collaborative effort between public health officials and the sex worker community. Through this unique model, sex workers have helped define public health priorities for their own community and public health officials have gained access to this hidden and stigmatised population.

ACKNOWLEDGEMENTS

We thank all the sex workers who participated in this study, the staff and volunteers at St James Infirmary, and the staff at City Clinic/ Department of Public Health, including Wendy Wolf, Bob Kohn, Charlotte Kent, and Dr Will Wong.

.....

Authors' affiliations

D Cohan, A Lutnick, UCSF, 1001 Potrero Avenue, Ward 6D San Francisco General Hospital Department of Obstetrics/Gynecology San Francisco, CA 94110 USA

P Davidson, University of California, San Francisco Department of Social and Behavioral Sciences, California, USA

C Cloniger, San Francisco Department of Public Health, California, USA

A Herlyn, University of California, San Francisco, California, USA

J Breyer, C Cobaugh, D Wilson, St James Infirmary and University of California, San Francisco, California, USA

J Klausner, San Francisco Department of Public Health and University of California, San Francisco Department of Medicine, California, USA

Funding: This study was supported by the National Institute for Drug Abuse (K23 DA01674), the Ford Foundation, the San Francisco Department of Public Health, and the University of California San Francisco School of Medicine Student Research Training Program.

REFERENCES

- 1 **Lutnick A**. The St James Infirmary: a history. *Sexuality Culture* 2006;**10**(2).
- 2 **Cohan DL, Breyer J, Cobaugh C, et al**. Social context and the health of sex workers in San Francisco, in: XV International AIDS Conference., 2004.
- 3 **Cohan DL, Cloniger C, Breyer J, et al**. Sex worker health, San Francisco style: a peer-based specialty clinic in San Francisco. In: National HIV Prevention Conference, 2001.
- 4 www.stjamesinfirmary.org.
- 5 **Rekart ML**. Sex-work harm reduction. *Lancet* 2005;**366**:2123-34.
- 6 **Breyer J, Cloniger C, Cohan DL, et al**. Training sex workers as disease prevention and health promotion and maintenance educators. In: National STD Prevention Conference, 2002.
- 7 **Alexander P**. Sex work and health: a question of safety in the workplace. *J Am Med Women's Ass* 1998;**53**:77-82.
- 8 **Bagley SC, White H, Golomb BA**. Logistic regression in the medical literature: standards for use and reporting, with particular attention to one medical domain. *J Clin Epidemiol* 2001;**54**:979-85.
- 9 **Harcourt C, Donovan B**. The many faces of sex work. *Sex Transm Infect* 2005;**81**:201-6.
- 10 **Harcourt C, van Beek I, Heslop J, et al**. The health and welfare needs of female and transgender street sex workers in New South Wales. *Aust N Z J Public Health* 2001;**25**:84-9.
- 11 **Cohan DL, Kim A, Ruiz J, et al**. Health indicators among low income women who report a history of sex work: the population based Northern California Young Women's Survey. *Sex Transm Infect* 2005;**81**:428-33.
- 12 **Day S, Ward H**. Violence towards female prostitutes. Violence in sex work extends to more than risks from clients. *BMJ* 2001;**323**:230.
- 13 **Potterat JJ, Brewer DD, Muth SQ, et al**. Mortality in a long-term open cohort of prostitute women. *Am J Epidemiol* 2004;**159**:778-85.
- 14 **Raphael J, Shapiro DL**. Violence in indoor and outdoor prostitution venues. *Violence Against Women* 2004;**10**:126-39.
- 15 **Ward H, Day S, Weber J**. Risky business: health and safety in the sex industry over a 9 year period. *Sex Transm Infect* 1999;**75**:340-3.
- 16 **Ward H, Day S, Green A, et al**. Declining prevalence of STI in the London sex industry, 1985 to 2002. *Sex Transm Infect* 2004;**80**:374-6.
- 17 **Wallace J, Porter J, Weiner A, Steinberg A**. Oral sex, crack smoking, and HIV infection among female sex workers who do not inject drugs. *Am J Public Health* 1997;**87**:470.

CORRECTION

doi: 10.1136/sti.2004.014662.corr1

In the February issue of the journal there was an error in the paper by Hamlyn E, McAllister J, Winston A *et al* (Is screening for sexually transmitted infections in men who have sex with men who receive non-occupational HIV post-exposure prophylaxis worthwhile? *Sex Transm Infect* 2006;**82**:21-3). In reference 10, the initial of the first author was incorrect. The correct name is K Manavi.