

Polymorphisms in the *VKORC1* gene are strongly associated with warfarin dosage requirements in patients receiving anticoagulation

T Li, L A Lange, X Li, L Susswein, B Bryant, R Malone, E M Lange, T-Y Huang, D W Stafford, J P Evans



J Med Genet 2006;43:740–744. doi: 10.1136/jmg.2005.040410

See end of article for authors' affiliations

Correspondence to:
Dr James P Evans, UNC
Chapel Hill, 103 Mason
Farm Road, Room 4200,
CB#7264, Chapel Hill,
NC 27599-7264, USA;
jpevans@med.unc.edu

Revised version received
16 March 2006
Accepted for publication
30 March 2006
Published Online First
12 April 2006

Background: Warfarin is a mainstay of therapy for conditions associated with an increased risk of thromboembolic events. However, the use of this common agent is fraught with complications and little is known regarding inter-individual variation in warfarin response.

Objective: We tested for association between single nucleotide polymorphisms (SNPs) in *VKORC1* and *CYP2C9* and average weekly warfarin dose required to maintain patients at their desired anticoagulation target.

Methods: The sample consisted of 93 European-American patients from anticoagulation clinics at the University of North Carolina at Chapel Hill. Data on mean weekly warfarin dose were collected over a mean treatment period of 20.6 months. ANCOVA models were used and haplotype analysis was performed.

Results: Three of six *VKORC1* SNPs were found to be very strongly associated with the average warfarin dose required to achieve the target international normalised ratio (INR; $p < 0.0001$). The mean weekly dose by genotype ranged from approximately 27 to 47 mg. There was no evidence for an association between either of the two *CYP2C9* polymorphisms studied, *CYP2C9*2* and *CYP2C9*3*. *CYP2C9*3* was significantly ($p = 0.05$) associated with average warfarin dosage after adjustment for *VKORC1*1173*.

Conclusions: These results are of considerable clinical interest and confirm recently published results regarding the role of these two genes in modifying warfarin metabolism and maintenance dosage. The consistent findings regarding the role of *VKORC1* and *CYP2C9* in warfarin metabolism and maintenance dosage represent a clinically useful proof of principal for the use of pharmacogenomic information in medicine and may lead to improved understanding of warfarin's actions.

Individuals vary widely in their response to pharmacologic agents and such variability can have profound health effects. Little is known about drug-specific genetic differences which lead to differential effects of commonly prescribed agents. Warfarin is a mainstay of therapy for conditions which result in an increased risk of thromboembolic events and is used by more than 1 million patients each year in the US. However, the use of this common agent is fraught with complications. Following initiation of warfarin therapy, major bleeding episodes occur in approximately 12% of patients and death results in as many as 2% of patients.^{1,2} Although computer aids, nomograms, and flexible protocols^{3,4} result in improved control of anticoagulation, determining a patient's optimal dose remains highly problematic. Guidance regarding how a given individual will respond to a standard dose of warfarin promises increased efficacy and decreased complications associated with the use of this widely prescribed drug.

Warfarin exerts its pharmacologic effect through its ability to disrupt the γ -glutamyl carboxylation of clotting factors II, VII, IX, and X. The γ -glutamyl carboxylase enzyme requires reduced vitamin K as a cofactor and warfarin inhibits regeneration of reduced vitamin K by targeting the enzyme vitamin K epoxide reductase. The vitamin K epoxide reductase gene (*VKORC1*) was cloned in 2004.^{5,6} The *VKORC1* gene (GI:41352830) is on human chromosome 16p11.2 and encodes a protein of 163 amino acids with a mass of 18.2 kDa. Although this gene has only recently been studied, several articles^{7–12} have already reported evidence for

an association between *VKORC1* genotypes and warfarin dosage.

A second gene, cytochrome P450, subfamily IIC, polypeptide 9 (*CYP2C9*), has been more widely studied in relation to warfarin sensitivity, due to its involvement in metabolism of the drug. Two commonly studied variants in *CYP2C9* (*CYP2C9*2* and *CYP2C9*3*) have consistently been shown to be associated with warfarin dosage. Sanderson *et al* presented a meta-analysis of these results from nine different studies.¹³ The results from the meta-analysis demonstrate that patients with the *CYP2C9*2* and *CYP2C9*3* variant alleles have lower mean daily warfarin dosage and greater risk of bleeding.

We sought to confirm whether genetic variation in the *VKORC1* and *CYP2C9* genes influences an individual's sensitivity to warfarin in a sample of European-American anticoagulation clinic patients, as well as to examine whether the *VKORC1* genetic effects were independent of the two *CYP2C9* polymorphisms that have been established to be associated with warfarin sensitivity.

METHODS

This study was approved by the Biomedical Institutional Review Board at the University of North Carolina at Chapel Hill. Patients attending anticoagulation clinics at UNC who were being treated with warfarin were approached for

Abbreviations: ANCOVA, analysis of covariance; INR, international normalised ratio; LD, linkage disequilibrium; SNP, single nucleotide polymorphism; VTE, venous thromboembolic events

participation in the current study. After obtaining informed consent, blood was obtained from 93 European-American patients for DNA analysis. An extensive database is maintained on these patients that includes the indication for treatment, duration of treatment, average dose, dosage adjustments, and international normalised ratio (INR). Mean weekly warfarin dose required to achieve each patient's target INR was gathered from the database over a mean period of treatment of 20.6 months. Genomic DNAs were extracted from whole blood using the QIAamp DNA Blood Mini Kit (Qiagen, Valencia, CA).

Genomic sequencing

Direct sequencing of 60 patients identified six single nucleotide polymorphisms (SNPs) in the *VKORC1* gene. Approximately 10 ng DNA was used for PCR reactions. The primers used to amplify the 5'-UTR and exon 1 region were: CCAATCGCCGAGTCAGAGG and CCCAGTCCCCAGCACTGTTCT; primers for exon 2 and flanking region were: AGGGGAGGATAGGGTTCAGTG and CCTGTAGTTACCTCCCACA; and primers for exon 3 and the 3'-UTR were: ATACGTGCGTAAGCCACCAC and ACCCAGATATGCCCTTAG. Sequencing was via high throughput capillary electrophoresis. Genotyping for these SNPs was then performed in the remaining patients.

Assay of known SNPs was via real time PCR. The assay reagents for SNP genotyping were obtained from the Assay-by-Design service (Applied Biosystems, Foster City, CA). The primers and probes (FAM and VIC dye-labelled) were designed using Primer Express software and were synthesised in Applied Biosystems (table 1). PCR reactions used 2x TaqMan Universal PCR Master Mix, No AmpErase UNG (Applied Biosystems). The real time PCR reactions were performed in an Opticon II system (Bio-Rad Laboratories/MJ Research, Waltham, MA). Conditions were as follows: 95°C 10 min preheat, 92°C 15 s, 60°C 1 min followed by a plate reading, 40 cycles. The results were read according to the signal value of FAM and VIC dye.

Statistical methods

Each of the SNPs was first assessed to determine if the observed genotype frequencies were consistent with expected Hardy-Weinberg proportions using Pearson's χ^2 tests. Pairwise marker-marker linkage disequilibrium was assessed using Lewontin's D' statistic¹⁴ and Devlin and Risch's Δ^2 statistic¹⁵ as implemented in the computer program GOLD (www.sph.umich.edu/csg/abecasis/GOLD).

To test for association between individual SNPs and the continuously distributed outcomes, average warfarin dose, and average INR, we performed analysis of covariance models (ANCOVA) using PROC GLM within the SAS software system (version 8.0, SAS Institute, Cary, NC). The outcomes were first examined for adherence to distributional assumptions (including approximate normality of error terms conditional on covariates and homoscedasticity); the natural log transformation was subsequently applied to average dosage to adhere to these assumptions. Covariate adjustment

was made for age, gender, and target INR in all analytic models. Genotype was tested for general association (no mode of inheritance assumption) using a 2 df F test. A 1 df F test was used to test the null hypotheses of no association for the *CYP2C9* genotypes due to a lack of observed homozygotes for the *CYP2C9**2 or *3 alleles. We additionally examined a two locus model, in which we selected the most statistically significant polymorphism from each of the two genes and tested the effects of each while controlling for the other. Finally, a model testing the interaction effects between the two polymorphisms was performed.

Tests of haplotype effects were performed using a score test developed by Schaid, as implemented in the computer program HAPLO.STAT (<http://mayoresearch.mayo.edu/mayo/research/biostat/schaid.cfm>). HAPLO.STAT utilises a weighting scheme based on EM derived haplotype frequency estimates and weights every haplotype rather than assigning a "most likely" haplotype to an individual. Statistical differences in overall haplotype frequencies (excluding haplotypes with extremely low frequencies, for example 0.01) were tested, after adjustment for covariates (age, gender, and target INR), for association with the outcomes. Specific individual haplotype effects were also tested.

RESULTS

The sample consisted of 93 European-American subjects with both genotype and phenotype data. The mean age was 63 years (SD 16, range 24–90 years) and 31% (n = 29) of the subjects were female. Mean treatment period for this study was 20.6 months (SD 11.0, range 12 days to 37.3 months). Indications for warfarin were predominantly atrial fibrillation or atrial flutter (45%, n = 47), with another large group (26%, n = 24) being treated or prophylaxed for venous thromboembolic events (VTE). Thirteen (14%) of the patients were receiving warfarin due to prosthetic mitral or aortic valves. The target INR for the majority of patients (83%, n = 77) was 2–3. Three of the prophylaxed VTE patients had a goal of 1.5–2.0. Thirteen patients had a goal of 2.5–3.5; this group included those with a prosthetic mitral valve as well as those with both a prosthetic atrial valve and atrial fibrillation. The positions and the corresponding observed allele frequencies of the identified SNPs in all 93 patients are given in table 2. The observed distributions of genotype data for the *VKORC1* and *CYP2C9* SNPs were consistent with expected Hardy-Weinberg proportions (data not shown).

Table 3 displays the association results of the six *VKORC1* polymorphisms with mean weekly warfarin dose and INR in these patients. As expected, the observed INRs cluster tightly around 2.5 since warfarin doses were continually adjusted clinically in order to achieve an appropriate level of anticoagulation, with an INR of 2–3 being the most common target. Five of the six *VKORC1* SNPs were found to be statistically associated with average warfarin dose required to achieve the target INR (p ranging from <0.0001 to 0.021). The *VKORC1**1173, *VKORC1**1542, and *VKORC1**2255 SNPs were found to be the most strongly associated with the average warfarin dose required to achieve the target INR

Table 1 The primers and probes used in real time PCR genotyping

VKOR SNPs	VIC probe sequence	FAM probe sequence	Forward primer	Reverse primer
VKOR 497 T>G	CCCCTTC _a CCTGCGC	CCCCTTC _c CCTGCGC	GCGGTAGAGATTGACGATGGT	GCAGCCATCGCCAACAC
VKOR 698 C>T	CAAGGCT _g GTATAACG	CAAGGCT _a GTATAACG	CTCTGATGCCAAAACCGAGTGAAC	GCCCGGCCCTTAAGTAATTCTT
VKOR 1173 C>T	CCTAGTCCAAG _g GTCTGAT	CTAGTCCAAG _a GTCTGAT	CCCCGGTGCCAGGAGATC	CACCTGGGCTATCCTCTGTTC
VKOR 1542 G>C	TCATCAC _g GAGCGTC	TCATCAC _c GAGCGTC	GGTGATCCACACAGCTGACA	CCTGTAGTTACCTCCCCACATC
VKOR 2255 T>C	CCAGGACC _a TGGTGC	CCAGGACC _g TGGTGC	GCTCCAGAGAAGGCATCACT	GCCAAGTCTGAACCATGTGTCA
VKOR 3730 G>A	ATACCC _g CACATGAC	CATACCC _a CACATGAC	GTCCCTAGAAGGCCCTAGATGT	GTGTGGCACATTTGGTCCAIT

Table 2 *VKORC1* and *CYP2C9* SNP major allele frequencies

SNPs	Position	NCBI dbSNP rs#	Frequency (%) of wild type allele
<i>VKORC1</i> *497 T>G	Intron 1	Rs2884737	68.8
<i>VKORC1</i> *698 C>T	Intron 1	Rs17708472	79.6
<i>VKORC1</i> *1173 C>T	Intron 1	Rs9934438	54.8
<i>VKORC1</i> *1542 G>C	Intron 2	Rs8050894	56.5
<i>VKORC1</i> *2255 T>C	Intron 2	Rs2359612	44.1
<i>VKORC1</i> *3730 G>A	3'-UTR	Rs7294	64.5
<i>CYP2C9</i> *2	Exon 3, C144R	Rs1799853	92.5
<i>CYP2C9</i> *3	Exon 7, L359I	Rs1057910	93.0

Six SNPs were identified in the *VKORC1* gene upon sequencing and real time PCR among 93 individuals. The location and allelic frequency of each SNP is shown.

($p < 0.0001$). The trend in mean warfarin dosage by all three genotypes is consistent with an additive mode of inheritance, where the mean for the heterozygote is roughly midway between the two homozygotes

The *VKORC1**1173, *1542, and *2255 SNPs were found to be in very strong linkage disequilibrium (LD) with each other, with estimated D' values of 1.00 and estimated Δ^2 values ranging from 0.94 to 0.98. Estimated D' values ranged from 0.84 to 1.00 and estimated Δ^2 values ranged from 0.11 to 0.56 for all other SNP pair combinations. Haplotype analysis results are reported in table 4. Four haplotypes were observed with an estimated frequency greater than 3%. The permutation based global p value for any difference between

Table 3 Individual *VKORC1* gene SNP association analysis results

<i>VKORC1</i> gene SNPs	n	Average (least square means) weekly warfarin dose (mg)	INR
497 T>G			
TT	43	43.6	2.5
TG	42	34.4	2.7
GG	8	24.9	2.7
p value		0.0015	0.12
698 C>T			
CC	59	35.2	2.7
CT	30	40.0	2.5
TT	4	54.7	2.4
p value		0.056	0.18
1173 C>T			
CC	31	47.1	2.5
CT	40	35.8	2.6
TT	22	26.9	2.7
p value		<0.0001	0.10
1542 G>C			
GG	32	46.4	2.5
GC	41	36.0	2.6
CC	20	26.5	2.7
p value		<0.0001	0.11
2255 T>C			
TT	20	26.5	2.7
TC	42	35.7	2.6
CC	31	47.1	2.5
p value		<0.0001	0.11
3730 G>A			
GG	40	33.3	2.6
GA	40	39.7	2.6
AA	13	46.4	2.5
p value		0.021	0.63

Least-square means are on the untransformed data and p values were obtained using the log transformation. All models adjusted for age and gender. Models for average weekly dose of warfarin additionally adjusted for target INR.

haplotypes was 0.0004. All four of the individual haplotype specific tests were found to be statistically significant ($p < 0.05$). The two haplotypes (EA1 and EA2) characterised by the T allele for *VKOR* 1173, the C allele for *VKOR* 1542, and the T allele for *VKOR* 2255 were associated with negative score statistics (indicating lower mean dosage), while the other two haplotypes were associated with positive score statistics (indicating higher mean dosage). Overall, the haplotype results are consistent with the single SNP results and suggest that the causal polymorphism is one of the identified SNPs (*VKORC1**1173, *1542, or *2255) or another polymorphism in LD with these three SNPs.

The *CYP2C9* SNP association results are presented in table 5. Only 14 and 13 heterozygotes were observed for *CYP2C9**2 and *CYP2C9**3 polymorphisms, respectively. There was no evidence for an association between *CYP2C9* genotypes and average warfarin weekly dose before additional adjustment for *VKORC1* genotype. *CYP2C9**3 became marginally statistically significant ($p = 0.050$) after adjustment for *VKOR* 1173. The *CYP2C9**2 and *CYP2C9**3 SNPs were not found to be in significant LD with one another ($D' = 0.04$, $\Delta^2 = 0.00$, $p = 0.97$). Given the lack of LD between the two SNPs, haplotype analyses were not performed.

DISCUSSION

We have here shown that variants within *VKORC1* are strongly associated with the mean weekly dose of warfarin required to maintain a desired target INR in a sample of European-American outpatients undergoing anticoagulation treatment. In addition to their robust statistical significance, the results presented here are potentially clinically meaningful given the notoriously narrow therapeutic window¹⁶ of the drug in question. The mean weekly dose of warfarin required to maintain the desired target INR is almost doubled (from 27 to 47 mg) for those who are homozygous C/C at position 1173, G/G at position 1542, or C/C at position 2255, as compared with the homozygotes for the opposite allele at each site. Also bolstering the significance of these results is the dose-response curve observed in heterozygotes, who demonstrate intermediate warfarin sensitivity. Moreover, these alleles are common in this unselected sample with an estimated prevalence of 54.2% for the *VKORC1**1173 C allele, 55.7% for the *VKORC1**1542 G allele, and 54.7% for the *VKORC1**2255 C allele. These results are strongly consistent with recent findings regarding the effects of *VKORC1* genotype on warfarin dosage levels required to reach desired INR both in terms of overall statistical significance and mean warfarin dosage levels by genotype.⁷⁻¹² For example, D'Andrea *et al*⁷ report an observed warfarin mean weekly dosage of 49.0, 35.7, and 25.9 (calculated by multiplying their reported mean daily dosage by 7) for *VKORC1**1173 genotypes C/C, C/T, and T/T, respectively, as compared with our observed weekly mean dosages of 47.1, 35.8, and 26.9, respectively, for the same genotypes. In addition, there is considerable agreement with the direction of the haplotype based associations. In our study, EA1 and EA2 are associated with lower mean levels of warfarin dosage and haplotypes EA3 and EA4 are associated with higher mean levels of warfarin dosage. Our haplotypes EA1 and EA2 have identical alleles as do haplotypes H3a and H3b in Wadelius *et al*¹⁰ for *VKORC1**1173, *2255, and *3730. Our haplotype EA3 corresponds to H2 and our haplotype EA4 corresponds to H1 in Wadelius *et al*,¹⁰ where H1 and H2 are also associated with higher warfarin dosage. Finally, our haplotypes EA1, EA2, EA3, and EA4 correspond to the sets of haplotypes {H2, H5}, {H1, H3}, {H4, H6, H9}, and {H7, H8}, respectively, in Rieder *et al*⁸; the direction of results between the two studies with respect to which haplotypes are associated with higher or lower warfarin dosage levels is also entirely consistent.

Table 4 Association between VKORC1 gene haplotypes and log average dose

Haplotype	VKORC1 gene SNPs						Estimated frequency (%)	Score*	p†
	497	698	1173	1542	2255	3730			
EA1	G	C	T	C	T	G	30.1	-3.42	0.00063
EA2	T	C	T	C	T	G	12.9	-2.26	0.024
EA3	T	T	C	G	C	G	19.3	2.47	0.013
EA4	T	C	C	G	C	A	34.9	2.73	0.0063
Global‡									0.0004

*HAPLO.STAT (<http://mayoresearch.mayo.edu/mayo/research/biostat/schaid.cfm>) program statistic; †haplotype specific empirical p values comparing each specific haplotype to all others combined; ‡simulated p value for global association test.

In addition to our VKORC1 results, we did not detect a statistically significant association between the two CYP2C9 polymorphisms and warfarin dosage before adjustment for VKORC1; however, results for CYP2C9*3 did reach a marginal level of statistical significance (p = 0.05) after adjustment for VKORC1*1173. Because the CYP2C9 SNPs have lower minor allele frequencies than the VKORC1 SNPs examined, the power to detect significant departures from the null hypothesis of no association for CYP2C9 was less than that for VKORC1.

In previously reported work, Sconce *et al*⁹ demonstrated the best multivariable regression model for fitting warfarin dosage included both CYP2C9 and VKORC1. Our multivariable model including age, gender, target INR, and VKORC1*1173 genotype explained approximately 30% (R² = 0.30) of our total observed variation. Our multivariable model including age, gender, target INR, and CYP2C9*3 explained only 12% of the total variation. Including both CYP2C9*3 and VKORC1*1173 along with our covariates explained 34% of the total variation. Thus, our multivariable model appeared to explain less of the total variation in the mean warfarin dosage required to reach the target INR than the multivariable regression model of Sconce *et al*⁹ (R² = 0.55). It should be noted that their model included height, a measure that we did not have available to us for this study. In addition, our results suggest, consistent with the findings of Wadelius *et al*¹⁰, that VKORC1 polymorphisms explain a larger amount of the observed variation in mean warfarin dosage required to reach the target INR than do CYP2C9 polymorphisms in the European-American population.

During the course of our study, we also collected data on warfarin dosage requirements to reach the target INR for 18 African-Americans. Of these 18 patients, one had a target INR of 1.5–2.0, 14 had a target INR of 2.0–3.0, and three had a target INR of 2.5–3.5. Marginally statistically significant

results were observed for VKORC1*1173 (p = 0.03) and *2255 (p = 0.03) and CYP2C9*3 (p = 0.01) in this limited data set. These results could be entirely explained by the low observed warfarin dosages (average weekly dosage of 9.3 mg) of a single individual who was homozygous T/T at both VKORC1*1173 and *2255 and heterozygous at CYP2C9*3. This individual's genotypes at these three markers were consistent with the genotypes that were shown to be associated with low warfarin dosage in the European-American population. In addition, this individual had the lowest target INR (1.75) and was also the only African-American individual in our sample with these aforementioned genotypes. We note that in our limited sample, there were strong differences in allele frequencies (data not shown) between the European-American and African-American samples. For example, VKORC1*698 and CYP2C9*2 were uninformative (completely homozygous for the wild type allele) in this small African-American sample. We also note that we observed weaker marker-marker LD and a larger diversity in haplotypes in this limited African-American sample (data not shown). These findings are all consistent with the comparative results between the European and African-Yoruban samples genotyped at VKORC1 as part of the International HapMap Project (www.hapmap.org). While these results based on 18 African-American patients are limited and inconclusive, they do suggest that it is important to assess the role of VKORC1 SNPs in warfarin response in other populations and that studying different populations with different haplotype patterns may be critical in ultimately identifying the functionally important variants.

It remains undetermined whether one or more of the SNPs identified here or elsewhere are causal in producing differential warfarin sensitivity or whether they are in LD with the actual causative SNP. Strong LD between VKORC1 polymorphisms in the European-American population hinders our ability to fine map/identify the functional loci. Further analysis of warfarin related gene(s) and surrounding sequence and the inclusion of larger numbers of patients, ideally in different populations, may lead to elucidation of the specific genotypic factors which influence warfarin sensitivity. In addition, prospective studies will ultimately be critical to establishing the clinical utility of assessing VKORC1 genotype for determining optimal warfarin dosage. Nonetheless, combining genotype data from SNPs of the VKORC1 gene with SNPs of the cytochrome CYP2C9 gene in a clinical setting promises to result in a more robust assessment of warfarin sensitivity.

The results presented here support those from other recent reports⁷⁻¹² and further demonstrate the need for prospective studies to evaluate the utility of VKORC1 and CYP2C9 genotype information with respect to outcomes such as length of time until the target INR is reached and adverse events. This current demonstration of common polymorphisms that predict warfarin sensitivity represents a potentially clinically useful proof of principal for the use of pharmacogenomic information in medicine and will likely play a

Table 5 Association between CYP2C9 genotypes and log average dose in European-American subjects

CYP2C9 gene SNPs	n	Least square means		
		Average weekly dose (mg)	Average weekly dose (mg) after adjustment for VKORC1*1173	INR
CYP2C9*2				
11	79	38.8	37.3	2.6
12	14	32.1	31.0	2.6
22	0	–	–	–
p value		0.19	0.16	0.93
CYP2C9*3				
11	80	38.5	37.1	2.6
12	13	33.0	31.5	2.7
22	0	–	–	–
p value		0.24	0.050	0.12

All models adjusted for age and gender. Models for average weekly dose of warfarin were additionally adjusted for the target INR.

critical role in future predictive models for warfarin dosage. Such models will have strong potential to guide physicians in quickly prescribing appropriate dosages and, ultimately, avoid complications.

ELECTRONIC-DATABASE INFORMATION



Information about the computer program GOLD is at www.sph.umich.edu/csg/abecasis/GOLD, the computer program HAPLO.STAT at <http://mayoresearch.mayo.edu/mayo/research/biostat/schaid.cfm>, and the International HapMap Project at www.hapmap.org.

Authors' affiliations

T Li, Gene Expression Laboratory, The Salk Institute for Biological Studies, La Jolla, CA, USA

L A Lange, L Susswein, E M Lange, J P Evans, Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

X Li, Department of Immunology, The Scripps Research Institute, La Jolla, CA, USA

B Bryant, R Malone, Department of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

T-Y Huang, D W Stafford, Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

Competing interests: none declared

This work was carried out at the University of North Carolina at Chapel Hill

REFERENCES

- Shulman S**. Oral anticoagulation. In: Beutler E, Lichtman MA, Coller BS, Kipps TJ, Seligsohn U, eds. *Williams hematology*. 6th ed. New York: McGraw-Hill, 2001:1777–92.
- Levine MN**, Raskob G, Landefeld S, Kearon C. Hemorrhagic complications of anticoagulant treatment. *Chest* 2001;**119**:108S–21S.
- Poller L**, Shiach CR, MacCallum PK, Johansen AM, Munster AM, Magalhaes A, Jespersen J. Multicentre randomised study of computerised anticoagulant dosage. European Concerted Action on Anticoagulation. *Lancet* 1998;**352**:1505–9.
- Doecke CJ**, Cosh DG, Gallus AS. Standardised initial warfarin treatment: evaluation of initial treatment response and maintenance dose prediction by randomised trial, and risk factors for an excessive warfarin response. *Aust N Z J Med* 1991;**21**:319–24.
- Li T**, Chang CY, Jin DY, Lin PJ, Khvorova A, Stafford DW. Identification of the gene for vitamin K epoxide reductase. *Nature* 2004;**427**:541–4.
- Rost S**, Fregin A, Ivaskevicius V, Conzelmann E, Hartnagel K, Pelz HJ, Lappégard K, Seifried E, Scharer I, Tuddenham EG, Muller CR, Strom TM, Oldenburg J. Mutations in VKORC1 cause warfarin resistance and multiple coagulation factor deficiency type 2. *Nature* 2004;**427**:537–41.
- D'Andrea G**, D'Ambrosio RL, Di Perna P, Chetta M, Santacroce R, Brancaccio V, Grandone E, Margaglione M. A polymorphism in the VKORC1 gene is associated with an interindividual variability in the dose-anticoagulant effect of warfarin. *Blood* 2005;**105**:645–9.
- Rieder MJ**, Reiner AP, Gage BF, Nickerson DA, Eby CS, McLeod HL, Blough DK, Thummel KE, Veenstra DL, Rettie AE. Effect of VKORC1 haplotypes on transcriptional regulation and warfarin dose. *N Engl J Med* 2005;**352**:2285–93.
- Sconce EA**, Khan TI, Wynne HA, Avery P, Monkhouse L, King BP, Wood P, Kesteven P, Daly AK, Kamali F. The impact of CYP2C9 and VKORC1 genetic polymorphism and patient characteristics upon warfarin dose requirements: proposal for a new dosing regimen. *Blood* 2005;**106**(7):2329–33.
- Wadelius M**, Chen LY, Downes K, Ghori J, Hunt S, Eriksson N, Wallerman O, Melhus H, Wadelius C, Bentley D, Deloukas P. Common VKORC1 and GGCCX polymorphisms associated with warfarin dose. *Pharmacogenomics J* 2005;**5**(4):262–70.
- Yuan HY**, Chen JJ, Lee MT, Wung JC, Chen YF, Charng MJ, Lu MJ, Hung CR, Wei CY, Chen CH, Wu JY, Chen YT. A novel functional VKORC1 promoter polymorphism is associated with inter-individual and inter-ethnic differences in warfarin sensitivity. *Hum Mol Genet* 2005;**14**:1745–51.
- Lee SC**, Ng SS, Oldenburg J, Chong PY, Rost S, Guo JY, Yap HL, Rankin SC, Khor HB, Yeo TC, Ng KS, Soong R, Goh BC. Interethnic variability of warfarin maintenance requirement is explained by VKORC1 genotype in an Asian population. *Clin Pharmacol Ther* 2006;**79**:197–205.
- Sanderson S**, Emery J, Higgins J. CYP2C9 gene variants, drug dose, and bleeding risk in warfarin-treated patients: a HuGENet systematic review and meta-analysis. *Genet Med* 2005;**7**:97–104.
- Lewontin RC**. The interaction of selection and linkage. I. General considerations. *Genetics* 1964;**49**:49–67.
- Devlin B**, Risch N. A comparison of linkage disequilibrium measures for fine-scale mapping. *Genomics* 1995;**29**:311–22.
- Schulman S**. Clinical practice. Care of patients receiving long-term anticoagulant therapy. *N Engl J Med* 2003;**349**:675–83.