

SUPEROXIDE DISMUTASE, LIPID PEROXIDATION, AND BELL-SHAPED DOSE RESPONSE CURVES

Joe M. McCord □ Division of Pulmonary and Critical Care Medicine, University of Colorado Denver Health Sciences Center, Denver, CO

□ Cellular metabolism generates the cytotoxic superoxide free radical, $O_2^{\cdot-}$, and a family of enzymes called superoxide dismutases (SOD) protects us from $O_2^{\cdot-}$ by catalyzing its conversion to O_2 and H_2O_2 . Superoxide production increases in a wide variety of pathological states, especially those involving inflammation or ischemic injury. Most of the literature has described systems wherein added or over expressed SOD produced beneficial effects, yet in some circumstances SOD provided no benefit, or was clearly detrimental, exacerbating cell injury or death. When broad dose-response studies were finally possible in models of reperfusion injury in the isolated heart, hormesis became clear. We propose that the mechanisms underlying the hormesis are related to the paradoxical abilities of the superoxide radical to serve as both an initiator and a terminator of the free radical-mediated chain reaction that results in lipid peroxidation. Lipid peroxidation is a universal feature of oxidative stress, causing loss of cellular structure and function. Under any given conditions, the optimal concentration of SOD is that which decreases chain *initiation* without elimination of the chain *termination* properties of the radical, resulting in a minimum of net lipid peroxidation. Mathematical modeling of this hypothesis yields predictions fully consistent with observed laboratory data.

Keywords: superoxide dismutase, lipid peroxidation, dose-response, reperfusion injury, free radical, hormesis

INTRODUCTION

The increased production of the superoxide free radical has come to occupy a remarkably central role in a wide variety of diseases, as documented by thousands of published studies (McCord, 2000). Superoxide is not, as first thought, just a toxic but unavoidable byproduct of oxygen metabolism. Rather, we now appreciate that it is a carefully regulated metabolite capable of signaling and communicating important information to the cell's genetic machinery. Redox regulation of gene expression by superoxide and other related oxidants and antioxidants has become a mechanism of great interest in health and disease (Rushmore and Kong, 2002).

Surprisingly, about 98% of the oxygen we metabolize is handled by a single enzyme—the cytochrome oxidase in our mitochondria, which transfers four electrons to oxygen in a concerted reaction to produce two

Address correspondence to Joe M. McCord, Division of Pulmonary and Critical Care Medicine, University of Colorado Denver Health Sciences Center, 4200 E. Ninth Ave., Denver, CO, 80246, Telephone: 303 315-6257, Fax: 303 315-0470, Email: joe.mccord@uchsc.edu





FIGURE 1. A schematic diagram of the mitochondrial electron transport system illustrating that at least two specific “leakage sites” exist in injured mitochondrial allowing electrons to prematurely reach oxygen molecules, giving rise to the superoxide radical. SOD and catalase normally detoxify the radical. If superoxide concentration rises too high, oxidation of proteins, lipids, and DNA may occur and various signalling pathways may be triggered.

molecules of water as the product. The probable reason for this near monopoly in the reduction of oxygen is the chemical difficulty of carrying out this combustion reaction in a safe and controlled manner. If oxygen receives less than the full complement of four electrons, the result is the production of active oxygen species $O_2\cdot^-$ and H_2O_2 . Even in a healthy cell approximately 1 or 2% of all oxygen metabolized might be converted directly to superoxide and/or hydrogen peroxide (Chance *et al.*, 1979), reflecting the ability of molecular oxygen to “steal” electrons from a variety of cellular reductants via spontaneous (non-enzyme-catalyzed) oxidation. Thus, even though our electron-conducting metabolic circuits are generally *well-insulated* by evolutionary design to prevent the accidental development of short-circuits, our insulation is not perfect. At least two sites have been identified in the electron-transport chain (Complex I and ubiquinone) where electrons may leak out to oxygen molecules, resulting in the formation of superoxide (McCord and Turrens, 1994). These sites are shown in Fig. 1 as the primary sources of intracellular superoxide generation. In a healthy cell, the superoxide produced is detoxified by the antioxidant enzyme SOD (McCord and Fridovich, 1969), which catalyzes the reaction:



This dismutation or disproportionation reaction makes use of the fact that the superoxide radical can react either as an oxidant or a reductant, eager to get rid of its extra electron or to take on another. The enzyme uses one superoxide radical to oxidize another. Catalases work in much

the same way to detoxify the resulting H_2O_2 , because hydrogen peroxide can be a weak reductant as well as a fairly strong oxidant:




Thus, SOD and catalase working together convert superoxide to the innocuous products O_2 and H_2O (Fig. 1).

Working myocardium consumes oxygen at about 8 mM/min, and therefore the rate of superoxide production in a healthy cell might reach 0.1 mM/min. During exposure to hyperoxia, the rates of leakage from these sites in lung mitochondria are believed to increase in direct proportion to the increased oxygen tension. Healthy adult rats die after about 72 hours in an atmosphere of 100% oxygen—only five times the normal concentration at sea level (Crapo and Tierney, 1974). This accounts for the hormesis observed with increasing oxygen concentrations.

ISCHEMIA/REPERFUSION INJURY

In 1981 intravenously administered SOD was first shown to be protective in a model of feline intestinal partial ischemia/reperfusion injury (Granger *et al.*, 1981), implying that increased superoxide production might be responsible in large part for the post-ischemic destruction of tissues as occurs, for example, in stroke and myocardial infarction. In the intestine, the major source of superoxide was concluded to be from the enzyme xanthine dehydrogenase, normally an innocuous NAD^+ -reducing enzyme involved in the catabolism of purines. During ischemia, however, tissues are unable to maintain ATP levels, allowing Ca^{++} to enter the cytosol as there is too little energy available to power the divalent cation pump at the cell membrane. This intracellular Ca^{++} can activate proteases such as calpain, which in turn can partially proteolyze xanthine dehydrogenase converting it into xanthine oxidase. Xanthine oxidase no longer reduces NAD^+ , but rather transfers electrons to molecular oxygen to produce the superoxide radical:



During ischemia, an abundant amount of hypoxanthine builds up as AMP is catabolized. Upon reoxygenation of the tissue the missing substrate, O_2 , is supplied, allowing a burst of superoxide production (McCord, 1985). In tissues rich in xanthine dehydrogenase such as the intestine, this seems to be the major source of reoxygenation injury. In myocardial reperfusion injury of rodents and dogs, this also appears to be the predominant mechanism of injury. Hearts of humans (Eddy *et al.*, 1987) and rabbits (Terada *et al.*, 1991), however, contain only traces of xanthine oxidase activity, and this mechanism appears to make a much

smaller contribution to reoxygenation injury, but one that may still be significant (Terada *et al.*, 1991; Zhao *et al.*, 2008).

In tissues or species that contain little xanthine dehydrogenase, SOD is still dramatically protective against reoxygenation injury (Nelson *et al.*, 1994), implying that another damaging source of superoxide exists. This alternate source appears to be much more basic—the resumption of mitochondrial respiration itself (Ambrosio *et al.*, 1993). As intracellular calcium levels rise in energy-compromised ischemia tissues, the calcium is avidly taken up by the mitochondria leading to uncoupling, futile cycling, and increase production of superoxide (Paraidathathu *et al.*, 1992). This mechanism of reoxygenation-induced superoxide production is likely common to all mammalian cells.

CONTROVERSY AND HORMESIS

The possibility that reoxygenated tissues might be significantly protected by SOD quickly captured the interest of experimental cardiologists and myocardial physiologists. As various laboratories began studying various models, controversy soon appeared in the literature. There was little agreement as to whether SOD did (Ambrosio *et al.*, 1986; Aoki *et al.*, 1988; Burton, 1985; Chambers *et al.*, 1985; Gardner *et al.*, 1983; McCord *et al.*, 1989; Tamura *et al.*, 1988; Werns *et al.*, 1985; Woodward and Zakaria, 1985) or did not (Buchwald *et al.*, 1989; Menasche *et al.*, 1986; Myers *et al.*, 1985; Uraizee *et al.*, 1987) protect the reoxygenated heart. There were many differences among the many models, including *in vitro* versus *in vivo*, global versus regional ischemia, partial versus complete ischemia, buffer perfused versus blood perfused, and, importantly, the concentration of SOD used. Because of the expense and labor-intensity of the models, none of these cited studies examined dose-response relationships. One of the first studies to vary the dose of SOD was quite illuminating, as U-shaped curves were apparent when SOD was used to prevent the occurrence of reperfusion-induced arrhythmias (ventricular fibrillation and tachycardia) following 10 min of coronary artery occlusion in isolated rat hearts (Bernier *et al.*, 1989). We soon reported similarly shaped curves with regard to creatine kinase release from hypoxic/reoxygenated isolated rat hearts (Fig. 2A) and lactate dehydrogenase release from globally ischemic/reperfused rabbit hearts (Fig. 2B), or area of infarct in an *in vivo* model of coronary artery ligation in rabbits (Fig. 2C) (Omar *et al.*, 1990). It is remarkable to note the consistent loss of protection at higher doses of SOD, even though the models differ considerably with respect to species (rat versus rabbit), end point measured (infarct size, enzyme release, arrhythmogenesis, or recovery of developed ventricular pressure), source of the SOD used (yeast in the rat and human recombinant in the rabbit) and its schedule of administration (continuous versus a single bolus).




FIGURE 2. Bell-shaped dose-response curves have been reported in various models reflecting the ability of SOD to protect or exacerbate injury to the post-hypoxic heart. Data are re-plotted from Omar *et al.* (1990) and Omar and McCord (1990). In Panel A, the black square represents data from Menasche *et al.* (1986) using a similar model. Similarly, in Panel B the black square represents data from Myers *et al.* (1985) using a similar model. Both of these studies reported no significant protection from SOD at these single doses.

Furthermore, our curves could clearly have predicted the outcomes of some reported negative studies in the literature using the isolated rat (Menasche *et al.*, 1986) (see Fig. 2A) and rabbit (Myers *et al.*, 1985) hearts (see Fig. 2B).

HOW CAN TOO MUCH SOD BE DAMAGING?

Several potential explanations have been offered to explain the hormetic behavior of SOD. The most common explanation is that since SOD produces H_2O_2 as a product, more SOD would produce more H_2O_2 , which is toxic to cells (Kedziora and Bartosz, 1988; Mao *et al.*, 1993; Norris and Hornsby, 1990). This explanation does not hold when the SOD-catalyzed reaction is competing for superoxide with spontaneous (uncatalyzed) dismutation, for the net reaction *rates* are identical in the steady-state. The molar rate of H_2O_2 production is exactly one-half the molar





FIGURE 3. Bell-shaped dose-response curves produced by SOD in a simple system consisting of 3 mM linolenic acid in the presence of 4.65 μM FeCl_3 and 1 mM xanthine. Superoxide was generated by the indicated concentrations of xanthine oxidase, and inhibition of malondialdehyde formation was noted at the indicated concentrations of SOD. Data are re-plotted from Lazzarino *et al.* (1994).

rate of $\text{O}_2\cdot^-$ production, regardless of the concentration of SOD (McCord and Fridovich, 1969). SOD will dramatically lower the *steady-state concentration* of superoxide, but obviously cannot alter the steady-state rate of its conversion to H_2O_2 , as this is limited by the rate of superoxide production. It has also been suggested numerous times that the toxicity of high-dose SOD might be due to a weak peroxidase activity of Cu,Zn-SOD (Hodgson and Fridovich, 1975). As seen in Fig. 2D, however, Mn-SOD shows the same bell-shaped dose-response (Omar and McCord, 1990) as those produced by Cu,Zn-SOD (Fig. 2A, 2B, and 2C) (Omar *et al.*, 1990), despite its lack of such a peroxidase activity (Hodgson and Fridovich, 1975).



We hypothesized an alternate explanation based not upon the properties of SOD *per se*, but rather upon a major physiological process being modulated by changes in the concentration of superoxide in a cell—lipid peroxidation. Even in a very simple chemically defined system, SOD displays bell-shaped dose-response curves *if that system involves lipid peroxidation*. An example is the oxidation of pure linolenic acid by xanthine/xanthine oxidase plus ferric iron (Lazzarino *et al.*, 1994). Fig. 3 shows a series of hormetic curves as the concentration of SOD is varied together with the rate of $\text{O}_2\cdot^-$ generation in such a system. At each higher rate of $\text{O}_2\cdot^-$ generation a new optimally protective concentration is defined. This optimally protective SOD concentration varies linearly with rate of $\text{O}_2\cdot^-$ production over several orders of magnitude. In simple systems that do not involve lipid peroxidation such as the inactivation of creatine kinase by xanthine/xanthine oxidase, SOD shows nearly complete protection with no indication of hormesis (McCord and Russell, 1988).

SUPEROXIDE RADICAL INITIATES AND TERMINATES LIPID PEROXIDATION

Like all other free radical chain reactions, lipid peroxidation must be initiated by a free radical, (i.e., by a one-electron oxidation) in this case abstracting an allylic hydrogen from a polyunsaturated side chain to create a lipid radical ($L\cdot$). The protonated form of superoxide, the hydroperoxyl radical ($HO_2\cdot$), can serve as an initiating radical due to its lipophilicity and redox potential:



The lipid radical $L\cdot$ adds molecular oxygen extremely rapidly to produce a lipid dioxygen radical, which is a sufficiently good oxidant to abstract an allylic hydrogen from another unsaturated side chain, producing a lipid hydroperoxide (LOOH) and propagating the chain:



Typically *in vivo* these two reactions cycle about 15 times, such that each initiating event results in the oxidation of about 15 polyunsaturated fatty acid molecules. (The $LOO\cdot$ radical may alternatively rearrange and break down to produce malondialdehyde and other radical products, rather than yielding exclusively a lipid hydroperoxide product, but these options do not affect the present discussion.) Hence, it is well-established that superoxide can initiate lipid peroxidation. Ironically, the only way to eliminate a radical is by reacting it with another radical. (Because radicals have an odd number of electrons, any reaction with a non-radical must yield a product that is another radical.) Thus, from a teleological perspective, it would be useful if a cell had available a continuous supply of a relatively unreactive free radical for the purpose of promoting a radical-radical annihilation to eliminate the very dangerous chain-propagating lipid dioxygen radicals. We have suggested that superoxide meets these criteria (Nelson *et al.*, 1994). It is produced as a by-product of oxygen metabolism, and is therefore continuously available in all cells. As free radicals go, it is relatively unreactive. Its concentration is maintained at a very low, but non-zero value by the cellular SOD content. If lipid dioxygen radicals were scavenged by $O_2\cdot^-$, then entire chains of reactions would be prevented or terminated:






FIGURE 4. Relationships of net lipid peroxidation and functional recovery of isolated rabbit hearts to administered dosages of SOD following ischemia-reperfusion. Recovery of developed pressure correlates inversely with net lipid peroxidation as assessed by thiobarbituric acid-reactive substances in the coronary effluent. Maximal recovery of developed pressures was seen at a dosage of 5 mg SOD/L in the coronary perfusate. Data are re-plotted from (Nelson *et al.*, 1994).

Therefore, it was hypothesized that $O_2^{\cdot-}$, in addition to being able to liberate iron and to initiate lipid peroxidation, may also serve as a terminator of lipid peroxidation, such that over scavenging the radical may increase net lipid peroxidation (Nelson *et al.*, 1994; Omar *et al.*, 1990). A functionally similar termination reaction was proposed by Thom and Elbuken (Thom and Elbuken, 1991) whereby the protonated form of superoxide, HO_2^{\cdot} , reacts with LOO^{\cdot} to produce water, oxygen, and a carbonyl containing lipid peroxidation product. If either hypothesis is correct, then indices of lipid peroxidation should be high at both low concentrations of SOD (where initiation would be high) and at high concentrations of SOD (where termination would be low). At some intermediate SOD concentration, initiation by superoxide would be largely suppressed, but termination by superoxide would still be making a contribution, and net lipid peroxidation would be at a minimum. We found these predictions to be true (Nelson *et al.*, 1994), and experimental verification is shown in Fig. 4. In this model of the isolated perfused rabbit heart, 60 min of no-flow ischemia followed by reperfusion results in serious oxidative stress. As the concentration of SOD in the perfusate is increased, a new balance is restored when the concentration reaches 5 mg/l. Note that maximal recovery of preischemic function (developed pressure) occurs when lipid peroxidation is reduced to the minimum. These results suggest that for any given level of oxidative stress, a single optimal concentration of SOD exists that will restore the superoxide level to yield the best combination of initiation and termination (i.e., the one producing




FIGURE 5. The predicted mathematical relationships between SOD concentration (which is inversely related to superoxide concentration) and lipid peroxidation. Panel A shows the relationship between SOD and initiation events, while Panel B shows the relationship between SOD and termination events (in terms of chain length). The equations for the rectangular hyperbolic functions are shown in the respective panels. As net lipid peroxidation is reflected by the product of initiation events times chain length, that resulting equation is plotted in Panel C, showing the predicted minimum at a certain concentration of SOD.

minimal net lipid peroxidation). Any concentration of SOD other than the optimal leads to increased lipid peroxidation and therefore to increased oxidative stress.

THEORETICAL PREDICTIONS

Fig. 5 presents the expected mathematical relationships between SOD concentration (which is inversely related to superoxide concentration) and initiation, termination, and net lipid peroxidation. Panel A shows that as SOD concentration increases (and $O_2^{\cdot-}$ decreases) initiation (at least that fraction dependent upon superoxide) would decrease according to an inverse rectangular hyperbola, asymptotically approaching zero initiating events at infinite SOD. Panel B shows the expected effect on chain length. At the highest concentration of $O_2^{\cdot-}$ (no SOD) the chain length would approach unity, reflecting complete suppression of propagation reactions. At the lowest concentration of the radical (high SOD) the chain length would be maximal (limited only by other terminators such as vitamin E, or by the mutual annihilation of propagating




FIGURE 6. Reconciliation of literature reports describing transgenic mice overexpressing SOD1 to different degrees. Our empirical dose-response curve from the data in Fig. 2D is seen here as the dashed curve. If pathological stress doubles the rate of superoxide production, then the curve would shift to the right (solid curve), such that optimal protection requires twice as much SOD, and 1x wild-type (wt) expression becomes inadequate. Results from three published studies of Tg-mice are shown: Point “1”, a 2-fold overexpressing mouse (Deshmukh *et al.*, 1997); Point “2”, a 2.7-fold overexpressing mouse (Chen *et al.*, 2000); and Point “3”, a 4.2-fold overexpressing mouse (Levy *et al.*, 2002). The data are normalized to their respective wild-types.

radicals). This produces another inverse rectangular hyperbolic function in the opposite direction. Net lipid peroxidation, shown in Panel C, is represented by the *product* of initiation events times chain length. Panel C shows the curves from Panels A and B replotted on a semilog coordinate, along with the curve produced by multiplying initiation by chain length. Note that the net amount of lipid peroxidation displays a minimum at a specific, optimal concentration of SOD. It is an inverted bell-shaped curve. Fig. 5 shows the remarkable similarity in shape of this theoretically predicted curve for net lipid peroxidation and that obtained experimentally by monitoring lipid peroxidation products generated by an isolated perfused heart (Fig. 4). This convinces us that the superoxide radical can act both as initiator of lipid peroxidation and terminator of lipid peroxidation. This realization provides a rational basis for the bell-shaped dose-response curves observed for SOD in systems involving lipid peroxidation, and argues that, for any specific set of conditions, there is a single, optimally protective concentration of SOD. It underscores the importance of balance between oxidants and antioxidants.

An understanding of hormetic behavior has allowed us to reconcile seemingly contradictory early studies using single doses of *exogenous* SOD in isolated organ preparations as pointed out in Figs. 2A and 2B. We can also use this understanding to reconcile seemingly contradictory studies

of transgenic (Tg) mice, where varying but fixed levels of *endogenous* SOD expression have been achieved in Tg mouse lines depending on gene copy number and promoter efficiency. Fig. 6 examines results from three such published studies of Tg mice overexpressing SOD1 to different degrees, superimposed on our own empirical dose-response curve (from the data in Fig. 2D, seen here as the dashed curve in Fig 6). This curve assumes that in a healthy wild-type mouse, the normal 1X expression level of SOD1 provides near optimal biological outcome. If we further assume that a pathological stress such as ischemia followed by reperfusion might double the rate of superoxide production in a tissue, then we can imagine that the dose-response curve would be shifted to the right (solid curve in Fig. 6), such that the optimal biological outcome would require twice as much SOD expression. Let us first look at the results of a study examining intestinal reperfusion injury in a Tg mouse line overexpressing SOD1 by 2-fold (Deshmukh *et al.*, 1997). These authors observed that the Tg mice showed only 26% as much increase in lipid peroxidation as their wild-type counterparts. These data are normalized to wild-type and plotted as the point labeled “1” in Fig. 6, falling close to its predicted position on the solid curve. Another study of a Tg mouse line overexpressing SOD1 by 2.7-fold examined myocardial ischemia-reperfusion in isolated hearts, measuring lactate dehydrogenase release as a marker of tissue damage (Chen *et al.*, 2000). Here the Tg mice showed only 40% as much enzyme release as their wild-type counterparts. These data are normalized to wild-type and plotted as the point labeled “2” in Fig. 6, again falling at its predicted position on the solid curve. In a third study of a Tg mouse line, this time overexpressing SOD1 by 4.2-fold, transient *in utero* ischemia-reperfusion was produced in pregnant mothers. After 24 h of reperfusion *in utero* the fetal brains were analyzed for lipid peroxidation products (Levy *et al.*, 2002). Here the Tg mice showed 76% *more* lipid peroxidation than their wild-type counterparts. These data are normalized to wild-type and plotted as the point labeled “3” in Fig. 6, again falling near the predicted position on the solid curve. Remarkably, very similar and predictable results are obtained whether is SOD dosed *in vivo* by transgenic manipulation or *ex vivo* in isolated organs perfused with purified exogenous SOD at various concentrations.

CONCLUSIONS AND PRACTICAL IMPLICATIONS

An obvious and far-reaching conclusion to draw from these observations is that the optimally protective concentration of SOD under any set of circumstances is at the top of the bell-shaped curve, but that the curve is not static. Rather, the curve will shift to the right as the flux of superoxide production increases due to a variety of pathophysiological conditions, including ischemia-reperfusion, inflammation, redox-cycling drugs or xenobiotics, hyperoxia, and even normal aging. Adding to the complexity

is the likelihood that the curve may shift to the right in one particular organ if the pathology is organ-specific, as would be the case following a heart attack or stroke. It is precisely these bell-shaped curves, we believe, that have made clinical applications of SOD1 or SOD2 so problematic. To maintain plasma concentrations of therapeutically administered enzymes in a range that varies by no more than two- or three-fold becomes clinically difficult with proteins that are rapidly cleared by the kidneys. It might be pointed out here that the apparent toxicity of high-dose SOD appears to be manifest only in injured tissues where lipid peroxidation is increased, and superoxide may not be the exclusive initiator. If lipid peroxidation is not ongoing, elevated concentrations of SOD appear to do no harm to tissues until they reach 5- or 6-fold above wild type.

A genetically-engineered chimeric recombinant human SOD (SOD2/3) has been produced by combining portions of the SOD2 and SOD3 genes (Gao *et al.*, 2003). This chimeric molecule has better pharmacological properties than any of the naturally occurring enzymes, and may eliminate some of the dose-response problems that have been described here (Hernandez-Saavedra *et al.*, 2005). A comparison of the effects of SOD2/3 in a variety of *in vivo* models at varying dosages produced a dramatically flattened bell curve. This may reflect the unusual properties of SOD2/3 regarding its subcellular location (on the outer surfaces of cells or bound to intracellular matrix), its size (small enough to extravasate easily), and its near-neutral charge (which facilitates both both extravasation and binding). If these factors are considered with regard to the three natural forms of SOD (SOD1 in the cytosol, SOD2 in the mitochondrial matrix, SOD3 in the extracellular and nuclear compartments) one realizes that changes in the activities of the isozymes may not produce equivalent effects with regard to hormetic behavior. That is, changing the SOD activity in any given subcellular compartment and/or binding location within a compartment may affect the shape or magnitude of the resulting hormetic curve.

While this review has focused on hormetic behavior that is strictly related to the dual effects of superoxide as initiator and terminator of lipid peroxidation, it is interesting to consider other data suggesting that hormetic behavior might also result from the signal transducing roles of the radical. As primitive life forms evolved in an increasingly aerobic environment, living with the unavoidable production of superoxide became a fact of life. The concentration of this omnipresent yet evanescent free radical afforded the cell with a realtime indicator of such important information as the availability of a food supply, reflected by mitochondrial oxidative activity. An abundance of food signals a good time for a cell to divide. Not surprisingly, cells learned to undergo mitosis when mild oxidative stress signaled an abundance of energy. One might wonder what would happen to this signaling system if the cell produced so much

SOD that this signal to divide might be extinguished. One possibility is that the cell would simply cease to divide. When scientists began to produce SOD-transfected cell lines and SOD-transgenic mice it became apparent that there was an upper limit of perhaps eight- to ten-fold to the amount of SOD over expression that could be tolerated (Norris and Hornsby, 1990). There is a growing appreciation of the roles played by superoxide and its dismutation product hydrogen peroxide in the regulation of cell growth (Burdon, 1995), while other signal transduction pathways seem to generate superoxide specifically for signaling purposes. A family of NADPH oxidases, activated by such stimuli as angiotensin II, tumor necrosis factor- α , and insulin, regulate the activation of distinct protein kinases (Anilkumar et al., 2008) that have the potential to affect vascular smooth muscle and cardiac hypertrophy, endothelial activation, angiogenesis, and atherosclerosis. Hence, the aberrant regulation of these signals by too much or too little SOD activity could, in theory, give rise to many additional instances of hormetic behavior.

An exciting alternative approach to the therapeutic use of exogenous SODs is the controlled induction of the family of antioxidant enzymes regulated directly or indirectly by the Antioxidant Response Element (ARE). The ARE is a promoter element activated by the transcription factor Nrf2. A number of phytochemicals long known to traditional medicine appear to be a safe and effective means of upregulating these enzymes in humans via Nrf2-dependent mechanisms (Chen and Kong, 2004; Nelson *et al.*, 2006). Interestingly, these phytochemicals themselves display strong hormetic effects which reflect their abilities to stimulate or inhibit multiple signaling pathways, with different effects occurring at different ED₅₀ concentrations of the phytochemical. Unfortunately, much of the literature in this area has not utilized dose-response curves, but rather has examined fixed, high concentrations of the phytochemicals. The concentrations examined are often not pharmacologically achievable and therefore of questionable physiological relevance. The studies have often overlooked dramatic ARE activation at lower concentrations that disappears at the higher concentrations. An awareness of hormesis is essential for studies involving signal transduction pathways.

Finally, it should be noted that the antioxidant enzymes are largely coordinately regulated, directly or secondarily, by ARE-regulated promoters. It may be highly artificial to manipulate the concentration of SOD *alone*, observing effects on net lipid peroxidation, when *in vivo* the entire network of antioxidant enzymes would more likely be induced in concert. This means that glutathione levels would be raised via increased synthesis, glutathione reductase and peroxidases would be induced to eliminate lipid hydroperoxides, and various peroxiredoxins would also be induced. All of these actions would serve to limit lipid peroxidation. Hence, an animal's natural response to oxidative stress might be much

less likely to produce hormetic effects than laboratory experiments that manipulate any single antioxidant enzyme such as SOD.

REFERENCES

- Ambrosio G, Becker L C, Hutchins G M, Weisman H F and Weisfeldt M L. 1986. Reduction in Experimental Infarct Size by Recombinant Human Superoxide Dismutase: Insights into the Pathophysiology of Reperfusion Injury. *Circulation* **74**:1424-1433
- Ambrosio G, Zweier J L, Duilio C, Kuppusamy P, Santoro G, Elia P P, Tritto I, Cirillo P, Condorelli M, Chiariello M and Flaherty J T. 1993. Evidence That Mitochondrial Respiration Is a Source of Potentially Toxic Oxygen Free Radicals in Intact Rabbit Hearts Subjected to Ischemia and Reperfusion. *J Biol Chem* **268**:18532-18541
- Anilkumar N, Weber R, Zhang M, Brewer A and Shah A M. 2008. Nox4 and Nox2 NADPH Oxidases Mediate Distinct Cellular Redox Signaling Responses to Agonist Stimulation. *Arterioscler Thromb Vasc Biol* **28**:1347-1354
- Aoki N, Bitterman H, Brezinski M E and Lefer A M. 1988. Cardioprotective Actions of Human Superoxide Dismutase in Two Reperfusion Models of Myocardial Ischaemia in the Rat. *Br J Pharmacol* **95**:735-740
- Bernier M, Manning A S and Hearse D J. 1989. Reperfusion Arrhythmias: Dose-Related Protection by Anti-Free Radical Interventions. *Am J Physiol* **256**:H1344-H1352
- Buchwald A, Klein H H, Lindert S, Pich S, Nebendahl K, Wiegand V and Kreuzer H. 1989. Effect of Intracoronary Superoxide Dismutase on Regional Function in Stunned Myocardium. *J Cardiovasc Pharmacol* **13**:258-264
- Burdon RH. 1995. Superoxide and Hydrogen Peroxide in Relation to Mammalian Cell Proliferation. *Free Radic Biol Med* **18**:775-794
- Burton KP. 1985. Superoxide Dismutase Enhances Recovery Following Myocardial Ischemia. *Am J Physiol* **248**:H637-H643
- Chambers DE, Parks D A, Patterson G, Roy R S, McCord J M, Yoshida S, Parmley L and Downey J M. 1985. Xanthine Oxidase As a Source of Free Radical in Myocardial Ischemia. *J Mol Cell Cardiol* **17**:145-152
- Chance B, Sies H and Boveris A. 1979. Hydroperoxide Metabolism in Mammalian Organs. *Physiol Rev* **59**:527-605
- Chen C and Kong A N. 2004. Dietary Chemopreventive Compounds and ARE/EpRE Signaling. *Free Radic Biol Med* **36**:1505-1516
- Chen Z, Oberley T D, Ho Y, Chua C C, Siu B, Hamdy R C, Epstein C J and Chua B H. 2000. Overexpression of CuZnSOD in Coronary Vascular Cells Attenuates Myocardial Ischemia/Reperfusion Injury. *Free Radic Biol Med* **29**:589-596
- Crapo JD and Tierney D F. 1974. Superoxide Dismutase and Pulmonary Oxygen Toxicity. *Am J Physiol* **226**:1401-1407
- Deshmukh DR, Mirochnitchenko O, Ghole V S, Agnese D, Shah P C, Reddell M, Brodin R E and Inouye M. 1997. Intestinal Ischemia and Reperfusion Injury in Transgenic Mice Overexpressing Copper-Zinc Superoxide Dismutase. *Am J Physiol* **273**:C1130-C1135
- Eddy LJ, Stewart J R, Jones H P, Engerson T D, McCord J M and Downey J M. 1987. Free Radical-Producing Enzyme, Xanthine Oxidase, Is Undetectable in Human Hearts. *Am J Physiol* **253** (Heart **22**):H709-H711
- Gao B, Flores S C, Leff J A, Bose S K and McCord J M. 2003. Synthesis and Anti-Inflammatory Activity of a Chimeric Recombinant Superoxide Dismutase: SOD2/3. *Am J Physiol Lung Cell Mol Physiol* **284**:L917-L925
- Gardner TJ, Stewart J R, Casale A S, Downey J M and Chambers D E. 1983. Reduction of Myocardial Ischemic Injury With Oxygen-Derived Free Radical Scavengers. *Surgery* **94**:423-427
- Granger DN, Rutigli G and McCord J M. 1981. Superoxide Radicals in Feline Intestinal Ischemia. *Gastroenterology* **81**:22-29
- Hernandez-Saavedra D, Zhou H and McCord J M. 2005. Anti-Inflammatory Properties of a Chimeric Recombinant Superoxide Dismutase: SOD2/3. *Biomed Pharmacother* **59**:204-208
- Hodgson EK and Fridovich I. 1975. The Interaction of Bovine Erythrocyte Superoxide Dismutase With Hydrogen Peroxide: Chemiluminescence and Peroxidation. *Biochem* **14**:5299-5303

- Kedziora J and Bartosz G. 1988. Down's Syndrome: a Pathology Involving the Lack of Balance of Reactive Oxygen Species. *Free Radical Biol Med* **4**:317-330
- Lazzarino G, Tavazzi B, Di Pierro D, Giardina B and McCord J M (1994) Superoxide radical as both initiator and terminator of lipid peroxidation: the importance of oxidative balance, in *Frontiers of Reactive Oxygen Species in Biology and Medicine* (Asada K and Yoshikawa T eds) pp 523-524, Elsevier Sciences B.V..
- Levy R, Gluzman S, Milman D, Seruty C, Hagay Z, Yavin E and Groner Y. 2002. Ischemic Reperfusion Brain Injury in Fetal Transgenic Mice With Elevated Levels of Copper-Zinc Superoxide Dismutase. *J Perinat Med* **30**:158-165
- Mao GD, Thomas P D, Lopaschuk G D and Poznansky M J. 1993. Superoxide Dismutase (SOD)-Catalase Conjugates - Role of Hydrogen Peroxide and the Fenton Reaction in SOD Toxicity. *J Biol Chem* **268**:416-420
- McCord JM. 1985. Oxygen-Derived Free Radicals in Post-Ischemic Tissue Injury. *N Engl J Med* **312**:159-163
- McCord JM. 2000. The Evolution of Free Radicals and Oxidative Stress. *Am J Med* **108**:652-659
- McCord JM and Fridovich I. 1969. Superoxide Dismutase: an Enzymic Function for Erythrocyte (Hemocuprein). *J Biol Chem* **244**:6049-6055
- McCord JM, Omar B A and Russell W J. 1989. Sources of oxygen-derived radicals in ischemia-reperfusion, in *Medical, Biochemical, and Chemical Aspects of Free Radicals* (Hayaishi O, Niki E, Kondo M and Yoshikawa T eds) pp 1113-1118, Elsevier Science, Amsterdam.
- McCord JM and Russell W J. 1988. Inactivation of Creatine Phosphokinase by Superoxide During Reperfusion Injury. *Basic Life Sci* **49**:869-873
- McCord JM and Turrens J F. 1994. Mitochondrial Injury by Ischemia and Reperfusion. *Curr Topics Bioenerg* **17**:173-195
- Menasche P, Grousset C, Gauduel Y, Mouas C and Pivnic A. 1986. Enhancement of Cardioplegic Protection With the Free-Radical Scavenger Peroxidase. *Circulation* **74**:III138-III144
- Myers CL, Weiss S J, Kirsh M M and Shlafer M. 1985. Involvement of Hydrogen Peroxide and Hydroxyl Radical in the 'Oxygen Paradox': Reduction of Creatine Kinase Release by Catalase, Allopurinol or Deferoxamine, but Not by Superoxide Dismutase. *J Mol Cell Cardiol* **17**:675-684
- Nelson SK, Bose S K, Grunwald G K, Myhill P and McCord J M. 2006. The Induction of Human Superoxide Dismutase and Catalase in Vivo: A Fundamentally New Approach to Antioxidant Therapy. *Free Radic Biol Med* **40**:341-347
- Nelson SK, Bose S K and McCord J M. 1994. The Toxicity of High-Dose Superoxide Dismutase Suggests That Superoxide Can Both Initiate and Terminate Lipid Peroxidation in the Reperfused Heart. *Free Radical Biol Med* **16**:195-200
- Norris KH and Hornsby P J. 1990. Cytotoxic Effects of Expression of Human Superoxide Dismutase in Bovine Adrenocortical Cells. *Mutat Res* **237**:95-106
- Omar BA, Gad N M, Jordan M C, Striplin S P, Russell W J, Downey J M and McCord J M. 1990. Cardioprotection by Cu,Zn-Superoxide Dismutase Is Lost at High Doses in the Reoxygenated Heart. *Free Radical Biol Med* **9**:465-471
- Omar BA and McCord J M. 1990. The Cardioprotective Effect of Mn-Superoxide Dismutase Is Lost at High Doses in the Postschemic Isolated Rabbit Heart. *Free Radical Biol Med* **9**:473-478
- Paraidathathu T, Degroot H and Kehrer J P. 1992. Production of Reactive Oxygen by Mitochondria From Normoxic and Hypoxic Rat Heart Tissue. *Free Radical Biol Med* **13**:289-297
- Rushmore TH and Kong A N. 2002. Pharmacogenomics, Regulation and Signaling Pathways of Phase I and II Drug Metabolizing Enzymes. *Curr Drug Metab* **3**:481-490
- Tamura Y, Chi L, Driscoll E M, Jr., Hoff P T, Freeman B A, Gallagher K P and Lucchesi B R. 1988. Superoxide Dismutase Conjugated to Polyethylene Glycol Provides Sustained Protection Against Myocardial Ischemia/Reperfusion Injury in Canine Heart. *Circ Res* **63**:944-959
- Terada LS, Rubinstein J D, Lesnefsky E J, Horwitz L D, Leff J A and Repine J E. 1991. Existence and Participation of Xanthine Oxidase in Reperfusion Injury of Ischemic Rabbit Myocardium. *Am J Physiol Heart Circ Physiol* **260**:H805-H810
- Thom SR and Elbuken M E. 1991. Oxygen-Dependent Antagonism of Lipid Peroxidation. *Free Radical Biol Med* **10**:413-426
- Uraze A, Reimer K A, Murry C E and Jennings R B. 1987. Failure of Superoxide Dismutase to Limit Size of Myocardial Infarction After 40 Minutes of Ischemia and 4 Days of Reperfusion in Dogs. *Circulation* **75**:1237-1248

- Werns SW, Shea M J, Driscoll E M, Cohen C, Abrams G D, Pitt B and Lucchesi B R. 1985. The Independent Effects of Oxygen Radical Scavengers on Canine Infarct Size. Reduction by Superoxide Dismutase but Not Catalase. *Circ Res* **56**:895-898
- Woodward B and Zakaria M N. 1985. Effect of Some Free Radical Scavengers on Reperfusion Induced Arrhythmias in the Isolated Rat Heart. *J Mol Cell Cardiol* **17**:485-493
- Zhao L, Roche B M, Wessale J L, Kijtawornrat A, Lolly J L, Shemanski D and Hamlin R L. 2008. Chronic Xanthine Oxidase Inhibition Following Myocardial Infarction in Rabbits: Effects of Early Versus Delayed Treatment. *Life Sci* **82**:495-502