

Effects of Antibiotics on Quorum Sensing in *Pseudomonas aeruginosa*[∇]

Mette E. Skindersoe,¹† Morten Alhede,¹ Richard Phipps,¹ Liang Yang,¹ Peter O. Jensen,³
Thomas B. Rasmussen,² Thomas Bjarnsholt,¹ Tim Tolker-Nielsen,¹
Niels Høiby,³ and Michael Givskov^{1*}

Biomedical Microbiology, BioSys, Bldg. 227, Technical University of Denmark, Kgs. Lyngby DK-2800,¹ Physiology, Cultures & Enzymes Division, Chr-Hansen A/S, Bøge Allé 10-12, Hørsholm DK-2970,² and Department of Clinical Microbiology, Rigshospitalet, University of Copenhagen, Copenhagen DK-2100,³ Denmark

Received 19 September 2007/Returned for modification 23 October 2007/Accepted 11 June 2008

During infection, *Pseudomonas aeruginosa* employs bacterial communication (quorum sensing [QS]) to coordinate the expression of tissue-damaging factors. QS-controlled gene expression plays a pivotal role in the virulence of *P. aeruginosa*, and QS-deficient mutants cause less severe infections in animal infection models. Treatment of cystic fibrosis (CF) patients chronically infected with *P. aeruginosa* with the macrolide antibiotic azithromycin (AZM) has been demonstrated to improve the clinical outcome. Several studies indicate that AZM may accomplish its beneficial action in CF patients by impeding QS, thereby reducing the pathogenicity of *P. aeruginosa*. This led us to investigate whether QS inhibition is a common feature of antibiotics. We present the results of a screening of 12 antibiotics for their QS-inhibitory activities using a previously described QS inhibitor selector 1 strain. Three of the antibiotics tested, AZM, ceftazidime (CFT), and ciprofloxacin (CPR), were very active in the assay and were further examined for their effects on QS-regulated virulence factor production in *P. aeruginosa*. The effects of the three antibiotics administered at subinhibitory concentrations were investigated by use of DNA microarrays. Consistent results from the virulence factor assays, reverse transcription-PCR, and the DNA microarrays support the finding that AZM, CFT, and CPR decrease the expression of a range of QS-regulated virulence factors. The data suggest that the underlying mechanism may be mediated by changes in membrane permeability, thereby influencing the flux of *N*-3-oxo-dodecanoyl-L-homoserine lactone.

The opportunistic pathogen *Pseudomonas aeruginosa* causes severe infections, particularly in immunocompromised individuals and patients with cystic fibrosis (CF). The progressive deterioration of the lungs caused by chronic *P. aeruginosa* infections and, thus, persistent inflammation are currently the main causes of morbidity and mortality in patients with CF (55). Intensive antipseudomonal treatment (maintenance therapy) has greatly improved the prospects for patients with CF (26, 29, 72). A serious side effect of antibiotic therapy is the development of resistance to the antibiotics used (3, 15, 28, 54). *P. aeruginosa* forms biofilms during the infection process, which adds to the difficulties of eradicating infections by antibiotic intervention, since bacterial cells living as biofilms are much more tolerant to antibiotics than their planktonic counterparts (4, 14, 17). When the infection enters a chronic state, the mucoid, alginate-overproducing phenotype dominates (22). Macrolides such as erythromycin, clarithromycin, and the erythromycin derivative azithromycin (AZM) have been shown to inhibit the enzymatic activity of guanosine diphosphomannose dehydrogenase in the alginate biosynthetic pathway of mucoid *P. aeruginosa* strains at concentrations well below the MIC (44, 49). Alginate induces a continuous antigen-antibody reaction locally in the small airways. Thus, a lower level of

alginate production not only reduces the viscosity of the sputum but may also lead to less inflammation and, thus, improved lung function (8, 41, 49). Several clinical studies have shown that long-term treatment with AZM improves lung function and body weight in patients with CF (27, 43, 76). AZM is not usually considered to exhibit antipseudomonal activity due to its MIC for *P. aeruginosa* in the range of 128 to 512 µg/ml (43). The highest concentrations of AZM found in the sputum of patients receiving high-dose therapy (250 mg AZM daily) is 0.6 to 79.3 µg/ml, with the median AZM concentration being 9.5 µg/ml (6). It has been suggested that AZM treatment inhibits neutrophil recruitment to the lung by reducing the levels of expression of proinflammatory cytokines and inhibition of neutrophil migration, resulting in a significant reduction in airway-specific inflammation (88).

It has also been suggested that inhibition of cell-cell communication is the mode of action by which AZM exerts its activity in *P. aeruginosa* infections (60, 84). In gram-negative bacteria, cell-cell communication, also known as quorum sensing (QS), relies upon small diffusible signal molecules (*N*-acyl L-homoserine lactones [AHLs]), which interact with transcriptional activators to couple gene expression with cell population density. The QS system of *P. aeruginosa* is organized hierarchically, with the RhII-RhIR components being subordinate to the LasI-LasR components. LasI directs the synthesis of *N*-3-oxo-dodecanoyl-L-homoserine lactone (OdDHL), whereas RhII synthesizes *N*-butanoyl-L-homoserine lactone (BHL). *P. aeruginosa* cells are permeable to BHL, which freely diffuses across the cell membrane, whereas active efflux is required for the transportation of OdDHL (67). In addition to the AHL

* Corresponding author. Mailing address: Biomedical Microbiology, BioSys, Bldg 227, Technical University of Denmark, Kgs. Lyngby DK-2800, Denmark. Phone: 45 4525 2769. Fax: 45 4593 2809. E-mail: immg@pop.dtu.dk.

† Present address: ChemoMetec A/S, Gydevang 43, DK-3450 Alleroed, Denmark.

[∇] Published ahead of print on 21 July 2008.

signal molecules BHL and OdDHL, a third intercellular signal, 2-heptyl-hydroxy-4-quinolone (designated the *Pseudomonas* quinolone signal [PQS]), has been found to be part of the QS regulon in *P. aeruginosa* (70). They all function in concert to control the expression of an array of genes, including genes encoding tissue-damaging exoproducts (31, 51, 62, 66). In addition to controlling the production of virulence factors, the QS signal molecules PQS and OdDHL have been reported to possess immunomodulatory effects, and in this way, they contribute directly to the pathogenesis of *P. aeruginosa* (24, 42, 85).

Several studies have shown the effects of AZM on the expression of QS-regulated virulence genes in vitro (40, 60, 84). Applying a CF mouse model, we have recently demonstrated that AZM treatment significantly improved the clearance of pulmonary bacteria, reduced the extent of lung abscesses and decreased the severity of lung pathology, and reduced the level of alginate production in vivo and also in in vitro experiments, although it had no effect on the growth of infecting bacteria (40). Since QS seems to play a key role in the expression of virulence and the interaction with host protection, inhibitors of QS have been suggested to be important components of future antipseudomonal therapies (37). Examples of compounds that can block QS in *P. aeruginosa* are the halofuranones (32, 38), which originate from a benthic macroalgae, and the fungal metabolites patulin and penicillic acid (74). Furanone C-30 specifically inhibits QS over a concentration range from 1 to 10 μM ; but when it is administered in excess, i.e., at concentrations 10-fold higher, the compound impairs growth and exhibits antibiotic properties. In addition to QS-inhibitory (QSI) activity, the bioactive halofuranones exhibit a multitude of activities, including antibiotic and antifouling properties, and can therefore be considered multifunctional compounds engaged in communication as well as competition (for a recent review, see reference 21). It was recently demonstrated that at sub-MICs antibiotics broadly affect the patterns of transcription in bacteria and seem to function as signaling mediators rather than function by targeting the growth of rivals (35). This raises the question of whether other compounds previously identified for their antibiotic properties may turn out to possess properties that interfere with communication and, potentially, QSI activity. To try to answer that question, we present the results of a screening of 12 antibiotics for their QSI activities. We identified three antibiotics, AZM, ciprofloxacin (CPR), and ceftazidime (CFT), that exhibit strong QSI activities in the screening assay. The effects of these three antibiotics were further investigated by means of transcriptome analysis and phenotyping assays.

MATERIALS AND METHODS

Bacterial strains applied in this study. The sequenced *P. aeruginosa* PAO1 wild-type strain was obtained from the Pseudomonas Genetic Stock Center (strain PAO0001; www.pseudomonas.med.ecu.edu). The *lasI-rhlI* mutant was constructed by using a previously reported knockout system (7). The QSI selector strain QS inhibitor selector 1 (QSI1) was described previously (73).

Bacterial screen for QS inhibition. Strain QSI1 has previously been described in detail (73). The QS element is derived from the *Vibrio fischeri* LuxR-regulated QS system and was cloned in *Escherichia coli*, where it responds to a broad range of AHLs and QS inhibitors (2, 73, 74). QSI1 harbors plasmid pUC18Not-*luxR*-*P_{luxR}*-RBSII-*phlA* T₀-T₁ Ap^r maintained in CSH37 (59), an *E. coli* strain that constitutively expresses β -galactosidase. The general concept of the selector

system implies the presence of a QS regulator (*luxR* or a homologue) gene and a QS-controlled promoter fused to a gene encoding a toxic gene product. In the presence of signal molecules, the gene under QS control is expressed, causing cell death. However, in the presence of a QSI compound, production of the toxic gene product is switched off and the bacteria are rescued and able to grow normally. The toxic gene product of the QSI1 selector system is cytosolic phospholipase A, which originates from *Serratia liquefaciens* MG1 (33, 34). Because signal molecules are added to the assay from an external source, QSIs targeting the AHL synthetase are not identified, whereas compounds that enhance the degradation of AHL, impede AHL uptake, or block the interaction of AHL with the *luxR* protein are found.

Briefly, the preparation of QSI1 screens was performed as follows: BT medium (B medium [16] plus 2.5 mg thiamine per liter) containing 2% agar (wt/vol) was melted, 10% A10 buffer (16) was added, and the mixture was cooled to 45°C. To the resulting ABT agar were then added *N*-3-oxo-hexanoyl-L-homoserine lactone (Sigma-Aldrich, Germany), ampicillin (VepiDan, Denmark), 5-bromo-4-chloro-3-indoxyl- β -D-galactopyranoside (X-Gal; Apollo Scientific, United Kingdom), and 1-methyl-2-pyrrolidone (Merck, Germany) to final concentrations of 100 nM, 100 $\mu\text{g}/\text{ml}$, 40 $\mu\text{g}/\text{ml}$, and 0.2% (vol/vol), respectively; and the medium was supplemented with 0.5% (wt/vol) glucose and 0.5% (wt/vol) Casamino Acids. An overnight culture of QSI1 (grown in ABT medium supplemented with 0.5% [wt/vol] glucose, 0.5% [wt/vol] Casamino Acids, and 100 $\mu\text{g}/\text{ml}$ ampicillin) was added to a final concentration of 0.4%.

Plates for the screening assay were subsequently made by pouring the mixture into a box to give an agar thickness of 5 mm. Wells with diameters of 5 mm were made in the agar plates. Upon solidification, 50 μl of a 1- to 10- $\mu\text{g}/\text{ml}$ solution of the antibiotic to be tested for QSI activity was added to the wells. The plate was then incubated at 30°C for 16 h. The appearance of a circular zone of growth (which appeared blue due to the presence of hydrolyzed X-Gal) indicated the presence of compounds with QSI activity in the sample tested.

Sample preparation for *P. aeruginosa* GeneChip analysis. ABT medium supplemented with 0.5% Casamino Acids was inoculated with exponentially growing *P. aeruginosa* PAO1 cells to an optical density at 600 nm (OD₆₀₀) of 0.05, and the cells were grown at 37°C and 200 rpm in five 500-ml flasks containing 100 ml each. When the OD₆₀₀ was 0.3, AZM (8 and 2 $\mu\text{g}/\text{ml}$; Zithromax; Pfizer), CPR (0.04 $\mu\text{g}/\text{ml}$; Fluka), or CFT (0.25 $\mu\text{g}/\text{ml}$; Fortum; GlaxoSmithKline) was added. These concentrations of antibiotics did not affect the growth of *P. aeruginosa*. Samples were retrieved when the OD₆₀₀ reached 2.0, immediately transferred to 2 volumes of RNAlater (Ambion), and stored at -80°C. RNA was isolated with an RNeasy mini purification kit (Qiagen), according to the protocol provided by the manufacturer, including the on-column DNase treatment. The synthesis of cDNA was performed with 12 μg RNA, 300 ng/ μl random primers (Invitrogen), 1,500 U SuperScript III reverse transcriptase (Invitrogen), and 30 U SUPERase[®] · In RNase inhibitor (Ambion), according to the Affymetrix expression analysis protocol. The cDNA synthesized was purified with a QIAquick PCR purification kit (Qiagen), and 3 to 4 μg cDNA was fragmented with 0.2 U FPLC pure DNase I (Amersham Bioscience) per μg cDNA. The fragmented cDNA was labeled at the terminus with biotin-ddUTP (Enzo Bioarray terminal labeling kit) and hybridized for 18 h at 55°C to a *P. aeruginosa* genome microarray GeneChip (Affymetrix). The chips were washed and stained according to the Affymetrix protocol. The microarray hybridization signal intensity was scaled to an overall signal average of 500; and the microarray was evaluated for the presence, marginal presence, or absence of probe sets (representing different genes or open reading frames) by the use of Affymetrix GeneChip operating system (version 1.4) software. The expression values only for probe sets estimated as being "present" were included in the further analyses. The microarray experiments were conducted twice for the antibiotic-treated cultures (two arrays for 2 $\mu\text{g}/\text{ml}$ AZM, two arrays for 8 $\mu\text{g}/\text{ml}$ AZM, two arrays for 0.04 $\mu\text{g}/\text{ml}$ CPR, and two arrays for 0.25 $\mu\text{g}/\text{ml}$ CFT), while the untreated controls (references) were tested in four replicates. The data presented here are based on the average of each set.

Quantification of mRNA by real-time PCR. (i) Primer design. The primers used for mRNA amplification (Table 1) were designed by the use of Integrated DNA Technologies Primer Quest software (<http://www.idtdna.com>). Briefly, the sizes of the amplicon designed ranged from 80 to 200 bp, and the primer melting temperatures were designed for 60°C, with a melting temperature difference of less than 2°C for each primer pair. The primer sequences were also subjected to a BLAST analysis against the *P. aeruginosa* PAO1 genome sequence to eliminate the possibility of nonspecific binding.

(ii) Real-time PCR. Each real-time PCR mixture (final volume, 20 μl) contained 9 μl of cDNA, 10 μl of 2 \times SYBR green quantitative PCR master mix (Fermentas), and 0.5 μM of each forward and reverse primer. Real-time PCR was performed with a Chromo4 real-time detector (Bio-Rad [formerly MJ Re-

TABLE 1. Primers used for real-time PCR

Gene	Forward	Reverse
<i>rhIA</i>	5'-TCT GTT GGT ATC GGT TTG CAA GGG-3'	5'-ACA GCA CCA CGT TGA AAT GTT CGG-3'
<i>pprA</i>	5'-TGC TGT TGG GCA TGG ACA TCT-3'	5'-AGA CGC GAA AGG ATC AGC TT-3'
<i>pprB</i>	5'-TCA AGT ACG AGG TAC ACG GCA ACA-3'	5'-TAT CTG GTA GTT GGT CAG GCC CTT-3'
<i>rpoD</i>	5'-ACA TGC GTG AAA TGG GTA CCG T-3'	5'-ATG CGC TTG GCG ATT TCG ATC T-3'

search]) by using the following cycling parameters: an initial denaturation at 95°C for 10 min before 40 cycles of 95°C for 15 s, 60°C for 30 s, and 72°C for 30 s. The data were analyzed by the efficiency-corrected relative quantification method (71). Expression of the target genes was normalized to the expression of reference gene *rpoD*, as the DNA microarray analysis showed that the expression of this gene was stable under all conditions. Furthermore, Savli et al. (77) have found that among the housekeeping genes, this gene has the most stable expression. To control for variations between runs, the housekeeping gene and the various target genes for each individual treatment were amplified at the same time on a 96-well plate. The changes in the levels of expression of the target genes in the treated samples were calculated relative to the average level of expression of the respective gene in the nontreated sample. The efficiency of the reverse transcription-PCR (RT-PCR) for the gene pairs (under treated versus untreated conditions) varied less than 5%.

Accordingly, the average efficiencies of each gene in this study were very similar for the conditions compared, allowing accurate analysis.

Supernatants for virulence factor assays. *P. aeruginosa* PAO1 cultures were grown either for 16 to 20 h (for hemolysis assay) or to an OD₆₀₀ of 2.0 (for the protease, chitinase, and elastase assays), as described above, with 8 or 2 µg/ml AZM, 0.04 µg/ml CPR, 0.25 µg/ml CFT, or no antibiotic (control). The *P. aeruginosa* PAO1 QS mutant $\Delta lasI\text{-}\Delta rhlI$ was also grown along with the treated and untreated *P. aeruginosa* PAO1 cultures. Cells were harvested by centrifugation, and the supernatants were filtered sterilized (TPP syringe filter; pore size, 0.22 µm). The supernatants were either used immediately or stored at -20°C.

Hemolysis assay. The filter-sterilized supernatants were autoclaved to destroy enzymatic activity and mixed with venous blood from a healthy individual at a volumetric ratio of 30:1 (supernatant to blood), observed for hemolysis after 20 min, and left at room temperature for 4 h to allow the sedimentation of intact erythrocytes. Hemolytic activity was observed as a clearing and a nonprecipitating red coloring of the blood solution, while a lack of hemolytic activity was characterized by the precipitation of intact erythrocytes. The experiment was performed four times with supernatants from four individual growth experiments.

Protease assay. The filter-sterilized supernatants (300 µl each) were added to the wells of ABT agar plates (ABT medium supplemented with 2% agar) containing 5% skim milk, and the plates were incubated overnight at 37°C. The zones of clearance (radii) were measured on photographs of the plates by the use of a ruler. The experiment was performed three times as four replicates with supernatants from three individual growth experiments. The values from the four replicates were averaged and used as one value to represent each of the three experiments.

Elastase assay. The filter-sterilized supernatants were mixed 2:1 with phosphate buffer (0.1 M, pH 6.3), and 2 mg/ml elastin Congo red (Sigma) was added. The mixture was incubated at 37°C with shaking (200 rpm) for 1 week. After centrifugation, the absorbance of the supernatant at 495 nm was measured with a spectrophotometer zeroed on an elastin Congo red sample incubated with medium alone. The procedure was modified from that described previously (63). The experiment was performed three times in triplicate with supernatants from three individual growth experiments. The values from the triplicate experiments were averaged and used as one value to represent each of the three experiments.

Chitinase assay. Chitinase was measured by a modified chitin azure assay (30, 36). The filter-sterilized supernatants were mixed 2:1 with sodium citrate buffer (0.1 M, pH 4.8), and 0.5 mg/ml chitin azure (Sigma) was added. The supernatant-chitin azure mixtures were incubated at 37°C with shaking (200 rpm) for 1 week. The samples were then centrifuged at 15,000 × g for 10 min, and the absorbance at 570 nm was determined. The samples were compared to blanks incubated with medium only. The experiment was performed three times in triplicates with supernatants from three individual growth experiments. The values from the

triplicate experiments were averaged and used as one value to represent each of the three experiments.

Quantification of rhamnolipid. A standard curve for rhamnolipid B (concentration versus total ionization current) was calculated from liquid chromatography-electrospray ionization-mass spectrometry (MS) data. To minimize potential differences in the ionization levels of rhamnolipid between the samples, the rhamnolipid standards used to calculate the concentration curve were analyzed immediately prior to as well as after the samples were analyzed. The total ionization current was determined on the [M + NH₄]⁺ ion at 668.4 over the 7 s over which rhamnolipid B was eluted. High-pressure liquid chromatography (LC)-MS analysis was performed with an Agilent 1100 series high-pressure liquid chromatograph connected to a Micromass LCT time-of-flight mass spectrometer. The concentration of rhamnolipid in the untreated culture was set equal to an index value of 100.

Docking of AZM, CFT, and CPR against the LBD of the LasR protein. The X-ray crystallographic structure of the ligand binding domain (LBD) of the LasR protein (Protein Data Bank accession no. 2UV0) was used for the docking of AZM, CFT, and CPR to the LasR receptor site by using the program Molegro Virtual Docker (86) and the template docking model with default settings. OdDHL was set as the template ligand in the LBD of the LasR protein. The ligands were ranked by using the Molegro Virtual Docker rerank scoring function.

Data analysis and statistics. One-way analysis of variance followed by Dunnett's test was used to evaluate the effects of AZM, CFT, and CPR treatment and the effect of the *lasI-rhlI* mutation on the expression of elastases, chitinase, and protease compared to their expression in untreated wild-type strain *P. aeruginosa* PAO1. Tests were performed with the InStat (version 3) program from GraphPad Software (San Diego, CA). Differences were considered significant if the *P* value was <0.05 or less.

Microarray data accession number. The DNA microarray data (CHP, CEL, and EXP files) have been submitted to the Gene Expression Omnibus database (<http://www.ncbi.nlm.nih.gov/geo/>) and may be found under experiment series number GSE8953.

RESULTS

Screening for QSI activity. AZM has been shown to inhibit the expression of a subfraction of QS-controlled genes (60, 84). Along with AZM, we have investigated 11 other antibiotics (including one antifungal compound, griseofulvin) for their abilities to interfere with bacterial QS using a simple plate diffusion screen based on QSI1 (73). By employing our standard conditions (73), we found that chloramphenicol, gentamicin, griseofulvin, kanamycin, piperacillin, spectinomycin, tetracycline, tobramycin, and streptomycin showed low levels of or no QSI activity, whereas AZM, in accordance with previous investigations, was very active. The fluoroquinolone CPR and the cephalosporin CFT also showed strong QSI activities in the assay with QSI1 (Fig. 1). CPR and CFT are employed in the clinic as antipseudomonal treatments, whereas AZM exhibits a low level of, if any, bactericidal or bacteriostatic effect against *P. aeruginosa*. CPR targets DNA gyrase (topoisomerase II) and thereby inhibits bacterial DNA synthesis. CFT is a β-lactam and functions by disrupting the synthesis of the peptidoglycan layer of bacterial cell walls. The mode of action of AZM is to bind to the 50S subunit of the bacterial ribosome, thereby inhibiting the translation of mRNA. These three antibiotics therefore exhibit diverse mechanisms of antimicrobial action, and they are structurally very different. We chose to further investigate the effects on *P. aeruginosa* QS-regulated gene expression and virulence factor production with AZM, CPR, and CFT as representatives of these widely diverse groups of antibiotics. The growth curves for *P. aeruginosa* PAO1 grown in the presence or absence of AZM, CPR, and CFT at different concentrations (including concentrations with slightly growth-inhibiting effects) and for the $\Delta lasI\ \Delta rhlI$ mutant are presented




FIG. 1. Screening of antibiotics by the QSIS1 assay. (a) Piperacillin; (b) azithromycin; (c) chloramphenicol; (d) ciprofloxacin; (e) gentamicin; (f) spectinomycin; (g) griseofulvin; (h) kanamycin; (i) tetracycline; (j) tobramycin; (k) ceftazidime; (l) streptomycin.

in Fig. 2. We found that addition of 8 $\mu\text{g/ml}$ AZM, 0.25 $\mu\text{g/ml}$ CFT, or 0.04 $\mu\text{g/ml}$ CPR did not affect the growth of *P. aeruginosa* cultures when drug was added when the OD_{600} was 0.1. The MICs of the three antibiotics were found to be 800 $\mu\text{g/ml}$ (AZM), 8 $\mu\text{g/ml}$ (CFT), and 12.5 $\mu\text{g/ml}$ (CPR) under the specified growth conditions. Thus, for the following series of experiments, the concentrations applied were 30- to 300-fold lower than the MICs.

Heat-stable hemolysin. We investigated the effects of AZM, CPR, and CFT on the production of heat-stable hemolysin. *P. aeruginosa* produces at least two hemolysins, the heat-labile phospholipase C (*phlC*, PA0844) and the heat-stable rhamnolipid (*rhlA*, *rhlB*, *rhlC*, and *rhlG*; PA3479, PA3478, PA1130, and PA3387, respectively) (9, 46, 83). Rhamnolipid synthesis is turned on in early stationary phase of *P. aeruginosa* PAO1 cultures (62, 89). After 14 to 20 h of growth (under the conditions outlined in Materials and Methods), the concentration of rhamnolipid in the supernatant exceeds 50 $\mu\text{g/ml}$, which is sufficient to lyse erythrocytes within 20 min (unpublished observations). The lysis of erythrocytes by rhamnolipid can be used as a simple assay for the detection of rhamnolipid by




FIG. 2. Growth curves for the $\Delta\text{lasI}-\Delta\text{rhlI}$ mutant (\square) and strain PAO1 grown without antibiotics (\times) or PAO1 grown in the presence of 2 $\mu\text{g/ml}$ AZM (Δ), 8 $\mu\text{g/ml}$ AZM (\triangle), 16 $\mu\text{g/ml}$ AZM (\blacktriangle), 0.04 $\mu\text{g/ml}$ CPR (\circ), 0.08 $\mu\text{g/ml}$ CPR (\bullet), 0.25 $\mu\text{g/ml}$ CFT (\diamond), or 0.5 $\mu\text{g/ml}$ CFT (\blacklozenge).




FIG. 3. Hemolysis of autoclaved supernatants from *P. aeruginosa* PAO1 cultures grown for 20 h. PAO1 was grown without antibiotics (a) and in the presence of 2 $\mu\text{g/ml}$ AZM (b), 8 $\mu\text{g/ml}$ AZM (c), 0.25 $\mu\text{g/ml}$ CFT (d), or 0.04 $\mu\text{g/ml}$ CPR (e). The results for the $\Delta\text{lasI}-\Delta\text{rhlI}$ mutant of PAO1 are also shown (f).

mixing autoclaved supernatant with blood and incubating the mixture at room temperature for 20 min. The autoclaving eliminates the activity of the heat-labile enzyme phospholipase. A clearing and a nonprecipitating red coloring of the blood-supernatant solution indicates that rhamnolipid is present at a concentration well above 50 $\mu\text{g/ml}$, while a lack of hemolytic activity (characterized by the precipitation of intact blood cells) indicates that little (less than 50 $\mu\text{g/ml}$) or no rhamnolipid is present in the supernatant. Investigating autoclaved supernatants from *P. aeruginosa* grown in the presence or the absence of AZM, CPR, and CFT and supernatants from the QS mutant $\Delta\text{lasI}-\Delta\text{rhlI}$, we found that only the untreated wild type produced enough rhamnolipid to cause the rapid hemolysis of red blood cells. However, treatment with CFT did not fully abolish the hemolytic activity (Fig. 3).

AZM at sub-MICs has previously been reported to affect an *rhlAB::lacZ* gene fusion (84). We decided to directly quantify the effect of AZM as well as the effects of CFT and CPR on rhamnolipid production. We thus determined the concentration of rhamnolipid in the supernatants of treated and untreated *P. aeruginosa* cells by LC-MS. When the cells were grown in the presence of 8 $\mu\text{g/ml}$ AZM, the concentration of rhamnolipid was reduced by 85% compared to that in the untreated cultures, and 2 $\mu\text{g/ml}$ AZM reduced the rhamnolipid content of the supernatants by 80% compared to that in the untreated cultures, whereas CPR (0.04 $\mu\text{g/ml}$) and CFT (0.25 $\mu\text{g/ml}$) reduced the concentration of rhamnolipid to 55 to 65% of that in the untreated cultures. The QS mutant $\Delta\text{lasI}-\Delta\text{rhlI}$ did not produce any measurable amount of rhamnolipid (Fig. 4).

Protease. The production of extracellular protease in *P. aeruginosa* PAO1 cultures grown in the absence or the presence of AZM, CPR, or CFT and in $\Delta\text{lasI}-\Delta\text{rhlI}$ mutant cultures was assayed by adding filter-sterilized supernatants from overnight cultures to wells made in agar plates containing 5% skim milk. The protease activity was lowered by AZM, CPR, and CFT (Fig. 5).

Elastase. The extracellular elastase activities in the supernatants of *P. aeruginosa* PAO1 cultures (untreated or grown with AZM, CPR, or CFT) were quantified by use of a previously described procedure (63). The QS mutant $\Delta\text{lasI}-\Delta\text{rhlI}$ was assayed by the same procedure. AZM treatment reduced the elastase activity of PAO1 almost to the level of that for the




FIG. 4. Rhamnolipid contents, in relative units (RU), in supernatants from *P. aeruginosa* cultures grown either without antibiotics (black bar) or in the presence of 2 $\mu\text{g/ml}$ AZM (bar with horizontal lines), 8 $\mu\text{g/ml}$ AZM (bar with vertical lines), 0.25 $\mu\text{g/ml}$ CFT (dark gray bar), or 0.04 $\mu\text{g/ml}$ CPR (light gray bar). The $\Delta\text{lasI-}\Delta\text{rhII}$ mutant of PAO1 was also included in the experiment but did not produce a measurable concentration of rhamnolipid (last slot [no column visible]). The concentration of rhamnolipid in the untreated culture was set equal to an index value of 100. The graph is based on the average of the indexes of three independent experiments. *, $P < 0.05$; **, $P < 0.01$ (Dunnnett's post test).

$\Delta\text{lasI-}\Delta\text{rhII}$ mutant. CPR and CFT also reduced the elastase activity (Fig. 6).

Chitinase. The chitinase activities in the supernatants of *P. aeruginosa* PAO1 cultures (untreated or grown with AZM, CPR, or CFT) and QS mutant $\Delta\text{lasI-}\Delta\text{rhII}$ cultures were quantified by measuring the degradation of chitin azure. AZM and CPR reduced the chitinolytic activity of PAO1 almost to the level of that for the $\Delta\text{lasI-}\Delta\text{rhII}$ mutant. CFT did not decrease chitinase activity as much as AZM and CPR did, although it still had an effect (Fig. 7).

DNA microarray analysis of gene expression in the presence of AZM, CFT, and CPR. In order to obtain knowledge of the mechanism by which virulence factor production was altered by the presence of the three antibiotics, we performed DNA microarray analyses with *P. aeruginosa* cultures treated with AZM, CPR, and CFT at concentrations that did not affect




FIG. 5. Protease activities of supernatants from *P. aeruginosa* cultures grown either without antibiotics (black bar) or in the presence of 2 $\mu\text{g/ml}$ AZM (bar with horizontal lines), 8 $\mu\text{g/ml}$ AZM (bar with vertical lines), 0.25 $\mu\text{g/ml}$ CFT (dark gray bar), or 0.04 $\mu\text{g/ml}$ CPR (light gray bar). The $\Delta\text{lasI-}\Delta\text{rhII}$ mutant of PAO1 was also included in the experiment but did not produce a measurable concentration of protease (last slot [no column visible]). The graph is based on the average results of three independent experiments. *, $P < 0.05$; **, $P < 0.01$ (Dunnnett's test).




FIG. 6. Elastolytic activities of supernatants from PAO1 cultures grown to an OD_{600} of 2.0 either without antibiotics (black bar) or in the presence of: 2 $\mu\text{g/ml}$ AZM (bar with horizontal lines), 8 $\mu\text{g/ml}$ AZM (bar with vertical lines), 0.25 $\mu\text{g/ml}$ CFT (dark gray bar), or 0.04 $\mu\text{g/ml}$ CPR (light gray bar). The elastolytic activity of the $\Delta\text{lasI-}\Delta\text{rhII}$ mutant of PAO1 is also shown (white bar). The graph is based on the average results of three independent experiments. *, $P < 0.05$; **, $P < 0.01$ (Dunnnett's post test).

growth. The medium with exponentially growing *P. aeruginosa* cultures (OD_{600} , 0.1) was supplemented with either AZM (2 and 8 $\mu\text{g/ml}$), CFT (0.25 $\mu\text{g/ml}$), or CPR (0.04 $\mu\text{g/ml}$) or the medium was left untreated as a control. By applying the growth conditions described in Materials and Methods, an OD_{600} of 2.0 corresponds to the transition from exponential phase into stationary phase, and at this point a substantial fraction of QS-regulated genes are maximally regulated (induced or repressed) (80). An OD_{600} of 2.0 was therefore chosen as the sample point for the transcriptomic analysis. Total RNA was extracted from the samples and subjected to cDNA synthesis. The absolute expression values from cultures grown in the presence of antibiotics were compared with those from untreated cultures, and the changes in expression are reported. As described previously (38, 89), changes in expression of less than fivefold were disregarded. Only probe sets that were present, as validated with Affymetrix GeneChip operating system (version 1.4) software, were included in the analyses. According to these criteria, non-growth-inhibitory concentrations of AZM altered the expression of 477 (8 $\mu\text{g/ml}$) and 323 (2 $\mu\text{g/ml}$) genes, with most genes (419 and 277, respectively) being downregulated. Non-growth-inhibitory concentrations of




FIG. 7. Chitinase activities of supernatants from PAO1 cultures grown to an OD_{600} of 2.0 either without antibiotics (black bar) or in the presence of: 2 $\mu\text{g/ml}$ AZM (bar with horizontal lines), 8 $\mu\text{g/ml}$ AZM (bar with vertical lines), 0.25 $\mu\text{g/ml}$ CFT (dark gray bar), or 0.04 $\mu\text{g/ml}$ CPR (light gray bar). The chitinase activity of the $\Delta\text{lasI-}\Delta\text{rhII}$ mutant of PAO1 is also shown (white bar). The graph is based on the average results of three independent experiments. *, $P < 0.05$; **, $P < 0.01$ (Dunnnett's post test).

TABLE 2. Effect on gene expression

Compound (concn)	No. of genes downregulated more than fivefold	No. of genes upregulated more than fivefold	No. (%) of QS genes ^a affected more than fivefold	Specificity (%) ^b
AZM (8 μ g/ml, 11 μ M)	427	61	121 (70)	28
AZM (2 μ g/ml, 2.7 μ M)	227	49	81 (47)	35
CFT (0.25 μ g/ml, 0.4 μ M)	136	10	49 (28)	36
CPR (0.04 μ g/ml, 0.1 μ M)	223	58	89 (51)	40
C30 ^c (2.5 μ g/ml, 10 mM)	260	0	125 (72)	48

^a Genes belonging to the QS regulon, as defined elsewhere (74).

^b Specificity is indicated by the number of QS genes repressed/total number of genes repressed.

^c Data for the QSI furanone C30 were obtained at an OD₆₀₀ of 2.0 and are included for comparison. The data are from a previous report (38).

CFT (0.25 μ g/ml) and CPR (0.04 μ g/ml) affected the expression of 122 (114 repressed) and 271 genes (215 repressed), respectively (Table 2).

AZM, CFT, and CPR affect expression of QS-regulated genes. The QS regulon has been defined by (among others) Rasmussen et al. (74). That particular study (74) considered genes to be regulated by QS if their levels of expression in both $\Delta lasI$ - $\Delta rhII$ and $\Delta lasR$ - $\Delta rhIR$ mutants were consistently altered more than fivefold compared to the levels of expression in their wild-type counterpart. The majority of all QS-regulated genes defined in this way are inducible with exogenous AHLs in a $\Delta lasI$ - $\Delta rhII$ background. Only one gene (PA1556, a probable cytochrome *c* oxidase subunit) is consistently upregulated in both the $\Delta lasI$ - $\Delta rhII$ and the $\Delta lasR$ - $\Delta rhIR$ mutants and is thus QS repressible. Using this QS regulon, we analyzed the fraction of QS-regulated genes among the genes repressed by the three antibiotics. These analyses showed that AZM (8 μ g/ml) repressed 71% of the 174 QS-regulated genes in *P. aeruginosa* PAO1 and that 29% of all genes being repressed by AZM were members of the QS regulon. Lowering of the concentration of AZM to 2 μ g/ml caused a relief in the inhibition of QS-regulated genes (47% of all QS genes repressed), and the total number of genes downregulated fell to 277, while the specificity toward the QS regulon increased slightly to 30%. CPR had the highest specificity for the QS regulon (48%) and targeted 49% of all QS genes in *P. aeruginosa*, whereas the specificity for CFT was 39% and CFT repressed only 25% of all QS genes. For comparison, the specificity of C30 was 48%, and it downregulated 72% of all QS genes in *P. aeruginosa* (raw data for C30 are from a previous report [38] and have been subjected to the same data treatment and criteria as the raw data for AZM, CFT, and CPR). The effects of AZM, CFT, and CPR on the expression of genes belonging to the QS regulon are shown in Table 3.

The DNA array data show that members of the QS regulon are overrepresented among the genes repressed by sub-MICs of AZM, CFT, and CPR. This suggests that these antibiotics exhibit some specificity for QS-regulated gene expression through an interaction with or an effect on QS regulatory

components. Investigating the expression profiles of genes regulated by LasR, RhlR, or LasR-RhlR, we further aimed to elucidate the potential QS target(s) of the antibiotics. The QS-regulated genes can be divided into four groups. Group A contains 44 LasR-dependent genes, i.e., genes downregulated in both a $\Delta lasR$ mutant and a $\Delta lasR$ - $\Delta rhIR$ mutant. Group B consists of RhlR-dependent genes (14 genes), i.e., genes downregulated in both a $\Delta rhIR$ mutant and a $\Delta lasR$ - $\Delta rhIR$ mutant. Group C comprises the main fraction of QS-regulated genes (102 genes in total) and consists of genes which are downregulated in all three types of mutants: $\Delta lasR$, $\Delta rhIR$, and $\Delta lasR$ - $\Delta rhIR$. Finally, group D consists of 14 LasR-RhlR-dependent genes, which are downregulated only if both LasR and RhlR are knocked out, as in the $\Delta lasR$ - $\Delta rhIR$ mutant. Interestingly it seems that AZM, CFT, and CPR had less of a preference for the group B genes (the RhlR-specific genes) (Fig. 8).

Even though all three antibiotics exhibited a higher degree of specificity for the LasR-regulated genes than for the RhlR-dependent genes, none of these antibiotics downregulated the AHL synthetase genes. AZM, however, caused the downregulation of *lasR* sevenfold, which may also contribute to the QSI effect of AZM. This is in accordance with the observation that AZM (2 μ g/ml) has been reported to decrease the level of expression of a *lasR::lacZ* gene fusion (84).

Among the genes affected by AZM, CFT, and CPR, many encode previously identified QS-regulated genes such as chitinase (*chiC*, PA2300) and chitin-binding protein (*cbpD*, PA0852) (30, 80, 89). Other known QS-induced genes, such as the genes encoding the osmotically inducible protein (*osmC*, PA0059) and cytochrome *c* oxidase (*coxAB* and *coIII*, PA0105 to PA0108), are downregulated by the presence of the three antibiotics (38, 80). The endoprotease gene *piv* (also known as *prpL* [PA4175]) was also downregulated by AZM, CFT, and CPR. This protease is capable of degrading elastin, decorin, and the transferrin family of proteins, including lactoferrin (92). PA5099 (a probable transporter), which belongs to the QS regulon, was downregulated more than 300 times by AZM and CPR and 50 times by CFT. The regulon contains only one gene, PA1556, which is repressed by QS. This gene, encoding cytochrome *c* oxidase, was upregulated by all three antibiotics, and so was the neighboring gene PA1557, a probable cytochrome oxidase subunit. Many genes from the region from PA2134 to PA2190 of the genome were downregulated by AZM, CFT, and CPR. This region has been found to be associated with stationary growth and to be under QS/*rpoS* control (79, 80, 89, 90). Of the 57 genes in this 60.69-kbp region, 46 were found to be present in all the arrays. Of the 46 genes, 35 genes were repressed more than fivefold by AZM and CPR. CPR and especially AZM seemed to exhibit higher target specificities for QS genes than CFT, which is in accordance with the findings of the virulence factor assays described earlier in this section. Many QS-regulated virulence factors, such as the alkaline metalloproteinase (*aprA*, PA1249), the PfpI protease (*pfpI*, PA0355), the LasA protease (*lasA*, PA1871), elastase B (*lasB*, PA3724), the cytotoxic galactophilic lectin (PA-IL; *lecA*, PA2570), and rhamnolipid (*rhlA* and *rhlB*; PA3478 and PA3479, respectively), were repressed more than fivefold by AZM and CPR but not by CFT. These results were confirmed by RT-PCR, which showed that AZM downregulated *rhlA* expression 45-fold,

TABLE 3. Effects of sub-MICs of AZM, CFT, and CPR on *P. aeruginosa* genes belonging to the QS regulon^a

Gene in <i>P. aeruginosa</i>	Gene	<i>pprB</i> ^b	Regulation by AZM ^c	Reg. ^d	Relative expression					Description			
					AZM (2 µg/ml)	AZM (8 µg/ml)	CFT	CPR	<i>last-rhlI</i>		<i>lasR-rhlR</i>	<i>lasR</i>	<i>rhlR</i>
PA0059	<i>osmC</i>			C	-19.8	-63.4	-8.6	-16.8	-22.0	-31.8	-44.0	-25.8	Osmotically inducible protein OsmC
PA0105	<i>coxB</i>			C	-20.0	-25.3	-8.3	-10.5	-11.5	-13.5	-18.7	-7.9	Cytochrome <i>c</i> oxidase, subunit II
PA0107				C	-91.9	-29.0	-11.0	-17.0	-17.4	-14.1	-18.6	-11.4	Conserved hypothetical protein
PA0108	<i>collI</i>			C	-14.2	-33.8	-13.2	-12.2	-27.8	-8.5	-15.8	-6.3	Cytochrome oxidase, subunit III
PA0122				C	-3.0	-14.2	-1.5	-2.9	-44.8	-26.1	-7.1	-6.0	Conserved hypothetical protein
PA0188		-		C	-2.9	-32.5	-4.8	-4.6	-8.1	-21.3	-28.2	-8.8	Hypothetical protein
PA0355	<i>pfpl</i>			C	-7.5	-12.5	-3.6	-6.6	-11.8	-21.5	-17.7	-11.6	Protease Pfpl
PA0567				C	-9.2	-22.0	-3.5	-9.1	-249.4	-25.2	-22.1	-9.5	Conserved hypothetical protein
PA0572				A	-3.5	-6.0	-2.3	-3.4	-10.4	-13.7	-8.2	-1.1	Hypothetical protein
PA0737				C	-3.3	-3.4	-4.1	-4.3	-9.4	-7.7	-94.4	-5.8	Hypothetical protein
PA0843	<i>plcR</i>			A	-4.7	-7.8	-2.7	-3.3	-7.1	-6.2	-25.0	-4.6	Phospholipase accessory protein PlcR precursor
PA0852	<i>cbpD</i>		-	C	-7.0	-35.0	-5.0	-10.1	-63.7	-74.4	-33.2	-8.3	Chitin-binding protein CbpD precursor
PA0990				A	-7.0	-9.3	-3.3	-5.9	-5.7	-19.9	-6.6	-4.4	Conserved hypothetical protein
PA0996	<i>pqsA</i>		+	A	2.0	1.7	1.3	-1.3	-7.0	-20.8	-34.5	3.3	Probable coenzyme A ligase
PA0997	<i>pqsB</i>			A	1.2	-1.2	1.1	-1.0	-12.6	-15.4	-59.2	3.1	Homologous to beta-keto-acyl-acyl carrier protein synthase
PA0998	<i>pqsC</i>			A	1.2	-1.3	1.2	-1.1	-8.2	-13.9	-29.9	2.8	Homologous to beta-keto-acyl-acyl carrier protein synthase
PA0999	<i>pqsD</i>			A	1.6	-1.2	-1.0	-1.3	-8.2	-9.8	-20.2	2.6	3-Oxoacyl-[acyl carrier protein] synthase III
PA1000	<i>pqsE</i>			A	2.1	1.1	1.4	1.3	-7.9	-12.0	-28.9	3.6	Quinolone signal response protein
PA1001	<i>phnA</i>			A	1.9	1.0	-1.0	-1.6	-6.6	-9.7	-20.9	3.2	Anthraniolate synthase component I
PA1002	<i>phnB</i>			A	1.5	-1.1	1.2	-1.3	-7.3	-5.2	-6.3	5.5	Anthraniolate synthase component II
PA1131				B	-1.9	-5.2	-1.4	-2.4	-5.9	-11.1	-2.9	-6.6	Probable major facilitator superfamily transporter
PA1168				A	-6.5	-8.1	-3.8	-3.3	-21.5	-19.1	-26.6	-3.9	Hypothetical protein
PA1190				C	-19.2	-26.3	-9.8	-8.9	-14.6	-12.4	-12.5	-10.4	Conserved hypothetical protein
PA1242				C	-5.7	-10.1	-5.5	-18.7	-19.6	-21.3	-6.7	-9.2	Hypothetical protein
PA1249	<i>aprA</i>			D	-23.5	-54.2	-3.5	-5.3	-7.3	-8.4	-4.1	-2.7	Alkaline metalloproteinase precursor
PA1250	<i>aprI</i>			D	-1.2	-1.5	1.1	-1.5	-9.7	-5.5	-3.0	-1.0	Alkaline proteinase inhibitor AprI
PA1323				C	-11.2	-35.6	-4.3	-9.6	-25.3	-39.3	-56.5	-15.5	Hypothetical protein
PA1324				C	-10.5	-23.4	-4.8	-8.1	-22.1	-26.7	-39.2	-17.0	Hypothetical protein
PA1412				C	-5.5	-11.2	-22.3	-7.0	-5.7	-5.6	-8.8	-6.6	Hypothetical protein
PA1431	<i>rsaL</i>			A	-2.6	-2.2	-1.3	-2.2	-3,619.1	-3,441.9	-3,038.0	-1.9	Regulatory protein RsaL
PA1432	<i>lasI</i>			A	-1.9	1.2	1.8	1.7	-97.0	-133.1	-39.0	-1.5	Autoinducer synthesis protein LasI
PA1471				C	-4.5	-5.8	-2.0	-2.9	-7.8	-5.6	-10.3	-9.7	Hypothetical protein
PA1556				C	15.6	17.5	9.9	9.2	7.9	6.0	7.8	5.3	Probable cytochrome <i>c</i> oxidase subunit
PA1625				C	-3.7	-5.1	-3.7	-7.4	-6.6	-9.0	-6.5	-5.5	Conserved hypothetical protein
PA1657				C	1.6	-1.5	-1.5	3.5	-12.7	-15.4	-12.7	-5.2	Conserved hypothetical protein
PA1662				D	-1.5	-3.5	-2.3	-1.4	-5.7	-9.0	-4.3	-4.4	Probable ClpA/B-type protease
PA1664				C	1.6	-2.4	-1.3	1.1	-5.2	-11.0	-15.5	-8.7	Hypothetical protein
PA1665				A	1.1	-2.7	-2.5	1.1	-5.6	-46.9	-15.1	-2.8	Hypothetical protein
PA1667				A	-1.5	-4.1	-1.1	-1.1	-5.5	-5.1	-5.0	-3.1	Hypothetical protein
PA1669				B	-6.3	-2.8	-2.0	-1.4	-28.6	-12.7	-3.8	-10.3	Hypothetical protein
PA1784				A	-5.5	-14.6	-2.4	-4.4	-9.4	-11.2	-9.9	-1.4	Hypothetical protein
PA1869				C	1.2	-3.2	-1.0	-2.7	-137.1	-287.9	-23.6	-25.1	Probable acyl carrier protein
PA1870				C	-4.3	-11.8	-6.9	-4.8	-22.0	-12.7	-8.8	-13.8	Hypothetical protein
PA1871	<i>lasA</i>		-	C	-10.3	-32.9	-2.9	-9.0	-105.4	-147.8	-31.3	-7.1	LasA protease precursor

PA1874	A	-7.8	-9.3	-6.0	-6.6	-8.0	-6.4	-7.9	1.9	Hypothetical protein
PA1875	A	-7.5	-16.7	-12.3	-20.5	-34.2	-9.0	-6.7	1.1	Probable outer membrane protein precursor
PA1901	C	-1.4	-27.6	-3.1	-4.1	-21.8	-25.3	-19.1	-51.2	Phenazine biosynthesis protein PhzC
PA1902	C	-1.3	-42.9	-3.5	-4.5	-133.5	-132.0	-97.7	-91.0	Phenazine biosynthesis protein PhzD
PA1903	C	-1.3	-20.0	-2.7	-4.5	-17.2	-24.4	-18.6	-19.7	Phenazine biosynthesis protein PhzE
PA1904	C	1.1	-20.8	-3.5	-3.8	-134.0	-61.6	-77.4	-44.7	Probable phenazine biosynthesis protein
PA1905	C	1.2	-15.9	-2.6	-4.4	-14.1	-32.7	-20.8	-30.0	Probable pyridoxamine 5'-phosphate oxidase
PA1914	A	-24.0	-31.9	-5.5	-6.8	-92.8	-15.1	-100.0	-1.4	Conserved hypothetical protein
PA2021	C	-12.2	-14.1	-5.0	-11.5	-8.6	-12.1	-12.2	-13.2	Hypothetical protein
PA2046	C	-5.9	-25.2	-5.9	-8.6	-32.6	-22.1	-38.7	-45.5	Hypothetical protein
PA2067	A	-3.5	-13.4	-2.1	-5.9	-6.0	-5.7	-5.3	-4.2	Probable hydrolase
PA2068	C	-3.2	-13.4	-1.8	-5.5	-8.5	-19.4	-18.5	-25.0	Probable major facilitator superfamily transporter
PA2069	C	-4.2	-70.4	-2.3	-6.7	-29.3	-72.9	-26.1	-26.8	Probable carbamoyl transferase
PA2137	C	-29.3	-107.3	-50.3	-20.8	-11.4	-15.9	-46.4	-71.3	Hypothetical protein
PA2139	C	-28.3	-20.2	-14.1	-20.7	-9.5	-10.0	-7.7	-24.3	Hypothetical protein
PA2141	C	-31.2	-23.4	-8.2	-33.2	-5.8	-45.3	-80.1	-69.0	Hypothetical protein
PA2142	C	-27.1	-18.9	-12.3	-19.5	-13.4	-16.4	-18.2	-26.8	Probable short-chain dehydrogenase
PA2143	C	-82.1	-43.2	-23.0	-42.5	-67.1	-60.1	-191.3	-83.7	Hypothetical protein
PA2144	A	-11.0	-22.1	-6.9	-7.4	-7.3	-20.8	-11.4	-4.8	Glycogen phosphorylase
PA2146	C	-8.5	-11.4	-8.0	-7.4	-6.9	-10.8	-15.5	-9.7	Conserved hypothetical protein
PA2148	A	-5.3	-8.0	-4.5	-5.5	-8.1	-8.1	-6.4	-4.0	Conserved hypothetical protein
PA2149	C	-6.7	-10.6	-4.5	-12.3	-6.3	-7.0	-39.5	-11.2	Hypothetical protein
PA2151	C	-47.9	-71.1	-11.2	-13.5	-39.3	-124.4	-41.2	-30.6	Conserved hypothetical protein
PA2153	C	-15.0	-30.0	-4.2	-6.4	-6.9	-11.3	-19.1	-9.2	1,4-Alpha-glucan branching enzyme
PA2158	C	-23.8	-68.3	-12.3	-43.9	-65.0	-10.3	-35.9	-52.4	Probable alcohol dehydrogenase (Zn dependent)
PA2163	C	-9.0	-9.6	-3.9	-7.9	-8.2	-8.2	-6.9	-5.3	Hypothetical protein
PA2165	C	-16.5	-19.7	-7.5	-14.4	-8.7	-5.4	-6.2	-7.5	Probable glycogen synthase
PA2166	C	-15.8	-19.0	-6.7	-14.8	-15.4	-38.5	-24.4	-19.2	Hypothetical protein
PA2167	C	-11.0	-86.2	-7.4	-18.1	-8.7	-8.1	-7.1	-15.4	Hypothetical protein
PA2170	C	-4.1	-5.1	-3.1	-8.6	-8.2	-14.1	-5.4	-6.5	Hypothetical protein
PA2171	C	-10.1	-9.9	-7.0	-21.7	-27.6	-109.9	-58.6	-22.5	Hypothetical protein
PA2176	C	-50.5	-112.3	-10.0	-14.8	-19.2	-25.2	-30.0	-12.0	Hypothetical protein
PA2178	D	-7.0	-3.6	-3.6	-21.9	-11.2	-6.3	-4.3	-4.9	Hypothetical protein
PA2184	C	-5.0	-6.7	-4.7	-7.3	-15.5	-13.9	-17.5	-20.8	Conserved hypothetical protein
PA2190	C	-13.3	-60.5	-6.3	-11.3	-17.0	-14.9	-9.9	-6.9	Conserved hypothetical protein
PA2193	B	3.5	-1.5	1.7	1.4	-262.2	-256.6	-3.0	-9.1	Hydrogen cyanide synthase HcnA
PA2194	B	2.3	-3.9	-1.1	-1.3	-123.1	-43.1	-2.3	-8.3	Hydrogen cyanide synthase HcnB
PA2195	B	1.7	-4.3	1.1	-1.5	-20.5	-38.9	-2.7	-6.9	Hydrogen cyanide synthase HcnC
PA2300	C	-20.2	-45.4	-8.1	-20.7	-33.8	-66.7	-82.8	-34.0	Chitinase
PA2302	A	-3.3	-9.6	-1.8	-3.3	-13.2	-17.5	-10.1	-1.2	Probable nonribosomal peptide synthetase
PA2303	A	-3.1	-10.5	-1.5	-2.1	-87.9	-212.8	-59.2	-1.3	Hypothetical protein
PA2304	A	-2.8	-6.9	-1.5	-2.3	-18.1	-23.2	-9.2	1.1	Hypothetical protein
PA2305	A	-1.7	-2.8	1.1	-1.5	-6.5	-16.7	-8.8	1.1	Probable nonribosomal peptide synthetase
PA2414	C	-12.2	-51.7	-4.6	-11.3	-32.2	-17.6	-16.4	-7.7	L-Sorbose dehydrogenase
PA2415	C	-5.1	-8.6	-2.9	-6.4	-13.8	-9.3	-9.2	-5.2	Hypothetical protein

Continued on following page

TABLE 3—Continued

Gene in <i>P. aeruginosa</i>	Gene	<i>ppvB^b</i>	Regulation by AZM ^c	Reg. ^d	Relative expression							Description	
					AZM (2 µg/ml)	AZM (8 µg/ml)	CFT	CPR	<i>last-rhlI</i>	<i>lasR-rhlR</i>	<i>lasR</i>		<i>rhlR</i>
PA2423			+	A	-1.2	-1.2	-1.1	-1.6	-5.8	-5.9	-7.6	1.6	Hypothetical protein
PA2433				C	-25.6	-70.7	-6.4	-24.8	-25.9	-67.3	-70.0	-32.8	Hypothetical protein
PA2485		+		C	-7.9	-8.1	-2.3	-7.0	-12.0	-18.0	-15.5	-12.3	Hypothetical protein
PA2486				C	-9.3	-18.4	-2.3	-4.5	-8.9	-10.6	-8.5	-5.7	Hypothetical protein
PA2566				A	-4.2	-6.8	1.1	-3.3	-9.6	-7.1	-6.3	-1.7	Conserved hypothetical protein
PA2570	<i>lecA</i>			C	-25.9	-76.4	-4.5	-14.9	-45.4	-49.2	-27.2	-30.3	LecA
PA2587	<i>pqsH</i>	-		A	-1.6	-2.1	-1.6	-2.3	-46.1	-36.1	-20.4	1.3	Probable FAD-dependent mono-oxygenase
PA2588				A	-3.3	-4.5	-2.4	-4.3	-6.7	-34.3	-87.8	-2.4	Probable transcriptional regulator
PA2591				D	-1.1	-1.2	1.1	-1.3	-7.1	-14.8	-2.2	1.1	Probable transcriptional regulator
PA2592				D	1.3	-1.2	1.4	-1.1	-6.7	-17.9	-3.8	-1.6	Probable periplasmic spermidine/putrescine-binding protein
PA2708				C	-4.1	-4.7	-4.2	-6.1	-5.8	-7.8	-8.7	-12.5	Hypothetical protein
PA2747		+		C	-3.8	-7.2	-2.6	-7.7	-32.6	-37.3	-37.5	-10.3	Hypothetical protein
PA2751				C	-3.4	-4.4	-2.4	-4.3	-8.5	-7.0	-10.3	-6.0	Conserved hypothetical protein
PA2754				C	-5.0	-9.2	-2.4	-3.5	-11.5	-7.3	-9.1	-5.4	Conserved hypothetical protein
PA2777				C	-4.4	-26.5	-4.6	-10.1	-8.4	-6.2	-10.5	-8.9	Conserved hypothetical protein
PA2873				D	-1.2	-7.8	-1.3	-3.1	-13.2	-5.8	-1.6	-1.3	Hypothetical protein
PA2937				C	-23.3	-45.2	-5.2	-13.2	-6.9	-8.6	-9.1	-5.4	Hypothetical protein
PA2939				A	-10.6	-38.6	-3.5	-6.5	-39.8	-9.1	-12.7	1.8	Probable aminopeptidase
PA3231				C	-17.7	-21.9	-6.4	-41.9	-33.7	-13.2	-27.7	-10.0	Hypothetical protein
PA3273				C	-10.2	-20.7	-4.0	-19.7	-13.3	-13.2	-13.2	-27.1	Hypothetical protein
PA3274				C	-11.5	-49.1	-7.0	-27.4	-45.1	-37.5	-112.5	-17.5	Hypothetical protein
PA3326				D	-1.5	-4.2	1.3	-1.0	-12.3	-11.0	-2.4	-4.1	Probable Clp family ATP-dependent protease
PA3327		-		C	-1.4	-5.0	-1.0	1.3	-5.4	-12.7	-5.0	-9.8	Probable nonribosomal peptide synthetase
PA3328		-		B	-1.3	-4.5	-1.1	-1.2	-6.5	-26.0	-3.2	-12.4	Probable FAD-dependent mono-oxygenase
PA3329				C	1.3	-3.8	1.9	1.3	-54.9	-173.5	-6.1	-69.3	Hypothetical protein
PA3330				B	-4.3	-1.1	1.7	1.3	-62.4	-179.6	-4.4	-19.5	Probable short-chain dehydrogenase
PA3331				B	1.1	-4.1	1.8	1.6	-19.8	-62.0	-3.8	-23.8	Cytochrome P450
PA3332				B	1.3	-4.3	1.3	1.2	-10.2	-18.7	-5.0	-17.7	Conserved hypothetical protein
PA3333	<i>fabH2</i>			B	-1.3	-7.4	1.0	-1.2	-17.0	-33.4	-4.1	-10.4	3-Oxoacyl-[acyl carrier protein] synthase III
PA3334				C	1.2	-3.8	1.8	1.7	-42.8	-17.4	-5.1	-11.2	Probable acyl carrier protein
PA3335				D	1.1	-3.9	1.8	1.2	-18.4	-9.4	-2.3	-3.9	Hypothetical protein
PA3336				B	-9.3	-2.4	1.5	-1.3	-8.2	-9.3	-4.3	-22.8	Probable major facilitator superfamily transporter
PA3361	<i>lecB</i>			C	-1.4	-12.3	-1.6	-2.5	-22.4	-28.9	-13.2	-15.3	Fucose-binding lectin PA-III
PA3369		+		C	-7.1	-11.9	-4.2	-12.7	-19.9	-42.2	-36.5	-14.2	Hypothetical protein
PA3370		+		C	-12.9	-26.1	-8.7	-18.8	-39.0	-66.4	-87.1	-31.2	Hypothetical protein
PA3371		+		C	-6.7	-11.7	-7.4	-18.8	-23.8	-51.7	-69.2	-25.7	Hypothetical protein
PA3460				A	-12.9	-28.0	-2.0	-3.2	-5.0	-7.0	-4.7	-4.7	Probable acetyltransferase
PA3476	<i>rhlI</i>			D	-1.6	-2.0	-1.2	-1.6	-19.0	-41.4	-2.4	-1.8	Autoinducer synthesis protein RhlI
PA3478	<i>rhlB</i>			C	-7.3	-31.7	-2.4	-6.3	-120.2	-122.7	-19.5	-53.0	Rhamnosyltransferase chain B
PA3479	<i>rhlA</i>			C	-5.0	-28.3	-1.8	-5.3	-320.2	-196.0	-14.5	-55.0	Rhamnosyltransferase chain A
PA3520				C	-3.7	-8.9	-2.7	-4.4	-28.3	-18.3	-13.9	-7.4	Hypothetical protein

PA3581	<i>glpF</i>	C	-163.4	-43.8	-14.0	-99.3	-61.0	-88.9	-19.1	-35.1	Glycerol uptake facilitator protein
PA3584	<i>glpD</i>	C	-9.2	-12.2	-8.1	-6.3	-5.7	-20.4	-12.0	-8.4	Glycerol-3-phosphate dehydrogenase
PA3691		C	-5.1	-13.3	-2.0	-3.1	-11.6	-19.2	-28.8	-12.3	Hypothetical protein
PA3692		C	-7.9	-26.7	-3.1	-4.4	-30.7	-16.8	-37.5	-12.3	Probable outer membrane protein precursor
PA3724	<i>lasB</i>	C	-3.5	-22.8	-2.5	-6.1	-167.2	-224.3	-15.1	-5.2	Elastase LasB
PA3734		C	-5.0	-9.3	-3.5	-4.9	-7.2	-21.3	-6.6	-8.5	Hypothetical protein
PA3788		A	-4.7	-6.5	-2.7	-4.8	-9.2	-5.8	-6.1	-4.0	Hypothetical protein
PA3819		C	-3.5	-5.0	-3.7	-3.6	-6.3	-7.0	-9.7	-7.1	Conserved hypothetical protein
PA3888		A	-2.5	-3.2	-2.1	-3.5	-5.8	-10.2	-10.5	-4.6	Probable permealase of ABC transporter
PA3890		C	-3.8	-5.1	-2.6	-6.7	-17.8	-87.2	-64.7	-31.8	Probable permealase of ABC transporter
PA3904		A	1.9	1.5	1.5	1.6	-68.3	-40.7	-172.1	-1.0	Hypothetical protein
PA3906		A	1.1	-1.1	-1.2	-1.1	-7.5	-176.5	-9.5	-1.4	Hypothetical protein
PA3907		A	1.4	-1.1	1.2	2.0	-5.8	-13.4	-5.2	1.0	Hypothetical protein
PA3908		D	-1.0	-2.6	-1.5	1.2	-6.2	-49.8	-4.5	1.1	Hypothetical protein
PA4078	*	A	-6.2	-15.8	-4.3	-6.4	-5.6	-11.0	-7.4	-4.8	Probable nonribosomal peptide synthetase
PA4130		D	1.4	-1.9	-1.3	1.1	-9.6	-6.2	-3.3	1.5	Probable sulfite or nitrite reductase
PA4133		A	2.6	-1.8	-2.1	1.1	-57.4	-13.0	-6.1	1.6	Cytochrome <i>c</i> oxidase subunit (cbb3 type)
PA4134		A	1.9	-3.3	-3.3	-1.8	-12.0	-5.6	-18.0	-1.2	Hypothetical protein
PA4139		A	-9.2	-11.6	-4.3	-3.2	-6.1	-10.2	-7.0	-4.2	Hypothetical protein
PA4141		C	-2.0	-12.2	-1.1	-1.6	-73.1	-105.0	-27.6	-57.6	Hypothetical protein
PA4142		C	-2.1	-11.7	-2.1	-4.2	-27.2	-71.7	-25.6	-26.4	Probable secretion protein
PA4143		C	-2.6	-14.4	-2.1	-3.4	-8.1	-15.3	-7.3	-9.0	Probable toxin transporter
PA4171	<i>prpL</i>	C	-7.4	-10.0	-4.9	-6.3	-5.4	-15.5	-13.6	-9.5	Probable protease
PA4175		A	-12.0	-25.4	-6.7	-15.8	-28.4	-47.1	-30.4	-1.6	Pvds-regulated endoprotease, lysyl class
PA4209	<i>phzM</i>	C	1.8	-10.8	-1.9	-1.9	-9.8	-18.8	-11.9	-7.5	Probable phenazine-specific methyltransferase
PA4210	<i>phzA1</i>	C	-1.0	-11.8	-2.3	-5.6	-65.1	-123.9	-79.7	-20.2	Probable phenazine biosynthesis protein
PA4211	<i>phzB1</i>	C	1.4	-15.5	-2.2	-2.3	-51.0	-61.2	-59.7	-46.1	Probable phenazine biosynthesis protein
PA4217	<i>phzS</i>	C	1.4	-9.6	-2.2	-2.2	-11.4	-12.9	-8.9	-10.9	Flavin-containing mono-oxygenase
PA4290		C	-23.4	-19.0	-26.8	-25.3	-7.2	-6.0	-14.2	-15.9	Probable chemotaxis transducer
PA4345		C	-5.3	-7.4	-3.9	-6.9	-10.6	-9.1	-7.7	-6.6	Hypothetical protein
PA4377		A	-13.9	-12.2	-3.5	-6.2	-10.1	-7.0	-7.8	-3.2	Hypothetical protein
PA4677		A	-6.8	-4.2	-3.1	-5.1	-17.0	-40.0	-22.5	-1.1	Hypothetical protein
PA4738		C	-6.8	-22.5	-5.2	-11.5	-39.6	-54.4	-58.1	-16.6	Conserved hypothetical protein
PA4739		C	-4.8	-12.9	-3.4	-5.7	-34.3	-34.3	-43.8	-16.6	Conserved hypothetical protein
PA4876	<i>osmE</i>	C	-5.3	-10.4	-3.4	-4.8	-18.2	-9.1	-10.8	-6.7	Osmotically inducible lipoprotein OsmE
PA4880		C	-6.8	-17.3	-4.5	-4.4	-6.0	-29.6	-12.9	-12.9	Probable bacterioferritin
PA5097		D	-4.2	-18.2	-12.5	-5.8	-9.5	-5.2	-3.4	-4.8	Probable amino acid permease
PA5099		C	-188.0	-353.2	-54.1	-450.6	-16.2	-14.7	-40.5	-39.9	Probable transporter
PA5212		C	-3.1	-5.6	-1.7	-2.5	-8.8	-8.7	-12.1	-5.7	Hypothetical protein
PA5220		B	-2.1	-5.5	-1.7	-2.3	-7.7	-8.9	-4.2	-6.9	Hypothetical protein
PA5235	<i>glpT</i>	B	-4.3	-2.7	-2.6	-5.0	-15.2	-32.4	-2.8	-13.9	Glycerol-3-phosphate transporter
PA5297	<i>poxB</i>	D	-3.1	-9.4	-2.0	-3.5	-17.8	-6.0	-3.8	-2.8	Pyruvate dehydrogenase (cytochrome)
PA5480		B	-10.6	-2.6	1.0	-4.2	-14.2	-19.9	-2.5	-6.1	Hypothetical protein

Continued on following page

TABLE 3—Continued

Gene in <i>P. aeruginosa</i>	Gene	<i>pprB</i> ^b	Regulation by AZM ^c	Reg. ^d	Relative expression							Description	
					AZM (2 µg/ml)	AZM (8 µg/ml)	CFT	CPR	<i>lasI-rhlI</i>	<i>lasR-rhlR</i>	<i>lasR</i>		<i>rhlR</i>
PA5481			+	C	-9.6	-30.8	-7.3	-15.0	-45.6	-50.7	-96.6	-22.6	Hypothetical protein
PA5482			+	C	-5.1	-15.2	-3.5	-9.2	-59.6	-43.0	-81.1	-12.2	Hypothetical protein

^a The effects of sub-MICs of AZM, CFT and CPR on genes belonging to the QS regulon, which is defined elsewhere (74). This regulon consists of genes downregulated in both a $\Delta lasI-\Delta rhlI$ mutant and a $\Delta lasR-\Delta rhlR$ mutant of *P. aeruginosa* PAO1. Genes downregulated more than five times are in boldface, and genes upregulated more than five times are underlined.

^b *pprB* genes upregulated more than five times (+) or downregulated five times (-) in a $\Delta pprB$ mutant of PAO1, as described elsewhere (25).

^c Genes previously reported to be upregulated (+) or downregulated (-) by AZM, as described elsewhere (60).

^d Genes were assigned to one of the following four QS regulation (Reg.) groups: group A, Las-dependent genes; group B, Rhl-dependent genes; group C, genes requiring both a functional Las system and a functional Rhl system; group D, Las-Rhl-regulated genes. See the text for further explanations.

while CPR and CFT reduced the level of *rhlA* expression 10- and 3-fold, respectively.

AZM, CFT, and CPR alter the expression of a variety of genes. A few genes were consistently upregulated by AZM, CFT, and CPR. The QS repressible genes PA1556 (cytochrome *c* oxidase subunit) and PA1557 were induced more than five times. PA2260 (a hypothetical protein previously identified [89] to be a QS repressible gene) was induced by AZM and CFT but not by CPR.

Several genes were induced only by AZM. A high proportion of these genes have also previously been reported to be AZM inducible (60), although there were a number of differences in the experimental setup between the current study and the previous study (60). For example, the previous study (60) used the *P. aeruginosa* PAO1 strain DSM 1707 instead of the *P. aeruginosa* PAO1 strain from the Pseudomonas Genetic Stock Center used here. Moreover, the study by Nalca et al. (60) also applied the complex and rich brain heart infusion medium instead of minimal medium. Many of the genes induced by AZM encode ribosomal subunits, such as PA3742 (*rplS*) and PA4671 (probable ribosomal protein). The overexpression of ribosomal genes may be a mechanism that compensates for the effect of AZM, which targets the 50S ribosomal subunit. The initiation factor *infA* (PA2619) was also induced in the presence of AZM, in accordance with previous findings (60). In agreement with the findings of two recent studies (12, 53), we found that CPR induces the expression of the pyocin S genes (PA0985 for pyocin S5, PA1150 for pyocin S2, and PA3866 for the pyocin S3-like protein). The S-type pyocins are protease-sensitive bacteriocins and have colicin-like structures and modes of action.

The ribosome modulation factor (*rmf*, PA4296) was repressed by AZM but was not affected by treatment with CFT or CPR. Previous transcriptome analyses (60) also showed that *rmf* is repressed by AZM. The *rmf* of *P. aeruginosa* shows a high degree of homology to the *rmf* of *E. coli* (similarity, 66%). *rmf* mutants of *E. coli* have been shown to gradually lose their viability in stationary-phase cultures (61); however, a recent study concluded that *rmf* (in *E. coli*) may be required only under competitive growth conditions (13).

AZM induces expression of genes related to T3S. In accordance with previous findings (60), we also found that AZM induced the expression of genes involved in type III secretion (T3S). *P. aeruginosa* uses its T3S system to deliver several cytotoxic products into the cytosol of eukaryotic cells. The expression of T3S increases the pathogenicity of *P. aeruginosa* and is associated with more severe infections and higher rates of mortality in animal models (1, 75, 78). QS has been shown to negatively modulate the expression of the T3S system in *P. aeruginosa* (10), which means that the T3S system is downregulated under high-cell-density conditions and thus presumably plays a role primarily in the early stages of the infection. Genes in the T3S system, such as PA1700 and PA1701 (conserved hypothetical proteins in T3S), *pcrGV* (PA1705 regulator in T3S and PA1706), *exsB* (PA1712, exoenzyme S synthesis protein B), *exsD* (PA1714, part of the *pscB* to *pscL* operon), and *pscBCEO* (PA1715, T3S export apparatus protein; PA1716, T3S outer membrane protein; PA1718, T3S export protein; PA1696, translocation protein in T3S), were upregulated by




FIG. 8. LasR, RhlR, and LasR-RhlR specificities of AZM (white bars), CFT (black bars), CPR (light gray bars), and C30 (dark gray bars). See the text for the definitions of groups (Gr.) A, B, C, and D.

AZM. Neither CFT nor CPR, however, induced the genes in the T3S system gene cluster from PA1690 to PA1725.

AZM, CPR, CFT, and the two-component regulator system *pprAB*. The QS-induced chemotaxis transducer PA4290 was downregulated by AZM, CFT, and CPR. AZM and CPR also repressed the main part of the genes in the region from PA4290 to PA4306. Many of these genes have been shown in previous studies to be regulated by QS. In those studies, QS genes were identified by mapping genes downregulated in $\Delta lasI-\Delta rhlI$ mutants and restored to wild-type levels by addition of signal molecules (38, 80, 89). The two-component regulator system *pprAB* (PA4293, PA4296) was downregulated by AZM and CPR but not by CFT. *pprA* was downregulated 16 and 6 times by AZM and CPR, respectively, and *pprB* downregulated 10 and 5 times by AZM and CPR, respectively. These results obtained by DNA microarray analysis were confirmed by RT-PCR, which showed that AZM reduced the levels of *pprA* and *pprB* expression 19- and 15-fold, respectively, whereas CPR downregulated the level of *pprA* expression 11-fold and the level of *pprB* expression 18-fold. RT-PCR also confirmed that the effect of CFT on *pprAB* expression was insignificant.

Docking of AZM, CFT, and CPR against the LBD of the LasR protein. In order to explore the possibility that AZM, CFT, and CPR directly interact with the LasR protein, we performed in silico docking of the three antibiotics with the recently published LBD of the LasR protein (11). None of the three antibiotics gave a high-affinity score in the docking experiment. In contrast, previously reported QSI compounds (e.g., C30, patulin, furanone, and 4-nitropyridine-*N*-oxide) exhibited high-affinity scores (Table 4). The in silico analysis indicates that AZM, CFT, and CPR do not fit inside the binding site cavity of LasR because they exhibit steric clashes with the protein target. This suggests that the effects of AZM, CFT, and CPR on QS may be due to a mechanism of action other than a direct interaction with the LasR protein.

DISCUSSION

In the present paper, we present the results from the screening of 12 antibiotics for QSI activity obtained with QSIS1. Three of the 12 antibiotics, AZM, CFT, and CPR, showed strong positive results in the screening system and were further investigated for their effects on the production of virulence factors in *P. aeruginosa*. AZM, CFT, and CPR exhibit diverse

mechanisms of antimicrobial action and are structurally very different. It is thus intriguing that all three antibiotics were active in QSIS1. Both CFT and piperacillin are β -lactams that target penicillin binding protein 3 (PBP 3); however, only CFT showed actual activity in the QSI screening system, indicating that the QSI activity is likely not PBP 3 dependent. We carefully selected the concentrations of the three antibiotics which had no effect on growth. Our experience from the screening of large numbers of compounds from compound libraries is that toxic or growth-inhibitory compounds may show QSI-like activities (unpublished observations). This is likely to be caused by stress-like effects rather than “true” QSI activity (such as an interaction with the QS receptor), since the effect on QS-controlled gene expression disappears when the concentration of the compound is lowered to a level that allows the bacteria to attain the optimum growth rate in a given medium. Furthermore, since the obstruction of QS (either by mutation or by antagonistic drugs) does not inhibit growth, we consider it essential to investigate the potential QSI activities of the three antibiotics under conditions that do not affect growth. When they were administered at such concentrations, AZM, CPR, and CFT inhibited the production of the important virulence factors chitinase, protease, and elastase and the production of the heat-stable hemolysin rhamnolipid. Our recent data suggest that rhamnolipid functions as a shield to protect biofilms against the predatory behavior of neutrophils, and rhamnolipid may thus be a pivotal virulence factor (45). Investigations have shown that biofilms of *P. aeruginosa* QS mutants have reduced tolerance to antibiotics compared with that of their wild-type parents, and inhibition of QS has been shown to promote the eradication of biofilms by antimicrobial treatments and make the biofilm more susceptible to phagocytosis by neutrophils (20, 38, 73). Rhamnolipid and the adhesive lectin LecA were downregulated more than fivefold by AZM and CPR. Both products are regulated by QS and are involved in biofilm maturation (19, 23, 52, 65). Repression of these genes may contribute to the development of the less structured and more antibiotic sensitive biofilm seen in QS mutants. Moreover, rhamnolipid and lectin LecA have been reported to have cytotoxic effects on mammalian cells, including neutrophils and macrophages, and are also important in the *P. aeruginosa* infectious process (5, 45, 57, 58, 81, 82, 95).

Another important virulence component of *P. aeruginosa* is the T3S system, which is negatively regulated by QS in *P. aeruginosa*. The DNA microarray data presented here suggest that AZM increases the level of expression of T3S genes, in

TABLE 4. In silico docking

Name of ligand	Rerank score
OdDHL (template)	-107.3
3-Oxo-C ₁₂	-94.5
2-Heptylthioacetyl homoserine lactone	-88.8
BHL.....	-82.2
Patulin.....	-69.8
4-Nitropyridine- <i>N</i> -oxide.....	-62.9
C30.....	-50.4
CPR.....	317.3
CFT.....	373.5
AZM.....	3,316.2

accordance with the QS-inhibiting properties of AZM reported here and by others. Interestingly, a previous study (50) showed that treatment of *P. aeruginosa* with sub-MICs of macrolides (AZM, clarithromycin, erythromycin) prior to the intranasal challenge of mice significantly enhanced the mortality rate (from 0% to 80 to 100%). In the same study, however, macrolide antibiotics were also administered to mice after the inoculation of *P. aeruginosa* PAO1, and in that case, the rate of mortality among the mice did not increase (50). Thus, it seems that the activation of genes by macrolides causes enhanced virulence only if the macrolides are applied prophylactically and not as a part of the treatment. Nevertheless, the potential risk, along with the potential benefits, of inducing virulence and cytotoxicity through the T3S system should be considered when patients infected with *P. aeruginosa* are treated with AZM (and possibly other macrolides).

As mentioned above, the two-component regulator system *pprAB* (PA4293 and PA4296, respectively) was downregulated more than fivefold by AZM and CPR but not by CFT; these results were confirmed by RT-PCR. The expression of *pprAB* has been shown to be regulated by QS (80, 89), but the gene products have also been identified to have activities that modulate QS, representing yet another autofeedback system of QS in *P. aeruginosa* (25). The knockout of *pprB* leads to reduced influx of the *P. aeruginosa* QS signal OdDHL, thereby globally influencing the expression of QS-regulated genes (25); but PprB is also involved in the regulation of sensitivity to antibiotics. The overexpression of *pprB* results in increased sensitivity to antibiotics, especially aminoglycosides, probably due to a decrease in membrane permeability (91). In vivo and in vitro studies have shown that *P. aeruginosa* develops increasing tolerance to aminoglycosides following pretreatment with these antimicrobials (18, 47, 93). Interestingly, cells from *P. aeruginosa* biofilms exposed to sub- and suprainhibitory concentrations of CPR have also recently been demonstrated to display transient increased levels of resistance to CPR (64). We speculate that the underlying mechanism by which AZM and CPR (and perhaps also CFT) exert their action on QS is by means of an adaptive response which downregulates *pprB* and perhaps also other membrane regulators or transporters, such as PA5099. This causes the membrane permeability to decrease, thereby providing the cells with transient protection against the toxic effects of the antibiotics. However, at the same time, the lower membrane permeability decreases the level of OdDHL influx, which results in a reduction in the levels of expression of QS-regulated genes. Support for this hypothesis can be found in the observation that the three antibiotics have higher specificities for *las*-regulated genes than *rhl*-regulated genes (group B). The *rhl* signal molecule BHL is capable of freely diffusing over the membrane and is, unlike OdDHL, not dependent on active transport.

A previous study (25) identified 53 genes that were downregulated more than fivefold by a Tn5 insertion in *pprB*. In comparison, as described in the present report, we found that the levels of expression of 38 of those *pprB*-regulated genes that were downregulated more than fivefold in the previously mentioned study (25) were reduced to half that level or less in the presence of AZM. This included 21 genes which were downregulated more than fivefold. We found that CPR downregulated 23 of the 53 genes more than twofold, including 6




FIG. 9. Inhibitors of and signal molecules for *P. aeruginosa* QS. (a) 4-Nitropyridine-*N*-oxide (73); (b) C30 (38); (c) 2-heptylthioacetyl homoserine lactone (69a); (d) 3-oxo-*C*₁₂-2-aminophenol (83a); (e) BHL; (f) patulin (74); (g) OdDHL.

genes that were downregulated more than fivefold. CFT caused a downregulation of 15 of the 53 genes downregulated by the *pprB* mutation, but none of the genes were downregulated more than fourfold. Genes reported to be affected by PprB include virulence factor genes such as *lasB*, *rhlAB*, *lecA*, and *prpL* (25), which are also downregulated by AZM and CPR. The levels of expression of the QS transcriptional regulator RhlR and the AHL synthetase genes *lasI* and *rhlI* were also decreased by the *prpB*-knockout mutation (25). This is not surprising, as *rhl* is subordinate to the *las* system and the AHL synthetase genes are under QS control, which may provide a negative feedback that enhances the QS-repressing effect of the *pprB* mutation.

QSIs constructed with AHL scaffolds are expected to bind to the LuxR homologue without activating it, thereby blocking the binding and activation by AHLs. Thus, these QS antagonists have a high degree of structural similarity to AHLs. Other QSIs which share the 2(5*H*)-furanone moiety, such as the natural occurring halogenated furanones, and the mycotoxins patulin and penicillic acid seem to exert their effects by destabilizing LuxR and thereby inducing the turnover of QS regulator proteins (56, 74). Figure 9 shows the structures of BHL, OdDHL, and previously published QSI compounds.

The three antibiotics AZM, CFT, and CPR are structurally very different from all QSIs described until now; moreover, like many other antibiotics, they are very bulky molecules (Fig. 10). In silico docking of AZM, CFT, and CPR to the LBD of the LasR protein showed that the antibiotics have a low affinity for the LasR receptor site, mainly due to spatial penalties. This supports the idea that these antibiotics may exert their QS regulatory effects through mechanisms different from those of the other QSIs, thereby opening new and interesting possibilities for combination therapy with QSIs with different modes of action. The usefulness of the combination of QSIs and traditional bactericidal compounds has been proved by several in vitro experiments (20, 38, 74). The understanding of some antibiotics' dual course of action may have the potential to




FIG. 10. Structures of the three antibiotics investigated for potential quorum sensing inhibitory activities. (A) AZM; (B) CFT; (C) CPR.

further add to therapies against pathogens in which the key element attenuates rather than kills the bacteria directly. This strategy seeks to allow the host defense system to eliminate the attenuated bacteria.

Most antibiotics originate from natural sources, such as soil microorganisms, or are derivatives of compounds originating from nature. A general point of view has been that the ecological purpose of antimicrobials in the environment is to fight competitors. However, the concentrations of free antibiotics in soil are likely to be well below the MICs.

As described in this paper and by others (53, 94), sub-MICs of antibiotics may have various effects on bacteria which are totally different from the effects of high doses; for example, they may enhance biofilm formation, alter motility patterns, increase cytotoxicity, and cause changes in the expression of virulence factors (39, 53, 60). It has been suggested that antibiotics in nature may function as intermicrobial signals rather than as bullets aimed at killing enemies in order to defend niches or food (53). This hypothesis offers an evolutionary explanation of why some antibiotics may be capable of interfering with QS-controlled gene expression. In line with the findings presented in the present report, it has recently been shown that OdDHL (and other 3-oxo-AHLs) as well as its tetramic acid degradation product exhibits activities against gram-positive bacteria (48), suggesting that the signal molecules, like some antibiotics, possess dual activities. Furthermore, our data suggest that antibiotics which function to down-regulate QS in *P. aeruginosa* may be administered at sub-MICs (where there is a reduced selection pressure for the development of resistance) to block QS and thereby attenuate the activity of the pathogen. Such interesting properties highlight the huge unexplored potential of using compounds derived from natural scaffolds in the search for new pharmaceuticals with a variety of targets (i.e., multifunctional antimicrobials). It seems that we have not fully explored and exploited the potential of the drugs that we know today.

The dual activities of some antibiotics (e.g., AZM, CFT, and CPR, investigated here) may help explain why CFT has been shown to be superior to comparative antipseudomonal antibiotic regimens for patients with CF in the Danish CF center (69). Caution must be taken, however. The subsequent shift to CFT as the predominant antipseudomonal antibiotic in the Danish CF center led to the epidemic spread of a multiresistant nonmucoid strain (68). A previous study (35) showed that sub-MICs of antibiotics may still modulate transcriptional patterns in members of the beneficial human flora, causing a

variety of unwanted effects. Likewise, it has been shown that the long-term use of AZM by patients with CF and chronic *P. aeruginosa* infections may lead to macrolide resistance in *Staphylococcus aureus* in these patients (87). This underlines the requirement for the further development of QSIs without classical bactericidal or bacteriostatic activity.

ACKNOWLEDGMENTS

This work was supported by grants from the German Mukoviszidose e.v., Deutsche Forschungsgemeinschaft, and the Danish Research Council (STF grant no. 2052-03-0013) to M.G.

REFERENCES

- Ader, F., R. Le Berre, K. Faure, P. Gosset, O. Epaulard, B. Toussaint, B. Polack, E. Nowak, N. B. Viget, E. Kipnis, and B. P. Guery. 2005. Alveolar response to *Pseudomonas aeruginosa*: role of the type III secretion system. *Infect. Immun.* **73**:4263–4271.
- Andersen, J. B., A. Heydorn, M. Hentzer, L. Eberl, O. Geisenberger, B. B. Christensen, S. Molin, and M. Givskov. 2001. *gfp*-based *N*-acyl-homoserine-lactone monitors for detection of bacterial communication. *Appl. Environ. Microbiol.* **67**:575–585.
- Andrade, S. S., R. N. Jones, A. C. Gales, and H. S. Sader. 2003. Increasing prevalence of antimicrobial resistance among *Pseudomonas aeruginosa* isolates in Latin American medical centres: 5 year report of the SENTRY Antimicrobial Surveillance Program (1997–2001). *J. Antimicrob. Chemother.* **52**:140–141.
- Anwar, H., and J. W. Costerton. 1990. Enhanced activity of combination of tobramycin and piperacillin for eradication of sessile biofilm cells of *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **34**:1666–1671.
- Bajolet-Laudinat, O., S. Girod-de Bentzmann, J. M. Tournier, C. Madoulet, M. C. Plotkowski, C. Chippaux, and E. Puchelle. 1994. Cytotoxicity of *Pseudomonas aeruginosa* internal lectin PA-I to respiratory epithelial cells in primary culture. *Infect. Immun.* **62**:4481–4487.
- Baumann, U., M. King, E. M. App, S. Tai, A. König, J. J. Fischer, T. Zimmermann, W. Sextro, and H. von der Hardt. 2004. Long term azithromycin therapy in cystic fibrosis patients: a study on drug levels and sputum properties. *Can. Respir. J.* **11**:151–155.
- Beatson, S. A., C. B. Whitchurch, A. B. T. Semmler, and J. S. Mattick. 2002. Quorum sensing is not required for twitching motility in *Pseudomonas aeruginosa*. *J. Bacteriol.* **184**:3598–3604.
- Berger, M., and M. Konstan. 1999. Immunopathogenesis of cystic fibrosis lung disease, p. 115–143. *In* J. R. Yankaskas and M. R. Knowles (ed.), *Cystic fibrosis in adults*. Lippincott-Raven, Philadelphia, PA.
- Berka, R. M., and M. L. Vasil. 1982. Phospholipase C (heat-labile hemolysin) of *Pseudomonas aeruginosa*: purification and preliminary characterization. *J. Bacteriol.* **152**:239–245.
- Bleves, S., C. Soscia, P. Nogueira-Orlandi, A. Lazdunski, and A. Filloux. 2005. Quorum sensing negatively controls type III secretion regulon expression in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **187**:3898–3902.
- Bottomley, M. J., E. Muraglia, R. Bazzo, and A. Carfi. 2007. Molecular insights into quorum sensing in the human pathogen *Pseudomonas aeruginosa* from the structure of the virulence regulator LasR bound to its auto-inducer. *J. Biol. Chem.* **282**:13592–13600.
- Brazas, M. D., and R. E. Hancock. 2005. Ciprofloxacin induction of a susceptibility determinant in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **49**:3222–3227.
- Bubunenko, M., T. Baker, and D. L. Court. 2007. Essentiality of ribosomal and transcription antitermination proteins analyzed by systematic gene replacement in *Escherichia coli*. *J. Bacteriol.* **189**:2844–2853.

14. Ceri, H., M. E. Olson, C. Stremick, R. R. Read, D. Morck, and A. Buret. 1999. The Calgary biofilm device: new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms. *J. Clin. Microbiol.* **37**:1771–1776.
15. Ciofu, O., V. Fussing, N. Bagge, C. Koch, and N. Hoiby. 2001. Characterization of paired mucoid/non-mucoid *Pseudomonas aeruginosa* isolates from Danish cystic fibrosis patients: antibiotic resistance, beta-lactamase activity and RiboPrinting. *J. Antimicrob. Chemother.* **48**:391–396.
16. Clark, D. J., and O. Maaloe. 1967. DNA replication and the division cycle in *Escherichia coli*. *J. Mol. Biol.* **23**:99–112.
17. Costerton, J. W., P. S. Stewart, and E. P. Greenberg. 1999. Bacterial biofilms: a common cause of persistent infections. *Science* **284**:1318–1322.
18. Daikos, G. L., G. G. Jackson, V. T. Lolans, and D. M. Livermore. 1990. Adaptive resistance to aminoglycoside antibiotics from first-exposure down-regulation. *J. Infect. Dis.* **162**:414–420.
19. Davey, M. E., N. C. Caiazza, and G. A. O'Toole. 2003. Rhamnolipid surfactant production affects biofilm architecture in *Pseudomonas aeruginosa* PAO1. *J. Bacteriol.* **185**:1027–1036.
20. Davies, D. G., M. R. Parsek, J. P. Pearson, B. H. Iglewski, J. W. Costerton, and E. P. Greenberg. 1998. The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science* **280**:295–298.
21. de Nys, R., M. Givskov, N. Kumar, S. Kjelleberg, and P. D. Steinberg. 2006. Furanones. *Prog. Mol. Subcell. Biol.* **42**:55–86.
22. DeVries, C. A., and D. E. Ohman. 1994. Mucoid-to-nonmucoid conversion in alginate-producing *Pseudomonas aeruginosa* often results from spontaneous mutations in *algT*, encoding a putative alternate sigma factor, and shows evidence for autoregulation. *J. Bacteriol.* **176**:6677–6687.
23. Diggle, S. P., R. E. Stacey, C. Dodd, M. Camara, P. Williams, and K. Winzer. 2006. The galactophilic lectin, LecA, contributes to biofilm development in *Pseudomonas aeruginosa*. *Environ. Microbiol.* **8**:1095–1104.
24. DiMango, E., H. J. Zar, R. Bryan, and A. Prince. 1995. Diverse *Pseudomonas aeruginosa* gene products stimulate respiratory epithelial cells to produce interleukin-8. *J. Clin. Investig.* **96**:2204–2210.
25. Dong, Y. H., X. F. Zhang, H. M. Soo, E. P. Greenberg, and L. H. Zhang. 2005. The two-component response regulator PprB modulates quorum-sensing signal production and global gene expression in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **56**:1287–1301.
26. Elborn, J. S., D. J. Shale, and J. R. Britton. 1991. Cystic fibrosis: current survival and population estimates to the year 2000. *Thorax* **46**:881–885.
27. Equi, A., I. M. Balfour-Lynn, A. Bush, and M. Rosenthal. 2002. Long term azithromycin in children with cystic fibrosis: a randomised, placebo-controlled crossover trial. *Lancet* **360**:978–984.
28. Flamm, R. K., M. K. Weaver, C. Thornsberrry, M. E. Jones, J. A. Karlowky, and D. F. Sahm. 2004. Factors associated with relative rates of antibiotic resistance in *Pseudomonas aeruginosa* isolates tested in clinical laboratories in the United States from 1999 to 2002. *Antimicrob. Agents Chemother.* **48**:2431–2436.
29. Fogarty, A., R. Hubbard, and J. Britton. 2000. International comparison of median age at death from cystic fibrosis. *Chest* **117**:1656–1660.
30. Folders, J., J. Algra, M. S. Roelofs, L. C. van Loon, J. Tommassen, and W. Bitter. 2001. Characterization of *Pseudomonas aeruginosa* chitinase, a gradually secreted protein. *J. Bacteriol.* **183**:7044–7052.
31. Gambello, M. J., and B. H. Iglewski. 1991. Cloning and characterization of the *Pseudomonas aeruginosa lasR* gene, a transcriptional activator of elastase expression. *J. Bacteriol.* **173**:3000–3009.
32. Givskov, M., R. de Nys, M. Manefield, L. Gram, R. Maximilien, L. Eberl, S. Molin, P. D. Steinberg, and S. Kjelleberg. 1996. Eukaryotic interference with homoserine lactone-mediated prokaryotic signaling. *J. Bacteriol.* **178**:6618–6622.
33. Givskov, M., and S. Molin. 1993. Secretion of *Serratia liquefaciens* phospholipase from *Escherichia coli*. *Mol. Microbiol.* **8**:229–242.
34. Givskov, M., L. Olsen, and S. Molin. 1988. Cloning and expression in *Escherichia coli* of the gene for extracellular phospholipase A1 from *Serratia liquefaciens*. *J. Bacteriol.* **170**:5855–5862.
35. Goh, E. B., G. Yim, W. Tsui, J. McClure, M. G. Surette, and J. Davies. 2002. Transcriptional modulation of bacterial gene expression by subinhibitory concentrations of antibiotics. *Proc. Natl. Acad. Sci. USA* **99**:17025–17030.
36. Hackman, R. H., and M. Goldberg. 1964. New substrates for use with chitinases. *Anal. Biochem.* **8**:397–401.
37. Hentzer, M., and M. Givskov. 2003. Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections. *J. Clin. Investig.* **112**:1300–1307.
38. Hentzer, M., H. Wu, J. B. Andersen, K. Riedel, T. B. Rasmussen, N. Bagge, N. Kumar, M. A. Schembri, Z. Song, P. Kristoffersen, M. Manefield, J. W. Costerton, S. Molin, L. Eberl, P. Steinberg, S. Kjelleberg, N. Hoiby, and M. Givskov. 2003. Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* **22**:3803.
39. Hoffman, L. R., D. A. D'Argenio, M. J. MacCoss, Z. Zhang, R. A. Jones, and S. I. Miller. 2005. Aminoglycoside antibiotics induce bacterial biofilm formation. *Nature* **436**:1171–1175.
40. Hoffmann, N., B. Lee, M. Hentzer, T. B. Rasmussen, Z. Song, H. K. Johansen, M. Givskov, and N. Hoiby. 2007. Azithromycin blocks quorum sensing and alginate polymer formation and increases the sensitivity to serum and stationary-growth-phase killing of *Pseudomonas aeruginosa* and attenuates chronic *P. aeruginosa* lung infection in Cfr^{-/-} mice. *Antimicrob. Agents Chemother.* **51**:3677–3687.
41. Hoiby, N. 1995. Microbiology of cystic fibrosis, p. 75–98. In M. E. Hodson and D. M. Geddes (ed.), *Cystic fibrosis*. Chapman & Hall Medical, London, United Kingdom.
42. Hooi, D. S., B. W. Bycroft, S. R. Chhabra, P. Williams, and D. I. Pritchard. 2004. Differential immune modulatory activity of *Pseudomonas aeruginosa* quorum-sensing signal molecules. *Infect. Immun.* **72**:6463–6470.
43. Howe, R. A., and R. C. Spencer. 1997. Macrolides for the treatment of *Pseudomonas aeruginosa* infections? *J. Antimicrob. Chemother.* **40**:153–155.
44. Ichimiya, T., K. Takeoka, K. Hiramatsu, K. Hirai, T. Yamasaki, and M. Nasu. 1996. The influence of azithromycin on the biofilm formation of *Pseudomonas aeruginosa* in vitro. *Chemotherapy* **42**:186–191.
45. Jensen, P. O., T. Bjarnsholt, R. Phipps, T. B. Rasmussen, H. Calum, L. Christoffersen, C. Moser, P. Williams, T. Pressler, M. Givskov, and N. Hoiby. 2007. Rapid necrotic killing of polymorphonuclear leukocytes is caused by quorum-sensing-controlled production of rhamnolipid by *Pseudomonas aeruginosa*. *Microbiology* **153**:1329–1338.
46. Johnson, M. K., and D. Boese-Marrazzo. 1980. Production and properties of heat-stable extracellular hemolysin from *Pseudomonas aeruginosa*. *Infect. Immun.* **29**:1028–1033.
47. Karlowky, J. A., M. H. Saunders, G. A. Harding, D. J. Hoban, and G. G. Zhanel. 1996. In vitro characterization of aminoglycoside adaptive resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **40**:1387–1393.
48. Kaufmann, G. F., R. Sartorio, S. H. Lee, C. J. Rogers, M. M. Meijler, J. A. Moss, B. Clapham, A. P. Brogan, T. J. Dickerson, and K. D. Janda. 2005. Revisiting quorum sensing: discovery of additional chemical and biological functions for 3-oxo-*N*-acylhomoserine lactones. *Proc. Natl. Acad. Sci. USA* **102**:309–314.
49. Kobayashi, H. 1995. Biofilm disease: its clinical manifestation and therapeutic possibilities of macrolides. *Am. J. Med.* **99**:26S–30S.
50. Kobayashi, T., K. Tateda, T. Matsumoto, S. Miyazaki, A. Watanabe, T. Nukiwa, and K. Yamaguchi. 2002. Macrolide-treated *Pseudomonas aeruginosa* induces paradoxical host responses in the lungs of mice and a high mortality rate. *J. Antimicrob. Chemother.* **50**:59–66.
51. Latifi, A., M. K. Winson, M. Foglino, B. W. Bycroft, G. S. Stewart, A. Lazdunski, and P. Williams. 1995. Multiple homologues of LuxR and LuxI control expression of virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Mol. Microbiol.* **17**:333–343.
52. Lequette, Y., and E. P. Greenberg. 2005. Timing and localization of rhamnolipid synthesis gene expression in *Pseudomonas aeruginosa* biofilms. *J. Bacteriol.* **187**:37–44.
53. Linares, J. F., I. Gustafsson, F. Baquero, and J. L. Martinez. 2006. Antibiotics as intermicrobial signaling agents instead of weapons. *Proc. Natl. Acad. Sci. USA* **103**:19484–19489.
54. Livermore, D. M. 2002. Multiple mechanisms of antimicrobial resistance in *Pseudomonas aeruginosa*: our worst nightmare? *Clin. Infect. Dis.* **34**:634–640.
55. Lyczak, J. B., C. L. Cannon, and G. B. Pier. 2002. Lung infections associated with cystic fibrosis. *Clin. Microbiol. Rev.* **15**:194–222.
56. Manefield, M., T. B. Rasmussen, M. Hentzer, J. B. Andersen, P. Steinberg, S. Kjelleberg, and M. Givskov. 2002. Halogenated furanones inhibit quorum sensing through accelerated LuxR turnover. *Microbiology* **148**:1119–1127.
57. McClure, C. D., and N. L. Schiller. 1992. Effects of *Pseudomonas aeruginosa* rhamnolipids on human monocyte-derived macrophages. *J. Leukoc. Biol.* **51**:97–102.
58. McClure, C. D., and N. L. Schiller. 1996. Inhibition of macrophage phagocytosis by *Pseudomonas aeruginosa* rhamnolipids in vitro and in vivo. *Curr. Microbiol.* **33**:109–117.
59. Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
60. Nalca, Y., L. Jansch, F. Bredenbruch, R. Geffers, J. Buer, and S. Haussler. 2006. Quorum-sensing antagonistic activities of azithromycin in *Pseudomonas aeruginosa* PAO1: a global approach. *Antimicrob. Agents Chemother.* **50**:1680–1688.
61. Niga, T., H. Ito, Y. Oyamada, J. Yamagishi, M. Kadono, T. Nishino, N. Gotoh, and M. Inoue. 2005. Cooperation between alteration of DNA gyrase genes and over-expression of MexB and MexX confers high-level fluoroquinolone resistance in *Pseudomonas aeruginosa* strains isolated from a patient who received a liver transplant followed by treatment with fluoroquinolones. *Microbiol. Immunol.* **49**:443–446.
62. Ochsner, U. A., and J. Reiser. 1995. Autoinducer-mediated regulation of rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. USA* **92**:6424–6428.
63. Ohman, D. E., S. J. J. Cryz, and B. H. Iglewski. 1980. Isolation and characterization of *Pseudomonas aeruginosa* PAO mutant that produces altered elastase. *J. Bacteriol.* **142**:836–842.
64. Oldak, E., and E. A. Trafny. 2005. Secretion of proteases by *Pseudomonas*

- aeruginosa* biofilms exposed to ciprofloxacin. Antimicrob. Agents Chemother. **49**:3281–3288.
65. Pamp, S. J., and T. Tolker-Nielsen. 2007. Multiple roles of biosurfactants in structural biofilm development by *Pseudomonas aeruginosa*. J. Bacteriol. **189**:2531–2539.
 66. Passador, L., and B. H. Iglewski. 1995. Quorum sensing and virulence gene regulation in *Pseudomonas aeruginosa*, p. 65–78. In J. A. Roth (ed.), Virulence mechanisms of bacterial pathogens, 2nd ed. American Society for Microbiology, Washington, DC.
 67. Pearson, J. P., C. Van Delden, and B. H. Iglewski. 1999. Active efflux and diffusion are involved in transport of *Pseudomonas aeruginosa* cell-to-cell signals. J. Bacteriol. **181**:1203–1210.
 68. Pedersen, S. S., C. Koch, N. Hoiby, and K. Rosendal. 1986. An epidemic spread of multiresistant *Pseudomonas aeruginosa* in a cystic fibrosis centre. J. Antimicrob. Chemother. **17**:505–516.
 69. Permin, H., C. Koch, N. Hoiby, H. O. Christensen, A. F. Moller, and S. Moller. 1983. Ceftazidime treatment of chronic *Pseudomonas aeruginosa* respiratory tract infection in cystic fibrosis. J. Antimicrob. Chemother. **12**(Suppl A):313–323.
 - 69a. Persson, T., T. H. Hansen, T. B. Rasmussen, M. E. Skindersoe, M. Givskov, and J. Nielsen. 2005. Rational design and synthesis of new quorum-sensing inhibitors derived from acylated homoserine lactones and natural products from garlic. Org. Biomol. Chem. **21**:253–262.
 70. Pesci, E. C., J. B. Milbank, J. P. Pearson, S. McKnight, A. S. Kende, E. P. Greenberg, and B. H. Iglewski. 1999. Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. USA **96**:11229–11234.
 71. Pfaffl, M. W. 2001. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res. **29**:e45.
 72. Ramsey, B. W. 1996. Management of pulmonary disease in patients with cystic fibrosis. N. Engl. J. Med. **335**:179–188.
 73. Rasmussen, T. B., T. Bjarnsholt, M. E. Skindersoe, M. Hentzer, P. Kristofersen, M. Kote, J. Nielsen, L. Eberl, and M. Givskov. 2005. Screening for quorum-sensing inhibitors (QSI) by use of a novel genetic system, the QSI selector. J. Bacteriol. **187**:1799–1814.
 74. Rasmussen, T. B., M. E. Skindersoe, T. Bjarnsholt, R. K. Phipps, K. B. Christensen, P. O. Jensen, J. B. Andersen, B. Koch, T. O. Larsen, M. Hentzer, L. Eberl, N. Hoiby, and M. Givskov. 2005. Identity and effects of quorum-sensing inhibitors produced by *Penicillium* species. Microbiology **151**:1325–1340.
 75. Roy-Burman, A., R. H. Savel, S. Racine, B. L. Swanson, N. S. Revadigar, J. Fujimoto, T. Sawa, D. W. Frank, and J. P. Wiener-Kronish. 2001. Type III protein secretion is associated with death in lower respiratory and systemic *Pseudomonas aeruginosa* infections. J. Infect. Dis. **183**:1767–1774.
 76. Saiman, L., B. C. Marshall, N. Mayer-Hamblett, J. L. Burns, A. L. Quittner, D. A. Cibene, S. Coquillette, A. Y. Fieberg, F. J. Accurso, and P. W. Campbell III. 2003. Azithromycin in patients with cystic fibrosis chronically infected with *Pseudomonas aeruginosa*: a randomized controlled trial. JAMA **290**:1749–1756.
 77. Savli, H., A. Karadenizli, F. Kolayli, S. Gundes, U. Ozbek, and H. Vahaboglu. 2003. Expression stability of six housekeeping genes: a proposal for resistance gene quantification studies of *Pseudomonas aeruginosa* by real-time quantitative RT-PCR. J. Med. Microbiol. **52**:403–408.
 78. Sawa, T., M. Ohara, K. Kurahashi, S. S. Twining, D. W. Frank, D. B. Doroques, T. Long, M. A. Gropper, and J. P. Wiener-Kronish. 1998. In vitro cellular toxicity predicts *Pseudomonas aeruginosa* virulence in lung infections. Infect. Immun. **66**:3242–3249.
 79. Schuster, M., A. C. Hawkins, C. S. Harwood, and E. P. Greenberg. 2004. The *Pseudomonas aeruginosa* RpoS regulon and its relationship to quorum sensing. Mol. Microbiol. **51**:973–985.
 80. Schuster, M., C. P. Lostroh, T. Ogi, and E. P. Greenberg. 2003. Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. J. Bacteriol. **185**:2066–2079.
 81. Sharabi, Y., and N. Gilboa-Garber. 1979. Mitogenic stimulation of human lymphocytes by *Pseudomonas aeruginosa* galactosephilic lectins. FEMS Microbiol. Lett. **5**:273–276.
 82. Shryock, T. R., S. A. Silver, M. W. Banschbach, and J. C. Kramer. 1984. Effect of *Pseudomonas aeruginosa* rhamnolipid on human neutrophil migration. Curr. Microbiol. **10**:323–328.
 83. Sierra, G. 1960. Hemolytic effect of a glycolipid produced by *Pseudomonas aeruginosa*. Antonie van Leeuwenhoek **26**:189–192.
 - 83a. Smith, K. M., Y. Bu, and H. Suga. 2003. Library screening for synthetic agonists and antagonists of a *Pseudomonas aeruginosa* autoinducer. Chem. Biol. **10**:563–571.
 84. Tateda, K., R. Comte, J. C. Pechere, T. Kohler, K. Yamaguchi, and C. Van Delden. 2001. Azithromycin inhibits quorum sensing in *Pseudomonas aeruginosa*. Antimicrob. Agents Chemother. **45**:1930–1933.
 85. Telford, G., D. Wheeler, P. Williams, P. T. Tomkins, P. Appleby, H. Sewell, G. S. Stewart, B. W. Bycroft, and D. I. Pritchard. 1998. The *Pseudomonas aeruginosa* quorum-sensing signal molecule *N*-(3-oxododecanoyl)-L-homoserine lactone has immunomodulatory activity. Infect. Immun. **66**:36–42.
 86. Thomsen, R., and M. H. Christensen. 2006. MolDock: a new technique for high-accuracy molecular docking. J. Med. Chem. **49**:3315–3321.
 87. Tramper-Stranders, G. A., T. F. Wolfs, A. Fleer, J. L. Kimpfen, and C. K. van der Ent. 2007. Maintenance azithromycin treatment in pediatric patients with cystic fibrosis: long-term outcomes related to macrolide resistance and pulmonary function. Pediatr. Infect. Dis. J. **268**:8–12.
 88. Tsai, W. C., M. L. Rodriguez, K. S. Young, J. C. Deng, V. J. Thannickal, K. Tateda, M. B. Hershenson, and T. J. Standiford. 2004. Azithromycin blocks neutrophil recruitment in *Pseudomonas* endobronchial infection. Am. J. Respir. Crit. Care Med. **170**:1331–1339.
 89. Wagner, V. E., D. Bushnell, L. Passador, A. I. Brooks, and B. H. Iglewski. 2003. Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. J. Bacteriol. **185**:2080–2095.
 90. Waite, R. D., A. Paccanaro, A. Papakonstantinou, J. M. Hurst, M. Saqi, E. Littler, and M. A. Curtis. 2006. Clustering of *Pseudomonas aeruginosa* transcriptomes from planktonic cultures, developing and mature biofilms reveals distinct expression profiles. BMC Genomics **7**:162.
 91. Wang, Y., U. Ha, L. Zeng, and S. Jin. 2003. Regulation of membrane permeability by a two-component regulatory system in *Pseudomonas aeruginosa*. Antimicrob. Agents Chemother. **47**:95–101.
 92. Wilderman, P. J., A. I. Vasil, Z. Johnson, M. J. Wilson, H. E. Cunliffe, I. L. Lamont, and M. L. Vasil. 2001. Characterization of an endoprotease (PrpL) encoded by a PvdS-regulated gene in *Pseudomonas aeruginosa*. Infect. Immun. **69**:5385–5394.
 93. Xiong, Y. Q., J. Caillon, M. F. Kergueris, H. Drugeon, D. Baron, G. Potel, and A. S. Bayer. 1997. Adaptive resistance of *Pseudomonas aeruginosa* induced by aminoglycosides and killing kinetics in a rabbit endocarditis model. Antimicrob. Agents Chemother. **41**:823–826.
 94. Yim, G., H. H. Wang, and J. Davies. 2006. The truth about antibiotics. Int. J. Med. Microbiol. **296**:163–170.
 95. Zulianello, L., C. Canard, T. Kohler, D. Caille, J. S. Lacroix, and P. Meda. 2006. Rhamnolipids are virulence factors that promote early infiltration of primary human airway epithelia by *Pseudomonas aeruginosa*. Infect. Immun. **74**:3134–3147.