

The ABC-Type Multidrug Resistance Transporter LmrCD Is Responsible for an Extrusion-Based Mechanism of Bile Acid Resistance in *Lactococcus lactis*[∇]

Arsalan Haseeb Zaidi,¹ Patrick J. Bakkes,¹ Jacek Lubelski,² Herfita Agustiandari,¹ Oscar P. Kuipers,² and Arnold J. M. Driessen^{1*}

Department of Molecular Microbiology¹ and Department of Molecular Genetics,² Groningen Biomolecular Sciences and Biotechnology Institute and the Kluyver Centre for Genomics of Industrial Microorganisms, University of Groningen, Kercklaan 30, 9751 NN Haren, The Netherlands

Received 9 April 2008/Accepted 2 September 2008

Upon prolonged exposure to cholate and other toxic compounds, *Lactococcus lactis* develops a multidrug resistance phenotype that has been attributed to an elevated expression of the heterodimeric ABC-type multidrug transporter LmrCD. To investigate the molecular basis of bile acid resistance in *L. lactis* and to evaluate the contribution of efflux-based mechanisms in this process, the drug-sensitive *L. lactis* NZ9000 Δ lmrCD strain was challenged with cholate. A resistant strain was obtained that, compared to the parental strain, showed (i) significantly improved resistance toward several bile acids but not to drugs, (ii) morphological changes, and (iii) an altered susceptibility to antimicrobial peptides. Transcriptome and transport analyses suggest that the acquired resistance is unrelated to elevated transport activity but, instead, results from a multitude of stress responses, changes to the cell envelope, and metabolic changes. In contrast, wild-type cells induce the expression of *lmrCD* upon exposure to cholate, whereupon the cholate is actively extruded from the cells. Together, these data suggest a central role for an efflux-based mechanism in bile acid resistance and implicate LmrCD as the main system responsible in *L. lactis*.

Multidrug transporters are responsible for active efflux of structurally and functionally unrelated drugs and are ubiquitous in nature (21, 32). The majority of the described multidrug transporters in bacteria depend on ion motive forces for extrusion activity, while only few systems use ATP hydrolysis to drive efflux. However, recent reports and functional genomic predictions suggest that the role of these bacterial ABC-type transporters in multidrug resistance (MDR) might be underestimated (32).

Lactococcus lactis is a nonpathogenic lactic acid bacterium that serves as a model organism to study bacterial MDR (31). The genome of *L. lactis* contains 40 putative drug transporter genes of which several have been characterized in detail. These are LmrP (8), a member of the major facilitator superfamily (MFS) transporter, and the ABC-type transporters LmrA (48) and LmrCD (31). *L. lactis* MG1363 readily acquires a stable MDR phenotype when challenged with increasing concentrations of cholate (51) or other drugs (7). An ATP-dependent transporter has been implicated in cholate resistance, but its identity has remained obscure (51). Recent transcriptome analysis revealed that a common response of the selected MDR strains is the elevated expression of *lmrCD* compared to the wild-type strain (31). Furthermore, the deletion of the *lmrCD* genes makes *L. lactis* more susceptible to a wide range of toxic compounds, while the drug resistance phenotype can

be restored by the overexpression of *lmrCD* from a plasmid (33).

Although previous studies implicated LmrCD as a major determinant of drug resistance in *L. lactis*, it has remained unclear whether other putative MDR transporters contribute to resistance in the selected strains as the observed phenotype covers both anionic and cationic drugs such as daunomycin, rhodamine, and ethidium bromide. Moreover, apart from the *lmrCD* upregulation, the cholate-resistant *L. lactis* strain (51) showed a different gene expression profile from the strains selected for resistance against cationic drugs (31). The cholate-resistant strain also showed strong upregulation of the *arc* operon genes involved in the arginine deiminase pathway and of a gene encoding a putative MDR transporter of the MFS, i.e., lmg2513 (formerly named *yxvD*).

Cholate resistance is of particular interest as lactic acid bacteria like *Lactobacillus* are important constituents of the human intestinal microflora and are widely used as probiotics in food supplements (22, 42, 50). Probiotic survival depends on resistance against inhibitory compounds present in the intestine such as bile acids (29). These compounds are synthesized from cholesterol in the liver and have an important role in the intestine, where they are involved in the absorption of dietary fats and lipid soluble vitamins (43). Bile acids are released as conjugates of glycine or taurine in the intestine, where indigenous microbes such as *Bifidobacteria* employ bile acid hydrolases that enzymatically liberate the free bile acids (FBA) from their conjugated forms (45). FBA are toxic, weakly acidic, and hydrophobic molecules that strongly inhibit the growth of various intestinal bacteria (6). Cholate, which is one of the most frequently encountered FBA in the gastrointestinal tract, is produced by the gut microflora from conjugated bile salts, such

* Corresponding author. Mailing address: Department of Molecular Microbiology, University of Groningen, Kercklaan 30, 9751 NN, Haren, The Netherlands. Phone: 31 50 3632164. Fax: 31 50 3632154. E-mail: a.j.m.driessen@rug.nl.

[∇] Published ahead of print on 12 September 2008.

as taurocholate and glycocholate (29). The exact mechanism of inhibition is still not clear but may involve permeabilization of the cytoplasmic membrane (28). The autochthonous microbiota as well as many gram-negative pathogens have evolved to survive the bactericidal effects of FBA (49). The mechanisms underlying bile acid resistance have been attributed to diverse physiological and cellular responses such as changes in cell envelope (11) and increased activity of bile acid modifying enzymes in many gram-positive bacteria (4) while MDR efflux pumps have been implicated in gram-negative bacteria (35). In contrast, little is known about the role of MDR efflux pumps in bile acid extrusion in gram-positive organisms.

Here, we have investigated the adaptive response of *L. lactis* cells that lack the major MDR transporter, LmrCD, when challenged with cholate. The data suggest that in the absence of LmrCD, cholate resistance relies on a multitude of transport-unrelated cellular responses. This investigation also establishes for the first time that LmrCD-mediated extrusion of cholate is the primary mechanism of bile acid resistance in *L. lactis*.

MATERIALS AND METHODS

Bacterial strains, medium, and growth conditions. *L. lactis* NZ9000 is a derivative of the plasmid-free *L. lactis* MG1363 strain containing *pepN:nisRK* (13, 14) and is referred to as the wild-type strain. The derivative *L. lactis* NZ9000 Δ lmrCD strain lacks the MDR transporter LmrCD (31). In all cases, cells were grown at 30°C in M17 medium (Difco) containing 0.5% (wt/vol) glucose (GM17).

Cholate adaptation assay. Cholate-resistant cells were selected by growth of the *L. lactis* NZ9000 Δ lmrCD strain in GM17 medium containing increasing concentrations of cholate. Exponentially growing cells were diluted 1:100 in 5 ml of fresh GM17 containing cholate and grown overnight. This procedure was repeated several times with a concomitant stepwise increase of the cholate concentration until a significant increase in MIC had occurred. The final concentration of cholate used was 4 mM. To obtain single colonies, the adapted culture was spread-plated on GM17 medium with 1.8% (wt/vol) agar containing 3 mM cholate. Thirty colonies were selected and subcultured in fresh GM17 broth without cholate. Cultures were supplemented with glycerol at a final concentration of 10% (vol/vol) and stored at -80°C. Growth analysis of the individual colonies indicated essentially identical 50% inhibitory concentrations (IC₅₀s) and MICs for cholate. One of the obtained cholate-adapted NZ9000 Δ lmrCD colonies was selected and used for further characterization and is referred to as the Δ LmrCD' strain.

Growth studies. Overnight cultures of *L. lactis* wild-type, Δ lmrCD and Δ LmrCD' strains were diluted in fresh GM17 medium and grown to an optical density at 600 nm (OD₆₀₀) of 0.6. In the case of strains bearing a plasmid, the GM17 medium was supplemented with chloramphenicol at a final concentration of 5 µg/ml. Cells were then diluted to an OD₆₀₀ of 0.05, and aliquots of 150 µl were transferred to 96-well microtiter plates that contained 50 µl of GM17 medium containing a range of drugs at various concentrations. Sterile silicon oil (50 µl) was pipetted on top of the samples to prevent evaporation. Growth was monitored at 30°C every 6 min for 8 to 12 h at 660 nm using a multiscan photometer (spectraMax 340; Molecular Devices). The maximum specific growth rate μ was calculated from the exponential growth phase (52) and plotted against the concentration of the different drugs. Concentrations that inhibited growth by 50% (IC₅₀s) and 100% (MIC) were determined. The experiments were carried out in triplicate, and the data shown are averaged to obtain the standard error of the mean.

Cholate transport assay. Exponentially growing cells were harvested at an OD₆₀₀ of ~1, washed once with 50 mM potassium phosphate (KP_i), pH 7.0, containing 1 mM MgSO₄, and resuspended in this buffer to an OD₆₀₀ of ~2.0. The cells were de-energized by incubation with 10 mM 2-deoxyglucose for 30 min at 30°C, washed three times with KP_i buffer, and finally resuspended in this buffer to an OD₆₀₀ of ~8. Aliquots (3 ml) of the cell suspension were dispensed in glass tubes and preincubated for 5 min at 30°C with gentle stirring. Next, 8 µl of 1.82 mM [¹⁴C]cholate (55 mCi/mmol) was mixed with 4 µl of 500 mM nonradioactive cholate in 1 ml of MilliQ water, and 150 µl of the mix was added to the cells (final cholate concentration is ~100 µM), followed by incubation for 14 min after which glucose (230 mM) was added as a source of metabolic energy. At the time

TABLE 1. Oligonucleotide primers used for RT-PCR analysis

Primer name ^a	Primer sequence (5'→3')
<i>lmrC</i> RT-PCR FW.....	GTTGAAGAACGTGGGAATAATTT CTCAGGTGG
<i>lmrC</i> RT-PCR RV.....	CCTCCTGTGCTTTCTGTGTATCGT AGATTTTC
<i>lmrD</i> RT-PCR FW.....	CGTTTCTGATGATGAATCAGTCT TCTCAGTTGG
<i>lmrD</i> RT-PCR RV.....	CAAAACGAATTGATTATGATAA AGTTCAGAG
<i>secY</i> RT-PCR FW.....	TACAACCTGCTCCAGCTACGA
<i>secY</i> RT-PCR RV.....	GTTCTCCAAGAGCGACAAT

^a FW, forward; RV, reverse.

points indicated in the figures, the amount of [¹⁴C]cholate associated with cells was determined by a filtration method. Herein, aliquots of 200 µl were passed over 0.2-µm-pore-size cellulose-acetate filters that were prewetted in 100 mM KP_i, pH 7.0. Retained cells were then washed two times with 2 ml of 100 mM LiCl. Finally, the radioactivity associated with the cells on the filter was measured by liquid scintillation counting. Values were corrected for the background level of radioactivity obtained for control incubations without cells. The amount of accumulated cholate was related to the quantity of cells. Due to morphological differences (see results) the cellular dry weight rather than the OD₆₀₀ was used. A similar experiment was performed using an *L. lactis* NZ9000 Δ lmrCD strain carrying the pLLmrCD plasmid (*lmrCD* expression under the control of native promoter) and control plasmid pIL252 (31).

Scanning electron microscopy. A drop of a bacterial culture was placed on a freshly cleaved mica surface pretreated with a 0.5% poly-L-lysine solution. Next, cells were washed with 0.1 M cacodylate buffer, fixed with 2% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.4) for 30 min, and washed with the cacodylate buffer. The specimen was dehydrated with ethanol in a sequence of 30, 50, 70, and 100% (10 min each), followed by two times in 100% for 30 min. Finally, the specimen was dried in a Bal-Tec Critical Point Dryer with CO₂ and sputter-coated with 2 to 3 nm of Au/Pd (Bal-Tec sputter coater). Observation was done with a Jeol FE-SEM 6301F (cold-field emission scanning electron microscope).

Expression analysis by RT-PCR. Overnight cultures of *L. lactis* NZ9000 were diluted in GM17 medium to an OD₆₀₀ of 0.05 and grown at 30°C until they reached an OD₆₀₀ of ~0.6. Cholate (1 mM) was added to the cultures, and at various time points, aliquots were removed, and cells were harvested by centrifugation (3,500 rpm at 4°C for 5 min). Total RNA was isolated using TRIzol reagent (Invitrogen). The RNA concentration was determined, and equal amounts of RNA were transferred into Illustra Ready To-Go reverse transcription-PCR (RT-PCR) tubes to generate cDNA from the RNA templates. The first synthesis of a cDNA strand was performed at 42°C for 40 min, followed by standard PCR conditions. RT-PCR products of the *lmrC*, *lmrD*, and *secY* genes were obtained with gene-specific primer pairs (Table 1), and samples were analyzed on a 2% (wt/vol) agarose gel.

DNA microarray analysis. DNA microarray experiments were essentially performed as described previously (14, 47). In brief, RNA was isolated from two replicate cultures of both *L. lactis* Δ lmrCD and cholate-adapted *L. lactis* Δ LmrCD'. Cultures were grown at 30°C in GM17 medium in the absence of cholate, and cells were harvested at an OD₆₀₀ of ~1. Next, single-strand RT (amplification) and indirect labeling of 20 µg of total RNA, with either Cy3 or Cy5 dye, were performed (including samples in which the dyes were swapped to correct for dye-specific effects). Labeled cDNA samples were hybridized to microarray slides containing probes representing 2,496 open reading frames (ORFs) of *L. lactis* MG1363 spotted in duplicate. After overnight hybridization, slides were washed for 5 min at 37°C in 2× SSC (1× SSC is 0.15 M NaCl plus 0.015 M sodium citrate) containing 0.5% sodium dodecyl sulfate followed by 5 min in 1× SSC containing 0.25% sodium dodecyl sulfate to remove nonspecifically hybridized cDNAs. Slides were scanned using a GenePix 4200AL instrument (Westburg). Subsequently, individual spot intensities were determined using ArrayPro, version 4.5 (Media Cybernetics Inc., Silver Spring, MD). Slide data were processed and normalized using MicroPrep, which yielded average gene expression ratios of the mutant to the control strain. Expression of a gene was considered to be significantly altered at a Cyber-T Bayesian *P* value of ≤0.001. From the set of genes that exhibited a significant change in expression, only genes that exhibited a strong, i.e., ≥1.8-fold, change in expression are discussed.




FIG. 1. Bile acid resistance of *L. lactis* cells harboring or lacking the *lmrCD* genes. (A) Cholate resistance of *L. lactis* wild-type (\square), Δ *lmrCD* (\triangle), and Δ *LmrCD*^r (\blacktriangle) strains. (B) Cholate resistance of *L. lactis* Δ *lmrCD* (\circ) and Δ *lmrCD* containing plasmid pIL*lmrCD* (\bullet) or control plasmid pIL252 (\triangle). (C) Cholate resistance of cholate-adapted Δ *LmrCD*^r cells containing pIL*lmrCD* (\triangle) or pIL252 (\blacktriangle). (D) Glycodeoxycholate resistance of Δ *LmrCD*^r cells containing pIL*lmrCD* (\blacksquare) or pIL252 (\square). Note that the plasmid-encoded *lmrCD* genes are under the control of their native promoter. Cells were grown for 8 h in GM17 medium in the absence or presence of various concentrations of bile acid, and the maximum specific growth rate, μ , was determined.

Microarray data accession number. The array data reported in this publication have been deposited in the NCBI Gene Expression Omnibus (GEO; <http://www.ncbi.nlm.nih.gov/geo/>) and are accessible through GEO Series accession number GSE10203.

RESULTS

***L. lactis* can acquire resistance to cholate in the absence of LmrCD.** Previously, we have demonstrated that *L. lactis* cells lacking the genes encoding the heterodimeric ABC-type MDR transporter LmrCD become about twofold more susceptible to cholate than wild-type cells (31). Transcriptome analysis of the MDR strain of *L. lactis* selected for cholate resistance showed, among other things, a marked induction of *lmrCD* expression (31) that could be related to inactivation of the transcriptional regulator LmrR (1). These data suggest an important role of LmrCD in resistance to cholate but do not exclude an involvement of other transporters or non-efflux-based mechanisms in cholate resistance. Therefore, the cholate-sensitive *L. lactis* Δ *lmrCD* strain was repeatedly exposed to stepwise increasing

sublethal concentrations of cholate to induce resistance. In this manner a strain with significantly improved resistance to cholate was obtained (Δ *LmrCD*^r) (Fig. 1A and Table 2). The observed phenotype was found to be stable as the Δ *LmrCD*^r strain remained resistant after growth for 7 days in the absence of cholate (data not shown). Cholate-adapted Δ *LmrCD*^r cells were found to be 1.7-fold more resistant to cholate than the parental Δ *lmrCD* strain, as judged from the MICs (1.7 and 1 mM, respectively). However, in the absence of cholate, the Δ *LmrCD*^r cells showed a lower growth rate than parental and wild-type *L. lactis* cells. For reasons that remain unclear, the IC₅₀ of cholate obtained for the Δ *lmrCD* strain was 2.5-fold lower than that reported previously (1 and 2.5 mM, respectively) (25). However, these IC₅₀s were found consistently in at least six independent experiments, while the relative resistance levels between the wild-type and Δ *lmrCD* strains obtained in both studies were very similar. Plasmid-based overexpression of *lmrCD* in the Δ *lmrCD* strain increased cholate resistance up to approximately fourfold (Fig. 1B), which greatly exceeded

TABLE 2. Susceptibility of *L. lactis* wild-type, $\Delta lmrCD$, and $\Delta LmrCD^f$ strains to various bile acids and drugs

Drug	IC ₅₀ (μM) of the indicated strain ^a			Relative resistance ^f
	Wild type	$\Delta lmrCD$	$\Delta LmrCD^f$	
Lithocholate ^c	85	85	80	0.9
Deoxycholate ^{b,d}	125 ± 15	120 ± 18	190 ± 18	1.6
Chenodeoxycholate ^{b,d}	115	105	160	1.5
Cholate ^{b,e}	1650 ± 273	1000 ± 146	1650 ± 261	1.7
Glycodeoxycholate ^b	6800 ± 566	1063 ± 53	1625 ± 35	1.5
Taurocholate ^b	NS	NS	NS	
Glycocholate ^b	NS	NS	NS	
Hoechst 33342	1.55 ± 0.13	0.3 ± 0.21	0.35 ± 0.17	1.2
Daunomycin	25.5	2.8	2.8	1.0
Rhodamine 6G	5.7	4.9	4.5	0.9
Ethidium bromide	4.8	3.7	3.4	0.9
Quinine	850	775	350	0.5
Tetracycline	0.26	0.2	0.18	0.9
Erythromycin	0.065	0.067	0.061	0.9
Kanamycin	35 ± 9	33 ± 9.5	41 ± 6.8	1.2
Chloramphenicol	4.0 ± 0.5	4.2 ± 0.9	4.5 ± 0.6	1.1
Puromycin	27.5	26.5	27	1.0

^a Values are means and standard errors of IC₅₀s obtained from at least three independent experiments. NS, not sensitive (no appreciable growth inhibition at 6 mM).

^b Sodium salts (anionic detergent).

^c One hydroxyl group.

^d Two hydroxyl groups.

^e Three hydroxyl groups.

^f Calculated as (IC₅₀ of $\Delta LmrCD^f$ cells)/(IC₅₀ of $\Delta lmrCD$ cells).

the cholate resistance level of both the $\Delta LmrCD^f$ and the wild-type strain (Fig. 1A and B). Overall, these data demonstrate that LmrCD provides resistance to cholate.

Exposure to one specific drug may evoke cross-resistance to other toxic compounds in bacteria (18, 41). Therefore, the resistance to structurally and functionally diverse toxic compounds was determined. To this end, $\Delta LmrCD^f$ cells were grown in the presence of a variety of drugs, among which a range of bile acids, several commonly used antibiotics, and structurally unrelated fluorescent dyes like rhodamine 6G, ethidium bromide, and Hoechst 33342 (Table 2). Compared to the wild-type cells, *L. lactis* $\Delta lmrCD$ cells proved to be susceptible to glycodeoxycholate in addition to cholate, Hoechst 33342, daunomycin, rhodamine 6G, and ethidium bromide. Compared to the $\Delta lmrCD$ strain, the cholate-adapted $\Delta LmrCD^f$ cells showed a significant increase in resistance to the unconjugated bile acids deoxycholate (1.6-fold) and chenodeoxycholate (1.5-fold) and the glycoconjugate of deoxycholate (1.5-fold). Interestingly, the acquired resistance to several of these cholate derivatives even exceeded that of the wild-type cells. However, this was certainly not the case for glycodeoxycholate. Compared to the wild-type strain, $\Delta LmrCD^f$ cells were 4.2-fold more susceptible to glycodeoxycholate (Table 2). Note that the deletion of the *lmrCD* genes from the wild-type strain results in a more than sixfold increase in sensitivity to glycodeoxycholate (Table 2.). These data suggest that glycodeoxycholate is an important substrate for LmrCD.

In contrast to the observed enhanced resistance to bile acids, $\Delta LmrCD^f$ cells did not show a significant improvement in resistance to any of the antibiotics and fluorescent dyes tested. On the other hand, the $\Delta LmrCD^f$ strain was twofold more susceptible to quinine than the $\Delta lmrCD$ and wild-type strains. Taken together, the results indicate that the acquired resistance of the $\Delta LmrCD^f$ strain does not arise from MDR but is specific for cholate and related bile acids.

Next, we tested whether the resistance to bile acids of the $\Delta LmrCD^f$ strain could be enhanced by reintroducing the *lmrCD* genes. The $\Delta LmrCD^f$ strain carrying either the control plasmid or the plasmid harboring the *lmrCD* genes (under the control of their own promoter) was grown in the presence of cholate (Fig. 1C) or glycodeoxycholate (Fig. 1D). Expression of *lmrCD* in $\Delta LmrCD^f$ cells resulted in a relatively small increase in cholate resistance, i.e., the IC₅₀ increased from 2.4 to 3.2 mM (Fig. 1C). For glycodeoxycholate (Fig. 1D), the IC₅₀ essentially did not change upon expression of *lmrCD* (i.e., it remained at ~1.5 mM). In contrast, the MICs, differed greatly; $\Delta LmrCD^f$ cells readily die in the presence of 2.4 mM glycodeoxycholate, but when expressing LmrCD they can withstand concentrations up to 8 mM (Fig. 1D), which is consistent with the notion that glycodeoxycholate is an excellent substrate for LmrCD. The $\Delta LmrCD^f$ strain carrying the *lmrCD* plasmid was also grown in the presence of ethidium bromide, rhodamine 6G, or Hoechst 33342. As expected, the $\Delta LmrCD^f$ strain gained the MDR phenotype associated with LmrCD expression (data not shown).

During growth *L. lactis* produces lactic acid, which results in acidification of the growth medium (40). Since the pH may affect the solubility (and, thus, toxicity) of the bile acids tested, the acidification of the growth medium caused by the wild-type, $\Delta lmrCD$, and $\Delta LmrCD^f$ strains, either in the absence or presence of cholate (0.7 mM and 2.2 mM), was monitored in time (data not shown). All strains showed similar degrees of acidification with a drop of the pH from pH 7.3 at the early stages of growth to pH 5.4 in the stationary growth phase. This shows that the bile acid resistance of the $\Delta LmrCD^f$ strain is not due to alteration in primary metabolism.

Cholate-adapted *L. lactis* $\Delta lmrCD$ cells exhibit an altered cell morphology. In the absence of drugs, $\Delta LmrCD^f$ cells had a slower growth rate than the parental $\Delta lmrCD$ and wild-type strains (Fig. 1A and 3). Interestingly, $\Delta LmrCD^f$ cells exhibited an unusual flaky morphology when cultivated in GM17 medium, and they sedimented more readily than $\Delta lmrCD$ and wild-type cells. Scanning electron microscopy revealed that $\Delta LmrCD^f$ cells were similar in size to the $\Delta lmrCD$ and wild-type cells but appeared to grow in unusually long strings of cocci clumped together as large aggregates (Fig. 2). Although multiplication of the $\Delta LmrCD^f$ cells seemed unaffected, the final stage of cell division, i.e., cell separation, was clearly impaired. Such an altered overall cellular morphology likely results from changes in the cell envelope.

To investigate the latter possibility, the sensitivity of the cells to various bacteriocins was tested. This included peptides that specifically act on the cell membrane either by a combination of pore formation and inhibition of peptidoglycan synthesis (nisin) or by exclusively inhibiting peptidoglycan synthesis (bacitracin and vancomycin) (Fig. 3). Wild-type and $\Delta lmrCD$ cells displayed very similar sensitivities to the tested bacteriocins. In contrast, $\Delta LmrCD^f$ cells were found to be highly susceptible to the bacteriocin nisin (fourfold lower IC₅₀), while improved resistance to bacitracin (twofold higher IC₅₀) was observed. The sensitivity to vancomycin was similar to that of the control strains. The altered sensitivity of $\Delta LmrCD^f$ cells to membrane- and peptidoglycan-acting compounds likely arises from an altered cell envelope. Such altered properties of the




FIG. 2. Morphological changes in *L. lactis* Δ lmrCD cells following cholate adaptation. Scanning electron micrographs of Δ LmrCD^r cells at magnifications of $\times 6,000$ (A) or $\times 45,000$ (B) and of the parental Δ lmrCD strain at a magnification of $\times 45,000$ (C). Bars, 1 μ m (A) and 100 nm (B and C).




FIG. 3. Sensitivity of *L. lactis* wild-type (\square), Δ lmrCD (\triangle), and the cholate-adapted Δ LmrCD^r (\blacktriangle) strain to bacteriocins. Cells were grown for 8 h in GM17 medium containing the bacteriocins bacitracin (A), vancomycin (B), or nisin (C) at various concentrations. The maximum specific growth rate, μ , is plotted against the bacteriocin concentration.

cell surface may also account for the increased susceptibility to quinine (Table 2).

Transcriptome analysis of cholate-adapted *L. lactis* Δ lmrCD cells. To investigate the underlying adaptive mechanism(s) of the cholate-adapted strain, DNA microarray analysis was performed on the global gene expression profiles of exponentially growing Δ lmrCD and Δ LmrCD^r cells in the absence of cholate. Compared to the parental Δ lmrCD strain, 124 genes of the Δ LmrCD^r strain showed ≥ 1.8 -fold change in expression with a Bayesian *P* value of ≤ 0.001 (the full array data set is available under accession number GSE10203 at <http://www.ncbi.nlm.nih.gov/geo/accessible>). Of these genes, 87 and 37 showed an increased and decreased expression, respectively. The maximum level of gene induction and repression was 5-fold (*matR*) and 11.6-fold (*llmg1356*), respectively. The differentially expressed genes were grouped into functional classes (Table 3), taking into account the categories defined earlier for the related *L. lactis* strain IL1403 (9). Responsive genes were related to cell envelope biogenesis, stress response and chaperones, general metabolism, housekeeping functions, and sex factor. Remarkably, the expression of none of the 40 genes encoding putative MDR transporters, among which are the well-characterized LmrA and LmrP, was significantly altered in the Δ LmrCD^r strain. This also includes the *llmg2513* (*yxbD*) gene that was upregulated in the MDR strain of *L. lactis* selected by cholate (31). Expression of a large number of hypothetical ORFs was also significantly changed including *llmg1960*, a putative di- and tricarboxylate transporter. However, none of the remaining upregulated hypothetical ORFs could be related to a (drug) transport function (see full array set).

Genes related to cell envelope. Various genes associated with cell envelope biosynthesis or morphology, namely, *cdsA*, *murC*, *rgpE*, *llmg0215*, *llmg0538*, and *llmg1148*, showed increased expression levels in the Δ LmrCD^r strain. Decreased expression levels were observed for *cfa*, which is involved in membrane lipid biosynthesis, and *mvk*, a key gene of the mevalonate biosynthetic pathway needed for isoprenoid synthesis.

Genes related to stress response and chaperones. A distinct level of overexpression of the molecular chaperone genes *groES*, *groEL*, and *hslO* was observed in the Δ LmrCD^r strain, as well the gene encoding a serine protease, *htrA*, implicated in protein folding stress. Also *nah*, encoding an Na⁺/H⁺ antiporter involved in sodium toxicity and intracellular pH regulation, was expressed at a higher level. In contrast, oxidative stress response genes *trxA*, *sodA*, and *gshR* and the general

TABLE 3. Differential expression of genes in cholate-adapted *L. lactis* Δ LmrCD^f cells versus Δ lmrCD cells

Functional class and gene name	Bayesian <i>P</i> value	Fold change	Proposed name and/or description
Sex factor/conjugation			
<i>matR</i>	1.38E-10	5.0	Retron-type reverse transcriptase/LtrA
<i>cluA</i>	2.73E-09	3.2	Cell surface antigen I/II precursor
<i>llmg1399</i>	8.14E-09	3.2	Putative cell surface antigen
<i>ltrC</i>	7.05E-08	2.7	Relaxosome formation
<i>ltrB</i>	1.51E-06	2.5	Group II intron-interrupted relaxase LtrB (<i>mobA</i>)
<i>ltrD</i>	5.47E-07	2.4	Relaxosome formation
<i>ltrE</i>	1.46E-06	2.0	Relaxosome formation
<i>traD</i>	5.81E-06	1.8	Conjugal transfer protein TraD
<i>llmg1353</i>	6.39E-11	-11.0	Putative tellurite resistance protein
<i>telB</i>	1.73E-13	-10.8	Putative tellurite resistance protein
<i>telC</i>	5.43E-11	-7.1	Putative tellurite resistance protein
<i>telA</i>	1.32E-10	-6.6	Putative tellurite resistance protein
Stress and chaperones			
<i>groES</i>	7.77E-09	2.9	GroES/Hsp10 chaperone
<i>nah</i>	3.96E-09	2.8	Na ⁺ /H ⁺ antiporter
<i>htrA</i>	4.91E-08	2.7	Housekeeping protease
<i>groEL</i>	2.83E-08	2.5	GroEL/Hsp60 chaperone
<i>hslO</i>	5.55E-08	2.4	Heat shock protein; 33-kDa chaperone
<i>llmg2047</i>	1.51E-08	-2.8	Universal stress protein E
<i>uspA2</i>	2.67E-06	-2.1	Universal stress protein A2
<i>uspA</i>	2.02E-07	-2.1	Universal stress protein A
<i>sodA</i>	1.32E-07	-2.1	Superoxide dismutase
<i>clpE</i>	6.57E-07	-2.0	ATP-dependent Clp protease ATP-binding subunit E
<i>gshR</i>	5.95E-07	-2.0	Glutathione reductase
<i>trxA</i>	3.66E-07	-1.9	Thioredoxin
Cell envelope			
<i>chiC</i>	8.22E-09	2.8	Acidic endochitinase precursor
<i>llmg1148</i>	1.65E-07	2.5	Putative cell surface antigen
<i>llmg2420</i>	1.24E-07	2.3	Putative glycosyltransferase
<i>chb</i>	2.43E-07	2.2	Chitin binding protein, putative
<i>cdsA</i>	7.95E-07	2.1	Phosphatidate cytidyltransferase
<i>llmg0538</i>	4.95E-07	2.0	(3 <i>R</i>)-Hydroxymyristoyl-(acyl-carrier-protein) dehydratase
<i>lplL</i>	1.29E-05	1.9	Lipoate-protein ligase A
<i>murC</i>	1.93E-04	1.9	UDP- <i>N</i> -acetyl muramyl-alanine ligase
<i>llmg2421</i>	1.67E-05	1.9	Putative glycosyltransferase
<i>llmg0215</i>	6.65E-04	1.8	Predicted membrane protein
<i>rgpE</i>	2.80E-04	1.8	Glycosyltransferase RgpE
<i>cfa</i>	1.82E-06	-2.0	Cyclopropane-fatty-acyl-phospholipid synthase
<i>mvk</i>	7.80E-07	-2.0	Mevalonate kinase
<i>apbE</i>	1.82E-06	-1.9	Thiamine biosynthesis lipoprotein ApbE precursor
General metabolism			
<i>rma</i>	5.78E-09	3.1	Putative rRNA (guanine-N1-)-methyltransferase
<i>glx</i>	3.69E-06	2.6	Glutamyl-tRNA synthetase
<i>llmg1361</i>	8.96E-06	2.6	Putative tyrosine recombinase
<i>polC</i>	4.66E-05	2.2	DNA polymerase III alpha subunit
<i>ackA1</i>	5.52E-07	2.1	AckA1 protein (Acetate kinase)
<i>lacX</i>	5.87E-07	2.1	Galactose mutarotase related enzyme
<i>purA</i>	8.09E-07	2.1	Putative adenylosuccinate synthetase
<i>hisS</i>	4.30E-08	2.0	Histidyl-tRNA synthetase, class IIa
<i>llmg1089</i>	1.49E-06	2.0	Carbamoyl-phosphate synthase, large subunit
<i>ihvB</i>	6.31E-05	2.0	Acetolactate synthase large subunit
<i>butA</i>	6.56E-08	2.0	Acetoin reductase
<i>llmg2209</i>	4.46E-07	1.9	tRNA-dihydrouridine synthase B
<i>pheS</i>	2.84E-06	1.9	Phenylalanyl-tRNA synthetase alpha chain
<i>pfl</i>	9.71E-06	1.9	Pyruvate formate lyase
<i>recF</i>	3.12E-06	1.9	DNA replication and repair protein RecF
<i>proS</i>	2.16E-06	1.8	Prolyl-tRNA synthetase
<i>llmg2205</i>	5.25E-06	1.8	Conserved hypothetical protein
<i>polA</i>	2.84E-06	1.8	DNA polymerase I
<i>metA</i>	1.32E-06	1.8	Homoserine <i>O</i> -succinyltransferase
<i>glmS</i>	7.67E-09	-2.7	Glucosamine-fructose-6-phosphate aminotransferase
<i>add</i>	2.59E-06	-2.6	Adenosine deaminase
<i>ihvE</i>	5.20E-04	-2.2	Branched-chain amino acid aminotransferase
<i>rtrG</i>	8.25E-07	-2.2	Transcriptional regulator, LysR family
<i>pepC</i>	4.72E-06	-2.1	Aminopeptidase C
<i>serB</i>	2.74E-06	-2.0	Phosphoserine phosphatase
<i>llmg1086</i>	1.91E-06	-1.9	MgtA-like cation transporting ATPase
<i>nadE</i>	5.04E-06	-1.9	NAD ⁺ synthase
<i>cysD</i>	1.70E-06	-1.9	<i>O</i> -acetylhomoserine sulfhydrylase
<i>fbaA</i>	8.76E-07	-1.8	Fructose-bisphosphate aldolase
<i>dtpT</i>	2.09E-06	-1.8	Di-/tripeptide transporter

^a Statistically significant changes in expression are given as the ratio of the cholate-adapted Δ LmrCD^f strain versus the control Δ lmrCD strain (fold change). For clarity, most hypothetical ORFs (56) were omitted.

stress response genes *uspA* and *uspA2* were distinctly repressed.

Genes related to general metabolism and housekeeping functions. A variety of genes related to various biosynthetic pathways and DNA replication and repair are differentially expressed. Increases in expression of the genes *rma*, *polC*, *polA*, and *recF* involved in DNA replication occurred in the Δ LmrCD^r strain. Likewise, an increase in expression of genes involved in amino acid biosynthesis (i.e., *ilvB*, *metA*, *proS*, *llmg1089*, *hisS*, *gltX*, and *pheS*) and sugar metabolism (i.e., *lacX*, *ackA1*, *pfl*, and *butA*) was observed, whereas several other genes were repressed, such as *serB* and *cysD* involved in glycine and serine biosynthesis, respectively, as well as genes involved in nucleotide biosynthesis. Of the extensive proteolytic system of *L. lactis*, only *pepC*, an amino acid peptidase, and *dtpT*, a di- and tripeptide transporter, exhibited significant repression.

Sex factor. The sex factor is a chromosomally located ~55-kb element (llmg1411-llmg1348) involved in conjugation. Forty-one out of 57 sex factor genes were significantly differentially expressed. One of the most prominently upregulated genes is *chuA*. This gene confers a cell aggregation phenotype in *L. lactis* (34), which would explain the clumping phenotype observed. A similar role for the product of llmg1399 can be envisaged as it lies in close proximity to *chuA* and shares high homology with it. In addition, *traD*, involved in conjugal transfer, was distinctly overexpressed. A group of 10 genes located at the distal end of the element is significantly repressed. Among these, *telA*, *telB*, and *telC* have been implicated in tellurite resistance. In line with this observation, the Δ LmrCD^r cells were found to be significantly more susceptible to tellurite than cells of the Δ lmrCD and wild-type strains, with MICs of 1.5, 2.5, and 2.5 mM, respectively (data not shown).

LmrCD is responsible for cholate extrusion. The transcriptome analysis of the cholate-adapted Δ LmrCD^r cells showed no evidence of an altered expression of any of the putative MDR transporters. To further analyze the mechanism of cholate resistance, the ability of cholate to enter cells was examined by means of a [¹⁴C]cholate transport assay. Wild-type, Δ lmrCD, and Δ LmrCD^r cells showed a similar level of [¹⁴C]cholate association when tested under energy-deprived conditions (Fig. 4A). Upon the addition of glucose, Δ lmrCD cells accumulated cholate in a time-dependent fashion, while wild-type cells showed a strong extrusion of cholate under identical conditions. For energized Δ LmrCD^r cells, initially a minor efflux is observed, followed by an extended cholate accumulation phase similar to that of the Δ lmrCD strain. The Δ LmrCD^r cells accumulated cholate to a lesser extent (approximately twofold) than the parental Δ lmrCD strain. The cholate extrusion activity of *L. lactis* Δ lmrCD cells could be restored by the reintroduction of the *lmrCD* genes on a plasmid under the control of their native promoter (Fig. 4B). It should be noted that since cholate is a weak acid, the observed energy-dependent uptake of cholate by the Δ lmrCD and Δ LmrCD^r cells (Fig. 4A) likely results from passive permeation and a Δ pH-dependent partitioning of the cholate across the membrane. Taken together, these data unequivocally show that in *L. lactis*, LmrCD efficiently extrudes cholate from the cell in an energy-dependent fashion. Furthermore, the apparent lack of major extrusion activity in the Δ LmrCD^r cells suggests that LmrCD is




FIG. 4. Accumulation of cholate by *L. lactis* cells harboring or lacking *lmrCD* genes. (A and B) De-energized cells were preloaded with [¹⁴C]cholate, and after 14 min cells were energized with glucose at a final concentration of 230 mM. The arrow indicates when glucose was added. (A) Cholate accumulation in the wild-type (□), Δ lmrCD (△), and Δ LmrCD^r (▲) strains. (B) Cholate accumulation in the Δ lmrCD strain containing plasmid pILlmrCD (■) or pIL252 (△).

the major contributor in cholate extrusion and lends further support for the notion that the cholate resistance of the Δ LmrCD^r strain is unrelated to a transport phenomenon.

The expression of the *lmrCD* genes in *L. lactis* is controlled by the transcriptional regulator LmrR (1). Previous studies have demonstrated that binding of drugs like Hoechst 33342 and daunomycin to LmrR relieves the repression of the *lmrCD* genes, thus leading to the manifestation of the MDR phenotype. Since growth and transport studies suggest that bile acids are natural substrates of the LmrCD transporter, the ability of cholate to induce the expression of the *lmrCD* genes was investigated. RT-PCR-based detection of mRNA revealed a transient increase of both the *lmrC* and *lmrD* transcripts upon a challenge of the cells with cholate (Fig. 5). On the other hand, the mRNA levels of the constitutively expressed *secY* gene remained unaltered. This indicates that cholate is an inducer of *lmrCD* expression.

DISCUSSION

The most widely used probiotic bacteria are *Lactobacillus* and *Bifidobacteria*, which are natural inhabitants of the human gastrointestinal tract. Lactococci are not considered part of the gastrointestinal tract, and only a few studies on their probiotic activity are available. However, lactococci do exhibit probiotic properties, like the ability to (i) survive in the gut, (ii) lower the cholesterol level, and (iii) modulate the immune response of the host (26, 46). Lactococci also tolerate gut secretions such as




FIG. 5. Cell-based RT-PCR analysis of *lmrCD* expression in *L. lactis* wild-type cells following cholate induction. PCR products were generated through the use of gene-specific primers for *lmrCD*. Amplified products were separated on 2.0% agarose gels and were identified by ethidium bromide staining. For each time sample, an RT-PCR with primers specific for the *secY* gene was run as a control.

bile acids (24, 25). The purpose of this study was to examine the role of the ABC-type MDR transporter LmrCD in bile acid resistance. In gram-negative bacteria, drug extrusion-based mechanisms across the outer membrane are major determinants of bile acid resistance. This resistance involves MDR efflux pumps that belong to resistance nodulation division and MFS (30, 38), both of which utilize the proton motive force as an energy source. In gram-positive bacteria such as *L. lactis*, the exact mechanisms of cholate resistance are less clear, but the involvement of efflux-based mechanisms based on secondary transport have been suggested (20).

Here, we demonstrate that in *L. lactis*, LmrCD is responsible for an extrusion-based mechanism of resistance against cholate and also provides resistance against the physiologically relevant conjugate glycodeoxycholate. Cells lacking the *lmrCD* genes are sensitive to these compounds. However, when challenged with increasing concentrations of cholate, these cells regain resistance. Remarkably, this resistance is no longer based on an extrusion mechanism but relates to secondary responses such as cell envelope changes, stress responses, and alterations in metabolism. On the other hand, in wild-type cells the short-term response to a challenge with cholate is the upregulation of the *lmrCD* genes, resulting in an increased extrusion of cholate. This upregulation involves the transcriptional repressor LmrR that likely interacts directly with cholate, whereupon repression of *lmrCD* is relieved (1). To our knowledge, this report shows for the first time a central role of an ABC-type transporter in bile acid resistance in prokaryotes. In this respect, ABC-type transporters are generally involved in bile extrusion and transport within the liver (15).

To investigate possible alternative mechanisms of cholate resistance, the cholate-adapted strain, Δ LmrCD^r, and parental strain were subjected to transcriptome profiling. About 100 genes involved in cellular metabolism and morphology were found to be differentially expressed in the transcriptome of the cholate-adapted strain. This is in sharp contrast to what was observed in cholate-adapted wild-type *L. lactis* cells, which showed a prominent defect in the *lmrR* gene that encodes a transcriptional repressor of *lmrCD* expression, which resulted in the constitutive expression of the *lmrCD* genes (1, 32). Interestingly, neither the well-characterized transporters LmrA and LmrP nor any of the other remaining putative MDR transporters were found to be upregulated in the cholate-

adapted Δ LmrCD^r strain, lending further support to the notion that bile acid extrusion is a key activity of LmrCD.

One of the most strongly induced genes in Δ LmrCD^r cells is *nah*, which encodes the Na⁺/H⁺ antiporter that has been implicated in intracellular pH homeostasis and Na⁺ toxicity. Due to passive permeation of cholic acid, high concentrations of cholate might interfere with intracellular pH regulation, which may explain this cellular response. Alternatively, upregulation of *nah* is a means to counteract Na⁺ toxicity as cholate is added as a sodium salt. One of the other remarkable responses is the upregulation of the genes associated with the chromosomally embedded sex factor. This 55-kb region comprises a unique mobile genetic element in *L. lactis* that can be excised in a closed circular form and is readily lost from cells (17). The genes contained in the proximal region (*ltrA-matR*, *ltrB*, *ltrC*, *ltrD*, *ltrE*, and *traD*) show strongly elevated expression. *ltrA-matR* codes for a protein with reverse transcriptase, endonuclease, and RNA maturase activity (37) that facilitates retro-homing of *ltrB* into intronless alleles. The *ltrC*, *ltrD*, and *ltrE* gene products are involved in relaxosome formation during conjugation (12) while *traD* encodes a coupling protein that links to the DNA transfer intermediate and perhaps leads the DNA through the mating channel (19). It has been shown for *L. lactis* MG1363 that the sex factor element can mobilize chromosomal genes (16). Possibly during cholate stress, genetic traits that confer resistance to cholate may be transferred to the recipient cells. Another sex factor gene, *cluA*, associated with cell aggregation phenotype also increased in expression and is involved in the cell-to-cell contact necessary for conjugal transfer (44). CluA is a 136-kDa surface-bound protein covalently linked to the cell wall peptidoglycan. This protein is not only responsible for a constitutive aggregation phenotype in *L. lactis* MG1363 but also linked to high-frequency conjugation and transfer of the sex factor (34, 42). The upregulation of the sex factor genes might thus be responsible for a major morphological change such as the clumping of the cells that may provide a certain level of protection to the inner cells in the aggregate to cholate. Interestingly, *Lactobacillus plantarum* also showed morphological changes in response to bile acid stress (10). Challenged cells clumped together (but did not form long strings) and showed elevated expression levels of several genes involved in membrane- and cell wall-associated functions. Thus, altering the properties of the bacterial cell surface may be a common response to bile acid stress.

In the Δ LmrCD^r strain several upregulated genes appear to be associated with cell envelope biogenesis such as *murC*, which is involved in the biosynthesis of the peptidoglycan murein, which catalyzes the addition of L-alanine to the nucleotide precursor UDP-N-acetylmuramoyl, and genes such as *rgpE*, *llmg2420*, and *llmg2421*, encoding putative glycosyltransferases that catalyze the formation of linear glycan chains. The upregulation of these genes may result in an alteration of the cell envelope composition and thus indirectly affect cholate permeation and susceptibility. This is further supported by the altered responses of the cholate-adapted cells to the activity of three peptide antibiotics, i.e., nisin, bacitracin, and vancomycin. These antimicrobials affect the cell envelope by different mechanisms of action (2, 3, 23). The increased nisin sensitivity in the cholate-adapted strain suggests that either the levels of lipid II, the binding site for nisin, have increased (2, 36) or that

lipid II is more accessible (27, 36). Nisin sensitivity frequently links to bacitracin resistance in bacteria (36), and this also appears to be the case in the cholate-adapted strain. On the other hand, the vancomycin sensitivity did not change, demonstrating that vancomycin binding sites have not changed in the mutant, thus suggesting that the different responses are due to antimicrobial-dependent differences in cell envelope permeation.

Although the cholate-adapted Δ LmrCD^r cells show a greater resistance to bile acids than wild-type cells, this is not the case for all derivatives tested (Table 2). The cholate-adapted cells remained very susceptible to the hydrophobic bile acid lithocholate and are substantially more sensitive to glycodeoxycholate than the wild-type. Our data show that the unconjugated bile acids (cholate, deoxycholate, chenodeoxycholate, and lithocholate) are more toxic to *L. lactis* than the conjugated forms (glycodeoxycholate, glycocholate, and taurocholate) as is the case with bacteria in general (39). The lack of toxicity of taurocholate and glycocholate may be due to the low pK_a values (~1.4 and ~2.4, respectively) which render these compounds fully ionized at neutral pH. Thus, these molecules are likely highly membrane impermeable. Overall, the susceptibility of *L. lactis* for bile acids seems to be directly related to their hydrophobicity. Indeed, membrane permeability decreases with the number of the hydroxyl group additions (28). However, this assessment does not take into account the extrusion-based resistance mechanism that prevails in wild-type cells. Notably, the toxicity of glycodeoxycholate is strongly dependent on the activity of LmrCD. This is intriguing since glyco-forms of bile acid conjugates are more toxic than tauro-conjugates while the former also represent the predominant form of bile salt conjugate in human bile (5). Therefore, the observation that LmrCD renders cells highly resistant to glycodeoxycholate suggests that bile acids are natural substrates of this transporter. As homologs of LmrCD are widely distributed among gut bacteria, ABC-type MDR transporters may be important factors in colonization and survival in the intestine.

ACKNOWLEDGMENTS

We gratefully acknowledge the contribution of I. Stokroos of the Department of Cell Biology, Molecular Imaging and Electron Microscopy Section (UMCG), University of Groningen, for the scanning electron microscopy work.

A.Z. was supported by a scholarship from the Higher Education Commission, Government of Pakistan.

REFERENCES

- Agustiandari, H., J. Lubelski, H. B. van den Berg van Saparoea, O. P. Kuipers, and A. J. Driessen. 2008. LmrR is a transcriptional repressor of expression of the multidrug ABC transporter LmrCD in *Lactococcus lactis*. *J. Bacteriol.* **190**:759–763.
- Arthur, M., and P. Courvalin. 1993. Genetics and mechanisms of glycopeptide resistance in enterococci. *Antimicrob. Agents Chemother.* **37**:1563–1571.
- Barna, J. C., and D. H. Williams. 1984. The structure and mode of action of glycopeptide antibiotics of the vancomycin group. *Annu. Rev. Microbiol.* **38**:339–357.
- Begley, M., C. Hill, and C. G. Gahan. 2003. Identification and disruption of btlA, a locus involved in bile tolerance and general stress resistance in *Listeria monocytogenes*. *FEMS Microbiol. Lett.* **218**:31–38.
- Begley, M., C. Hill, and C. G. Gahan. 2006. Bile salt hydrolase activity in probiotics. *Appl. Environ. Microbiol.* **72**:1729–1738.
- Binder, H. J., B. Filburn, and M. Floch. 1975. Bile acid inhibition of intestinal anaerobic organisms. *Am. J. Clin. Nutr.* **28**:119–125.
- Bolhuis, H., D. Molenaar, G. Poelarends, H. W. Van Veen, B. Poolman, A. J. Driessen, and W. N. Konings. 1994. Proton motive force-driven and ATP-dependent drug extrusion systems in multidrug-resistant *Lactococcus lactis*. *J. Bacteriol.* **176**:6957–6964.
- Bolhuis, H., G. Poelarends, H. W. Van Veen, B. Poolman, A. J. Driessen, and W. N. Konings. 1995. The lactococcal *lmrP* gene encodes a proton motive force-dependent drug transporter. *J. Biol. Chem.* **270**:26092–26098.
- Bolotin, A., B. Quinquis, A. Sorokin, and D. S. Ehrlich. 2004. Recent genetic transfer between *Lactococcus lactis* and enterobacteria. *J. Bacteriol.* **186**:6671–6677.
- Bron, P. A., M. Marco, S. M. Hoffer, M. E. Van, W. M. de Vos, and M. Kleerebezem. 2004. Genetic characterization of the bile salt response in *Lactobacillus plantarum* and analysis of responsive promoters in vitro and in situ in the gastrointestinal tract. *J. Bacteriol.* **186**:7829–7835.
- Bron, P. A., D. Molenaar, W. M. de Vos, and M. Kleerebezem. 2006. DNA micro-array-based identification of bile-responsive genes in *Lactobacillus plantarum*. *J. Appl. Microbiol.* **100**:728–738.
- Chen, I., P. J. Christie, and D. Dubnau. 2005. The ins and outs of DNA transfer in bacteria. *Science* **310**:1456–1460.
- den Hengst, C. D., S. A. van Hijum, J. M. Geurts, A. Nauta, J. Kok, and O. P. Kuipers. 2005. The *Lactococcus lactis* CodY regulon: identification of a conserved cis-regulatory element. *J. Biol. Chem.* **280**:34332–34342.
- de Ruyter, P. G., O. P. Kuipers, and W. M. de Vos. 1996. Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl. Environ. Microbiol.* **62**:3662–3667.
- Faber, K. N., M. Muller, and P. L. Jansen. 2003. Drug transport proteins in the liver. *Adv. Drug Deliv. Rev.* **55**:107–124.
- Gasson, M. J., J. J. Godon, C. J. Pillidge, T. J. Eaton, K. Jury, and C. A. Shearman. 1995. Characterization and exploitation of conjugation in *Lactococcus lactis*. *Int. Dairy J.* **5**:757–762.
- Gasson, M. J., S. Swindell, S. Maeda, and H. M. Dodd. 1992. Molecular rearrangement of lactose plasmid DNA associated with high-frequency transfer and cell aggregation in *Lactococcus lactis* 712. *Mol. Microbiol.* **6**:3213–3223.
- George, A. M. 1996. Multidrug resistance in enteric and other gram-negative bacteria. *FEMS Microbiol. Lett.* **139**:1–10.
- Grohmann, E., G. Muth, and M. Espinosa. 2003. Conjugative plasmid transfer in gram-positive bacteria. *Microbiol. Mol. Biol. Rev.* **67**:277–301, table.
- Gunn, J. S. 2000. Mechanisms of bacterial resistance and response to bile. *Microbes Infect.* **2**:907–913.
- Higgins, C. F. 2007. Multiple molecular mechanisms for multidrug resistance transporters. *Nature* **446**:749–757.
- Hord, N. G. 2008. Eukaryotic-microbiota crosstalk: potential mechanisms for health benefits of prebiotics and probiotics. *Annu. Rev. Nutr.* **28**:215–231.
- Hyde, A. J., J. Parisot, A. McNichol, and B. B. Bonev. 2006. Nisin-induced changes in *Bacillus* morphology suggest a paradigm of antibiotic action. *Proc. Natl. Acad. Sci. USA* **103**:19896–19901.
- Kimoto, H., J. Kurisaki, N. M. Tsuji, S. Ohmomo, and T. Okamoto. 1999. Lactococci as probiotic strains: adhesion to human enterocyte-like Caco-2 cells and tolerance to low pH and bile. *Lett. Appl. Microbiol.* **29**:313–316.
- Kimoto, H., M. Nomura, M. Kobayashi, K. Mizumachi, and T. Okamoto. 2003. Survival of lactococci during passage through mouse digestive tract. *Can. J. Microbiol.* **49**:707–711.
- Kimoto-Nira, H., K. Mizumachi, M. Nomura, M. Kobayashi, Y. Fujita, T. Okamoto, I. Suzuki, N. M. Tsuji, J. Kurisaki, and S. Ohmomo. 2006. *Lactococcus* sp. as potential probiotic lactic acid bacteria. *Jpn. Agric. Res. Q.* **41**:181–189.
- Kramer, N. E., E. J. Smid, J. Kok, K. B. de, O. P. Kuipers, and E. Breukink. 2004. Resistance of gram-positive bacteria to nisin is not determined by lipid II levels. *FEMS Microbiol. Lett.* **239**:157–161.
- Kurdi, P., K. Kawanishi, K. Mizutani, and A. Yokota. 2006. Mechanism of growth inhibition by free bile acids in lactobacilli and bifidobacteria. *J. Bacteriol.* **188**:1979–1986.
- Kurdi, P., H. Tanaka, H. W. Van Veen, K. Asano, F. Tomita, and A. Yokota. 2003. Cholic acid accumulation and its diminution by short-chain fatty acids in bifidobacteria. *Microbiology* **149**:2031–2037.
- Li, X. Z., and H. Nikaido. 2004. Efflux-mediated drug resistance in bacteria. *Drugs* **64**:159–204.
- Lubelski, J., J. A. de, M. R. van, H. Agustiandari, O. P. Kuipers, J. Kok, and A. J. Driessen. 2006. LmrCD is a major multidrug resistance transporter in *Lactococcus lactis*. *Mol. Microbiol.* **61**:771–781.
- Lubelski, J., W. N. Konings, and A. J. Driessen. 2007. Distribution and physiology of ABC-type transporters contributing to multidrug resistance in bacteria. *Microbiol. Mol. Biol. Rev.* **71**:463–476.
- Lubelski, J., P. Mazurkiewicz, M. R. van, W. N. Konings, and A. J. Driessen. 2004. *ydaG* and *ydbA* of *Lactococcus lactis* encode a heterodimeric ATP-binding cassette-type multidrug transporter. *J. Biol. Chem.* **279**:34449–34455.
- Luo, H., K. Wan, and H. H. Wang. 2005. High-frequency conjugation system facilitates biofilm formation and pAMβ1 transmission by *Lactococcus lactis*. *Appl. Environ. Microbiol.* **71**:2970–2978.
- Ma, D., D. N. Cook, J. E. Hearst, and H. Nikaido. 1994. Efflux pumps and drug resistance in gram-negative bacteria. *Trends Microbiol.* **2**:489–493.
- Mantovani, H. C., and J. B. Russell. 2001. Nisin resistance of *Streptococcus bovis*. *Appl. Environ. Microbiol.* **67**:808–813.

37. Matsuura, M., R. Saldanha, H. Ma, H. Wank, J. Yang, G. Mohr, S. Cavanagh, G. M. Dunny, M. Belfort, and A. M. Lambowitz. 1997. A bacterial group II intron encoding reverse transcriptase, maturase, and DNA endonuclease activities: biochemical demonstration of maturase activity and insertion of new genetic information within the intron. *Genes Dev.* **11**:2910–2924.
38. Nishino, K., and A. Yamaguchi. 2001. Analysis of a complete library of putative drug transporter genes in *Escherichia coli*. *J. Bacteriol.* **183**:5803–5812.
39. Okoli, A. S., T. Wadstrom, and G. L. Mendz. 2007. Minireview: bioinformatic study of bile responses in *Campylobacterales*. *FEMS Immunol. Med. Microbiol.* **49**:101–123.
40. Rallu, F., A. Gruss, and E. Maguin. 1996. *Lactococcus lactis* and stress. *Antonie van Leeuwenhoek* **70**:243–251.
41. Russell, A. D., U. Tattawasart, J. Y. Maillard, and J. R. Furr. 1998. Possible link between bacterial resistance and use of antibiotics and biocides. *Antimicrob. Agents Chemother.* **42**:2151.
42. Simon, G. L., and S. L. Gorbach. 1984. Intestinal flora in health and disease. *Gastroenterology* **86**:174–193.
43. Stagers, J. E., S. C. Frost, and M. A. Wells. 1982. Studies on fat digestion, absorption, and transport in the suckling rat. III. Composition of bile and evidence for enterohepatic circulation of bile salts. *J. Lipid Res.* **23**:1143–1151.
44. Stentz, R., K. Jury, T. Eaton, M. Parker, A. Narbad, M. Gasson, and C. Shearman. 2004. Controlled expression of CluA in *Lactococcus lactis* and its role in conjugation. *Microbiology* **150**:2503–2512.
45. Tanaka, H., K. Doesburg, T. Iwasaki, and I. Mierau. 1999. Screening of lactic acid bacteria for bile salt hydrolase activity. *J. Dairy Sci.* **82**:2530–2535.
46. Timmerman, H. M., L. E. Niers, B. U. Ridwan, C. J. Koning, L. Mulder, L. M. Akkermans, F. M. Rombouts, and G. T. Rijkers. 2007. Design of a multispecies probiotic mixture to prevent infectious complications in critically ill patients. *Clin. Nutr.* **26**:450–459.
47. van Hijum, S. A., J. A. de, R. J. Baerends, H. A. Karsens, N. E. Kramer, R. Larsen, C. D. den Hengst, C. J. Albers, J. Kok, and O. P. Kuipers. 2005. A generally applicable validation scheme for the assessment of factors involved in reproducibility and quality of DNA-microarray data. *BMC Genomics* **6**:77.
48. van Veen, H. W., K. Venema, H. Bolhuis, I. Oussenko, J. Kok, B. Poolman, A. J. Driessen, and W. N. Konings. 1996. Multidrug resistance mediated by a bacterial homolog of the human multidrug transporter MDR1. *Proc. Natl. Acad. Sci. USA* **93**:10668–10672.
49. van Velkinburgh, J. C., and J. S. Gunn. 1999. PhoP-PhoQ-regulated loci are required for enhanced bile resistance in *Salmonella* spp. *Infect. Immun.* **67**:1614–1622.
50. Vaughan, E. E., H. G. Heilig, K. Ben-Amor, and W. M. de Vos. 2005. Diversity, vitality and activities of intestinal lactic acid bacteria and bifidobacteria assessed by molecular approaches. *FEMS Microbiol. Rev.* **29**:477–490.
51. Yokota, A., M. Veenstra, P. Kurdi, H. W. Van Veen, and W. N. Konings. 2000. Cholate resistance in *Lactococcus lactis* is mediated by an ATP-dependent multispecific organic anion transporter. *J. Bacteriol.* **182**:5196–5201.
52. Zwietering, M. H., I. Jongenburger, F. M. Rombouts, and R. K. van't. 1990. Modeling of the bacterial growth curve. *Appl. Environ. Microbiol.* **56**:1875–1881.