

## Nematicidal Activity of Cassia and Cinnamon Oil Compounds and Related Compounds toward *Bursaphelenchus xylophilus* (Nematoda: Parasitaphelenchidae)

JEONG-OK KONG,<sup>1</sup> SANG-MYUNG LEE,<sup>2</sup> YIL-SEONG MOON,<sup>2</sup> SANG-GIL LEE,<sup>2</sup> YOUNG-JOON AHN<sup>1</sup>

**Abstract:** The nematicidal activity of two cassia, *Cinnamomum cassia*, oils (Especial and true), four cinnamon, *Cinnamomum zeylanicum*, oils (technical, #500, bark and green leaf), and their compounds (e.g., trans-cinnamaldehyde and trans-cinnamic acid) toward adult *Bursaphelenchus xylophilus* was examined by a direct contact bioassay. Results were compared with those of 34 related compounds. As judged by 24-hour LC<sub>50</sub> values, two cassia oils (0.084–0.085 mg/ml) and four cinnamon oils (0.064–0.113 mg/ml) were toxic toward adult *B. xylophilus*. Of 45 test compounds, trans-cinnamaldehyde (0.061 mg/ml) was the most active nematicide, followed by ethyl cinnamate,  $\alpha$ -methyl-trans-cinnamaldehyde, methyl cinnamate and allyl cinnamate (0.114–0.195 mg/ml). Potent nematicidal activity was also observed with 4-methoxycinnamitrile, trans-4-methoxycinnamaldehyde, trans-2-methoxycinnamaldehyde, ethyl  $\alpha$ -cyanocinnamate, cinnamitrile and cinnamyl bromide (0.224–0.502 mg/ml). Structure-activity relationships indicate that structural characteristics, such as types of functional groups, saturation and carbon skeleton, appear to play a role in determining the toxicities to adult *B. xylophilus*. Cassia and cinnamon oils and test compounds described merit further study as potential nematicides or leads for the control of pine wilt disease caused by *B. xylophilus*.

**Key words:** *Bursaphelenchus xylophilus*, pine wood nematode, pine wilt disease, botanical nematicide, cassia oil, *Cinnamomum cassia*, cinnamon oil, *Cinnamomum zeylanicum*, cinnamaldehyde, structure-activity relationship.

Pinewood nematode, *Bursaphelenchus xylophilus* (Stern and Buhner, 1934) Nickle, 1970, is associated with dead and dying conifers, particularly pines, and recently cut conifer logs throughout North America and Japan. The nematode is the causal agent of pine wilt disease (PWD) but is also associated with trees stressed or killed by other mortality factors, both biotic and abiotic (Linit, 1988). Among the cerambycid beetle species of the genus *Monochamus*, *M. alternatus* (Hope) and *M. carolinensis* (Olivier) are primary vectors of *B. xylophilus* in East Asia (Kobayashi et al., 1984) and the US (Linit, 1988), respectively. Control of PWD in East Asia has been principally provided by killing the vector beetles by aerial and ground applications of conventional insecticides, by fumigation of pine trees killed by the disease, or by eradicating *B. xylophilus* by trunk implantation of nematicides such as levamisol hydrochloride and emamectin benzoate (Kishi, 1995; Kosaka et al., 2001; Lee et al., 2003a, 2003b). Certain trunk-implantation nematicides often caused phytotoxicity (Kishi, 1995). Growing public concern for the environmental effects of pesticides, human health effects and undesirable effects on nontarget organisms become more critical with repeated applications of pesticides (Kishi, 1995; Mun and Woo, 2001). Additionally, factors such as increased costs of nematicides and labor have made PWD control difficult in Korea. These problems substantiate the need for biorational *B. xylophilus* control alternatives.

Plant essential oils and their constituents have been suggested as alternative sources for nematode control products because some are selective, degrade to nontoxic products and have few harmful effects on nontarget organisms and the environment (Isman, 2000; Chitwood, 2002). These potential new nematicides can be applied to pine trees in the same manner as the trunk-implanted nematicides currently used. Plant essential oils are widely available, and some are relatively inexpensive, compared with plant extracts (Isman, 2000). The US Environmental Protection Agency (EPA) lists only 17 different plant oils as being exempt from toxicity data requirements as minimum risk pesticides under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) (US EPA, 1996). Thus, much effort has been focused on these compounds and their constituents as potential sources of commercial nematode control agents. Previously the present authors reported that cinnamon, *Cinnamomum zeylanicum* Blume (Lauraceae), bark oil had potent nematicidal activity toward adult *B. xylophilus* (Kong et al., 2006).

This paper describes a laboratory study aimed at isolating nematicidal constituents of oils of cassia, *Cinnamomum cassia* Blume, and cinnamon active toward adult *B. xylophilus*. Also, the structure-activity relationships of (E)-cinnamic acid and its related compounds toward adult *B. xylophilus* are discussed.

### MATERIALS AND METHODS

**Materials:** Chemicals used in this study were as follows: benzaldehyde, benzyl cinnamate,  $\beta$ -caryophyllene, (E)-cinnamaldehyde, (E)-cinnamic acid, (E)-cinnamide, cinnamitrile, cinnamoyl chloride, cinnamyl acetate, cinnamyl alcohol, cinnamyl bromide, cinnamyl chloride, (E)-1-cinnamylpiperazine, p-cymene, ethyl cinnamate, hydrocinnamaldehyde, hydrocinnamoyl chloride, (E)-2-hydroxycinnamic acid, isopropyl cinnamate, linalool, (E)-2-methoxycinnamaldehyde, (Z)-2-methoxycinnamic acid, (E)-2-methoxy-

Received for publication July 11, 2006

<sup>1</sup> School of Agricultural Biotechnology, Seoul National University, Seoul 151–921, Republic of Korea

<sup>2</sup> Southern Forest Research Institute, Korea Forest Research Institute, Jinju 660–300, Republic of Korea

Supported by grants from the Ministry of Agriculture and Forestry through the R&D Promotion Center for Agriculture and the Ministry of Education & Human Resources Development for Brain Korea 21 Project of the Korean Government to YJA.

E-mail: yjahn@snu.ac.kr

This paper was edited by Ed Lewis.

cinnamic acid, (E)-4-methoxycinnamic acid, 4-methoxycinnamionitrile,  $\alpha$ -methyl-(E)-cinnamaldehyde, methyl cinnamate, (E)-2-methylcinnamic acid, (E)-4-methylcinnamic acid and  $\alpha$ -terpinene purchased from Aldrich (Milwaukee, WI);  $\alpha$ -copaene and eugenol from Fluka (Bucks, Switzerland); allyl cinnamate, cinnamyl cinnamate,  $\alpha$ -cyanocinnamic acid, ethyl  $\alpha$ -cyanocinnamate, ethyl 4-methoxycinnamate, hydrocinnamic acid, hydrocinnamionitrile, hydrocinnamyl acetate, hydrocinnamyl alcohol, (E)-2-hydroxycinnamaldehyde, (E)-4-methoxycinnamaldehyde and vinyl cinnamate from Tokyo Kasei (Tokyo, Japan); and limonene and  $\alpha$ -terpineol from Wako (Osaka, Japan). Hydrophobic parameter values for these compounds were calculated using ACD/Log P v8.02 (ACD/I-Lab, Montreal, Canada) (Table 1). The results of ACD/Log P v8.02 were obtained using the ACD/I-Lab service. Cassia oils Especial and true and cinnamon oils technical, #500 and green leaf oil were provided by Berjé (Bloomfield, NJ). Cinnamon bark oil was supplied by Jin-Ah Aromatics (Anyang, Gyeonggi, Korea). Polyoxyethylene hydrogenated castor oil (HCO-50), a solubilizer, was a gift from Nikko Chemical (Tokyo). Potato dextrose agar (PDA) was purchased from Becton, Dickinson and Company (Sparks, MD). All other chemicals were of reagent grade and available commercially.

*Nematodes:* *Bursaphelenchus xylophilus* isolated from chips of PWD-infected *P. densiflora* were cultured for 6 d on *Botrytis cinerea* Persoon cultured on PDA plates at  $25 \pm 1^\circ\text{C}$  in darkness as previously described (Kong et al., 2006). The cultured nematodes were separated from the culture medium by the Baermann funnel technique (Viglierchio and Schmitt, 1983). The standard nematode suspension was prepared by appropriate dilution with distilled water, and the number of nematodes was counted under a binocular microscope ( $\times 20$ ).

*Gas chromatography (GC):* A Shimadzu GC-2010 gas chromatograph (Tokyo, Japan), equipped with split injector, was used. Analytes were separated with a 0.32 mm i.d.  $\times$  60 m DB-1 MS bonded-phase fused-silica capillary column (J&W Scientific, Ringoes, NJ) with a film thickness of 0.25  $\mu\text{m}$ . The oven temperature was programmed from  $35^\circ\text{C}$  (4 min isothermal) to  $220^\circ\text{C}$  at  $2^\circ\text{C}/\text{min}$  and then to  $300^\circ\text{C}$  (held for 5 min at final temperature) at  $30^\circ\text{C}/\text{min}$ . The linear velocity of the helium carrier gas was 25.4 cm/sec ( $30^\circ\text{C}$ ) at a split ratio of 1:10. Constituents were identified based on retention time and by co-injection with authentic compounds.

*Gas chromatography-mass spectroscopy (GC-MS):* GC-MS analysis was performed using a gas chromatograph (Shimadzu GC-2010)-mass spectrometer (Shimadzu GC-MS QP-2010). The capillary column and temperature conditions for the GC-MS analyses were the same as described above for GC analysis. Helium carrier gas was used at a column head pressure of 26.9 kPa. The ion source temperature was  $200^\circ\text{C}$ , and the injector temperature was  $170^\circ\text{C}$ . The interface was kept at  $280^\circ\text{C}$ , and mass spectra were obtained at 70 eV. The effluent of the capillary column was introduced directly into the ion source of the mass spectrometer. The sector mass analyzer was set to scan from 50 to 500 amu every 0.5 sec. Components of cassia and cinnamon oils were identified by comparison of mass spectra of each peak with those of authentic samples in a mass spectra library (Anonymous, 2000).

*Bioassays:* A direct contact bioassay (Kong et al., 2006) was used to evaluate the toxicity to adult *B. xylophilus* of cassia and cinnamon oils and compounds tested. An aqueous nematode suspension (approx. 3,000 nematodes/ml) was prepared by diluting the standard nematode suspension. The nematode suspension (99  $\mu\text{l}$ ) was added separately to each well of a

TABLE 1. Hydrophobic parameter values for test compounds.

Compound	Log $P^a$	Compound	Log $P^a$
Allyl cinnamate	2.93	Hydrocinnamionitrile	1.72
(E)-Cinnamaldehyde	1.90	Hydrocinnamoyl chloride	2.65
(E)-Cinnamamide	1.35	Hydrocinnamyl acetate	2.87
(E)-Cinnamic acid	2.41	Hydrocinnamyl alcohol	1.88
Cinnamionitrile	2.32	2-Hydroxycinnamaldehyde	2.14
Cinnamoyl chloride	3.19	(E)-2-Hydroxycinnamic acid	2.43
Cinnamyl acetate	2.62	Isopropyl cinnamate	3.06
Cinnamyl alcohol	1.70	(E)-2-Methoxycinnamaldehyde	2.13
Cinnamyl bromide	3.29	(E)-4-Methoxycinnamaldehyde	2.07
Cinnamyl chloride	3.16	(Z)-2-Methoxycinnamic acid	2.42
(E)-1-Cinnamylpiperazine	2.05	(E)-4-Methoxycinnamic acid	2.36
$\alpha$ -Cyanocinnamic acid	2.06	4-Methoxycinnamionitrile	2.27
Ethyl cinnamate	2.71	$\alpha$ -Methyl-(E)-cinnamaldehyde	2.68
Ethyl $\alpha$ -cyanocinnamate	2.36	Methyl cinnamate	2.18
Ethyl 4-methoxycinnamate	2.65	(E)-2-Methylcinnamic acid	2.87
Hydrocinnamaldehyde	2.04	(E)-4-Methylcinnamic acid	2.87
Hydrocinnamic acid	1.84	Vinyl cinnamate	2.51

<sup>a</sup> Partition coefficient.

96-well plate prior to toxicity tests. Six to eight concentrations of each test material in 1  $\mu$ l of polyoxyethylene hydrogenated castor oil-ethanol solution (1 mg/ml) (Takai et al., 2000) were added to the wells. The castor oil-ethanol solution did not affect adversely adult *B. xylophilus*. Each well plate was then covered with a solid lid. Controls received 1  $\mu$ l of the castor oil-ethanol solution.

Treated and castor oil-ethanol-treated control nematodes were held at the same conditions used for nematode maintenance. Nematicidal activity was evaluated 24 hr post-treatment under a binocular microscope. Nematodes were considered to be dead if their bodies showed straight shapes without movement when they were prodded with fine wooden dowels (Kong et al., 2006). The straight-shaped nematodes never recovered, even when transferred to untreated distilled water. All treatments were replicated four to six times.

**Data analysis:** Data were corrected for control mortality using Abbott's formula (Abbott, 1925). The LC<sub>50</sub> values were calculated by probit analysis (SAS Online-Doc®, Version 8.01, Statistical Analysis System Institute, Cary, NC). Nematicidal activity was considered to be significantly different among treatments when 95% confidence limits of the LC<sub>50</sub> values failed to overlap.

## RESULTS

**Nematicidal activity of cassia and cinnamon oils:** The nematicidal activities of two cassia oils and four cinnamon oils were evaluated by comparing the LC<sub>50</sub> values estimated from the bioassay toward adult *B. xylophilus* (Table 2). As judged by 24-hr LC<sub>50</sub> values, cinnamon oil technical (0.064 mg/ml) was the most toxic oil, followed by cassia oil true and cassia oil Especial (approx. 0.084 mg/ml). Potent nematicidal activity was also observed with cinnamon oil #500, cinnamon bark oil and cinnamon green leaf oil (0.097–0.113 mg/ml). Mortality in the castor oil-ethanol-treated controls was less than 3%.

**Chemical constituents of cassia and cinnamon oils:** Cassia and cinnamon oils used were mainly composed of (E)-cinnamaldehyde (Table 3). The content of this compound was more pronounced in cinnamon oil technical (99%) and cassia oil Especial (91%) than in cinna-

mon oil #500 (82%), cassia oil true (80%), cinnamon bark oil (61%) or cinnamon green leaf oil (53%). The other major constituents, eugenol,  $\alpha$ -terpineol and p-cymene, comprised 13, 10 and 8% of cinnamon bark oil, respectively. The other major constituents, limonene, cinnamyl alcohol and eugenol, comprised 17, 16 and 13% of cinnamon green leaf oil, respectively.

**Nematicidal activity of cassia and cinnamon oil compounds:** The toxicity of cassia and cinnamon oil compounds to adult *B. xylophilus* is given in Table 4. On the basis of 24-hr LC<sub>50</sub> values, (E)-cinnamaldehyde was the most toxic compound toward adult *B. xylophilus*, followed by (E)-2-methoxycinnamaldehyde and (E)-cinnamic acid.

**Nematicidal activity of cinnamic acid derivatives:** Toxic effects on adult *B. xylophilus* in the direct contact bioassay of (E)-cinnamaldehyde, (E)-cinnamic acid and their related compounds are recorded in Table 5. Potencies varied according to the compound tested. (E)-Cinnamaldehyde was the most potent compound, followed by ethyl cinnamate,  $\alpha$ -methyl-(E)-cinnamaldehyde, methyl cinnamate, cinnamoyl chloride and allyl cinnamate (LC<sub>50</sub>, 0.114–0.195 mg/ml). Potent nematicidal activity (0.224–0.502 mg/ml) was also observed with six compounds. Moderate toxicity (0.750–1.896 mg/ml) was obtained from 16 compounds, whereas weak nematicidal activity (2.193–5.084 mg/ml) was produced from six compounds. The body of a nematode treated with these compounds usually showed a straight shape without movement when prodded with a wooden dowel. The castor oil-ethanol-treated control nematodes moved actively without being prodded.

**Structure-activity relationship:** Comparisons were made to determine nematicidal activity differences involving the skeletal structure, saturation and functional groups of cinnamaldehyde, cinnamic acid and their related compounds using the toxicity data obtained (Table 6). Aldehydes were more effective nematicides than their corresponding acids (e.g., cinnamaldehyde vs. cinnamic acid; hydrocinnamaldehyde vs. hydrocinnamic acid) and alcohols (cinnamaldehyde vs. cinnamyl alcohol; hydrocinnamaldehyde vs. hydrocinnamyl alcohol). Introduction of a functional group, such as hydroxy, methoxy or methyl, in cinnamaldehyde or cinnamic acid reduced its nematicidal activity. Allyl, ethyl and methyl cinnamates were significantly more toxic than isopropyl and vinyl cinnamates. Benzyl and cinnamyl cinnamates were ineffective. The toxicity of cinnamaldehyde, cinnamic acid, cinnamyl alcohol and cinnamionitrile was more pronounced than that of their corresponding saturated compounds.

Linear regression analysis of the toxicities of the test compounds toward adult *B. xylophilus* was performed using their LC<sub>50</sub> values and the values of the hydrophobicity parameter for the test compounds. Compounds showing LC<sub>50</sub> values > 20 mg/ml were excluded for the

TABLE 2. Nematicidal activity of cassia and cinnamon oils toward *Bursaphelenchus xylophilus* using the direct contact bioassay during a 24 hr exposure.

Material	Slope $\pm$ SE	LC <sub>50</sub> (95% CL <sup>a</sup> ), mg/ml
Cassia oil Especial	5.0 $\pm$ 0.52	0.085 (0.072–0.100)
Cassia oil true	3.9 $\pm$ 0.30	0.084 (0.072–0.095)
Cinnamon oil technical	3.7 $\pm$ 0.43	0.064 (0.053–0.075)
Cinnamon oil #500	4.3 $\pm$ 0.22	0.097 (0.090–0.107)
Cinnamon bark oil	4.5 $\pm$ 0.35	0.107 (0.093–0.125)
Cinnamon green leaf oil	6.1 $\pm$ 0.20	0.113 (0.110–0.115)

<sup>a</sup> CL denoted confidence limit.

TABLE 3. Chemical constituents of cassia and cinnamon oils identified by GC and GC-MS.<sup>a</sup>

Compound	RT <sup>b</sup> (min)	% Area					
		1 <sup>c</sup>	2	3	4	5	6
Benzaldehyde	22.35	0.8	0.9	1.3	2.2	— <sup>d</sup>	—
α-Terpinene	28.48	—	—	—	0.3	—	—
p-Cymene	28.52	—	—	—	—	8.0	—
Limonene	29.56	—	—	—	—	—	17.0
Linalool	34.69	—	—	—	—	3.0	—
α-Terpineol	41.14	—	—	—	—	9.6	—
(E)-Cinnamaldehyde	45.46	91.0	80.2	98.7	81.7	60.7	52.9
Cinnamyl alcohol	49.17	—	—	—	—	—	16.4
Eugenol	51.78	—	—	—	—	12.6	13.3
Unknown	52.05	—	—	—	—	4.8	—
Unknown	52.24	—	—	—	—	0.8	—
α-Copaene	54.98	—	1.5	—	—	—	—
(E)-Cinnamic acid	56.64	1.0	—	—	—	—	—
Cinnamyl acetate	56.97	1.8	4.2	—	9.6	—	—
β-Caryophyllene	57.13	—	—	—	—	0.5	0.4
2-Methoxycinnamaldehyde	61.67	5.4	13.2	—	6.2	—	—

<sup>a</sup> Gas chromatography-mass spectroscopy.<sup>b</sup> Retention time.<sup>c</sup> 1, Cassia oil Especial; 2, Cassia oil true; 3, Cinnamon oil technical; 4, Cinnamon oil #500; 5, Cinnamon bark oil; 6, Cinnamon green leaf oil.<sup>d</sup> Not detected.

analysis. Hydrophobicity parameter ( $r^2 = 0.024$ ,  $n = 34$ ) was not significantly related to the observed toxicities of the test compounds in the contact toxicity bioassay.

#### DISCUSSION

Very little work has been done to consider the potential of cassia and cinnamon oils to manage *B. xylophilus*, although extracts of some higher plants, particularly asteraceous and euphorbiaceous plants, have been found to possess good nematocidal activity against *B. xylophilus* (Kawazu et al., 1980a). In the present study, two cassia oils and four cinnamon oils exhibited good nematocidal activity toward adult *B. xylophilus*. Cassia and cinnamon oils might be good candidates for natural *B. xylophilus* control agents.

Mono- and sesquiterpenoids and related phenols exist in essential oils (Isman, 2000; Lawless, 2002), and

TABLE 4. Toxicity of cassia and cinnamon oil compounds to adult *Bursaphelenchus xylophilus* using the direct contact bioassay during a 24 hr exposure.

Compound	Slope ± SE	LC <sub>50</sub> (95% CL <sup>a</sup> ), mg/ml
Benzaldehyde		>20
β-Caryophyllene		>20
(E)-Cinnamaldehyde	5.7 ± 0.79	0.057 (0.042–0.069)
Cinnamyl acetate	1.1 ± 0.04	2.766 (2.560–2.994)
(E)-Cinnamic acid	1.3 ± 0.04	0.750 (0.698–0.808)
Cinnamyl alcohol	1.8 ± 0.11	1.278 (1.103–1.488)
Copaene		>20
p-Cymene		>20
Eugenol	4.3 ± 0.24	1.212 (1.073–1.360)
Limonene		>20
Linalool	1.6 ± 0.13	6.441 (5.752–7.081)
α-Terpinene		>20
α-Terpineol	2.6 ± 0.15	3.385 (3.123–3.635)

<sup>a</sup> CL denotes confidence limit.

jointly or independently they contribute to a variety of biological activities, including repelling or attracting nematodes, stimulating or inhibiting egg hatch, and nematocidal activity (Chitwood, 2002). Much effort has been focused on determining the distribution, nature and practical use of plant-derived chemicals that have nematocidal activity. Naturally occurring compounds nematocidal against *B. xylophilus* include the acetylenes (Z)-dehydromatricaria ester, 9,10-epoxyheptadec-16-en-4,6-diyn-8-ol, 1-phenylhepta-1,3,5-triyn-2-ene, 2-phenyl-5-(1-propynyl)-thiophene and tridec-1-en-3,5,7,9,11-pentayne (Kawazu et al., 1980b); the quinolizidine alkaloids, such as aloperine, cytisine, N-methylcytisine and matrine (Matsuda et al., 1989, 1991; Zhao, 1999); the lignans (–)-nortrachelogenin and (+)-pinoselinol, the phenylpropanoid methyl ferulate and the stilbene pinosylvin monomethyl ether (Suga et al., 1993); and the phenolics 3-undecylphenol and 3-(8Z-tridecenyloxy)phenol (Alen et al., 2000). In this study, the nematocidal principles of cassia and cinnamon oils were the phenylpropanoids (E)-cinnamaldehyde, cinnamyl acetate and eugenol, and the monoterpene α-terpineol. The nematocidal activity of these essential oils appeared to be associated with cinnamaldehyde content. Of the test compounds, (E)-cinnamaldehyde was the most active nematocide. Potent nematocidal activity was also observed with ethyl cinnamate, α-methyl-(E)-cinnamaldehyde, methyl cinnamate, cinnamoyl chloride and allyl cinnamate. Structural characteristics such as types of functional groups, saturation and carbon skeleton appear to play a role in determining the toxicity of these chemicals to adult *B. xylophilus*. The hydrophobicity parameter was not significantly related to the observed toxicities.

Available information is limited because immobile



TABLE 5. Toxicity of cinnamaldehyde, cinnamic acid and their related compounds to adult *Bursaphelenchus xylophilus* using the direct contact bioassay during a 24 hr exposure.

Compound	Slope (± SE)	LD <sub>50</sub> (95% CL <sup>a</sup> ), mg/ml
(E)-Cinnamaldehyde	5.7 ± 0.79	0.057 (0.042–0.069)
Hydrocinnamaldehyde	2.0 ± 0.23	0.779 (0.636–0.942)
2-Hydroxycinnamaldehyde	1.6 ± 0.08	0.877 (0.832–0.925)
(E)-2-Methoxycinnamaldehyde	3.0 ± 0.24	0.270 (0.214–0.339)
(E)-4-Methoxycinnamaldehyde	2.2 ± 0.14	0.262 (0.217–0.314)
α-Methyl-(E)-cinnamaldehyde	4.4 ± 0.16	0.131 (0.127–0.134)
(E)-Cinnamic acid	1.3 ± 0.04	0.750 (0.698–0.808)
α-Cyanocinnamic acid	1.3 ± 0.12	0.805 (0.582–1.140)
Hydrocinnamic acid	1.5 ± 0.06	1.896 (1.786–2.012)
(E)-2-Hydroxycinnamic acid	2.8 ± 0.28	0.854 (0.773–0.924)
(Z)-2-Methoxycinnamic acid	1.1 ± 0.11	1.449 (0.958–1.975)
(E)-2-Methoxycinnamic acid		>20
(E)-4-Methoxycinnamic acid	2.0 ± 0.12	5.084 (4.638–5.611)
(E)-2-Methylcinnamic acid	1.0 ± 0.08	2.940 (2.467–3.436)
(E)-4-Methylcinnamic acid	1.2 ± 0.06	1.500 (1.298–1.711)
Allyl cinnamate	1.2 ± 0.10	0.195 (0.144–0.244)
Benzyl cinnamate		>20
Cinnamyl acetate	1.1 ± 0.04	2.766 (2.560–2.994)
Cinnamyl cinnamate		>20
Ethyl cinnamate	5.8 ± 0.19	0.114 (0.111–0.117)
Ethyl α-cyanocinnamate	1.4 ± 0.11	0.333 (0.280–0.389)
Ethyl 4-methoxycinnamate	0.6 ± 0.05	0.909 (0.655–1.199)
Hydrocinnamyl acetate	1.1 ± 0.12	4.209 (2.967–5.861)
Isopropyl cinnamate	0.9 ± 0.04	1.018 (0.918–1.136)
Methyl cinnamate	3.4 ± 0.10	0.163 (0.158–0.169)
Vinyl cinnamate	1.1 ± 0.08	3.771 (2.960–4.683)
Cinnamyl alcohol	1.8 ± 0.11	1.278 (1.103–1.488)
Hydrocinnamyl alcohol	1.0 ± 0.09	2.193 (1.619–2.951)
(E)-Cinnamamide	1.6 ± 0.17	1.195 (0.968–1.467)
Cinnamionitrile	5.8 ± 0.38	0.448 (0.405–0.493)
Hydrocinnamionitrile	0.9 ± 0.05	1.687 (1.401–2.036)
4-Methoxycinnamionitrile	2.6 ± 0.10	0.224 (0.210–0.237)
Cinnamoyl chloride	1.8 ± 0.07	0.184 (0.172–0.195)
Hydrocinnamoyl chloride	0.8 ± 0.04	0.754 (0.674–0.839)
Cinnamyl bromide	1.1 ± 0.07	0.502 (0.381–0.628)
Cinnamyl chloride	2.8 ± 0.22	(0.789 (0.695–0.894)
(E)-1-Cinnamylpiperazine	1.8 ± 0.18	1.429 (1.159–1.803)

<sup>a</sup> CL denotes confidence limit.

nematodes intoxicated by some plant-derived materials and chemicals were described as dead (Jourand et al., 2004). Jourand et al. (2004) reported that an aqueous extract of *Crotalaria virgulata* subsp. *grantiana* Harvey leaves caused paralysis of *Meloidogyne incognita* (Kofoid & White) Chitwood juveniles, but the effect was reversible, i.e., nematostatic. They suggested that pyrrolizidine alkaloids such as grantianine and grantaline of *C. grantiana* (Hartmann and Witte, 1995) might be responsible for the nematostatic effect on the basis that Fabaceae (formerly Leguminosae) plant alkaloids possess anthelmintic properties (Fassuliotis and Skucas, 1969; Zhao, 1999). It has been reported that nematode bodies treated with the muscle activity blockers levamisole hydrochloride and morantel tartrate usually exhibit semicircular and coiling shapes, respectively (Kong et al., 2006). In the current study with adult *B. xylophilus*, the active compounds tested had rapid nematocidal action. The bodies of nematodes killed by these com-

TABLE 6. Structure-activity relationships of test chemicals against adult *Bursaphelenchus xylophilus*.

	Very highly active	Highly active	Moderately active	Poorly active	Inactive
Aldehydes	(E)-4-Methoxycinnamaldehyde	(E)-4-Methoxycinnamaldehyde	Hydrocinnamaldehyde	(E)-2-Methoxycinnamic acid	
Acids	α-Methyl-(E)-cinnamaldehyde	(E)-2-Methoxycinnamaldehyde	2-Hydroxycinnamaldehyde	(E)-Cinnamic acid	
			α-Cyanocinnamic acid	(E)-2-Hydroxycinnamic acid	
			(E)-2-Hydroxycinnamic acid	(Z)-2-Methoxycinnamic acid	
			(e)-4-Methylcinnamic acid	(E)-4-Methoxycinnamic acid	
			Hydrocinnamic acid		
Esters		Ethyl α-cyanocinnamate	Ethyl 4-methoxycinnamate	Cinnamyl acetate	Benzyl cinnamate
			Isopropyl cinnamate	Vinyl cinnamate	Cinnamyl cinnamate
Alcohols			Cinnamyl alcohol	Hydrocinnamyl acetate	
Others			Hydrocinnamoyl chloride	Hydrocinnamyl alcohol	
		4-Methoxycinnamionitrile	Cinnamyl chloride		
		Cinnamionitrile	Cinnamide		
		Cinnamyl bromide	(E)-1-Cinnamylpiperazine		
			Hydrocinnamionitrile		

<sup>a</sup> Very highly active, LC<sub>50</sub> 0.114–0.195 mg/ml; highly active, LC<sub>50</sub> 0.224–0.502 mg/ml; moderately active, LC<sub>50</sub> 0.750–1.896 mg/ml; poorly active, LC<sub>50</sub> 2.193–5.084 mg/ml; inactive, LC<sub>50</sub> > 20 mg/ml.

pounds usually showed straight shapes. Our present and previous other studies suggest that the nematocidal mode of action of the test phenylpropanoids might be different from that of either levamisole hydrochloride and morantel tartrate or plant alkaloids.

Results of this study indicate that the cassia and cinnamon oils and test compounds described could be useful as potential nematocides for *B. xylophilus*. Cinnamon oil is listed on the US Food and Drug Administration's GRAS (Generally Recognized as Safe) list and is exempt from toxicity data requirements by the US EPA (2004). For the practical use of the oils and test compounds as novel nematocides to proceed, further research is necessary on their systemic action, because the injected oils and compounds must be translocated and diffused to every part of the pine tree at an effective concentration. Other areas requiring attention are nematocidal mode of action, phytotoxicity and formulations for improving nematocidal potency and stability and for reducing cost.

#### LITERATURE CITED

- Abbott, W. S. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18:265–267.
- Alen, Y., Nakajima, S., Nitoda, T., Baba, N., Kanzaki, H., and Kawazu, K. 2000. Two antinematodal phenolics from *Knema hookeriana*, a Sumatran rainforest plant. *Zeitschrift für Naturforschung* 55C: 300–303.
- Anonymous. 2000. The Wiley registry of mass spectral data, 7th edition. New York: John Wiley & Sons, Inc.
- Chitwood, D. J. 2002. Phytochemical based strategies for nematode control. *Annual Review of Phytopathology* 40:221–249.
- Fassuliotis, G., and Skucas, G. P. 1969. The effect of a pyrrolizidine alkaloid ester and plants containing pyrrolizidine on *Meloidogyne incognita acrita*. *Journal of Nematology* 1:287–288.
- Hartmann, T., and Witte, L. 1995. Chemistry, biology and chemocology of the pyrrolizidine alkaloids. Pp. 155–233 in S. W. Pelletier, ed. *Alkaloids: Chemical & biological perspectives*, vol. 9. Oxford, UK: Elsevier Science.
- Isman, M. B. 2000. Plant essential oils for pest and disease management. *Crop Protection* 19:603–608.
- Jourand, P., Rapior, S., Fargette, M., and Mateille, T. 2004. Nematostatic effects of a leaf extract from *Crotalaria virgulata* subsp. *grantiana* on *Meloidogyne incognita* and its use to protect tomato roots. *Nematology* 6:79–84.
- Kawazu, K., Nishii, Y., Ishii, K., and Tada, M. 1980a. A convenient screening method for nematocidal activity. *Agricultural and Biological Chemistry* 44:631–635.
- Kawazu, K., Nishii, Y., and Nakajima, S. 1980b. Two nematocidal substances from roots of *Cirsium japonicum*. *Agricultural and Biological Chemistry* 44:903–906.
- Kishi, Y. 1995. The pine wood nematode and the Japanese pine sawyer. Tokyo: Thomas Company.
- Kobayashi, F., Yamane, A., and Ikeda, T. 1984. The Japanese pine sawyer beetle as a vector of pine wilt disease. *Annual Review of Entomology* 29:115–135.
- Kong, J. O., Lee, S. M., Moon, Y. S., Lee, S. G., and Ahn, Y. J. 2006. Nematocidal activity of plant essential oils against *Bursaphelenchus xylophilus* (Nematoda: Aphelenchoididae). *Journal of Asia-Pacific Entomology* 9:173–178.
- Kosaka, H., Aikawa, T., Ogura, N., Tabata, K., and Kiyohara, T. 2001. Pine wilt disease caused by the pine wood nematode: The induced resistance of pine trees by the avirulent isolate of nematode. *European Journal of Plant Pathology* 107:667–675.
- Lawless, J. 2002. The encyclopedia of essential oils. London, UK: Thorsons.
- Lee, S. M., Chung, Y. J., Lee, S. G., Lee, D. W., Choo, H. Y., and Park, C. G. 2003a. Toxic effects of some insecticides on the Japanese pine sawyer, *Monochamus alternatus*. *Journal of Korean Forestry Society* 92:305–312 (in Korean; English summary).
- Lee, S. M., Chung, Y. J., Moon, Y. S., Lee, S. G., Lee, D. W., Choo, H. Y., and Lee, C. K. 2003b. Insecticidal activity and fumigation conditions of several insecticides against Japanese pine sawyer (*Monochamus alternatus*) larvae. *Journal of Korean Forestry Society* 92:191–198 (in Korean; English summary).
- Linit, M. J. 1988. Nematode-vector relationships in the pine wilt disease system. *Journal of Nematology* 20:227–235.
- Matsuda, K., Kimura, M., Komai, K., and Hamada, M. 1989. Nematocidal activities of (–)-*N*-methylcytisine and (–)-anagyrene from *Sophora flavescens* against pine wood nematodes. *Agricultural and Biological Chemistry* 53:2287–2288.
- Matsuda, K., Yamada, K., Kimura, M., and Hamada, M. 1991. Nematocidal activity of matrine and its derivatives against pine wood nematodes. *Journal of Agricultural and Food Chemistry* 39:181–191.
- Mun, S. G., and Woo, Y. T. 2001. The ecological study for the rest period of Mt. Gumjeong. Busan, Korea: Busan Metropolitan City (in Korean).
- Suga, T., Ohta, S., Munesada, K., Ide, N., Kurokawa, M., Shimizu, M., and Ohta, E. 1993. Endogenous pine wood nematocidal substance in pines, *Pinus massoniana*, *P. strobes* and *P. palustris*. *Phytochemistry* 33:1395–1401.
- Takai, K., Soejima, T., Suzuki, T., and Kawazu, K. 2000. Emamectin benzoate as a candidate for a trunk-injection agent against the pine wood nematode, *Bursaphelenchus xylophilus*. *Pest Management Science* 56:937–941.
- US EPA. 1996. Exemption of certain pesticide substances from federal insecticide, fungicide, and rodenticide act requirements. U.S. Environmental Protection Agency, Final Rule, 40 CFR 152.25 (g), May 6, 1996.
- US EPA. 2004. Biopesticides—25b Minimum risk pesticides. [http://www.epa.gov/opppbd1/biopesticides/regtools/25b\\_list.htm](http://www.epa.gov/opppbd1/biopesticides/regtools/25b_list.htm).
- Viglierchio, D. R., and Schmitt, R. V. 1983. On the methodology of nematode extraction from field samples: Baermann funnel modifications. *Journal of Nematology* 15:438–444.
- Zhao, B. G. 1999. Nematocidal activity of quinolizidine alkaloids and the functional group pairs in their molecular structure. *Journal of Chemical Ecology* 25:2205–2214.