

Short report

Open Access

Molecular characterization of Umbre virus (Bunyaviridae)

Pragya D Yadav, Akhilesh C Mishra and Devendra T Mourya*

Address: Microbial Containment Complex, National Institute of Virology, 130/1 Sus Road, Pashan, Pune 411021, India

Email: Pragya D Yadav - yadavpd@icmr.org.in; Akhilesh C Mishra - acm1750@rediffmail.com; Devendra T Mourya* - mouryadt@vsnl.net.in

* Corresponding author

Published: 8 October 2008

Received: 9 September 2008

Virology Journal 2008, 5:115 doi:10.1186/1743-422X-5-115

Accepted: 8 October 2008

This article is available from: <http://www.virologyj.com/content/5/1/115>

© 2008 Yadav et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Umbre (UMB) virus was first isolated from India in 1955 and classified as Orthobunyavirus (Turlock serogroup). Eight isolates of this virus, isolated from *Culex* mosquitoes were characterized on the basis of partial glycoprotein (G2) gene. Twenty-six percent differences at nucleotide level while 17% differences at amino acid level were noted within different isolates. Phylogenetic data shows that this virus represents a distinct group within the genus Orthobunyavirus.

Findings

The viruses of *Bunyaviridae* family are spherical particles, range 80 to 120 nm in diameter and share a common genetic organization of three predominantly negative stranded RNA segments (S, M and L). Based on antigenic, genetic and ecological relatedness, the Bunyaviruses are divided into five genera. The genus Orthobunyavirus includes approximately 60 viruses, which are known to cause disease in humans (Elliot, 1996). Virological surveillance of these viruses depends primarily on detecting the viruses in arthropod vector populations in nature. Although, serological test like immunoassays are available for antigen detection for a few viruses, cross-reaction in closely related viruses cannot be ignored (Artsob *et. al.*, 1984; Hildreth *et. al.*, 1982).

Umbre virus (UMB) [strain G-1424] was first isolated from *Culex bitaeniorhynchus* mosquitoes, collected in 1955 at Umbre, Kolaba district, Maharashtra State, India. The virus has been registered in the International Arbovirus Catalogue No. 43 (Dandwate *et. al.*, 1969). During further field investigations, seven more strains of virus were isolated from *Cx. vishnui* mosquitoes. Recent reports on Bunyaviruses via. Ganjam virus isolation from Maharashtra (Joshi *et. al.*, 2004) and antibody detection of Hantan

virus in India (Chandi *et. al.*, 2005), has provided evidences that Bunyaviruses are circulating in this country but their involvement in causing human and animal disease are not known yet. In the Gene Bank, only one sequence of Turlock serogroup i.e. N gene of M'poke virus is available.

UMB viruses used in this study are listed in (Table 1) along with their geographical origin, host source and year of isolation. The available eight strains of this virus was procured from the virus registry of National Institute of Virology, Pune and propagated in *VeroE-6* cells. Cytopathic effect (CPE) was observed during 4th – 6th post infection day. Infected cells were harvested, centrifuged and supernatant was used for molecular characterization of the virus.

RNAs were isolated using chloroform, isoamylalcohol and further purified using RNAaid kit (Biogene), according to the manufacturer's instructions. RNAs were dissolved in 50 µl nuclease free water. Different sets of primers were used to amplify partial N, L and M gene. Partial M gene of 570 bp could be amplified using primer pair M14C and M619R, as described by Bowen *et. al.*, (2001), represents the nucleotide sequences of the N-terminal half

Table 1: Details of the virus strains used in the current study

Strain no.	Year of isolation	Host association	Place of isolation	Accession No.
G-1424	1955	<i>Culex bitaeniorynchus</i>	Umbre, Maharashtra	EU697948
G-7441	1956	<i>Cx. vishnui</i>	Kammavanpet, Tamil Nadu	EU697945
G-8335	1956	<i>Cx. vishnui</i>	Minnal, Tamil Nadu	EU697946
G-9601	1956	<i>Cx. vishnui</i>	Sulari, Tamil Nadu	EU697947
G-16283	1957	<i>Cx. vishnui</i>	Sathuperi, Tamil Nadu	EU678356
G-16310	1957	<i>Cx. vishnui</i>	Sathuperi, Tamil Nadu	EU697942
631308	1963	<i>Cx. vishnui</i>	Vellore, Tamil Nadu	EU697944
809365	1980	<i>Cx. vishnui</i>	Muduvadi, Karnataka	EU697943

of the G2 glycoprotein. Superscript III single step RT-PCR with Platinum Taq DNA polymerase kit (Invitrogen) was used for amplification of partial M gene according to the manufacturer's instructions.

Amplified products were detected in 2% agarose gel after staining with ethidium bromide in Tris/acetate/EDTA buffer (TAE). A desired size of 575 bp product was purified using QIAquick gel extraction kit (Qiagen), as per manufacturer's instructions. The sequences of amplified products were determined by using ABI PRISM BigDye Terminator V3.1 cycle sequencing ready reaction kit (Applied Biosystems). Amplification primers were used to sequence the amplified products. Cycle sequencing PCR program was used for 96°C-1 min, 96°C-10 sec, 50°C-5 sec and extension of 2 min at 60°C for 30 cycles.

The partial M gene sequence was curreted with the help of KODON Software and aligned with known Gene Bank sequences of Bunyamwera serogroup, California serogroup, and Kaeng Khoi viruses using clastal W program. Phylogenetic analysis was performed using Mega 3.0 by using neighbor-joining algorithm with thousand bootstrap values.

Partial M gene sequences showed maximum homology with Bunyamwera serogroup virus. Nucleotide and amino acid similarity within eight isolates varied from 74–100% and 83–99% respectively. Isolate G1424 and 809365 come together with 6% and 1% difference of nucleotide and amino acid respectively, while other six isolates club together with 5% nucleotide and 4% amino acid differences (Figure 1). Nucleotide and amino acid homology in UMB viruses ranged from 49–75% and 25–84% respectively with other Orthobunyaviruses. There were 26 amino acid conserved sites as compared to other Orthobunyaviruses.

UMB virus was isolated from two different species of *Culex* mosquitoes i.e. *Cx. bitaeniorhynchus* and *Cx. vishnui* which were collected from three different states. Both the species have different host range and breeding habitats, but the viruses obtained from these mosquito species were not different on the genomic bases. The first UMB

isolate [strain no. G 1424] and the last isolate [strain no. 809365] from India showed maximum similarity while other six isolates club together. Phylogenetic analysis of genetically characterized member of the genus Orthobunya (345 bp of glycoprotein gene G2, data not shown) are divides into six distinct lineage viz. Bunyamwera, Simbu, California encephalitis, Group C, Kaeng Khoi (KK) and UMB virus. Comparison of 550 bp of partial M gene showed four lineages, excluding Group C viruses and Simbu group of virus (Figure 1). The genetic divergence between these lineages suggests that UMB virus represent a distinct group within the genus *Orthobunya*.

UMB virus strain G 1424 and 809365 not only highly homologues on the genomic level but also in their cell infectivity pattern. These two strains took more time to show CPE in comparison of other six isolates. Complete genome sequencing may shed light, why these two isolates are separate from other six, which club together despite isolated from two different mosquito species.

Turlock and Umbre virus are distinct from each other based on neutralization test (Calisher *et al.*, 1984). Availability of more sequences of Turlock group may answer about placement of this group of viruses. Bunyaviruses being three segmented RNA viruses have the capacity to reassort their segments into new genetically distinct viruses, if the target cells are subject to dual infection. The possibility of drift, shift and UMB virus evolution towards an emerging disease pathogen cannot be predicted based on partial sequences. Complete genome sequencing of UMB virus can possibility suggest whether there is any reassortment between three genes of this virus as known for Ngeri, Batai and Jatobal virus (Briese *et al.*, 2006; Yanase *et al.*, 2006; Saeed *et al.*, 2001).

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PDY performed the PCR and sequencing. ACM helped in preparation of manuscript. DTM and PDY designed, coordinated the study and prepared the manuscript.




Figure 1
Phylogenetic comparison of partial M RNA segments of the Umbre virus with other Orthobunyaviruses. Using Mega 3.0 software Neighbor-joining analysis performed with 1000 bootstrap replicates. Umbre virus forms a separate group within *Orthobunyaviruses*. Partial M segment source are: Umbre virus strain no G-16310 (EU697942), 809365 (EU697943), 631308 (EU697944), G-7441 (EU697945), G-8335 (EU697946), G-9601 (EU697947), G-1424 (EU697948) and G-16283 (EU678356).

Acknowledgements

The authors are thankful to Department of Science and Technology, Delhi for providing the funds and thanks to Mr. Rajen Lakra for technical help during the study.

References

- Artsob H, Spence LP, Thing C: **Enzyme-linked immunosorbent assay typing of California serogroup viruses isolated in Canada.** *J Clin Microbiol* 1984, **20**:276-280.
- Bowen MD, Trappier SG, Sanchez AJ, Meyer RF, Goldsmith CS, Zeki SR, Dunster LM, Peters CJ, Ksiazek TG, Nichol ST: **A reassortant bunyavirus isolated from acute hemorrhagic fever cases in Kenya and Somalia RVF task Force.** *Virology* 2001, **291**:185-190.
- Briese T, Bird B, Kapoor K, Nichol ST, Lipkin WI: **Batai and Ngeri viruses, M segment Reassortment and association with severe febrile disease outbreaks in East Africa.** *J Virol* 2006, **80**(11):5627-5630.
- Calisher CH, Lazuick JS, Wolf KL, Muth DJ: **Antigenic relationships among Turlock serogroup Bunyaviruses as determined by neutralization test.** *Acta Virol* 1984, **28**:148-151.

5. Chandy S, Mitra S, Sathish N, Vijayakumar TS, Abraham OC, Jesudason MV, Abraham P, Yoshimatsu K, Arikawa J, Sridharan G: **A pilot study for serological evidence of hantavirus infection in human population in south India.** *Indian J Med Res* 2005, **122**:211-215.
6. Dandawate CN, Rajagopalan PK, Pavri KM, Work TH: **Virus isolations from mosquitoes collected in North Arcot District, Madras State and Chittoor district, Andhra Pradesh between November 1955 and October 1957.** *Indian J Med Res* 1969, **57**:1420-1426.
7. Elliott RM, The Bunyaviridae: **Concluding remarks and future prospects.** In *The Bunyaviridae* Edited by: Elliott RM. New York: Plenum Press; 1996:295-333.
8. Hildreth SW, Beaty BJ, Meegan JM, Frazier CL, Shope RE: **Detection of La Crosse arbovirus antigen in mosquito pools, application of chromogenic and fluorogenic enzyme immunoassay systems.** *J Clin Microbiol* 1982, **15**:879-884.
9. Joshi MV, Geevarghese G, Joshi GD, Ghodke YS, Mourya DT, Mishra AC: **Isolation of Ganjam virus from ticks collected off domestic animals around Pune, Maharashtra.** *J Med Entomol* 2005, **42**:204-206.
10. Saeed MF, Li L, Wang H, Weaver SC, Barrett ADT: **Phylogeny of the Simbu serogroup of the genus Bunyavirus.** *J Gen Virol* 2001, **82**:2173-2181.
11. Yanase T, Kato T, Yamakawa , Takayoshi K, Nakamura K, Kokuba T, Tsuda TM: **Genetic characterization of Batai virus indicates a genomic reassortment between orthobunyaviruses in nature.** *Arch virol* 2006, **151**:2253-2260.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

