

## Proteasomal Components Required for Cell Growth and Stress Responses in the Haloarchaeon *Haloferax volcanii*<sup>∇†</sup>

Guangyin Zhou, David Kowalczyk, Matthew A. Humbard, Sunil Rohatgi, and Julie A. Maupin-Furlow\*

Department of Microbiology and Cell Science, University of Florida, Gainesville, Florida 32611-0700

Received 20 August 2008/Accepted 6 October 2008

Little is known regarding the biological roles of archaeal proteases. The haloarchaeon *Haloferax volcanii* is an ideal model for understanding these enzymes, as it is one of few archaea with an established genetic system. In this report, a series of *H. volcanii* mutant strains with markerless and/or conditional knockouts in each known proteasome gene was systematically generated and characterized. This included single and double knockouts of genes encoding the 20S core  $\alpha 1$  (*psmA*),  $\beta$  (*psmB*), and  $\alpha 2$  (*psmC*) subunits as well as genes (*panA* and *panB*) encoding proteasome-activating nucleotidase (PAN) proteins closely related to the regulatory particle triple-A ATPases (Rpt) of eukaryotic 26S proteasomes. Our results demonstrate that 20S proteasomes are required for growth. Although synthesis of 20S proteasomes containing either  $\alpha 1$  or  $\alpha 2$  could be separately abolished via gene knockout with little to no impact on growth, conditional depletion of either  $\beta$  alone or  $\alpha 1$  and  $\alpha 2$  together rendered the cells inviable. In contrast, the PAN proteins were not essential based on the robust growth of the *panA panB* double knockout strain. Deletion of genes encoding either  $\alpha 1$  or PanA did, however, render cells more sensitive to growth on organic versus inorganic nitrogen sources and hypo-osmotic stress and limited growth in the presence of L-canavanine. Abolishment of  $\alpha 1$  synthesis also had a severe impact on the ability of cells to withstand thermal stress. This contrasted with what was seen for *panA* knockouts, which displayed enhanced thermotolerance. Together, these results provide new and important insight into the biological role of proteasomes in archaea.

Detailed three-dimensional structures of archaeal proteasomes have provided a firm foundation for understanding how these elaborate nanocompartmentalized complexes mediate protein degradation (3, 11). While their thermostable properties make archaeal proteasomes ideal for structural studies, the optimum growth requirements of archaea, which often include extreme pH, salt, and/or temperature, have limited our fundamental knowledge of how these proteolytic enzymes work in the ubiquitin-free archaeal cell.

Halophilic archaea or haloarchaea have developed into model organisms that are used to study many biological processes. Recent advances in haloarchaeal genome sequencing (4, 6, 14, 15) and the development of new genetic tools, such as the targeted and markerless deletion of chromosomal genes (1), have made this group of microbes ideal for providing insight into cell physiology.

*Haloferax volcanii* is a haloarchaeon which encodes at least five protein components associated with the proteasome system. These include the  $\alpha 1$ ,  $\beta$ , and  $\alpha 2$  proteins, encoded by *psmA*, *psmB*, and *psmC*, respectively, which form at least two 20S proteasome subtypes of differing subunit compositions ( $\alpha 1\beta$  and  $\alpha 1\alpha 2\beta$ ) (8, 20). *H. volcanii* also encodes two proteasome-activating nucleotidase (PAN) proteins, PanA (*panA*) and PanB (*panB*), closely related to the regulatory particle triple A-ATPase (Rpt) proteins of eukaryal 26S proteasomes

(17). These five proteasomal genes are separately dispersed throughout the major chromosome (17, 20) and, based on Northern blotting, are all translated from single-gene transcripts (7). The  $\alpha 1$  and  $\alpha 2$  proteins are also translated from the most distal and proximal ends of 2.1- and 1.6-kb polycistronic transcripts, respectively (7).

The only proteasome mutant strains generated in an archaeon to date are those of *H. volcanii*. These mutant strains include one with a novobiocin resistance marker inserted into the coding region of the *psmC* gene, encoding the  $\alpha 2$  protein (8), and a second strain with a deletion of  $\sim 100$  bp and an insertion of a mevinolin resistance marker into the *panA* gene, encoding the Rpt-like PanA protein (9). The former strain (WFD11 [*psmC1::gyrB\*-gyrA'*]) was used to purify 20S proteasomes of  $\alpha 1\beta$  composition, and the (phospho)proteome of the latter strain (GG102 [ $\Delta panA::hmgA^*$ ]) was investigated by tandem mass spectrometry based the robust number of phosphoproteins detected by two-dimensional electrophoresis in its proteome compared to what is the case for the parent strain, DS70 (9). Together, these mutant strains have provided insight into proteasome structure and function, but they are limited in number and scope.

In this report, we present the first systematic generation and phenotypic characterization of mutants with markerless knockouts and/or conditional promoter fusions of each gene associated with the proteasome system in an archaeal genome (*H. volcanii psmA*, *psmB*, *psmC*, *panA*, and *panB* genes). The same parent strain H26 ( $\Delta pyrE$ ) was used for the construction of all mutant strains to minimize nonrelated variables and enable comparison. Double mutants were also generated and characterized, including those deficient in both  $\alpha$ -type subunits ( $\alpha 1$  and  $\alpha 2$ ), in both Pan proteins (PanA and PanB), and in the

\* Corresponding author. Mailing address: Department of Microbiology and Cell Science, University of Florida, Gainesville, FL 32611-0700. Phone: (352) 392-4095. Fax: (352) 392-5922. E-mail: jmaupin@ufl.edu.

† Supplemental material for this article may be found at <http://jb.asm.org/>.

<sup>∇</sup> Published ahead of print on 17 October 2008.

TABLE 1. List of strains and plasmids used in this study

Strain or plasmid	Description <sup>b</sup>	Source or reference
<b>Strains</b>		
<i>E. coli</i> strains		
DH5 $\alpha$	F <sup>-</sup> <i>recA1 endA1 hsdR17</i> (r <sub>K</sub> <sup>-</sup> m <sub>K</sub> <sup>+</sup> ) <i>supE44 thi-1 gyrA relA1</i>	Life Technologies
GM2163	F <sup>-</sup> <i>ara-14 leuB6 fhuA31 lacY1 tsx78 glnV44 galK2 galT22 mcrA dcm-6 hisG4 rfbD1 rpsL136 dam13::Tn9 xylA5 ml-1 thi-1 mcrB1 hsdR2</i>	New England Biolabs
<i>H. volcanii</i> strains <sup>a</sup>		
DS70	Wild-type isolate DS2 cured of plasmid pHV2	19
H26	DS70 <i>pyrE2</i>	2
GZ108	H26 <i>panB</i> (devoid of PanB)	This study
GZ109	H26 <i>panA</i> (devoid of PanA)	This study
GZ114	H26 <i>psmC</i> (devoid of $\alpha$ 2)	This study
GZ120	GZ114 <i>panB</i> (devoid of $\alpha$ 2 and PanB)	This study
GZ130	H26 <i>psmA</i> (devoid of $\alpha$ 1)	This study
GZ131	GZ114 <i>panA</i> (devoid of $\alpha$ 2 and PanA)	This study
GZ132	GZ108 <i>panA</i> (devoid of PanA and PanB)	This study
GZ133	GZ108 <i>psmA</i> (devoid of $\alpha$ 1 and PanB)	This study
GZ134	GZ130 <i>panA</i> (devoid of $\alpha$ 1 and PanA)	This study
GZ136	H26 P <sub><i>maA</i></sub> - <i>psmA</i> (devoid of $\alpha$ 1 in the absence of Trp)	This study
GZ137	GZ114 ( <i>psmC</i> ) P <sub><i>maA</i></sub> - <i>psmA</i> (devoid of $\alpha$ 1 in the absence of Trp and $\alpha$ 2)	This study
GZ138	H26 P <sub><i>maA</i></sub> - <i>psmB</i> (devoid of $\beta$ in the absence of Trp)	This study
GZ112	H26 <i>psmB</i> -pJAM202	This study
<b>Plasmids</b>		
pTA131	Ap <sup>r</sup> ; pBluescript II containing P <sub><i>fdx</i></sub> - <i>pyrE2</i>	2
pET24b	Km <sup>r</sup> ; <i>E. coli</i> expression vector	Novagen
pJAM202c	Ap <sup>r</sup> Nv <sup>r</sup> ; control plasmid derived from pBAP5010	This study
pJAM202	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>psmB</i> - <i>His6</i>	8
pJAM204	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>psmA</i> - <i>His6</i>	8
pJAM205	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>psmC</i> - <i>His6</i>	8
pJAM650	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>panA</i> - <i>His6</i>	Maupin-Furlow, unpublished
pJAM648	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>panA</i>	Maupin-Furlow, unpublished
pJAM503	Ap <sup>r</sup> Nv <sup>r</sup> ; pBAP5010 containing P <sub><i>2mm</i></sub> - <i>His6</i> - <i>panA</i>	Maupin-Furlow, unpublished
pJAM2027	Ap <sup>r</sup> ; pTA131 containing <i>psmA</i> with ~500 bp of genomic DNA flanking 5' and 3' of the <i>psmA</i> coding region	This study
pJAM2030	Ap <sup>r</sup> ; pTA131 containing <i>psmB</i> with ~500 bp of genomic DNA flanking 5' and 3' of the <i>psmB</i> coding region	This study
pJAM2024	Ap <sup>r</sup> ; pTA131 containing <i>psmC</i> with ~500 bp of genomic DNA flanking 5' and 3' of the <i>psmC</i> coding region	This study
pJAM2022	Ap <sup>r</sup> ; pTA131 containing <i>panB</i> with ~500 bp of genomic DNA flanking 5' and 3' of the <i>panB</i> coding region	This study
pJAM2029	Ap <sup>r</sup> ; pTA131-derived <i>psmA</i> suicide plasmid	This study
pJAM2031	Ap <sup>r</sup> ; pTA131-derived <i>psmB</i> suicide plasmid	This study
pJAM2025	Ap <sup>r</sup> ; pTA131-derived <i>psmC</i> suicide plasmid	This study
pJAM2026	Ap <sup>r</sup> ; pTA131-derived <i>panA</i> suicide plasmid	This study
pJAM2023	Ap <sup>r</sup> ; pTA131-derived <i>panB</i> suicide plasmid	This study
pJAM2051	Ap <sup>r</sup> ; Nv <sup>r</sup> ; pJAM204 containing 323-bp P <sub><i>maA</i></sub>	This study
pJAM2053	Ap <sup>r</sup> ; pTA131-derived P <sub><i>maA</i></sub> - <i>psmA</i> suicide plasmid	This study
pJAM2052	Ap <sup>r</sup> ; Nv <sup>r</sup> ; pJAM202 containing 323-bp P <sub><i>maA</i></sub>	This study
pJAM2054	Ap <sup>r</sup> ; pTA131-derived P <sub><i>maA</i></sub> - <i>psmB</i> suicide plasmid	This study

<sup>a</sup> Parent strain, gene deletions, P<sub>*maA*</sub> promoter fusions, and/or replicating plasmids (pJAM202) are included for *H. volcanii* mutant strains.

<sup>b</sup> Km<sup>r</sup>, kanamycin resistance; Ap<sup>r</sup>, ampicillin resistance; Nv<sup>r</sup>, novobiocin resistance.

various combinations of PAN proteins with the  $\alpha$ -type subunits. The phenotypes of these strains were investigated under defined growth conditions in the presence and absence of stress agents including the addition of the amino acid analogue L-canavanine, low salt, and high temperature. The results provide new and important insight into proteasome function in archaeal cells.

#### MATERIALS AND METHODS

**Materials.** Biochemicals were purchased from Sigma-Aldrich (St. Louis, MO). Other organic and inorganic analytical-grade chemicals were from Fisher Scientific (Atlanta, GA) and Bio-Rad (Hercules, CA). Desalted oligonucleotides were from Integrated DNA Technologies (Coralville, IN). 2'-Deoxyuridine-5'-triphosphate coupled by an 11-atom spacer to digoxigenin (DIG-11-dUTP), alkaline

phosphatase-conjugated antibody raised against DIG, disodium 3-(4-methoxy-spiro{1,2-dioxetane-3,2'-(5'-chloro)tricyclo[3.3.1.1<sup>3,7</sup>]decan}-4-yl) phenyl phosphate (CSPD), and other DIG-related biochemicals were from Roche Molecular Biochemicals (Indianapolis, IN). Positively charged membranes for Southern hybridization were from Ambion (Austin, TX). Phusion and Vent DNA polymerases, restriction enzymes, T4 polynucleotide kinase, and T4 DNA ligase were from New England Biolabs (Ipswich, MA). AccuPrime GC-rich DNA polymerase was from Invitrogen (Carlsbad, CA). Standard agarose used for the separation of DNA for Southern blotting and routine analysis was from Bio-Rad Laboratories (Hercules, CA). SeaKem GTG agarose used for the separation and isolation of DNA fragments for ligation was from FMC Bioproducts (Rockland, ME).

**Strains, media, and plasmids.** Strains, oligonucleotide primers used for cloning, and plasmids are summarized in Table 1 and also in Table S1 in the supplemental material. *Escherichia coli* DH5 $\alpha$  was used for routine recombinant DNA experiments. *H. volcanii* strains were transformed (5) using plasmid DNA

isolated from *E. coli* GM2163. Liquid cultures were aerated at 200 rpm, and solid cultures were grown on 15% (wt/vol) agar plates. *E. coli* strains were grown at 37°C in Luria-Bertani medium. *H. volcanii* strains were grown at 42°C in various media including yeast extract-peptone-Casamino Acids (YPC), Casamino Acids, glycerol-minimal medium (GMM), and GMM with alanine (GMM-Ala). Medium formulae were according to *The Halohandbook* (5) with the following modifications: 20 mM glycerol served as the sole carbon source for GMM and 25 mM alanine replaced the 5 mM NH<sub>4</sub>Cl in GMM to generate GMM-Ala. Media were supplemented as needed with novobiocin (0.1 µg per ml), 5-fluoroorotic acid (5-FOA) (50 µg per ml), and uracil (10 and 50 µg per ml for growth in the presence and absence of 5-FOA, respectively). Uracil was solubilized in 1 M NaOH or 100% dimethyl sulfoxide at 50 mg per ml prior to addition to the growth medium.

For growth assays, cells were grown in YPC, GMM-Ala, and GMM as indicated. Cells were freshly inoculated from -80°C glycerol stocks onto appropriate agar-based media on plates. Cells were grown twice to log phase in 2-ml portions of media and used as an inoculum for final analysis of growth under various conditions as described below. Each subculture was inoculated to a final optical density at 600 nm (OD<sub>600</sub>) of 0.01 to 0.02. For analysis of growth rate and cell yield, cells were grown in 20 ml of media in 250-ml baffled Erlenmeyer flasks. For stress studies, cells were grown in 2 ml of media in capped 13- by 100-mm<sup>2</sup> culture tubes. For hypo-osmotic stress studies, cells were grown in GMM-Ala with the standard 2.46 M NaCl as well as concentrations of NaCl reduced as low as 1.0 M. GMM-Ala with no NaCl was added to standard medium for these assays. For canavanine sensitivity assays, cells were grown in GMM-Ala supplemented with sterile-filtered L-canavanine sulfate solution (10 mg/ml in water) at final concentrations of 0, 11.4, 22.7, 34.1, 45.4, 56.8, and 85.15 µM. For heat stress, cells were diluted to OD<sub>600</sub> of 0.04 units and transferred in aliquots (200 µl per 1.7-ml Eppendorf tube) to a 65°C water bath for 0 to 7 h. Aliquots were removed, diluted, and grown on GMM-Ala agar at 42°C. For liquid cultures, growth was monitored by an increase in OD<sub>600</sub> (where 1 OD<sub>600</sub> unit equals approximately 1 × 10<sup>9</sup> CFU per ml for all strains used in this study). For solid cultures, growth was determined by CFU per ml by spotting 20-µl aliquots of dilution series. All experiments were performed at least three times with triplicate cultures for each.

**DNA isolation and analysis.** DNA was separated by electrophoresis using 0.8% (wt/vol) agarose gels in 1× TAE electrophoresis buffer (40 mM Tris acetate, 2 mM EDTA, pH 8.5). Plasmid DNA was isolated from *E. coli* strains by use of the QIAprep spin miniprep kit (Qiagen, Valencia, CA). DNA fragments were isolated from agarose gels by use of the QIAquick gel extraction kit (Qiagen) as needed. PCRs were purified by MinElute (Qiagen) prior to modification by restriction enzymes (BamHI, HindIII, SpeI, XbaI, or NdeI) or T4 DNA polynucleotide kinase. For rapid PCR screening, template DNA was released from *H. volcanii* mutant strains and recombinant *E. coli* DH5α strains as follows. Isolated colonies were transferred to 30 µl of sterile deionized H<sub>2</sub>O by use of toothpicks, boiled (5 min), chilled on ice (10 min), and centrifuged (14,000 × g; 10 min at room temperature). Supernatant (5 to 10 µl) was used as the template for PCR. For Southern blotting, *H. volcanii* genomic DNA was isolated by DNA spooling (5).

**PCRs.** High-fidelity double-stranded DNA, used for the construction of plasmids listed in Table 1, was amplified by PCR using Phusion, Vent, or a mixture of AccuPrime GC-rich and Vent DNA polymerase at a 9:1 ratio. *Taq* DNA polymerase was used for screening mutant strains and for the generation of the DIG-labeled double-stranded DNA probes that were used for Southern blotting. All PCRs were performed according to the instructions of the suppliers with the following modifications: 3% (vol/vol) dimethyl sulfoxide was included as needed and 0.1 mM deoxyribonucleoside triphosphate mix was added to the standard DIG-labeling reaction mixture, which included 1× DIG deoxyribonucleoside triphosphate (catalog no. 1277065; Roche). Primer pairs and template DNA used for the PCRs are outlined in Table S1 in the supplemental material. PCR was performed using an iCycler or GeneCycler (Bio-Rad Laboratories), and products were analyzed on 0.8% (wt/vol) agarose gels in TAE buffer. Gels were photographed after being stained with ethidium bromide at 0.5 µg · ml<sup>-1</sup> with a Mini visionary imaging system (Fotodyne, Hartland, WI). Sizes of the fragments were estimated using the Hi-Lo DNA molecular weight marker (Minnesota Molecular, Minneapolis, MN).

**DNA sequencing.** Fidelity of all PCR amplified products was confirmed by sequencing the DNA of the plasmid inserts listed in Table 1. Sequence fidelity was also confirmed for the DNA fragments amplified by PCR from the genomic DNA of *H. volcanii* proteasomal mutant strains using 'BamHI Forward' and 'HindIII Reverse' primers. Both strands of DNA were sequenced by Sanger automated DNA sequencing using an Applied Biosystems model 3130 genetic analyzer (ICBR Genomics Division, University of Florida).

**Southern blotting.** Genomic DNA isolated from *H. volcanii* parent and mutant strains (~10 µg) was digested for 6 to 8 h with restriction enzymes, which generated DNA fragments of 1.1 to 2.7 kb, based on the most recent version of the genome sequence (<http://archaea.ucsc.edu/>, April 2007 version). The cleaved DNA was separated by agarose gel electrophoresis (20 V, 15 to 16 h) and capillary transferred and UV cross-linked to nylon membranes. Double-stranded DNA probes specific for either of the ~500-bp regions flanking 5' or 3' of the coding region of the target gene were labeled with DIG by PCR (see above and Table S1 in the supplemental material for details). DIG-labeled probes were hybridized to the Southern blots and detected by CSPD-mediated chemiluminescence as recommended by the supplier (Roche) with the following modification. An increase in stringency from 0.5× SSC (1× SSC is 0.15 M NaCl plus 0.015 M sodium citrate), 0.1% sodium dodecyl sulfate to 0.1× SSC, 0.1% sodium dodecyl sulfate was included in the washing of the membranes at 65°C after hybridization as needed. Sizes of the fragments were estimated by methylene blue staining of the Hi-Lo DNA molecular weight markers (Minnesota Molecular) on the membrane.

**Chromosomal knockout of proteasomal genes.** Proteasomal genes were targeted for markerless deletion from the chromosome of *H. volcanii* H26, GZ114, GZ108, or GZ130 by the *pyrE2*-based "pop-in/pop-out" method described in reference 2. In brief, primer pairs (see Table S1 in the supplemental material for details) were designed to PCR amplify ~500 bp of genomic DNA flanking the 5' and 3' ends of the coding region for each target gene. BamHI and HindIII sites were included in the primers ('BamHI Forward' and 'HindIII Reverse') to facilitate cloning of the PCR-generated DNA fragments into plasmid vector pTA131. For all genes with the exception of *panA*, an intermediate plasmid which carried the target gene in addition to the ~500 bp 5' and 3' of this gene was generated and used as a template in inverse PCRs with 'Inverse Forward' and 'Inverse Reverse' primer pairs. Products from the inverse PCRs were phosphorylated using T4 polynucleotide kinase, ligated, and transformed into *E. coli* DH5α for the generation of the suicide plasmid. For the generation of the *panA* suicide plasmid, the 'BamHI Forward' and 'Inverse Reverse' primer pairs and 'Inverse Forward' and 'HindIII Reverse' primer pairs were used in two separate PCRs, respectively. These ~500-bp PCR products were purified, phosphorylated, and cloned into the BamHI and HindIII sites of pTA131 by three-way ligation to generate the suicide plasmid. Suicide plasmids which carried the target gene deletion were identified by PCR screening of isolated colonies, sequenced to confirm fidelity, and used for homologous recombination with the *H. volcanii* genome by the pop-in/pop-out method with 5-FOA (2). Colonies were screened for the absence of a readily generated ~500-bp PCR product by use of primers specific for the coding region of the gene ('Negative-Forward' and 'Negative-Reverse' primer pairs). Strains which did not generate this PCR product were confirmed to be mutant strains by Southern blotting (as described above) and PCR with 'Confirm-Forward' and 'Confirm-Reverse' primer pairs. These latter PCR products were separated by agarose gel electrophoresis and sequenced to confirm DNA fidelity (as described above).

***P<sub>maA</sub>* promoter fusions of proteasomal genes.** The tryptophan-dependent tryptophanase (*P<sub>maA</sub>*) promoter region was inserted upstream of *psmA* and *psmB* on the genome of H26 and upstream of *psmA* on the genome of GZ114 Δ*psmC* by use of the following approach. A 337-bp XbaI-to-NdeI DNA fragment carrying the *P<sub>maA</sub>* promoter region was isolated from the genome of *H. volcanii* DS70 by PCR using the XbaI-*P<sub>maA</sub>* up and NdeI-*P<sub>maA</sub>* down primer pairs as listed in Table S1 in the supplemental material. This PCR product was inserted in the NdeI and XbaI sites of plasmids pJAM202 and pJAM204 to replace the *P<sub>2<sub>mm</sub></sub>* with the *P<sub>maA</sub>* promoter upstream of *psmB* and *psmA* and generate plasmids pJAM2052 and pJAM2051, respectively (Table 1). The 1.0- to 1.7-kb XbaI-to-StuI fragments of pJAM2051 and pJAM2052 were inserted into the XbaI to EcoRV sites of pTA131 to generate "half-suicide" plasmids, which were completed by respectively cloning (into the XbaI and AleI sites) the ~500- to 800-bp region immediately flanking the 5' end of *psmA* and *psmB* on the genome. These 5' regions were generated by PCR using the following primer pairs: SpeI-*P<sub>maA</sub>*-*psmB* and 00375-Confirm-Forward (for *psmB*) and XbaI-*P<sub>maA</sub>*-*psmA* and 00857-Confirm-Forward (for *psmA*) (see Table S1 in the supplemental material for details). The PCR products were treated with T4 polynucleotide kinase and SpeI (for *psmB*) or XbaI (for *psmA*) prior to cloning into the half-suicide plasmids. The final suicide plasmids for insertion of the *P<sub>maA</sub>* promoter upstream of *psmA* and *psmB* on the *H. volcanii* genome were pJAM2053 and 2054, respectively. Colonies were screened by patching onto GMM plates plus and minus tryptophan. Those strains exhibiting growth only in the presence of tryptophan were confirmed by Southern blotting and PCR using primer pairs listed in Table S1 in the supplemental material.

## RESULTS

**Generation of *H. volcanii* strains deficient in the synthesis of proteasomal proteins with a genetic approach.** The proteasome genes of *H. volcanii* were targeted for markerless deletion to examine the phenotypic consequences of proteasome dysfunction with genetic methods. The genes targeted for deletion included those encoding the  $\alpha 1$  (*psmA*),  $\beta$  (*psmB*), and  $\alpha 2$  (*psmC*) subunits of 20S core proteasomes and the PanA (*panA*) and PanB (*panB*) components of PAN complexes. The *pyrE2* mutant strain H26 served as the parent for this initial series of knockouts to facilitate the use of the “pop-in/pop-out” method described by Allers et al. (2). Up to 205 colonies per target gene were isolated and screened for generation of the desired chromosomal deletion. Initial screening was performed using PCRs with primers specific for the coding region of the target gene (‘Negative-Forward’ and ‘Negative-Reverse’ primer pairs; see Table S1 in the supplemental material). This enabled robust and reproducible high-throughput screening to detect colonies no longer able to generate the ~500-bp PCR product in contrast to the parent strain. Putative mutant strains were confirmed by Southern blotting using a probe specific for the regions flanking the gene deletion and PCR (see Fig. S1 and S2 in the supplemental material) using primer pairs annealing ~200 bp outside (5’ and 3’) of the genomic DNA region carried on the suicide plasmid (‘Confirm-Forward’ and ‘Confirm-Reverse’; see Tables S1 and S2 in the supplemental material). With exception of *psmB*, all four proteasomal genes were deleted singly from the chromosome, with knockout frequencies ranging from 1.6 to 14% with an average of ~6% (see Table S3 in the supplemental material). The *psmB* gene, encoding the sole  $\beta$  subunit which harbors the central proteolytic active site of the 20S core of the proteasome system, was deleted only in part when a wild-type copy of the gene was included in *trans* on the self-replicating plasmid pJAM202 (see Table S3 and Fig. S1 in the supplemental material).

By use of a similar “pop-in/pop-out” strategy, markerless double mutations of the proteasome system were also generated and confirmed by Southern blotting and PCR (see Fig. S1 and S2 in the supplemental material). Using the  $\Delta$ *psmC* strain GZ114, double mutant strains no longer encoding  $\alpha 2$  and either PanA or PanB were generated (GZ131 or GZ120, respectively). Likewise, using the  $\Delta$ *panB* strain GZ108 as the parent, double mutants no longer encoding PanB and either  $\alpha 1$  or PanA were generated (GZ133 or GZ132, respectively). With *psmA* mutant GZ130 as the parent strain, a double knockout devoid of  $\alpha 1$  and PanA was also generated (GZ134). The knockout frequencies for these double mutations ranged from ~2 to 16% with an average of ~7% (see Table S3 in the supplemental material) and were similar to those frequencies obtained when generating the single-proteasomal-gene mutations, as described above. In contrast, the construction of a double mutant strain deficient in  $\alpha 1$  and  $\alpha 2$  production was not successful when using either GZ114 ( $\Delta$ *psmC*) or GZ130 ( $\Delta$ *psmA*) as the parent strain in combination with suicide vectors designed for the deletion of either the *psmA* or the *psmC* gene, respectively. Over 350 isolated colonies in total were screened for a *psmA psmC* double mutation by the high-throughput PCR method (described above and in Materials and Methods). All of these colonies generated PCR products

which included 500 bp of the target gene and thus were similar to their respective parent strains.

To generate *H. volcanii* strains deficient in the production of all potential 20S proteasome subtypes (*i.e.*,  $\alpha 1\beta$ ,  $\alpha 1\alpha 2\beta$ , and  $\alpha 2\beta$ ), a new genetic strategy was used in which the tryptophanase promoter  $P_{trpA}$ , which is tightly controlled by tryptophan, was inserted upstream of the *psmA* and *psmB* genes (Table 1). This type of  $P_{trpA}$  promoter fusion strategy was recently used to demonstrate the consequences of depletion of the essential chaperonin protein CCT1 from *H. volcanii* (10). In the presence of tryptophan, the  $P_{trpA}$  promoter inserted upstream of the CCT1-encoding gene was transcribed and cells were viable. In contrast, when *cct1* gene transcription was off, cells were no longer able to grow in the absence of tryptophan (10). For our work, the  $P_{trpA}$  promoter was fused upstream of the *psmA* ( $\alpha 1$ ) genes of both H26 parent and GZ114 ( $\Delta$ *psmC*) strains with efficiencies ranging from ~3 to 4% (see Table S3 in the supplemental material). Likewise, the  $P_{trpA}$  promoter was fused upstream of the H26 *psmB* gene, encoding the sole  $\beta$  subunit of *H. volcanii*, at a frequency of ~7%. All strains were confirmed by Southern blotting and PCR (see Fig. S1 and S2 in the supplemental material).

**Growth of proteasome mutant strains on defined media.** To initiate phenotypic analysis, growth of the proteasome gene knockout strains was analyzed and compared to that of the parent strain in various media under standard laboratory conditions (42°C and 200 rpm). All of the single and double deletion strains displayed relatively normal growth rates and overall cell yield compared to the H26 parent strain on YPC medium (Fig. 1) and/or GMM (Table 2). Thus, singular deficiencies in the synthesis of  $\alpha 1$ ,  $\alpha 2$ , PanA, or PanB did not significantly limit the growth of *H. volcanii* under these conditions. Likewise, *psmA panA* (GZ134), *psmA panB* (GZ133), *psmC panA* (GZ131), *psmC panB* (GZ120), and *panA panB* (GZ132) double knockouts did not show impaired growth under these conditions. When the source of nitrogen in GMM (ammonium chloride) was replaced with alanine (GMM-Ala), strains with deficiencies in  $\alpha 1$  and/or PanA displayed a two- to threefold reduction in their growth rates, which revealed that these strains are sensitive to an organic versus an inorganic source of nitrogen. However, the  $\alpha 1$  and/or PanA deficiencies did not significantly impact the overall cell yield of these strains (Table 2). In a previous study, we reported a  $\Delta$ *panA* strain (GG102) that displayed a reduced growth rate and cell yield compared to its parent, DS70, in ATCC 974 medium (9). ATCC 974 is a complex medium that contains a reduced level of sodium chloride, *i.e.*, 2.1 M rather than to the 2.5 M present in YPC and GMM used in this study.

In contrast to the limited influence on cell growth of the proteasomal mutations described above (including knockout of all PAN protein synthesis, as demonstrated by  $\Delta$ *panA*  $\Delta$ *panB* of GZ132), when *H. volcanii* strains were depleted of all 20S proteasome subtypes, cell growth and overall cell yield were severely impaired (Fig. 2). This was demonstrated by insertion of the tryptophanase promoter,  $P_{trpA}$ , immediately upstream of the *psmB* gene in “wild-type” cells as well as upstream of *psmA* in a  $\Delta$ *psmC* strain (GZ138 and GZ137, respectively; Table 1). Since transcription from the  $P_{trpA}$  promoter requires tryptophan, cells can be depleted of the downstream gene product by growth on GMM, a medium devoid of amino acids.




FIG. 1. *H. volcanii* proteasomal mutant strains display growth rates and cell yields similar to that of parent strain H26 on complex YPC medium. Growth at 42°C (200 rpm) was monitored by an increase in OD<sub>600</sub>, where 1 unit was approximately  $1 \times 10^9$  CFU per ml for all strains.

The GZ137 and GZ138 strains were constructed to deplete all  $\alpha$ -type and  $\beta$ -type subunits of the 20S core particles, respectively, when grown on GMM alone. To analyze the phenotypic consequences of these gene fusions, cells were grown on GMM plus tryptophan and transferred to GMM plus or minus Trp. In the presence of tryptophan (GMM plus Trp), which turns on

the transcription of the  $P_{maA}$ -controlled genes, all strains tested were fully viable and displayed robust growth (Fig. 2). When tryptophan was omitted from the medium (GMM alone), the growth of strains depleted of all 20S proteasome core particle subtypes was severely impaired (GZ137 and GZ138). Control strains including H26 (parent) and GZ114 ( $\Delta psmC$ ) as well as GZ130 ( $\Delta psmA$ ) and GZ136 ( $P_{maA}$ - $psmA$ ) were all viable and displayed robust growth in GMM whether or not tryptophan was present. Likewise, the growth of the 20S proteasome core particle-depleted cells was complemented by copies of the proteasome genes provided in *trans*. That is, GZ137 ( $\Delta psmC$   $P_{maA}$ - $psmA$ ) was complemented for growth on tryptophan minus medium by the self-replicating pHV2-based plasmid pJAM204, which carries the *psmA* gene in *trans*. In addition, GZ138 ( $P_{maA}$ - $psmB$ ) cells were fully viable on tryptophan-minus medium when provided a copy of *psmB* (pJAM202). Both GZ137 and GZ138 strains were not complemented by the plasmid vector alone (pJAM202c).

Based on these results, production of at least a subgroup of the 20S core proteasome population is essential for *H. volcanii* growth. Although only  $\alpha 1\beta$  and  $\alpha 1\alpha 2\beta$  20S proteasome subtypes have been purified and characterized to date (8, 20), the viability of the  $\Delta psmA$  strain compared to that of the  $P_{maA}$ - $psmB$  fusion strain suggests that an  $\alpha 2\beta$  complex is also present. The essential role of 20S proteasome particles in cell growth and/or division provides substantial support toward our working model in which proteasomes are proposed to be central to the quality control and regulated turnover of proteins in archaea and thus are major players in posttranscriptional control within this group of organisms (12). This premise was originally based on the nanocompartmentalized structure and energy-dependent proteolytic mechanism conserved between archaeal proteasomes and known regulatory proteases (e.g., bacterial Clp and Lon proteases, eukaryal 26S proteasome) as well as the universal distribution of the 20S core particle and absence of other predictable, energy-dependent proteases in the cytosol of archaeal cells.

The essential function of the 20S core particles of proteasomes contrasts with what is seen for PAN complexes, which do not appear essential based on the robust growth of GZ132 ( $\Delta panA$   $\Delta panB$ ). These results, however, are consistent with the sporadic distribution of PAN (Rpt-like) coding

TABLE 2. Growth of proteasomal mutant strains

Strain no.	Proteasome gene(s) deleted	Proteasome protein deficiency	GMM <sup>a</sup>		GMM-Ala <sup>a</sup>	
			Growth rate ( <i>k</i> )	Cell yield	Growth rate ( <i>k</i> )	Cell yield
H26	None	None	0.239 ± 0.034	1.61 ± 0.12	0.183 ± 0.070	1.35 ± 0.01
GZ130	<i>psmA</i>	$\alpha 1$	0.152 ± 0.069	1.58 ± 0.01	0.086 ± 0.016	0.92 ± 0.08
GZ114	<i>psmC</i>	$\alpha 2$	0.237 ± 0.062	1.65 ± 0.02	0.189 ± 0.044	1.27 ± 0.01
GZ109	<i>panA</i>	PanA	0.225 ± 0.038	1.56 ± 0.02	0.052 ± 0.002	1.22 ± 0.09
GZ108	<i>panB</i>	PanB	0.205 ± 0.047	1.56 ± 0.05	0.158 ± 0.063	1.36 ± 0.20
GZ131	<i>psmC</i> , <i>panA</i>	$\alpha 2$ , PanA	0.182 ± 0.026	1.51 ± 0.02	0.110 ± 0.020	1.23 ± 0.05
GZ120	<i>psmC</i> , <i>panB</i>	$\alpha 2$ , PanB	0.170 ± 0.004	1.61 ± 0.03	0.120 ± 0.007	1.29 ± 0.08
GZ132	<i>panA</i> , <i>panB</i>	PanA, PanB	0.166 ± 0.014	1.58 ± 0.06	0.076 ± 0.012	1.25 ± 0.03
GZ134	<i>psmA</i> , <i>panA</i>	$\alpha 1$ , PanA	ND <sup>b</sup>	ND	0.087 ± 0.018	1.22 ± 0.22
GZ133	<i>psmA</i> , <i>panB</i>	$\alpha 1$ , PanB	ND	ND	0.095 ± 0.022	0.93 ± 0.13

<sup>a</sup> Strains were grown on GMM or GMM-Ala at 42°C (200 rpm) as indicated. Growth rate (*k*) is represented as generations per h; cell yield is represented as  $10^9$  CFU per ml.

<sup>b</sup> ND, not determined.




FIG. 2. 20S proteasomes are required for growth of *H. volcanii*. *H. volcanii* strains were grown on solid GMM supplemented with 2.5 mM tryptophan. Cells were transferred by loop to solid GMM with or without 2.5 mM tryptophan (Hv-GMM + Trp or Hv-GMM, respectively) and grown at 42°C as indicated. See Table 1 for strain genotype details.

sequences among archaea, including their absence from the genomes of *Thermoplasma* and *Pyrobaculum* spp. (12). This also lends support to our working model in which other triple-A ATPase proteins (in addition to PAN) facilitate the proteasome-mediated degradation of proteins in archaea. Likely candidates include Cdc48-like proteins, which appear universal among archaea and are linked to proteasome function in eukaryotes (12).

**Proteasome components are important for survival during and after stress.** To further investigate the phenotypes of the *H. volcanii* proteasome mutant strains, cells were grown under a variety of environmental stresses. These included growth under a low-salt condition (Fig. 3), growth in the presence of the amino acid analogue L-canavanine (Fig. 4), and growth after exposure to high temperature (Fig. 5).

*H. volcanii* is a halophilic archaeon that is sensitive to hypo-




FIG. 3. Strains deficient in 20S proteasomal  $\alpha 1$  or PanA are sensitive to hypo-osmotic stress. Growth of *H. volcanii* parent (H26) and proteasomal mutant strains in GMM-Ala with NaCl concentrations as indicated and described in Materials and Methods was determined. Growth is represented as a percentage relative to growth for that strain at 2.46 M NaCl (100% growth was  $0.92$  to  $1.36 \times 10^9$  CFU  $\cdot$  ml $^{-1}$ ). Experiments and plating were performed in triplicate and the mean  $\pm$  standard deviation (SD) was calculated.

osmotic stress, with requirements for sodium chloride in the range of 1.7 to 2.5 M for optimal growth (13). The GMM-Ala used in this study includes a variety of salts, most notably sodium chloride at 2.46 M and magnesium salts totaling 0.23 M (88 mM MgCl<sub>2</sub> and 142 mM MgSO<sub>4</sub>). GMM-Ala also includes 20 mM glycerol, which is a common osmolyte of bacteria and eukaryotes, as a carbon source. To assess the ability of *H. volcanii* parent and mutant strains to grow under a low-salt condition, the sodium chloride concentration of GMM-Ala was varied. Consistent with the salt requirements of this halophilic archaeon, little to no growth was observed for all *H. volcanii* strains when the NaCl was reduced to 1.25 M and below, even in the presence of the 20 mM glycerol used in this medium (Fig. 3). When the sodium chloride concentrations of GMM-Ala were reduced to 1.6 to 1.8 M, a reduced yet comparable level of growth was observed for parent H26 and *psmC* and/or *panB* deletion strains (GZ108, GZ114, and GZ120). Likewise, all four of these strains grew to comparable levels in medium with 2 to 2.46 M concentrations of sodium chloride. Thus, deficiencies in either  $\alpha 2$  or PanB appeared to have little to no influence on the ability of *H. volcanii* to grow under a

low-salt condition (Fig. 3A). In contrast, strains with deficiencies in  $\alpha 1$  and/or PanA were more sensitive to hypo-osmotic stress than their parent, H26 (Fig. 3B). All strains carrying deletions in *psmA* and/or *panA* displayed an approximately twofold reduction in overall cell yield when grown in low-salt medium (1.6 to 2 M NaCl) compared to what was seen for parent strain H26 (Fig. 3B). Growth of the single  $\Delta psmA$  and  $\Delta panA$  deletion strains was similar to that of the double  $\Delta psmA \Delta panA$  knockout strain under all sodium chloride concentrations examined. Thus, mutations in *psmA* and *panA* were not additive, suggesting PanA and  $\alpha 1$  are associated in the same hypo-osmotic stress response pathway.

Both *psmA* and *panA* deletion strains were complemented to wild-type levels of growth under a low-salt condition by gene copies provided in *trans* (Fig. 3C). The *psmA* deletion strain GZ130 was readily complemented to wild-type levels of growth under a low-salt condition by transformation of plasmid pJAM204, encoding the  $\alpha 1$  protein with a C-terminal polyhistidine tag (His<sub>6</sub>) and was not complemented by plasmid vector pJAM202c alone. Thus, the hypo-osmotic stress phenotype observed for the *psmA* deletion strain is due to an  $\alpha 1$  defi-




FIG. 4. Strains deficient in 20S proteasomal  $\alpha$ 1 or PanA are sensitive to the amino acid analogue L-canavanine. Growth of *H. volcanii* parent (H26) and proteasomal mutant strains in GMM-Ala supplemented with L-canavanine as indicated and described in Materials and Methods was determined. Growth is represented as a percentage relative to growth at 0  $\mu$ M canavanine (100% growth similar to that for Fig. 3). Experiments were performed in triplicate and the mean  $\pm$  SD was calculated.

ciency. Furthermore, the His<sub>6</sub> tag, which is fused to the C terminus of  $\alpha$ 1 and predicted to be localized to the external wall of the 20S core cylinder (8), does not impair the ability to complement the *psmA* knockout strain. Similar to the *psmA* strain, the *panA* deletion strain GZ109 was complemented in *trans* by plasmid pJAM648, which encodes the PanA protein, and GZ109 was not complemented by plasmid vector pJAM202c alone (Fig. 3C). In contrast to what was seen for the *psmA* strain, however, plasmids pJAM650 and pJAM503, encoding PanA with C- and N-terminal His<sub>6</sub> tags, respectively, were unable to fully restore the growth of the *panA* mutant to wild-type levels under low-salt conditions (Fig. 3C). Both His<sub>6</sub>-tagged PanA proteins synthesized from these plasmids are purified as soluble complexes from *H. volcanii* cells (unpub-

lished results). The N terminus of PanA is predicted to form a coiled-coil structure important in substrate recognition (17), and the C terminus of PanA includes the hydrophobic tyrosine X (HbYX) motif, which triggers 20S proteasome gate opening after ATP binding (16). These functions may be disrupted by the addition of the His<sub>6</sub> tags. Thus, although speculative, the ability to complement the *panA* mutant with only the unmodified PanA protein suggests that the recognition of substrates and 20S gate opening are important functions of PanA during hypo-osmotic stress.

Growth of *H. volcanii* strains was also examined in the presence of L-canavanine (Fig. 4), an amino acid analogue related to arginine that is incorporated into nascent polypeptides during translation and promotes protein unfolding. In general, all




FIG. 5. Strains deficient in 20S proteasomal  $\alpha$ 1 are sensitive to heat stress, whereas strains deficient in PanA are “superthermotolerant.” Parent (H26) and proteasomal mutant strains were grown to log phase at 42°C in GMM-Ala. Cells were diluted to an OD<sub>600</sub> of 0.04 units and the effects of heat on cell viability were measured by a shift to 65°C (see Materials and Methods for details). Aliquots of cells were removed at 0-, 2-, 4-, 6-, and 8-h intervals. Cells were diluted, plated on GMM-Ala, and incubated at 42°C for 5 days. Experiments were performed in triplicate and the mean  $\pm$  SD was calculated.

*H. volcanii* strains examined, including parent H26, were sensitive to this amino acid analogue with little to no growth at concentrations of  $\geq 34 \mu\text{M}$  L-canavanine (Fig. 4). Deletions in *psmC* and/or *panB* had little to no influence on this sensitivity. In contrast, strains with deletions in either *psmA* or *panA* were hypersensitive to L-canavanine and displayed impaired growth at concentrations as low as  $\sim 10 \mu\text{M}$ , in contrast to what was seen for parent strain H26, the growth of which was not impaired at this concentration. The  $\Delta\textit{psmA} \Delta\textit{panA}$  double mutant grew to a cell density similar to that of the single  $\Delta\textit{psmA}$  or  $\Delta\textit{panA}$  mutant strains at all concentrations of L-canavanine examined. Thus, similar to hypo-osmotic stress, mutations in *psmA* and *panA* were not additive in the L-canavanine-hypersensitive response.

The viability of *H. volcanii* proteasomal mutant strains was also compared to that of parent H26 after exposure to heat stress at  $65^\circ\text{C}$  (Fig. 5). Strains with deletions in *panB* and/or *psmC* genes displayed reductions in viability similar to that seen for parent H26 throughout the 6.5-h time course, with an average reduction of 4 log units from  $\sim 4 \times 10^7$  to  $3 \times 10^3$  CFU per ml (Fig. 5A). In contrast, all strains which harbored a deletion in the *psmA* gene were severely impaired in their thermotolerance (Fig. 5B). Few if any of the  $\Delta\textit{psmA}$  strains were viable after the 6.5-h exposure to  $65^\circ\text{C}$ . Surprisingly, strains deficient in *panA*, whether in the presence or in the absence of a functional *psmC* or *panB* gene, displayed greater thermotolerance than parent strain H26 (Fig. 5C). Although a significant decrease in CFU per ml was observed for these  $\Delta\textit{panA}$  strains after the 6.5-h heat stress, this reduction in cell viability was less than 2 log units from an average of  $\sim 4 \times 10^7$  to  $4.1 \times 10^5$  CFU per ml. Introduction of the *psmA* deletion into the  $\Delta\textit{panA}$  background reversed this "superthermotolerance" and resulted in cells severely impaired in their ability to overcome heat stress, similar to what was seen for the other *psmA* deletion strains (Fig. 5B).

## DISCUSSION

The ability to generate markerless deletions or conditional promoter ( $P_{\textit{maA}}$ ) fusions in genes encoding all of the known components of the proteasome system of *H. volcanii* has provided tremendous insight into the central role that this proteolytic system plays in the biology of archaea. Our results demonstrate that *H. volcanii* requires 20S core particles for growth. Although the synthesis of 20S proteasome subtypes containing either  $\alpha 1$  or  $\alpha 2$  can be separately abolished with little to no impact on cell viability under standard "nonstressful" laboratory conditions, depletion of either the  $\beta$  subunit alone or  $\alpha 1$  and  $\alpha 2$  subunits together had a profound impact on cell growth. Synthesis of an  $\alpha$ -type or  $\beta$ -type 20S proteasome subunit was essential for growth. Thus, of all the individual components of the proteasomal system, the  $\beta$  subunit, which is needed for the generation of the active site of 20S core particles, is the most crucial for cell viability. The  $\alpha 1$  and  $\alpha 2$  proteins, which form the outer rings of the 20S core particles, provide some redundancy of function. The depletion of either one of these  $\alpha$ -type subunits did not impair growth under standard laboratory conditions; however, the double deletion was lethal.

Interestingly, the *psmA* and *panA* genes, each of which were

required to overcome the hypersensitive responses to L-canavanine and low salt, encode the respective  $\alpha 1$  and PanA proteasomal proteins, shown to be predominant throughout the growth of *H. volcanii* (17). The depletion of  $\alpha 1$  via the *psmA* mutation also had a severe impact on the ability of *H. volcanii* to withstand thermal stress. In contrast, the *psmC* and *panB* genes, which had little if any impact on growth under the various conditions examined, encode the  $\alpha 2$  and PanB proteasomal proteins, respectively, which are at relatively low levels during exponential or log-phase growth and increase several-fold during the stationary phase of *H. volcanii* (17). This is consistent with our model, which suggests that  $\alpha 2$  and PanB have ancillary roles in proteasome function, whereas  $\alpha 1$ ,  $\beta$ , and PanA are the central proteasome components of the growing cell.

In contrast to the approach and findings of our study, Ruepp et al. (18) used the tripeptide carboxybenzyl-leucyl-leucyl-leucine vinyl sulfone inhibitor (Z-L3VS) to study the *in vivo* function of 20S proteasomes in *Thermoplasma acidophilum*. This archaeon differs from *H. volcanii* in that it is thermoacidophilic and does not encode PAN proteins. The addition of Z-L3VS to cell culture irreversibly modified 75 to 80% of the  $\beta$  subunits of 20S proteasomes in *T. acidophilum*. Although the effects of this proteasome inhibition impaired thermotolerance, inhibition had only a marginal influence on cell growth under normal conditions. Whether these differences are biological remains to be determined. The use of Z-L3VS limited the ability of Ruepp et al. (18) to completely abolish proteasome function in *T. acidophilum*.

It was somewhat surprising that *H. volcanii* cells harboring deletions in the *panA* gene (with exception of  $\Delta\textit{panA} \Delta\textit{psmA}$  cells) displayed enhanced thermotolerance compared to that of parent strain H26 and other proteasome mutant strains. The reason for this finding remains to be determined. Our recent characterization of the (phospho)proteome of the related *panA* mutant strain GG102, which harbors a mevinolin resistance marker insertion and deletion within the *panA* gene, reveals not only a global increase in the number of different phosphoproteins but also an increase in the abundance of proteins related to stress responses as well as Cdc48-related triple-A ATPase homologs (9). Whether any of these proteins compensate in part for the loss of the PanA function and render the cell more resistant to thermal stress than wild-type cells remains speculative.

## ACKNOWLEDGMENTS

Thanks to P. Lund, J. Soppa, and T. Allers for providing strains, plasmids, and helpful discussion. Thanks also to Jonathan Eisen and The Institute of Genomic Research for completion and annotation of the *H. volcanii* genome sequence.

This research was funded in part by the following grants to J.A.M.-F.: from NIH, R01 GM057498; and from DOE, DE-FG02-05ER15650.

## REFERENCES

- Allers, T., and M. Mevarech. 2005. Archaeal genetics—the third way. *Nat. Rev. Genet.* **6**:58–73.
- Allers, T., H. P. Ngo, M. Mevarech, and R. G. Lloyd. 2004. Development of additional selectable markers for the halophilic archaeon *Haloflex volcanii* based on the *leuB* and *trpA* genes. *Appl. Environ. Microbiol.* **70**:943–953.
- Baumeister, W., J. Walz, F. Zühl, and E. Seemüller. 1998. The proteasome: paradigm of a self-compartmentalizing protease. *Cell* **92**:367–380.
- Bolhuis, H. H., P. P. Palm, A. A. Wende, M. M. Falb, M. M. Rampp, F. F.

- Rodriguez-Valera, F. F. Pfeiffer, and D. D. Oesterhelt. 2006. The genome of the square archaeon *Haloquadratum walsbyi*: life at the limits of water activity. *BMC Genomics* **7**:169.
5. Dyall-Smith, M. 2008. The Halohandbook: protocols for halobacterial genetics, p. 67. [http://www.haloarchaea.com/resources/halohandbook/Halohandbook\\_2008\\_v7.pdf](http://www.haloarchaea.com/resources/halohandbook/Halohandbook_2008_v7.pdf).
  6. Falb, M., F. Pfeiffer, P. Palm, K. Rodewald, V. Hickmann, J. Tittor, and D. Oesterhelt. 2005. Living with two extremes: conclusions from the genome sequence of *Natronomonas pharaonis*. *Genome Res.* **15**:1336–1343.
  7. Gil, M. A., K. E. Sherwood, and J. A. Maupin-Furlow. 2007. Transcriptional linkage of *Haloferax volcanii* proteasomal genes with non-proteasomal gene neighbours including RNaseP, MOSC domain and SAM-methyltransferase homologues. *Microbiology* **153**:3009–3022.
  8. Kaczowka, S. J., and J. A. Maupin-Furlow. 2003. Subunit topology of two 20S proteasomes from *Haloferax volcanii*. *J. Bacteriol.* **185**:165–174.
  9. Kirkland, P. A., M. A. Gil, I. M. Karadzic, and J. A. Maupin-Furlow. 2008. Genetic and proteomic analyses of a proteasome-activating nucleotidase A mutant of the haloarchaeon *Haloferax volcanii*. *J. Bacteriol.* **190**:193–205.
  10. Large, A., C. Stamme, C. Lange, Z. Duan, T. Allers, J. Soppa, and P. A. Lund. 2007. Characterization of a tightly controlled promoter of the halophilic archaeon *Haloferax volcanii* and its use in the analysis of the essential *cct1* gene. *Mol. Microbiol.* **66**:1092–1106.
  11. Löwe, J., D. Stock, B. Jap, P. Zwickl, W. Baumeister, and R. Huber. 1995. Crystal structure of the 20S proteasome from the archaeon *T. acidophilum* at 3.4 Å resolution. *Science* **268**:533–539.
  12. Maupin-Furlow, J. A., M. A. Humbard, P. A. Kirkland, W. Li, C. J. Reuter, A. J. Wright, and G. Zhou. 2006. Proteasomes from structure to function: perspectives from archaea. *Curr. Top. Dev. Biol.* **75**:125–169.
  13. Mullakhanbhai, M. S., and H. Larsen. 1975. *Halobacterium volcanii* spec. nov., a Dead Sea *Halobacterium* with a moderate salt requirement. *Arch. Microbiol.* **104**:207–214.
  14. Ng, W. V., S. P. Kennedy, G. G. Mahairas, B. Berquist, M. Pan, H. D. Shukla, S. R. Lasky, N. S. Baliga, V. Thorsson, J. Sbrogna, S. Swartzell, D. Weir, J. Hall, T. A. Dahl, R. Welti, Y. A. Goo, B. Leithauser, K. Keller, R. Cruz, M. J. Danson, D. W. Hough, D. G. Maddocks, P. E. Jablonski, M. P. Krebs, C. M. Angevine, and H. Dale. 2000. Genome sequence of *Halobacterium* species NRC-1. *Proc. Natl. Acad. Sci. USA* **97**:12176–12181.
  15. Pfeiffer, F., S. C. Schuster, A. Broicher, M. Falb, P. Palm, K. Rodewald, A. Ruepp, J. Soppa, J. Tittor, and D. Oesterhelt. 2008. Evolution in the laboratory: the genome of *Halobacterium salinarum* strain R1 compared to that of strain NRC-1. *Genomics* **91**:335–346.
  16. Rabl, J., D. M. Smith, Y. Yu, S. C. Chang, A. L. Goldberg, and Y. Cheng. 2008. Mechanism of gate opening in the 20S proteasome by the proteasomal ATPases. *Mol. Cell* **30**:360–368.
  17. Reuter, C. J., S. J. Kaczowka, and J. A. Maupin-Furlow. 2004. Differential regulation of the PanA and PanB proteasome-activating nucleotidase and 20S proteasomal proteins of the haloarchaeon *Haloferax volcanii*. *J. Bacteriol.* **186**:7763–7772.
  18. Ruepp, A., C. Eckerskorn, M. Bogyo, and W. Baumeister. 1998. Proteasome function is dispensable under normal but not under heat shock conditions in *Thermoplasma acidophilum*. *FEBS Lett.* **425**:87–90.
  19. Wendoloski, D., C. Ferrer, and M. L. Dyall-Smith. 2001. A new simvastatin (mevinolin)-resistance marker from *Haloarcula hispanica* and a new *Haloferax volcanii* strain cured of plasmid pHV2. *Microbiology* **147**:959–964.
  20. Wilson, H. L., H. C. Aldrich, and J. A. Maupin-Furlow. 1999. Halophilic 20S proteasomes of the archaeon *Haloferax volcanii*: purification, characterization, and gene sequence analysis. *J. Bacteriol.* **181**:5814–5824.