

# SOS1 is the second most common Noonan gene but plays no major role in cardio-facio-cutaneous syndrome

Martin Zenker, Denise Horn, Dagmar Wiczorek, Judith Allanson, Silke Pauli, Ineke van der Burgt, Helmuth-Guenther Doerr, Harald Gaspar, Michael Hofbeck, Gabriele Gillesen-Kaesbach, Andreas Koch, Peter Meinecke, Stefan Mundlos, Anja Nowka, Anita Rauch, Silke Reif, Christian von Schnakenburg, Heide Seidel, Lars-Erik Wehner, Christiane Zweier, Susanne Bauhuber, Verena Matejas, Christian P Kratz, Christoph Thomas, Kerstin Kutsche

*J Med Genet* 2007;44:651–656. doi: 10.1136/jmg.2007.051276

**Background:** Heterozygous gain-of-function mutations in various genes encoding proteins of the Ras-MAPK signalling cascade have been identified as the genetic basis of Noonan syndrome (NS) and cardio-facio-cutaneous syndrome (CFCS). Mutations of *SOS1*, the gene encoding a guanine nucleotide exchange factor for Ras, have been the most recent discoveries in patients with NS, but this gene has not been studied in patients with CFCS.

**Methods and results:** We investigated *SOS1* in a large cohort of patients with disorders of the NS–CFCS spectrum, who had previously tested negative for mutations in *PTPN11*, *KRAS*, *BRAF*, *MEK1* and *MEK2*. Missense mutations of *SOS1* were discovered in 28% of patients with NS. In contrast, none of the patients classified as having CFCS was found to carry a pathogenic sequence change in this gene.

**Conclusion:** We have confirmed *SOS1* as the second major gene for NS. Patients carrying mutations in this gene have a distinctive phenotype with frequent ectodermal anomalies such as keratosis pilaris and curly hair. However, the clinical picture associated with *SOS1* mutations is different from that of CFCS. These findings corroborate that, despite being caused by gain-of-function mutations in molecules belonging to the same pathway, NS and CFCS scarcely overlap genotypically.

Research data accumulated during the past few years have significantly contributed to our current understanding that constitutive activation of the Ras-MAPK signalling cascade is most likely the common pathogenetic mechanism underlying Noonan syndrome (NS; OMIM 163950) and two related disorders, cardio-facio-cutaneous syndrome (CFCS; OMIM 115150) and Costello syndrome (OMIM 218040).<sup>1–5</sup> These disorders share a common pattern of congenital anomalies, including typical heart defects, craniofacial dysmorphism, short stature, skeletal anomalies and varying degrees of mental retardation.<sup>6–8</sup> *PTPN11*, encoding the protein tyrosine phosphatase SHP-2, which transduces signals from activated growth factor receptors to Ras and other signalling molecules, was first discovered as the major gene mutated in NS.<sup>1</sup> In contrast, CFCS was found to be associated with mutations in any of the genes *BRAF*, *MEK1* or *MEK2*, encoding components of the well-known B-Raf-MEK-ERK signalling cascade downstream from Ras.<sup>3,4</sup> Mutations in the *KRAS* gene have been found in patients with NS and CFCS,<sup>3,5,9,10</sup> and specific mutations in *HRAS* have been detected in the vast majority of people with Costello syndrome.<sup>2</sup> Nevertheless, no genetic defect has been found in ~30% of patients with CFCS and in ~50% of those with NS until recently.

To identify novel genes for these disorders, we screened additional functional candidates encoding proteins involved in the Ras-MAPK signalling pathway, including *SOS1*. While this work was in progress, two other groups reported gain-of-function mutations of *SOS1* in 17–20% of *PTPN11* mutation-negative patients with NS.<sup>11,12</sup> *SOS1* encodes a guanine exchange factor (GEF) for Ras catalysing the conversion of the inactive GDP-bound form of Ras to its active GTP-bound form. The precise mechanisms of *SOS1* activation are incompletely understood. Conformational changes within *SOS1* allowing Ras to access its allosteric binding site that, in the inactive state, is blocked by an intramolecular interaction involving the Dbl homology–pleckstrin homology (DH-PH) unit are presumed to play an important role.<sup>13–15</sup> NS-associated *SOS1* mutations have been suggested to result in a release of autoinhibition, followed by an increase in GEF activity subsequently leading to enhanced levels of active, GTP-bound Ras.<sup>11,12</sup> Indeed, by analysing representative mutant *SOS1* proteins, these NS-associated mutations were found to cause a gain-of-function effect, as shown by increased Ras and ERK activation in vitro.<sup>11,12</sup>

## PATIENTS AND METHODS

### Study population

Ethics approval for this study was obtained from the Ethics Committee of the University of Erlangen-Nuremberg, and informed consent for the genetic analyses was received from patients or their legal guardians.

Our initial study population (group 1) consisted of 85 clinically well-characterised patients with NS and CFCS, who were assessed by experienced clinical geneticists. In total, 53 patients (group 1A) were classified as having NS according to established diagnostic criteria,<sup>16,17</sup> and 21 patients (group 1B) were given the diagnosis of CFCS, supported by the published CFC index.<sup>18</sup> In 11 patients (group 1C), a clear-cut assignment to one of these two syndromes was not possible. This subgroup comprised six patients with normal mental development suggestive of NS but ectodermal anomalies similar to those seen in CFCS, and five infants classified as “borderline”, as they had a phenotype that could evolve into either NS or CFCS. All study participants had previously tested negative for mutations in the genes *PTPN11*, *KRAS* and *HRAS*. Patients in groups 1B and 1C had been screened for mutations in *BRAF*, *MEK1* and *MEK2* with normal results.

In addition, we included a second cohort (group 2), comprising 80 patients referred for molecular diagnosis of NS

**Abbreviations:** CFCS, cardio-facio-cutaneous syndrome; DH-PH, Dbl homology–pleckstrin homology; GEF, guanine exchange factor; NS, Noonan syndrome; OMIM, Online Mendelian Inheritance in Man

and tested negative for a *PTPN11* mutation. This group contained a considerable number of cases with a mild or atypical phenotype. A *PTPN11* mutation detection rate of 18% in the original cohort, from which group 2 was derived, reflects the clinical and genetic heterogeneity in this group.

### Molecular analysis

DNA specimens obtained from blood cells were screened for *SOS1* mutations by direct sequencing of all coding exons (ABI BigDye Terminator Sequencing Kit V.2.1; Applied Biosystems, Weiterstadt, Germany) using an automated capillary sequencer (ABI 3730, Applied Biosystems). Primer pairs and PCR conditions are available on request. In group 2, we restricted analysis to those exons in which mutations have been found in this and previous studies (exons 3, 6–8, 10–14, 16 and 19).<sup>11–12</sup> Where mutations were shown to have arisen de novo, we verified declared relationships by genotyping at 10 microsatellite loci for each patient and both parents. PCR products from one patient showing two sequence alterations in exon 10 were cloned in *Escherichia coli* (TOPO TA Cloning Kit; Invitrogen, Karlsruhe, Germany and One Shot, Invitrogen) to determine whether these alterations occurred on the same allele.

### Structural analysis of novel *SOS1* variations

The potential effects of novel *SOS1* variations detected in this study were analysed in more detail using the known three-dimensional structures of *SOS1*<sup>14–15–19–20</sup> and the computer program PyMOL (<http://www.pymol.org>). Figures were prepared using PyMOL and Adobe Photoshop (Adobe Systems Inc., USA).

## RESULTS

### *SOS1* variations and associated phenotypes

Overall, we found 18 different heterozygous *SOS1* sequence variations predicting amino acid changes of the encoded *SOS1* protein in 28 unrelated patients, including 25 sporadic cases and 3 patients with a positive family history for NS (table 1). Three of the observed sequence alterations represent possible polymorphisms: the variation c.1964C→T (P655L) which has previously been identified as a polymorphism<sup>12</sup> was found in 4 of the 28 patients. In line with the assumption that this is a non-pathogenic change, we also identified it in unaffected parents. One female patient with features of NS and significant ectodermal features (group 1C) was found to carry a novel sequence alteration, c.2999G→A (S1000N), inherited from her father, who lacked any clinical signs of NS. Based on this observation and the fact that S1000 is not highly conserved through evolution (fig 1), we regarded this variant as a probable neutral polymorphism. Another novel missense change, c.233T→G, affecting the evolutionarily conserved residue F78 (fig 1) was discovered in a male patient with many features of NS and in his mother. She was said to be clinically normal but could not be examined personally for the presence of possible minor signs of NS. Because we cannot be sure if F78C is a pathogenic mutation or polymorphism, we conservatively counted this patient among the *SOS1* mutation-negative group.

The remaining 15 *SOS1* sequence alterations observed in 22 unrelated patients were regarded as pathogenic mutations; 10 nucleotide variations identified in our cohort have been established as causative mutations previously,<sup>11–12</sup> whereas five are novel (table 1). Of the novel ones, mutation c.1300G→A has the same impact on the protein level as a previously published mutation (c.1300G→C; G434R).<sup>11</sup> The remaining four sequence variations predict novel amino acid changes. One of these, c.806T→C (M269T), affects the same amino acid residue as the previously reported mutation M269R.<sup>11–12</sup> It is likely that both exchanges of the neutral methionine by a polar residue have

similar consequences on *SOS1* protein function. The novel variations c.1433C→G (P478R) and c.1867T→A (F623I), detected in two sporadic cases, were regarded as causative mutations based on de novo occurrence, conservation (fig. 1) and function (see below). One affected person had two sequence changes, 1431G→T (Q477H) and 1433C→T (P478L). Cloning of the amplified product and sequencing of both parents demonstrated that both changes had occurred de novo on the same allele (not shown). De novo occurrence of the respective mutations was also confirmed in all other sporadic patients except two, for whom parental samples were unavailable. These two patients carried the known *SOS1* mutations c.1654A→G (R552G) and c.2536G→A (E846K). In the three patients with a positive family history of NS, previously reported *SOS1* mutations were found (table 1), which were in each case inherited from an affected mother. The nucleotide changes predicting F78C, Q477H, P478L/R, F623I and S1000N were not found in >150 population-matched controls.


Considering only those cases with bona fide mutations, 18 of 85 patients of group 1 were found to have a *SOS1* mutation. A causative *SOS1* alteration was identified in 14 of 53 (26%) patients in group 1A, 4 of 11 (36%) patients in group 1C, but in none of the 21 patients with CFCS (group 1B). Only four (5%) people with *SOS1* mutations were identified in group 2.

Clinically, most patients with *SOS1* mutations exhibited a clear NS phenotype (fig 2). One patient from group 2, a girl carrying the de novo P478R mutation, only had very mild facial features of NS and short stature, and would not have met the published strict clinical criteria.<sup>17</sup> Four patients with *SOS1* mutations had been assigned to group 1C; three of them exhibited significant ectodermal symptoms resembling those found in CFCS, but had normal psychomotor and mental development in keeping with the diagnosis of NS, and the fourth was a young infant defying a clear-cut diagnosis. Facial features of representative *SOS1* mutation-positive cases are illustrated in figure 2.

For a more detailed phenotypic analysis, we included the 22 index patients with bona fide mutations and the three affected mothers with a confirmed *SOS1* mutation. Comparison of clinical features in people carrying a *SOS1* mutation with those of our cohort and three other cohorts of *PTPN11* mutation-positive patients<sup>21–23</sup> revealed a similar spectrum of congenital heart defects, but a lower frequency of mental retardation or need for special education, easy bruising, and in males, cryptorchidism (table 2). These differences become even more obvious when combining findings of this study with data published recently.<sup>11–12</sup> However, a significantly lower prevalence of short stature in patients with a *SOS1* mutation, as indicated by previous studies,<sup>11–12</sup> could not be confirmed in our cohort. Remarkably, patients with *SOS1* mutations commonly had ectodermal manifestations, including keratosis pilaris of the face, sparse eyebrows, curly hair, and in one patient, ichthyosiform skin changes (fig 2). The incidence of ectodermal manifestations in patients with NS with *PTPN11* mutations has not yet been well documented. By reviewing a cohort (n = 42) of clinically well documented *PTPN11* mutation-positive people mainly derived from our previous study,<sup>17</sup> we could confirm a significantly higher prevalence of keratosis pilaris/hyperkeratotic skin and curly hair in patients with *SOS1* mutations compared with those with *PTPN11* alterations (58% vs 6% and 78% vs 34%, respectively). Moreover, ocular ptosis was observed more frequently in patients with NS with *SOS1* mutations than in patients with a *PTPN11* alteration (80% vs 54%).

### Structural assessment of novel *SOS1* variants

We analysed in more detail the potential effects of the novel predicted *SOS1* variants F78C and F623I, as well as those of



**Figure 1** (A) Domain organisation of the human SOS1 protein. Histone-like folds, Dbl homology (DH), pleckstrin homology (PH), helical PH-Rem linker (orange), Ras exchanger motif (Rem), Cdc25 and polyproline domains are shown along with previously described Noonan syndrome-associated mutations (black, top of the figure) and the novel missense changes identified in this study (red, bottom). Asterisks mark those variants also detected in one of the healthy parents. (B) Partial amino acid sequence alignments of human SOS1 with its orthologues of different species and human SOS2. Amino acids surrounding the four novel altered amino acids (indicated in red) are shown.

substitutions at residues Q477 and P478, using the known three-dimensional structures of SOS1.<sup>14 15 19 20</sup> The invariant residue F78 is located in the N-terminal histone folds domain. This residue is predicted to interact indirectly with R552, the most commonly mutated residue, and substitutions of F78 may

therefore have similar effects to those of mutations at R552. Q477 and P478 are not located in one of the known mutation clusters. We were unable to predict the precise consequences of substitutions of these residues based on structural modelling. Most likely, they do not directly affect the autoinhibited


**Table 1** Missense variants of *SOS1*

Exon*	Nucleotide change	Amino acid change	Domain	Pathogenic change or polymorphism	Number of affected people
3	c.233T→G	<b>F78C</b>	HF	Possible polymorphism	1 sporadic case†
6	c.806T→G	M269R	DH	Pathogenic	1 sporadic case
6	<b>c.806T→C</b>	<b>M269T</b>	DH	Pathogenic	1 sporadic case
7	c.925G→T	D309Y	DH	Pathogenic	1 sporadic case
10	c.1294T→C	W432R	PH	Pathogenic	1 sporadic case
10	c.1297G→A	E433K	PH	Pathogenic	1 sporadic case
10	<b>c.1300G→A‡</b>	<b>G434R</b>	PH	Pathogenic	1 sporadic case
10	c.1322G→A	C441Y	PH	Pathogenic	1 sporadic case
10	<b>c.[1431G→T; 1433C→T]§</b>	<b>[Q477H; P478L]</b>	PH	Pathogenic	1 sporadic case
10	<b>c.1433C→G</b>	<b>P478R</b>	PH	Pathogenic	1 sporadic case
10	c.1654A→G	R552G	PH-Rem linker	Pathogenic	2 sporadic cases, 1 familial observation¶
10	c.1655G→A	R552K	PH-Rem linker	Pathogenic	1 sporadic case, 1 familial observation¶
10	c.1656G→T	R552S	PH-Rem linker	Pathogenic	1 sporadic case, 1 familial observation¶
11	<b>c.1867T→A</b>	<b>F623I</b>	Rem	Pathogenic	1 sporadic case
12	c.1964C→T	P655L	Rem	Polymorphism	4 sporadic cases
13	c.2104T→C	Y702H	Rem	Pathogenic	1 sporadic case
16	c.2536G→A	E846K	Cdc25	Pathogenic	4 sporadic cases
19	<b>c.2999G→A</b>	<b>S1000N</b>	Cdc25	Probable polymorphism	1 sporadic case**

HF, histone-like folds; DH, Dbl homology domain; PH, pleckstrin homology domain; Rem, Ras exchanger motif. Novel variants are printed in bold type.

\*Exon 1 refers to the exon containing the ATG starting codon; †unaffected mother carries the same variant; ‡novel nucleotide exchange predicting a known missense mutation on protein level; §both sequence changes occurred de novo on the same allele; ¶affected mother-child duo; \*\*unaffected father carries the same variant.





**Figure 2** Facial appearance of 13 representative patients with Noonan syndrome (NS) carrying *SOS1* mutations. The respective missense mutation is indicated below and the clinical assignment above each photograph. The female patient classified as having “mild NS” presented only with short stature and mild facial anomalies. Ichthyosiform skin changes in a patient carrying the mutation E846K are illustrated in the lower right image. Parental/guardian informed consent was obtained for publication of this figure.

conformation. The highly conserved F623 residue is considered to play an important role in stabilising the Rem domain by forming an aromatic interaction with F958. Interestingly, substitution of F623 by glutamate in a *SOS1* mutant protein composed of the Rem-Cdc25 module has been shown to decrease exchange activity.<sup>24</sup> Substitution of F623 by isoleucine presumably also abolishes aromatic interaction between F958 and F623 and may therefore lead to a reduced guanine nucleotide exchange activity of *SOS1* (see supplementary information for further details; available online at <http://jmg.bmj.com/supplemental>).

## DISCUSSION

By screening genes encoding proteins involved in the Ras-MAPK signalling pathway we have independently discovered *SOS1* mutations in patients with classic features of NS (group

1A) and people with NS with more florid ectodermal symptoms and signs (group 1C). Taking these two subgroups of our core study population (group 1) together, our mutation detection rate reaches 28% among clinically well-characterised patients who lacked mutations in the previously reported NS and CFCS genes. This proportion exceeds those in previously published cohorts (Tartaglia *et al* 17%; Roberts *et al* 21%).<sup>11 12</sup> In aggregate, the two published studies and our study establish *SOS1* as the second major gene for NS. A *SOS1* mutation detection rate of 5% in group 2 corresponds to the low *PTPN11* mutation rate of 18% in this heterogeneous cohort. It is evident that established clinical criteria for NS are valid for patients with *SOS1* mutations, as only one patient with an *SOS1* mutation did not meet strict diagnostic criteria. Thus, our findings do not support *SOS1* as an important gene for people presenting with a mild or even atypical NS phenotype.

**Table 2** Clinical manifestations in patients with Noonan syndrome with *SOS1* and *PTPN11* mutations

	Patients with:				
	<i>SOS1</i> mutations				<i>PTPN11</i> mutations
	Present study	Tartaglia <i>et al.</i> <sup>12</sup>	Roberts <i>et al.</i> <sup>11</sup>	Total	Combined data†
Pulmonic stenosis	20/25	10/16	11/15	41/56 (73%)	125/171 (73%)
Hypertrophic cardiomyopathy	1/25	2/16	3/15	6/56 (11%)	10/171 (6%)
Atrial septal defect	4/25	4/16	2/15	10/56 (18%)	41/170 (24%)
Thorax deformity	17/24	16/16	ND	33/40 (83%)*	110/171 (64%)
Easy bruising	3/25*	5/16	5/15	13/56 (23%)*	46/90 (51%)‡
Mental retardation/need for special education	5/24*	1/16	3/11	9/51 (18%)*	71/164 (43%)
Short stature (<3rd centile)	13/25	2/15	4/13	19/53 (36%)*	122/171 (71%)
Cryptorchidism	5/11*	ND	5/10	10/21 (48%)*	75/94 (80%)

ND, no data.

\* $p < 0.05$  (Fisher's exact test); †combined data from four large studies: Tartaglia *et al.*,<sup>21</sup> Musante *et al.*,<sup>22</sup> Zenker *et al.*,<sup>17</sup> and Jongmans *et al.*<sup>23</sup>; ‡data available only from two of four studies.

## Key points

- We identified mutations of *SOS1* in 28% of patients with NS who had previously tested negative for mutations in *PTPN11* and *KRAS*, but not in patients with cardio-facio-cutaneous syndrome.
- Five novel mutations were detected, defining additional mutation hotspots and indicating possible new mechanisms of perturbed *SOS1* protein function.
- People with *SOS1* mutations commonly show ocular ptosis, curly hair and hyperkeratotic skin, but they seem to have a lower frequency of mental retardation than patients with Noonan syndrome and mutations of *PTPN11*.

People with *SOS1* mutations typically display a distinctive form of NS characterised by frequent ptosis, ectodermal symptoms and generally normal intelligence, an observation that has also been made by Tartaglia *et al.*<sup>12</sup> However, the clinical spectrum is broad (as shown in fig 2) making it difficult to predict the genetic defect of a person with NS based on clinical appearance, thus we will not change our diagnostic strategy of first analysing *PTPN11* in any patient with NS. In contrast to previous reports, we did not find a significantly lower prevalence of short stature in subjects with *SOS1* mutations compared with those with *PTPN11* mutations. Differences in patient selection may account for this apparent discrepancy, as several of our patients were recruited from a paediatric endocrinology department where they were seen for growth retardation.

One important finding of this study is the absence of *SOS1* mutations in patients with the diagnosis of CFCS, who had tested negative for mutations in the established CFCS genes. Although the ectodermal manifestations in patients with *SOS1* mutations are common and may be similar to those typically found in CFCS,<sup>12</sup> the *SOS1* phenotype in our study cohort is distinct from CFCS. The most important discriminating feature is the absence of significant mental retardation in patients with *SOS1* mutations. Nonetheless, patients with *SOS1* mutations may meet published criteria for CFCS.<sup>18</sup> This brings into question the usefulness of the "CFC index" and emphasises the significance of intellectual functioning level in discriminating between NS and CFCS. In young children, however, clinical discrimination between these two conditions, particularly NS caused by *SOS1* mutations, is difficult and may even be impossible. In such cases, determination of the molecular defect may have important prognostic implications.

*SOS1* mutations associated with NS have been shown to result in a gain of function,<sup>11–12</sup> consistent with the concept that NS and related disorders are caused by hyperactive Ras and aberrant activation of the B-Raf-MEK-ERK pathway.<sup>25</sup> However, to date, few *SOS1* mutant proteins have been assessed functionally. This study provides the first hint that NS-associated *SOS1* mutations may not uniformly lead to increased GEF activity, but instead may cause a decrease in exchange activity of *SOS1*, as already suggested by Shannon and Bollag.<sup>26</sup> The precise consequences of the F623I mutation remain to be determined by experimental means.

Taken together, our findings reinforce the concept that NS and CFCS scarcely overlap genotypically.<sup>27</sup> The pathogenetic basis of this obvious genotype phenotype correlation remains to be elucidated. It has been speculated that the influence of Ras effector pathways other than B-Raf-MEK-ERK or negative feedback mechanisms differentially affecting proximal and

## Electronic database information

- Online Mendelian Inheritance in Man (OMIM) (for NS, CFCS and CS). <http://www.ncbi.nlm.nih.gov/Omim/>
- Gene, for *SOS1* genomic (accession number NC\_000002.10) and cDNA (accession number NM\_005633.2) sequences. <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>.
- Protein Data Bank, for coordinates of DH-PH-Rem-Cdc25 construct of h*SOS1* (molecule B of code 1XD4), ternary Ras-*SOS1*-Ras-GppNHp complex (1NVW) and Ras-*SOS1* (1BKD). <http://www.rcsb.org/pdb/home/home.do>

distal components may account for the phenotypic differences in patients with NS and CFCS.<sup>11–25</sup> These emerging correlations between phenotypes and genotypes in this group of similar disorders is useful in defining new candidate genes for patients with NS or CFCS whose underlying genetic defect has yet to be discovered.

## ACKNOWLEDGEMENTS

We are grateful to the patients and families who participated in this study. We thank Angelika Diem and Inka Jantke for help with mutation screening. This work was supported by a grant from the German Research Foundation (DFG) to MZ.



Supplementary material is available online at <http://img.bmj.com/supplemental>.

## Authors' affiliations

**Martin Zenker, Anita Rauch, Christiane Zweier, Susanne Bauhuber, Verena Matejas**, Institute of Human Genetics, University Hospital Erlangen, Friedrich-Alexander University Erlangen-Nuremberg, Germany  
**Denise Horn, Stefan Mundlos**, Institute of Medical Genetics, Charité, University Medicine of Berlin, Berlin, Germany  
**Dagmar Wiczorek**, Institut für Humangenetik, Universität Duisburg-Essen, Essen, Germany  
**Judith Allanson**, Children's Hospital of Eastern Ontario, Ottawa, Ontario, Canada  
**Silke Pauli, Lars-Erik Wehner**, Institute of Human Genetics, University of Goettingen, Goettingen, Germany  
**Ineke van der Burgt**, Department of Human Genetics, University Medical Center St Radboud, Nijmegen, The Netherlands  
**Helmuth-Guenther Doerr**, Department of Pediatric Endocrinology, University Children's Hospital, Erlangen, Germany  
**Harald Gaspar**, Institute of Medical Genetics, University of Zurich, Zurich, Switzerland  
**Michael Hoffbeck**, University Children's Hospital, Pediatric Cardiology, Tuebingen, Germany  
**Gabriele Gillesen-Kaesbach**, Institut für Humangenetik, Universität zu Lübeck, Lübeck, Germany  
**Andreas Koch**, Department of Pediatric Cardiology, University Children's Hospital, Erlangen, Germany  
**Peter Meinecke**, Medizinische Genetik, Altonaer Kinderkrankenhaus, Hamburg, Germany  
**Anja Nowka, Kerstin Kutsche**, Institut für Humangenetik, Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany  
**Silke Reif**, Institut für Humangenetik und Medizinische Biologie, Universität Halle, Germany  
**Christian von Schnakenburg, Christian P Kratz**, Department of Pediatrics and Adolescent Medicine, University of Freiburg, Freiburg, Germany  
**Heide Seidel**, Institute of Human Genetics, Ludwig-Maximilian University, Munich, Germany

**Christoph Thomas**, Max Planck Institute of Molecular Physiology, Department of Structural Biology, Dortmund, Germany

Competing interests: None declared.

Parental/guardian informed consent was obtained for publication of figure 2.

Correspondence to: Martin Zenker, MD, Institute of Human Genetics, University of Erlangen-Nuremberg, Schwabachanlage 10, 91054 Erlangen, Germany; mzenker@humgenet.uni-erlangen.de

Received 2 May 2007

Revised 5 June 2007

Accepted 5 June 2007

Published Online First 23 June 2007

## REFERENCES

- Tartaglia M, Mehler EL, Goldberg R, Zampino G, Brunner HG, Kremer H, van der Burgt I, Crosby AH, Ion A, Jeffery S, Kalidas K, Patton MA, Kucherlapati RS, Gelb BD. Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nat Genet* 2001;**29**:465–8.
- Aoki Y, Niihori T, Kawame H, Kurosawa K, Ohashi H, Tanaka Y, Filocamo M, Kato K, Suzuki Y, Kure S, Matsubara Y. Germline mutations in HRAS proto-oncogene cause Costello syndrome. *Nat Genet* 2005;**37**:1038–40.
- Niihori T, Aoki Y, Narumi Y, Neri G, Cave H, Verloes A, Okamoto N, Hennekam RC, Gillissen-Kaesbach G, Wiczorek D, Kavamura MI, Kurosawa K, Ohashi H, Wilson L, Heron D, Bonneau D, Corona G, Kaname T, Naritomi K, Baumann C, Matsumoto N, Kato K, Kure S, Matsubara Y. Germline KRAS and BRAF mutations in cardio-facio-cutaneous syndrome. *Nat Genet* 2006;**38**:294–6.
- Rodriguez-Viciana P, Tetsu O, Tidyman WE, Estep AL, Conger BA, Cruz MS, McCormick F, Rauen KA. Germline mutations in genes within the MAPK pathway cause cardio-facio-cutaneous syndrome. *Science* 2006;**311**:1287–90.
- Schubbert S, Zenker M, Rowe SL, Boll S, Klein C, Bollag G, van der Burgt I, Musante L, Kalscheuer V, Wehner LE, Nguyen H, West B, Zhang KY, Sistermans E, Rauch A, Niemeyer CM, Shannon K, Kratz CP. Germline KRAS mutations cause Noonan syndrome. *Nat Genet* 2006;**38**:331–6.
- Allanson JE. Noonan syndrome. *J Med Genet* 1987;**24**:9–13.
- Hennekam RC. Costello syndrome: an overview. *Am J Med Genet C Semin Med Genet* 2003;**117**:42–8.
- Roberts A, Allanson J, Jadico SK, Kavamura MI, Noonan J, Opitz JM, Young T, Neri G. The cardio-facio-cutaneous (CFC) syndrome: a review. *J Med Genet* 2006.
- Carta C, Pantaleoni F, Bocchinfuso G, Stella L, Vasta I, Sarkozy A, Digilio C, Palleschi A, Pizzuti A, Grammatico P, Zampino G, Dallapiccola B, Gelb BD, Tartaglia M. Germline missense mutations affecting KRAS Isoform B are associated with a severe Noonan syndrome phenotype. *Am J Hum Genet* 2006;**79**:129–35.
- Zenker M, Lehmann K, Schulz AL, Barth H, Hansmann D, Koenig R, Korinthenberg R, Kreiss-Nachtsheim M, Meinecke P, Morlot S, Mundlos S, Quante AS, Raskin S, Schnabel D, Wehner LE, Kratz CP, Horn D, Kutsche K. Expansion of the genotypic and phenotypic spectrum in patients with KRAS germline mutations. *J Med Genet* 2007;**44**:131–5.
- Roberts AE, Araki T, Swanson KD, Montgomery KT, Schiripo TA, Joshi VA, Li L, Yassin Y, Tamburino AM, Neel BG, Kucherlapati RS. Germline gain-of-function mutations in SOS1 cause Noonan syndrome. *Nat Genet* 2007;**39**:70–4.
- Tartaglia M, Pennacchio LA, Zhao C, Yadav KK, Fodale V, Sarkozy A, Pandit B, Oishi K, Martinelli S, Schackwitz W, Ustaszewska A, Martin J, Bristow J, Carta C, Lepri F, Neri C, Vasta I, Gibson K, Curry CJ, Siguero JP, Digilio MC, Zampino G, Dallapiccola B, Bar-Sagi D, Gelb BD. Gain-of-function SOS1 mutations cause a distinctive form of Noonan syndrome. *Nat Genet* 2007;**39**:75–9.
- Corbalan-Garcia S, Margarit SM, Galron D, Yang SS, Bar-Sagi D. Regulation of Sos activity by intramolecular interactions. *Mol Cell Biol* 1998;**18**:880–6.
- Margarit SM, Sondermann H, Hall BE, Nagar B, Hoelz A, Pirruccello M, Bar-Sagi D, Kuriyan J. Structural evidence for feedback activation by Ras.GTP of the Ras-specific nucleotide exchange factor SOS. *Cell* 2003;**112**:685–95.
- Sondermann H, Soisson SM, Boykevich S, Yang SS, Bar-Sagi D, Kuriyan J. Structural analysis of autoinhibition in the Ras activator Son of sevenless. *Cell* 2004;**119**:393–405.
- van der Burgt I, Berends E, Lommen E, van Beersum S, Hamel B, Mariman E. Clinical and molecular studies in a large Dutch family with Noonan syndrome. *Am J Med Genet* 1994;**53**:187–91.
- Zenker M, Buheitel G, Rauch R, Koenig R, Bosse K, Kress W, Tietze HU, Doerr HG, Hofbeck M, Singer H, Reis A, Rauch A. Genotype-phenotype correlations in Noonan syndrome. *J Pediatr* 2004;**144**:368–74.
- Kavamura MI, Peres CA, Alchorne MM, Brunoni D. CFC index for the diagnosis of cardiofaciocutaneous syndrome. *Am J Med Genet* 2002;**112**:12–16.
- Boriack-Sjodin PA, Margarit SM, Bar-Sagi D, Kuriyan J. The structural basis of the activation of Ras by Sos. *Nature* 1998;**394**:337–43.
- Sondermann H, Nagar B, Bar-Sagi D, Kuriyan J. Computational docking and solution x-ray scattering predict a membrane-interacting role for the histone domain of the Ras activator son of sevenless. *Proc Natl Acad Sci U S A* 2005;**102**:16632–7.
- Tartaglia M, Kalidas K, Shaw A, Song X, Musat DL, van der Burgt I, Brunner HG, Bertola DR, Crosby A, Ion A, Kucherlapati RS, Jeffery S, Patton MA, Gelb BD. PTPN11 mutations in Noonan syndrome: molecular spectrum, genotype-phenotype correlation and phenotypic heterogeneity. *Am J Hum Genet* 2002;**70**:1555–63.
- Musante L, Kehl HG, Majewski F, Meinecke P, Schweiger S, Gillissen-Kaesbach G, Wiczorek D, Hinkel GK, Tinschert S, Hoeltzenbein M, Ropers HH, Kalscheuer VM. Spectrum of mutations in PTPN11 and genotype-phenotype correlation in 96 patients with Noonan syndrome and five patients with cardio-facio-cutaneous syndrome. *Eur J Hum Genet* 2003;**11**:201–6.
- Jongmans M, Sistermans EA, Rikken A, Nillesen WM, Tamminga R, Patton M, Maier EM, Tartaglia M, Noordam K, van der Burgt I. Genotypic and phenotypic characterization of Noonan syndrome: new data and review of the literature. *Am J Med Genet A* 2005;**134**:165–70.
- Hall BE, Yang SS, Boriack-Sjodin PA, Kuriyan J, Bar-Sagi D. Structure-based mutagenesis reveals distinct functions for Ras switch 1 and switch 2 in Sos-catalyzed guanine nucleotide exchange. *J Biol Chem* 2001;**276**:27629–37.
- Bentires-Alj M, Kontaridis MI, Neel BG. Stops along the RAS pathway in human genetic disease. *Nat Med* 2006;**12**:283–5.
- Shannon K, Bollag G. Sending out an SOS. *Nat Genet* 2007;**39**:8–9.
- Kratz CP, Schubbert S, Bollag G, Niemeyer CM, Shannon KM, Zenker M. Germline mutations in components of the ras signaling pathway in Noonan syndrome and related disorders. *Cell Cycle* 2006;**5**:1607–11.