

cause this technology is available in relatively few clinical laboratories, cases of infection with *M. massiliense* may be mistakenly attributed to *M. abscessus*. Although infections with *M. massiliense* may be underrecognized, reports of these infections are raising concern. The capacity of this bacteria to infect different body sites is further evidence for the pathogenic potential of a rapidly growing mycobacteria in human infections (10).

**Enrico Tortoli,* Rita Gabini,†
Irene Galanti,†
and Alessandro Mariottini***

*Careggi Hospital, Florence, Italy; and †San Donato Hospital, Arezzo, Italy

References

1. Böttger H, Rogall T, Flohr T, Blöcker H, Böttger EC. Detection and identification of mycobacteria by amplification of rRNA. *J Clin Microbiol.* 1990;28:1751–9.
2. Adékambi T, Colson P, Drancourt M. *rpoB*-based identification of nonpigmented and late-pigmenting rapidly growing mycobacteria. *J Clin Microbiol.* 2003;41:5699–708.
3. Delaney V, Sumrani N, Hong JH, Sommer B. Mycobacterial infections in renal allograft recipients. *Transplant Proc.* 1993;25:2288–9.
4. Branger B, Gouby A, Oulès R, Barducci JP, Mourad G, Fourcade J, et al. *Mycobacterium haemophilum* and *Mycobacterium xenopi* associated infection in a renal transplant patient. *Clin Nephrol.* 1985;23:46–9.
5. Adékambi T, Reynaud-Gaubert M, Greub G, Gevaudan MJ, La Scola B, Raoult D, et al. Amoebal coculture of “*Mycobacterium massiliense*” sp. nov. from the sputum of a patient with hemoptoic pneumonia. *J Clin Microbiol.* 2004;42:5493–501.
6. Simmon KE, Pounder JI, Greene JN, Walsh F, Anderson CM, Cohen S, et al. Identification of an emerging pathogen, *Mycobacterium massiliense*, by *rpoB* sequencing of clinical isolates collected in the United States. *J Clin Microbiol.* 2007;45:1978–80.
7. Kim HY, Yun YJ, Park CG, Lee BH, Cho YK, Park BJ, et al. Outbreak of *Mycobacterium massiliense* infection associated with intramuscular injections. *J Clin Microbiol.* 2007;45:3127–30.
8. Viana-Niero C, Lima KV, Lopes ML, da Silva Rabello MC, Marsola LR, Brilhante VC, et al. Molecular characteriza-

tion of *Mycobacterium massiliense* and *Mycobacterium bolletii* in outbreaks of infections after laparoscopic surgeries and cosmetic procedures. *J Clin Microbiol.* 2008;46:850–5.

9. Clinical and Laboratory Standards Institute (CLSI). Susceptibility testing of *Mycobacteria*, *Nocardia*, and other aerobic actinomycetes: approved standard M24-A. Wayne (PA): The Institute; 2003.
10. Brown-Elliott BA, Wallace RJ Jr. Clinical and taxonomic status of pathogenic nonpigmented or late-pigmenting rapidly growing mycobacteria. *Clin Microbiol Rev.* 2002;15:716–46.

Address for correspondence: Enrico Tortoli, Regional Reference Center for Mycobacteria, Microbiology Laboratory, Careggi Hospital, Viale Morgagni 85, 50134 Florence, Italy; email: enrico.tortoli@unifi.it

Bovine Kobuviruses from Cattle with Diarrhea

To the Editor: A new species of kobuvirus, named U-1 strain, was first recognized in 2003 as a cytopathic contaminant in a culture medium of HeLa cells that had been used for >30 years in the laboratory (1). The RNA genome of the U-1 strain comprises 8,374 nucleotides; the genome organization is analogous to that of picornaviruses. Morphologically, the U-1 strain resembles the Aichi virus, but genetically it is distinct (1). Therefore, the U-1 strain is classified as a new species of genus *Kobuvirus* in the family *Picornaviridae*, and it is called *bovine kobuvirus* (1). To date, the genus *Kobuvirus* consists of 2 species, *Aichi virus* and *bovine kobuvirus* (2). The Aichi virus is associated with acute gastroenteritis in humans (3–5); bovine kobuvirus infection has been detected only in cattle (1).

Only 1 report has described the discovery and epidemiologic features

of bovine kobuvirus (1). Of serum samples from 72 healthy cattle, 43 (59.7%) were positive for neutralizing antibody against bovine kobuvirus U-1 standard strain at a titer of ≥ 16 . In addition, 12 (16.7%) of 72 stool samples collected from the cattle were positive for the bovine kobuvirus genome by reverse transcription-PCR (RT-PCR) (1). This finding suggested that bovine kobuvirus is common and that the virus particles could be detected in the stool samples of infected cattle. We therefore conducted an epidemiologic survey of bovine kobuvirus and report detection of this virus in stool samples from calves with diarrhea during 2001–2004 in Chiang Mai Province, Thailand.

From November 2001 to July 2004, a total of 72 fecal specimens were collected. The age of the calves ranged from 7 to 49 days. The presence of bovine kobuvirus in fecal specimens was detected by using RT-PCR with a protocol modified from the method described by Yamashita et al. (1). All the bovine kobuvirus strains detected in our study were analyzed further by direct sequencing of their PCR amplicons with the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems, Foster City, CA, USA) on an automated sequencer (ABI 3100, Applied Biosystems). The nucleotide sequences of these portions were compared with those of reference strains available in the GenBank database by using BLAST (6). Phylogenetic and molecular evolutionary analyses were conducted by using MEGA version 3.1 (7). The nucleotide sequences of bovine kobuvirus strains described in this study were deposited in GenBank under accession nos. EF659450–EF659455.

The bovine kobuvirus was detected by the RT-PCR screening method in 6 (8.3%) of the 72 fecal specimens collected. The partial 3D regions of all 6 bovine kobuviruses exhibited highly conserved sequences of 99.3%–100% nucleotide and 100% amino acid




Figure. Phylogenetic analysis of the partial nucleotide sequence encoding the 3D region of bovine kobuviruses isolated in this study and other reference strains recognized to date. The tree was generated on the basis of the neighbor-joining method using the MEGA 3.1 program (7). Scale bar indicates branch length for a 5% nucleotide difference.

identities to each other. In addition, by searching for the closest sequences in the databank, we found that these sequences were closely related to all 13 bovine kobuvirus reference strains available in the GenBank database (U-1, K-35, K-36, K-49, K-38, N-5, K-3, K-4, K-6, K-60, N-2, K-44, and K-55); nucleotide and amino acid sequence identities ranged from 91.0% to 95.0% and 97.4% to 99.4%, respectively. Phylogenetic analysis of partial 3D nucleotide sequences of our bovine kobuvirus strains, together with those of all published bovine kobuvirus reference strains available in the GenBank database, is shown in the Figure. The Aichi virus standard strain was included in the tree as an outlier virus. The phylogenetic tree confirmed that all 6 virus strains belonged to bovine kobuvirus and formed a tight cluster in a monophyletic branch with other published bovine kobuvirus reference strains, but they were distantly related to the Aichi virus standard strain.

Detection and characterization of bovine kobuvirus strains from different geographic areas are important for understanding the worldwide distribu-

tion, heterogeneity, and association of bovine kobuvirus with enteric disease in cattle. Our findings indicate the role of kobuviruses in diarrheal disease in cattle and provide additional information on their relationship to bovine kobuviruses reported previously.

This research was supported by the Core University System Exchange Program under the Japan Society for the Promotion of Science, coordinated by the Graduate School of Medicine, the University of Tokyo, Japan, and Mahidol University, Thailand. The study was also supported in part by the Research Fund from the Faculty of Medicine, Chiang Mai University, Thailand.

**Pattara Khamrin,*†
Niwat Maneekarn,‡
Supatra Peerakome,‡
Shoko Okitsu,†
Masashi Mizuguchi,*
and Hiroshi Ushijima*†**

*University of Tokyo, Tokyo, Japan; †Aino Health Science Center, Tokyo; and ‡Chiang Mai University, Chiang Mai, Thailand

References

1. Yamashita T, Ito M, Kabashima Y, Tsuzuki H, Fujiura A, Sakae K. Isolation and characterization of a new species of kobuvirus associated with cattle. *J Gen Virol*. 2003;84:3069–77.
2. Racaniello VR. Picornaviridae: the viruses and their replication. In: Knipe DM and Howley PM, editors. *Fields virology*, vol. 1. 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2006. p. 795–838.
3. Oh DY, Silva PA, Hauroeder B, Diedrich S, Cardoso DD, Schreier E. Molecular characterization of the first Aichi viruses isolated in Europe and in South America. *Arch Virol*. 2006;151:1199–206.
4. Pham NT, Khamrin P, Nguyen TA, Kanti DS, Phan TG, Okitsu S, et al. Isolation and molecular characterization of Aichi viruses from fecal specimens collected in Japan, Bangladesh, Thailand, and Vietnam. *J Clin Microbiol*. 2007;45:2287–8.
5. Yamashita T, Sakae K, Kobayashi S, Ishihara Y, Miyake T, Mubina A, et al. Isolation of cytopathic small round virus (Aichi virus) from Pakistani children and Japanese travelers from Southeast Asia. *Microbiol Immunol*. 1995;39:433–5.
6. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215:403–10.
7. Kumar S, Tamura K, Nei M. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform*. 2004;5:150–63.

Address for correspondence: Hiroshi Ushijima, Department of Developmental Medical Sciences, Institute of International Health, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan; email: ushijima-hiroshi@jcom.home.ne.jp

Letters

Letters commenting on recent articles as well as letters reporting cases, outbreaks, or original research are welcome. Letters commenting on articles should contain no more than 300 words and 5 references; they are more likely to be published if submitted within 4 weeks of the original article's publication. Letters reporting cases, outbreaks, or original research should contain no more than 800 words and 10 references. They may have one Figure or Table and should not be divided into sections. All letters should contain material not previously published and include a word count.