

Fungal endosymbionts of plants reduce lifespan of an aphid secondary parasitoid and influence host selection

Simone A. Härri^{1,*}, Jochen Krauss^{1,2} and Christine B. Müller^{1,†}

¹*Institute of Environmental Sciences, University of Zürich, Winterthurerstrasse 190, 8057 Zürich, Switzerland*

²*Department of Animal Ecology I, Population Ecology, University of Bayreuth, Universitätsstrasse 30, 95447 Bayreuth, Germany*

Complex biotic interactions shape ecological communities of plants, herbivores and their natural enemies. In studies of multi-trophic interactions, the presence of small, invisible micro-organisms associated with plants and those of a fourth above-ground trophic level have often been neglected. Incorporating these neglected factors improves our understanding of the processes within a multi-trophic network. Here, we ask whether the presence of a fungal endosymbiont, which alters plant quality by producing herbivore-toxic substances, trickles up the food chain and affects the performance and host-selection behaviour of aphid secondary parasitoids. We simultaneously offered hosts from endophyte-free and endophyte-infected environments to secondary parasitoids. Older and more experienced parasitoid females discriminated against hosts from the endophyte-infected environment. Developing in lower quality hosts from the endophyte-infected environment reduced the lifespan of secondary parasitoids. This indicates that aphid secondary parasitoids can perceive the disadvantage for their developing offspring in parasitoids from the endophyte environment and can learn to discriminate against them. In the field, this discrimination ability may shift the success of primary parasitoids to endophyte-infected plants, which co-occur with endophyte-free plants. Ultimately, the control of aphids depends on complex interactions between primary and secondary parasitoids and their relative sensitivity to endophytic fungi.

Keywords: *Asaphes vulgaris*; bottom-up cascade; endophytic fungi; mummy parasitoids; *Neotyphodium lolii*; oviposition strategy

1. INTRODUCTION

The mediating effect of plants on the interaction between herbivores and their natural enemies plays a crucial role in structuring natural communities (Ohgushi *et al.* 2007). However, natural communities often consist of more than three trophic levels: plants; herbivores; and natural enemies (Pimm & Lawton 1977). For example, in insect communities, primary parasitoids are commonly attacked by a range of secondary parasitoids (Sullivan 1987). The activity of secondary parasitoids can limit the success of primary parasitoids and may disrupt the effectiveness of essential biological control (Rosenheim 1998). In aphid communities, secondary parasitoids are ubiquitous, diverse and may have a strong impact on interactions among plants, herbivores and primary parasitoids (Müller *et al.* 1999; Sullivan & Völkl 1999). Nevertheless, studies on multi-trophic interactions in insect communities often neglect the presence of this fourth trophic level (Brodeur 2000; but see Harvey *et al.* 2003; Soler *et al.* 2005).

Aside from ignoring consumers at the top of food webs, the presence of micro-organisms associating with plants at the bottom of food webs has also been frequently overlooked (Polis & Strong 1996). For example,

endophytic fungi (=endophytes) are endosymbionts of a large variety of plant species (Arnold & Lutzoni 2007). Endophytes of the genus *Neotyphodium* are associated with temperate grasses and produce herbivore-toxic alkaloids (Clay 1990). These alkaloids can change food plant quality for herbivores and the effects can cascade up the food chain and propagate to higher trophic levels (Omacini *et al.* 2001; Müller & Krauss 2005; de Sassi *et al.* 2006; Härri 2007). The potential of endophytic fungi to affect secondary parasitoids, however, has never been investigated experimentally. Secondary parasitoids are intimately linked to primary parasitoids with their larvae feeding directly on the pre-pupae and pupae of the primary parasitoid (Quicke 1997). In addition, adult females of many species also feed directly on the hosts (host feeding; Jarvis & Kidd 1986; Jarvis *et al.* 2008). It is therefore likely that the presence of endophytes affects the behaviour and the performance of secondary parasitoids.

Spatial heterogeneity in plant quality can affect multi-trophic interactions (van Nouhuys & Hanski 2002). Spatial mosaics in endophyte infection in European grasslands (Saikkonen *et al.* 2000; Zabalgoceazcoa *et al.* 2003) leave the secondary parasitoids with the choice of hosts from either endophyte-free or endophyte-infected grasses. Such a choice confronts the parasitoids with a sequence of decisions when encountering a host: should the host be accepted for oviposition, ignored or used as

* Author and address for correspondence: Department of Environmental Biology, University of Guelph, Guelph, Ontario N1G 2W1, Canada (shaerri@uoguelph.ca).

† Deceased 7 March 2008. We dedicate this paper to Christine, our co-author and dear friend, who died of cancer. She is sadly missed.

protein-rich food? Host choice by parasitoids has been studied experimentally and theoretically, with the majority of the experimental studies focusing on host choice by primary parasitoids, whereas theoretical models often ignore important differences between primary and secondary parasitoids (Godfray 1994). If eggs were cheap and not limited, and the time required for host attack minimal, parasitoids would be predicted to attack all hosts they encounter. However, as many secondary parasitoids are either time and/or egg limited, oviposition becomes more costly and a precise host choice is important (Heimpel *et al.* 1996). Most aphid secondary parasitoids are synovigenic, which means that they continue to mature eggs during the adult stage (Quicke 1997). For many synovigenic species, the process of egg maturation requires proteins that are obtained by host feeding (Harvey 2008) and slows down with progressing age (Godfray 1994). Synovigenic parasitoids can therefore become egg limited either temporarily when deprived of hosts or after encountering a large number of hosts within a short time period or permanently with increasing age (Rosenheim *et al.* 2000). With progressive, age-dependent egg limitation, females should preferentially oviposit into the highest quality hosts (Iwasa *et al.* 1984) and use lower quality hosts for host feeding (Kidd & Jervis 1991). The ability to distinguish between hosts of different qualities may depend on the parasitoid's experience (Vet *et al.* 1990). Older and more experienced females are predicted to make more precise host choice decisions than younger ones. If host choice by secondary parasitoids indeed occurs, it may define the impact of the endophyte on secondary parasitoids and the structure of the associated aphid–parasitoid community.

Here, we investigate the cascading effect of primary parasitoid hosts of the endophyte-infected and endophyte-free environments on secondary parasitoid choices. We assumed that primary parasitoids that develop in aphids from endophyte-infected grasses are of reduced quality and predicted that this reduced host quality should influence the host choice behaviour and offspring performance of aphid secondary parasitoids, by (i) increasing larval development time, (ii) reducing adult lifespan, (iii) resulting in a male-biased sex ratio as theory predicts that fertilized eggs (=females) are laid preferentially into high-quality hosts whereas unfertilized eggs (=males) are laid into low-quality hosts (Charnov *et al.* 1981) and (iv) reducing the weight of adults that had developed in such low-quality hosts. We further tested whether females are able to distinguish between mummies from endophyte-free (E–) and endophyte-infected (E+) environments, depending on age and experience, predicting that (v) older females should be more selective, (vi) host experience would lead to increased choice of E– mummies for oviposition and (vii) the lower quality hosts (E+ mummies) would preferentially be used for host feeding.

2. MATERIAL AND METHODS

(a) Model system

The system used to address our hypotheses consisted of the endophytic fungi *Neotyphodium lolii* Glen, Bacon, Hanlin, a specialist on the grass *Lolium perenne* L. The aphid species *Metopolophium festucae* Theobald served as hosts for the primary parasitoids *Aphidius ervi* Haliday whose mummies

served as hosts for the secondary parasitoids *Asaphes vulgaris* Wlk. The aphid species *M. festucae* is relatively insensitive to the presence of *N. lolii*, whereas the fecundity of *A. ervi* is reduced when developing within *M. festucae* feeding on *L. perenne* infected by *N. lolii* (Härrı 2007).

Seeds of *L. perenne* were provided by Brian Tapper, AgResearch, New Zealand. The seeds were either infected with *N. lolii* (wild-type; E+) or endophyte free (E–). All of the seeds belonged to the agricultural cultivar Grassland Samson. Endophyte infection was lost by selectively choosing plants with unsuccessful endophyte transmission (B. Tapper 2004, personal communication). After the experiment, the endophyte infection status of the plants was verified with the Phytoscreen *Neotyphodium* immunoblot assays (Agrinostics Ltd., Watkinsville, GA, USA). Infection level of E– plants was 3 per cent, whereas 80 per cent of the E+ plants were infected.

The stock culture of *M. festucae* was started in summer 2005 with a few individuals collected close to the University of Zurich, Switzerland. Since then, the culture was kept on *L. perenne* Arion (commercially available endophyte-free fodder grass). The stock culture of *A. ervi* was started with 250 individuals from Andermatt Biocontrol AG, Switzerland, in Spring 2006. *Aphidius ervi* was maintained on *M. festucae* feeding on *L. perenne* Arion. The stock culture of *A. vulgaris* was started with a few individuals obtained from the collected aphid mummies close to the University of Zurich, Switzerland, in Summer 2006. The stock culture was maintained on *A. ervi*.

Asaphes vulgaris (Pteromalidae, Chalcidoidea) is an ectoparasitic, solitary mummy parasitoid. Females oviposit on the surface of the pupae of the primary parasitoid and the hatched larva feeds on the primary parasitoid prepupa or pupa within the mummified aphid (Christiansen-Weniger 1992). When ovipositing or host feeding, females inject venom that kills the primary parasitoid pupae immediately (=idiobionts). Proteins gained from host feeding are allocated to the maturation of eggs. In the absence of hosts, the females can reallocate nutrients from the eggs into the somatic maintenance by resorption (=oosorption; Godfray 1994). The process of oosorption can lead to egg limitation in synovigenic species (Rosenheim *et al.* 2000).

(b) Experimental set-up

To obtain 1-day-old mummies from the E– and E+ environments, aphids from the stock culture were placed on either E– or E+ pots. After 10 days, *A. ervi* females were added, and a few days later, the freshly formed mummies were collected over four consecutive days. The explanatory factors were the presence of endophytes ('endophyte'), age of *A. vulgaris* females ('age') and *A. vulgaris* female experience ('experience'). In the endophyte choice treatment, 14 E– and 14 E+ mummies were offered simultaneously to one female *A. vulgaris* that had no previous experience with the E+ or E– environments. The choice experiments were conducted in Petri dishes and one Petri dish represented one replicate. The 28 mummies represented ad libitum number of hosts, as in all trials some primary parasitoids were emerging from the mummies. The mummies were glued to the Petri dish with a honey–water solution in a checkerboard pattern to ensure that individuals had the same probability of encountering E– and E+ mummies. To each Petri dish, one female and one male *A. vulgaris* were added. The mating partners

had the same age. The factor age was manipulated using mating pairs of different ages. From the nine mating pairs used in the experiment, the youngest was 5 days old and the oldest 14 days old with 1 day difference between each of the pairs (age class 8 was missing). Each of the mating pairs was kept without aphid hosts after emergence until they entered the experimental arena where they were left together with the 14 E− and 14 E+ mummies for 24 hours. The factor experience was obtained by offering each mating pair fresh mummies (14 E− and 14 E+) in a new Petri dish over the four consecutive days.

Three days after female and male *A. vulgaris* were removed from the Petri dishes, the mummies were put individually into gelatine capsules. The gelatine capsules were checked daily for the emergence of *A. vulgaris*. The time until emergence (=development time) was recorded. Emerged *A. vulgaris* were sexed and put singly into a plastic vial (5×2 cm) closed with a foam plug. A fresh piece of apple was provided as a sugar source every second day, which reduced the overall lifespan and workload compared with daily feedings (S. A. Härrä 2007, personal observation). The relative differences in lifespan between individuals developing with E− and E+ hosts are most likely not affected by the feeding regime. Time until death (=adult lifespan) was recorded in days. After death, each individual was weighed. The remaining mummies were dissected to detect host feeding or the death of the secondary parasitoid larvae. All stock cultures and experiments were conducted under controlled climatic conditions at 22°C with an 16 L : 8 D light regime.

(c) Statistical analyses

All statistical analyses were performed with R (v. 2.7.0 for Mac OS X). Values are represented as mean ± s.e. The choice between E− and E+ mummies was analysed with a generalized mixed-effects model (glmmPQL—function) with ‘experience’, ‘day’, ‘endophyte’ and all their interactions as fixed effects and ‘female identity’ as random effect using the quasi-Poisson error structure to account for overdispersion of the count data (Venables & Ripley 2002). To account for the repeated measurements over the four consecutive days, experience was nested within female identity for the random factor.

The life-history traits, development time, lifespan and weight, of the emerged *A. vulgaris* offspring were analysed with linear mixed-effects model (lme—function) with ‘age’, ‘experience’, ‘sex’, ‘endophyte’ and including all interactions with the exception of sex for which only the interaction with endophyte was included. Female identity was included as a random factor with experience nested within female identity to account for the repeated measurements. Differences in the sex of the emerged *A. vulgaris* were analysed using a generalized mixed-effects model (glmmPQL—function) including the same fixed and random effects as described above, but excluding sex and its interaction with endophyte. For the four life-history traits, *p*-values were adjusted using Bonferroni corrections.

For all analyses, experience and day were treated as continuous factors and the first-order autocorrelation structure (corAR1) was applied to account for the dependency of the repeated measures (Pinheiro & Bates 2000). For all linear mixed-effects models, the maximum-likelihood method was used for model fit and the general positive-definite symmetric variance-covariance structure for the random effects (Pinheiro & Bates 2000). Owing to the

Table 1. The influence of age, experience, endophyte and the interactions of interest on oviposition and host feeding, for the adult female generation that had a choice between mummies from E+ and E− environments. (For host feeding, all interactions were not significant and therefore not included in the analysis. The significant results are presented in bold letters. For the random factor ‘female identity’, the percentage of variation occurring between females is presented.)

	oviposition	host feeding
<i>single factors</i>		
age	$F_{1,7}=0.52$ $p=0.492$	$F_{1,7}=0.09$ $p=0.767$
experience	$F_{1,25}=38.14$ $p<0.001$	$F_{1,26}=0.90$ $p=0.351$
endophyte	$F_{1,32}=4.64$ $p=0.039$	$F_{1,35}=0.49$ $p=0.491$
<i>interactions</i>		
age×experience	$F_{1,25}=7.06$ $p=0.014$	—
age×endophyte	$F_{1,32}=6.16$ $p=0.019$	—
experience ×endophyte	$F_{1,32}=1.71$ $p=0.200$	—
age×experience ×endophyte	$F_{1,32}=8.50$ $p=0.006$	—
<i>random factor</i>		
female identity	14.96%	39.93%

non-orthogonality of the analyses, non-significant interaction terms were removed and only the main effects were tested. If the interactions were significant, only the highest order interaction would be interpreted.

3. RESULTS

Test statistics and *p*-values for all measured traits are summarized in table 1 and table 2.

Seven of the nine *A. vulgaris* females reproduced successfully showing a preference for E− mummies (figure 1a). This resulted in a mean number of offspring of 2.50 ± 0.58 from E− mummies and 1.85 ± 0.34 from E+ mummies. The preference for E− mummies was especially pronounced for the older parasitoids and increased with experience (age×experience×endophyte; figure 1a; table 1). Contrary to our expectations, host feeding was not significantly influenced by any of the explanatory factors (figure 1b; table 1). A few mummies from which no primary or mummy parasitoid emerged and that were not used for host feeding contained dead *A. ervi* larvae that did not finish their development (E− : 4 mummies; E+ : 8 mummies). In contrast to the primary parasitoid *A. ervi*, we did not observe larval and/or pupal mortality among *A. vulgaris*.

A total of 40 parasitoid offspring emerged from E− mummies and 25 from E+ mummies. The development time of *A. vulgaris* larvae from egg at oviposition to adult emergence was not influenced by the presence of endophytes but was shorter for males (19.98 ± 0.09 days) than females (21.33 ± 0.24 days; table 2). *Asaphes vulgaris* emerging from E+ had a significant shorter lifespan than those emerging from E− (figure 2).




Figure 1. The presence and the quantity of (a) oviposition events and (b) host-feeding events on E- (white bars) and E+ (grey bars). From left to right are the 4 days ('experience') and from top to down are the different ages of the females (excluding age class 8). Data show the increase in oviposition probability over the 4 days and an increase in oviposition probability with increasing age (table 1). Two females (aged 7 and 9 days at the beginning of the experiment) were not ovipositing despite showing host-feeding behaviour.

After the conservative Bonferroni correction, the reduction in lifespan is only marginally significant (table 2). Independent of endophyte infection, females lived significantly longer than males (females: 14.11 ± 2.73 days, males: 9.37 ± 0.59 days; table 2). The only factor that influenced the weight of the emerged adult wasps was their sex (females: 84.67 ± 4.32 μg , males: 65.50 ± 2.27 μg ; table 2). Contrary to our expectation, the sex ratio of the progeny was not influenced by any of the experimental factors (table 2).

4. DISCUSSION

The adult lifespan of a generalist secondary parasitoid at the top of an aphid-parasitoid food web tended to decrease when oviposition and larval development took place in hosts experiencing the endophyte environment. Female *A. vulgaris* improved their host choice with progressing age and oviposition experience through selection of more hosts from the endophyte-free environment compared with younger and less experienced females. Offspring performance in endophyte-free hosts was only improved in terms of lifespan and no effects on developmental time, sex ratio or weight were detected. *Asaphes vulgaris* parasitoids are long lived and synovigenic, maturing their eggs during adulthood. Under such conditions, a reduced lifespan will most likely result in a shorter reproductive time and thus reduced fitness. Similar fitness penalties of endophytes are known for predators and primary parasitoids (Bultman *et al.* 1997;




Figure 2. The mean \pm s.e. lifespan for *A. vulgaris* offspring emerging from E- mummies (open square) and E+ mummies (filled square). The mean was calculated from the average of each Petri dish. The statistics are presented in table 2 (second column).

de Sassi *et al.* 2006; Härrä 2007). Ours is the first study showing experimentally that even the top trophic level in this aphid-parasitoid food web can suffer from ubiquitous endophytic fungi in agricultural grasses. It is also one of very few studies showing that variation in plant quality—here caused by a microbial endophyte—affects individual performance of the fourth trophic level (but see Harvey *et al.* 2003; Soler *et al.* 2005).

In the field, where endophyte infection often occurs in a heterogeneous pattern (e.g. Saikkonen *et al.* 2000), a fitness reduction caused by the presence of endophytes depends on the ability of secondary parasitoids to discriminate against such inferior hosts. We show that more secondary parasitoids offspring emerged from E- mummies than E+ mummies when females were given a choice. This increase in the number of offspring resulted from an increased number of oviposition events on E- mummies and not from increased larval mortality within E+ mummies as we detected no mortality among secondary parasitoid larvae. Thus, our experiment demonstrated that secondary parasitoids were able to discriminate against hosts from the endophyte-infected environment and this discrimination improved over time and with experience.

The exact cues that females used when selecting hosts from the endophyte-free environment must be linked to the aphid mummy as no plants or living aphids were present in the choice arenas. Aphid secondary parasitoids may be attracted to the presence of aphid honeydew (Buitenhuis *et al.* 2004), but this was not present in the choice arenas either. Host acceptance of secondary parasitoids often involves careful, time-consuming examination of a mummy by tapping it and probing the host inside with the ovipositor (Vinson 1976). As mummy size does not differ between E- and E+ (Härrä 2007), the ovipositor that examines the host may perceive signals about endophyte-produced chemicals that have accumulated in the primary parasitoid. Alternatively, the mummified aphid skin could carry clues on past plant

Table 2. The influence of age, experience, sex and endophyte on the measured life-history parameters of *A. vulgaris*. (All the interactions were not significant and were therefore not included in the analyses. The *p*-values were adjusted for multiple testing using Bonferroni corrections. The significant and marginally significant results are presented in bold letters. For the random factor 'female identity', the percentage of variation occurring between females is presented.)

	development	lifespan	weight	sex ratio
<i>single factors</i>				
age	$F_{1,5}=6.12$ $p=0.225$	$F_{1,5}=0.38$ $p=1.00$	$F_{1,5}=3.51$ $p=0.479$	$F_{1,5}=2.55$ $p=0.684$
experience	$F_{1,11}=5.47$ $p=0.157$	$F_{1,11}=2.10$ $p=0.700$	$F_{1,11}=0.99$ $p=1.00$	$F_{1,11}=1.16$ $p=1.00$
sex	$F_{1,40}=21.95$ $p<0.001$	$F_{1,40}=11.89$ $p=0.004$	$F_{1,40}=9.81$ $p=0.010$	—
endophyte	$F_{1,40}=0.14$ $p=1.00$	$F_{1,40}=6.74$ $p=0.053$	$F_{1,40}=0.20$ $p=1.00$	$F_{1,41}=1.64$ $p=0.830$
<i>random factor</i>				
female identity	<0.01%	0.09%	<0.01%	<0.01%

associations of the now dead aphid. *Asaphes vulgaris* appears to also respond to kairomones that occur in the silky cocoon of aphidiine wasp (Christiansen-Weniger 1992). Independent of the exact cue the female secondary parasitoid perceives, host preference has been shown previously for different host species (Chow & Mackauer 1999), and we provide further evidence that age and experience of *A. vulgaris* females play a decisive role for their reproductive success.

For primary parasitoids, it has been proposed that mothers deposit a chemical cue inside the silk surrounding the pupa, which may condition the offspring to the environment in which they developed (van Emden *et al.* 1996; Douloumpaka & van Emden 2003). The fact that, in our experiment, the degree of preference exhibited by female wasps for E- mummies was time dependent suggests that the mechanism is likely to be different. All *A. vulgaris* females concentrated on host feeding for the first couple of days, presumably to mature their eggs. The oviposition events increased with more experience, and increasing numbers of eggs were laid in E- mummies. For host-feeding behaviour, no such temporal patterns were observed and no evidence for preference was detected. Therefore, females could have learned during host feeding to distinguish high- versus low-quality hosts.

The lack of detection of host-feeding patterns may have been caused by the relatively small sample size of nine females and the short time design of the experiment. Experiments with secondary parasitoids are extremely difficult to conduct, which is probably why very little data exist for multi-trophic interactions including secondary parasitoids (Brodeur 2000; but see Harvey *et al.* 2003; Soler *et al.* 2005). We believe that our experiment provides strong evidence for the cascading effects of endophytes to the top fourth trophic level of the food web, despite the small number of females tested. Changes in secondary parasitoid number and diversity depending on endophyte infection are also supported by a field study (Omacini *et al.* 2001). In that study, the food chain based on endophyte-infected plants and the aphid *Rhopalosiphum padi* showed strong numerical responses of primary and secondary parasitoids. The co-occurring aphid *M. festucae* was less affected by the presence of endophytes, but its parasitoid community also showed decreased abundance and

diversity. Apart from numerical responses in aphids, the pattern of reduced aphid secondary parasitoid abundance and diversity in the endophyte environment could have arisen by discrimination against hosts containing endophytes. In our laboratory experiment, the herbivore aphid *M. festucae* was relatively insensitive to the presence of the endophyte, while the associated primary parasitoids (Härrä 2007), and as shown here a secondary parasitoid, showed reduced fitness in the presence of endophytes.

In conclusion, we showed that the fitness of aphid secondary parasitoids when developing within hosts from an endophyte-infected environment was reduced and females learned to discriminate against these hosts of lower quality. The reduction in fitness of aphid secondary parasitoids in the presence of endophytes limits their top-down control on primary parasitoids and may result in complex indirect effects for lower trophic levels in the food web.

We thank Dennis Hansen, Uli Steiner and Nancy Bunbury for their helpful discussions and comments. Jeff Harvey and three anonymous reviewers helped improving the manuscript. The project was funded by an SNSF grant to C.B.M. (631-065950).

REFERENCES

- Arnold, A. E. & Lutzoni, F. 2007 Diversity and host range of foliar fungal endophytes: are tropical leaves biodiversity hotspots? *Ecology* **88**, 541–549. (doi:10.1890/05-1459)
- Brodeur, J. 2000 Host specificity and trophic relationships of hyperparasitoids. In *Parasitoid population biology* (eds M. E. Hochberg & A. R. Ives), pp. 163–183. Princeton, NJ: Princeton University Press.
- Buitenhuis, R., McNeil, J. N., Boivin, G. & Brodeur, J. 2004 The role of honeydew in host searching of aphid hyperparasitoids. *J. Chem. Ecol.* **30**, 273–285. (doi:10.1023/b:joec.0000017977.39957.97)
- Bultman, T. L., Borowicz, K. L., Schneble, R. M., Coudron, T. A. & Bush, L. P. 1997 Effect of a fungal endophyte on the growth and survival of two *Euplectrus* parasitoids. *Oikos* **78**, 170–176. (doi:10.2307/3545812)
- Charnov, E. L., Los-den Hartogh, R. L., Jones, W. T. & van den Assem, J. 1981 Sex ratio evolution in a variable environment. *Nature* **289**, 27–33. (doi:10.1038/289027a0)
- Chow, A. & Mackauer, M. 1999 Host handling and specificity of the hyperparasitoid wasp, *Dendrocerus*

- carpenteri* (Curtis) (Hym., Megaspilidae): importance of host age and species. *J. Appl. Entomol.* **123**, 83–91. (doi:10.1046/j.1439-0418.1999.00322.x)
- Christiansen-Weniger, P. 1992 Wirt-Parasitoid-Beziehung zwischen Blattlausprimärparasitoiden und den Blattlaushyperparasitoiden *Asaphes vulgaris* Wlk. und *Asaphes suspensus* (Nees) (Hymenoptera: Pteromalidae). PhD thesis Christian-Albrechts-Universität zu Kiel, Kiel, Germany.
- Clay, K. 1990 Fungal endophytes of grasses. *Annu. Rev. Ecol. Syst.* **21**, 275–297. (doi:10.1146/annurev.es.21.110190.001423)
- de Sassi, C., Müller, C. B. & Krauss, J. 2006 Fungal plant endosymbionts alter life history and reproductive success of aphid predators. *Proc. R. Soc. B* **273**, 1301–1306. (doi:10.1098/rspb.2005.3442)
- Douloumpaka, S. & van Emden, H. F. 2003 A maternal influence on the conditioning to plant cues of *Aphidius colemani* Viereck, parasitizing the aphid *Myzus persicae* Sulzer. *Physiol. Entomol.* **28**, 108–113. (doi:10.1046/j.1365-3032.2003.00323.x)
- Godfray, H. C. J. 1994 *Parasitoids. Behavioral and evolutionary ecology*. Monographs in Behavior and Ecology. Princeton, NJ: Princeton University Press.
- Härril, S. A. 2007 Fungal endosymbionts of grasses, and their effects on multitrophic interactions. PhD thesis, University of Zürich, Zürich, Switzerland.
- Harvey, J. A. 2008 Comparing and contrasting development and reproductive strategies in the pupal hyperparasitoids *Lysibia nana* and *Gelis agilis* (Hymenoptera: Ichneumonidae). *Evol. Ecol.* **22**, 153–166. (doi:10.1007/s10682-007-9164-x)
- Harvey, J. A., van Dam, N. M. & Gols, R. 2003 Interactions over four trophic levels: foodplant quality affects development of a hyperparasitoid as mediated through a herbivore and its primary parasitoid. *J. Anim. Ecol.* **72**, 520–531. (doi:10.1046/j.1365-2656.2003.00722.x)
- Heimpel, G. E., Rosenheim, J. A. & Mangel, M. 1996 Egg limitation, host quality, and dynamic behavior by a parasitoid in the field. *Ecology* **77**, 2410–2420. (doi:10.2307/2265742)
- Iwasa, Y., Suzuki, Y. & Matsuda, H. 1984 Theory of oviposition strategy of parasitoids—1. Effect of mortality and limited egg number. *Theor. Popul. Biol.* **26**, 205–227. (doi:10.1016/0040-5809(84)90030-3)
- Jervis, M. A. & Kidd, N. A. C. 1986 Host-feeding strategies in hymenopteran parasitoids. *Biol. Rev. Camb. Phil. Soc.* **61**, 395–434. (doi:10.1111/j.1469-185X.1986.tb00660.x)
- Jervis, M. A., Eilers, J. & Harvey, J. A. 2008 Resource acquisition, allocation, and utilization in parasitoid reproductive strategies. *Annu. Rev. Entomol.* **53**, 361–385. (doi:10.1146/annurev.ento.53.103106.093433)
- Kidd, N. A. C. & Jervis, M. A. 1991 Host-feeding and oviposition strategies of parasitoids in relation to host stage. *Res. Popul. Ecol.* **33**, 13–28. (doi:10.1007/bf02514570)
- Müller, C. B. & Krauss, J. 2005 Symbiosis between grasses and asexual fungal endophytes. *Curr. Opin. Plant Biol.* **8**, 450–456. (doi:10.1016/j.pbi.2005.05.007)
- Müller, C. B., Adriaanse, I. C. T., Belshaw, R. & Godfray, H. C. J. 1999 The structure of an aphid-parasitoid community. *J. Anim. Ecol.* **68**, 346–370. (doi:10.1046/j.1365-2656.1999.00288.x)
- Ohgushi, T., Craig, T. P. & Price, P. W. 2007 Indirect interaction webs: an introduction. In *Ecological communities: plant mediation in indirect interaction webs* (eds T. Ohgushi, T. P. Craig & P. W. Price), pp. 3–15. Cambridge, UK: Cambridge University Press.
- Omacini, M., Chaneton, E. J., Ghersa, C. M. & Müller, C. B. 2001 Symbiotic fungal endophytes control insect host-parasite interaction webs. *Nature* **409**, 78–81. (doi:10.1038/35051070)
- Pimm, S. L. & Lawton, J. H. 1977 Number of trophic levels in ecological communities. *Nature* **268**, 329–331. (doi:10.1038/268329a0)
- Pinheiro, J. C. & Bates, D. M. 2000 *Mixed-effects models in S and S-PLUS. Statistics and computing*. New York, NY: Springer.
- Polis, G. A. & Strong, D. R. 1996 Food web complexity and community dynamics. *Am. Nat.* **147**, 813–846. (doi:10.1086/285880)
- Quicke, D. L. J. 1997 *Parasitic wasps*. London, UK: Chapman & Hall.
- Rosenheim, J. A. 1998 Higher-order predators and the regulation of insect herbivore populations. *Annu. Rev. Entomol.* **43**, 421–447. (doi:10.1146/annurev.ento.43.1.421)
- Rosenheim, J. A., Heimpel, G. E. & Mangel, M. 2000 Egg maturation, egg resorption and the costliness of transient egg limitation in insects. *Proc. R. Soc. B* **267**, 1565–1573. (doi:10.1098/rspb.2000.1179)
- Saikkonen, K., Ahlholm, J., Helander, M., Lehtimäki, S. & Niemeläinen, O. 2000 Endophytic fungi in wild and cultivated grasses in Finland. *Ecography* **23**, 360–366. (doi:10.1034/j.1600-0587.2000.d01-1645.x)
- Soler, R., Bezemer, T. M., van der Putten, W. H., Vet, L. E. M. & Harvey, J. A. 2005 Root herbivore effects on above-ground herbivore, parasitoid and hyperparasitoid performance via changes in plant quality. *J. Anim. Ecol.* **74**, 1121–1130. (doi:10.1111/j.1365-2656.2005.01006.x)
- Sullivan, D. J. 1987 Insect hyperparasitism. *Annu. Rev. Entomol.* **32**, 49–70. (doi:10.1146/annurev.en.32.010187.000405)
- Sullivan, D. J. & Völkl, W. 1999 Hyperparasitism: multitrophic ecology and behavior. *Annu. Rev. Entomol.* **44**, 291–315. (doi:10.1146/annurev.ento.44.1.291)
- van Emden, H. F., Sponagl, B., Wagner, E., Baker, T., Ganguly, S. & Douloumpaka, S. 1996 Hopkins 'host selection principle', another nail in its coffin. *Physiol. Entomol.* **21**, 325–328. (doi:10.1111/j.1365-3032.1996.tb00873.x)
- van Nouhuys, S. & Hanski, I. 2002 Multitrophic interactions in space: metacommunity dynamics in fragmented landscapes. In *Multitrophic level interactions* (eds T. Tscharrntke & B. A. Hawkins), pp. 124–147. Cambridge, UK: Cambridge University Press.
- Venables, W. N. & Ripley, B. D. 2002 *Modern applied statistics with S*. New York, NY: Springer.
- Vet, L. E. M., Lewis, W. J., Papaj, D. R. & van Lenteren, J. C. 1990 A variable-response model for parasitoid foraging behavior. *J. Insect Behav.* **3**, 471–490. (doi:10.1007/BF01052012)
- Vinson, S. B. 1976 Host selection by insect parasitoids. *Annu. Rev. Entomol.* **21**, 109–133. (doi:10.1146/annurev.en.21.010176.000545)
- Zabalgozcoa, I., Vázquez de Aldana, B. R., García Ciudad, A. & García Criado, B. 2003 Fungal endophytes in grasses from semi-arid permanent grasslands of western Spain. *Grass Forage Sci.* **58**, 94–97. (doi:10.1046/j.1365-2494.2003.00352.x)