

Review

Neural mechanisms underlying the vulnerability to develop compulsive drug-seeking habits and addiction

Barry J. Everitt*, David Belin, Daina Economidou, Yann Pelloux,
Jeffrey W. Dalley and Trevor W. Robbins

*Behavioural and Clinical Neuroscience Institute, Department of Experimental Psychology,
University of Cambridge, Downing Street, Cambridge CB2 3EB, UK*

We hypothesize that drug addiction can be viewed as the endpoint of a series of transitions from initial voluntary drug use through the loss of control over this behaviour, such that it becomes habitual and ultimately compulsive. We describe evidence that the switch from controlled to compulsive drug seeking represents a transition at the neural level from prefrontal cortical to striatal control over drug-seeking and drug-taking behaviours as well as a progression from ventral to more dorsal domains of the striatum, mediated by its serially interconnecting dopaminergic circuitry. These neural transitions depend upon the neuroplasticity induced by chronic self-administration of drugs in both cortical and striatal structures, including long-lasting changes that are the consequence of toxic drug effects. We further summarize evidence showing that impulsivity, a spontaneously occurring behavioural tendency in outbred rats that is associated with low dopamine D_{2/3} receptors in the nucleus accumbens, predicts both the propensity to escalate cocaine intake and the switch to compulsive drug seeking and addiction.

Keywords: vulnerability; compulsion; addiction; striatum; dopamine; habits

1. INTRODUCTION

The central hypothesis guiding our research during the past decade or more has been that drug addiction can be understood in terms of the operation of the brain's learning and memory systems (Robbins & Everitt 1999; Everitt *et al.* 2001; Everitt & Robbins 2005). In particular, that chronically self-administered drugs may in some way pathologically subvert these memory systems and so lead to the establishment of compulsive drug-seeking habits (Everitt & Robbins 2005). Initially, our approach to the understanding of drug addiction built upon the clinical insight embodied within the DSM-IV (1994) criteria for 'substance abuse' and 'substance dependence'. Therefore, we began experimentally to decompose and augment the DSM-IV diagnostic framework in terms of specified learning and cognitive processes deriving from animal learning theory and that increasingly have been attributed to the operation of specific neural, especially limbic cortical–striatal, systems (Everitt *et al.* 2001).

The early focus of much experimental drug addiction research was to understand the reinforcing, or

'rewarding', effects of abused drugs; this has led to great advances in defining the primary molecular targets of addictive drugs as well as, more recently, the adaptations in these targets that develop with chronic drug self-administration (Nestler 2004; Koob & Le Moal 2005). However, it has been appreciated for some time that the molecular and neurochemical correlates of acute and chronic drug administration must be interpreted in behavioural and cognitive terms if the psychological processes and neurobiological mechanisms determining human drug addiction are to be specified. Therefore, we and others have increasingly come to view drug addiction as the endpoint of a series of transitions from initial drug use—when a drug is voluntarily taken because it has reinforcing, often hedonic, effects—through the loss of control over this behaviour, such that it becomes habitual and ultimately compulsive. We have recently reviewed the evidence that these transitions depend upon interactions between Pavlovian and instrumental learning processes (Everitt & Robbins 2005). Furthermore, we have hypothesized that the 'switch' from voluntary drug use to habitual and progressively compulsive drug use represents a transition at the neural level from prefrontal cortical to striatal control over drug-seeking and drug-taking behaviours, as well as a progression from ventral to more dorsal domains of the striatum, mediated at least in part by its stratified dopaminergic innervation (Everitt & Robbins 2005).

* Author for correspondence (bje10@cam.ac.uk).

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rstb.2008.0089> or via <http://journals.royalsociety.org>.

One contribution of 17 to a Discussion Meeting Issue 'The neurobiology of addiction: new vistas'.

We summarize here recent evidence in support of the hypothesis that drug-seeking habits are associated with a shift from ventral to dorsal striatal control over behaviour. We also address the major issue of vulnerability to drug addiction—that some individuals are more likely than others to take drugs, lose control over and escalate their drug intake, and ultimately to seek drugs compulsively. Experimental models of addiction, rather than drug self-administration, must reflect such individual differences and also incorporate extended periods of drug taking if the underlying neural mechanisms of addiction are to be identified (Koob & Le Moal 2005).

2. FROM VOLUNTARY TO HABITUAL DRUG SEEKING: THE SHIFT FROM VENTRAL TO DORSAL STRIATUM

With drugs such as cocaine, there is wide, though not universal, agreement that the dopaminergic innervation of the nucleus accumbens shell (AcbS), and even more ventral regions of the striatum, such as the olfactory tubercle, underlies its primary reinforcing effects (Di Chiara *et al.* 2004; Wise 2004; Ikemoto *et al.* 2005) as measured in drug self-administration procedures. We term this 'drug taking' to distinguish it from 'drug-seeking' behaviour, which must often be maintained over long periods of time and is profoundly influenced by environmental stimuli associated with self-administered drugs through Pavlovian conditioning (Everitt *et al.* 2001). In humans, these conditioned stimuli (CSs) can induce subjective states such as craving, as well as drug seeking and relapse after abstinence (Grant *et al.* 1996; Childress *et al.* 1999; Garavan *et al.* 2000). We have therefore developed a general model of drug seeking—a second-order schedule of cocaine reinforcement—in which this behaviour is sensitive not only to the contingency between instrumental responses and drug administration, but also to the presence of drug-associated CSs that have a powerful effect on performance by acting as conditioned reinforcers (Everitt & Robbins 2000). We have established that the acquisition of this cocaine-seeking behaviour depends upon the integrity of the nucleus accumbens core (AcbC) and its afferents from the basolateral amygdala (BLA). Thus, selective lesions of the BLA prevented the acquisition of cocaine seeking under a second-order schedule (Whitelaw *et al.* 1996), as expected given its fundamental role in conditioned reinforcement (Cador *et al.* 1989; Cardinal *et al.* 2002). Similarly, selective lesions of the AcbC also greatly impaired the acquisition of cocaine seeking (Ito *et al.* 2004). Simple drug taking, on the other hand, was unimpaired by BLA or AcbC lesions (Whitelaw *et al.* 1996; Ito *et al.* 2004). As we have reviewed elsewhere, the AcbC region of the ventral striatum is also a key locus not only for conditioned reinforcement but also for other Pavlovian influences on appetitive behaviour, including approach, as measured in autoshaping procedures, and Pavlovian-instrumental transfer, the process by which Pavlovian CSs energize ongoing instrumental behaviour, an example of conditioned motivation,

each of which also depends upon processing in sub-regions of the amygdala (Cardinal *et al.* 2002; Cardinal & Everitt 2004).

These observations indicate that the BLA and AcbC may function together as nodes within a limbic cortical-ventral striatopallidal system that underlies the acquisition of drug seeking. This notion is further supported by the observation that disconnecting these structures by unilateral pharmacological blockade of dopamine and AMPA receptors in the BLA and AcbC, respectively, on opposite sides of the brain also greatly diminished cocaine seeking (Di Ciano & Everitt 2004). Thus, the acquisition and early performance of cocaine seeking that is maintained over protracted periods of time by the contingent presentation of drug-associated conditioned reinforcers depends upon the integrity of the AcbC and its afferent input from the BLA. It can be assumed that, at this stage, drug seeking is under the control of instrumental response–outcome contingencies in which animals respond in a voluntary and goal-directed way for intravenous cocaine infusions. Indeed, we have clear evidence that this is so from studies using a 'seeking–taking' chained schedule in which animals perform a drug-seeking response in the initial link of the chain, which then gives access to a drug-taking response in the second link, performance of which delivers cocaine (Olmstead *et al.* 2001). After a limited experience with the drug, cocaine seeking was shown to be a goal-directed action in that its performance was sensitive to devaluation of the drug-taking link (Olmstead *et al.* 2001). This devaluation effect demonstrates that cocaine seeking at early stages of acquisition and performance is mediated by the knowledge of the contingency between the seeking response and its outcome.

This response–outcome process can be contrasted with a second, stimulus–response (S–R) instrumental process in which seeking behaviour is a response habit triggered and maintained by drug-associated stimuli (Everitt *et al.* 2001). Tiffany (1990), O'Brien & McClellan (1996) and ourselves (Robbins & Everitt 1999; Everitt *et al.* 2001; Everitt & Robbins 2005) have all advanced the hypothesis that drug addiction encompasses changes in the associative structure underlying drug seeking such that it becomes 'automatic' or habitual. We have additionally hypothesized that the progressive engagement of dorsal striatal mechanisms underlies this transition (Everitt & Robbins 2005), based upon evidence that the dorsolateral (DL) striatum mediates S–R habit learning—evidence that has been considerably strengthened through the recent studies of instrumental responding for food and its resistance to reinforcer devaluation, a canonical test of the development of S–R habits (Yin *et al.* 2004, 2006). What then is the evidence that the dorsal striatum mediates the performance of well-established drug-seeking habits that depend initially upon ventral striatal mechanisms during acquisition? Correlative evidence came from *in vivo* microdialysis measurement of extracellular dopamine in rats that had attained over a two-month period stable responding under a second-order schedule of cocaine seeking. While self-administered cocaine increased dopamine release in the AcbS, AcbC and caudate–putamen, extracellular




Figure 1. Within striatum serial processing underlies the establishment of cocaine-seeking habits. (a) Schematic of the intrastriatal dopamine-dependent spiralling circuitry functionally connecting the ventral with the dorsal striatum in the rat (modified from Belin & Everitt 2008). The spiralling loop organization is depicted as the alternation of pink and black arrows from the ventral to the more dorsal parts of the circuit, i.e. from the AcbS (yellow) to the AcbC (light blue) via the ventral tegmental area (pink) and from the AcbC, via the substantia nigra to the dorsal striatum (dark blue). (b) Cocaine seeking is dose dependently impaired by bilateral infusions of the DA receptor antagonist α -flupenthixol (depicted as green dots) into the DL striatum. α -Flupenthixol infusions into the DL striatum dose dependently decreased responding on the active lever under a second-order schedule of cocaine reinforcement, but had no effect on responding on the inactive lever (Belin & Everitt 2008). (c) Disconnecting the AcbC from the dopaminergic innervation of the dorsal striatum impairs habitual cocaine seeking. In unilateral AcbC-lesioned rats, the AcbC relay of the loop is lost on the one side of the brain. However, on the non-lesioned side, the spiralling circuitry is intact and functional. When α -flupenthixol (green dots) is infused in the DL striatum contralateral to the lesion it blocks the DAergic innervation from the midbrain, impairing the output structure of the spiralling circuitry on the non-lesioned side of the brain. Therefore, this asymmetric manipulation disconnects the core of the nucleus accumbens from the DL striatum bilaterally and greatly diminishes cocaine seeking (figure adapted with permission from Belin & Everitt (2008)).

dopamine was increased selectively in the AcbC in response to unexpected (i.e. non-response contingent) presentations of a cocaine-associated stimulus. However, during a prolonged period of cocaine seeking maintained by contingent presentations of the same cocaine-associated CS, dopamine release was increased only in the dorsal striatum, not in the AcbC or AcbS (Ito *et al.* 2000, 2002). Furthermore, dopamine release in the dorsal striatum was subsequently shown to be causally important for the *maintenance* of drug seeking, since it was greatly reduced by dopamine receptor blockade at this site, whereas this treatment was without effect in the AcbC (Vanderschuren *et al.* 2005). This apparent shift from ventral to dorsal striatal control over drug seeking provides support for the hypothesis that it is under S–R, or ‘habit’, control owing to accumulated evidence implicating the dorsal striatum in habit learning (Yin *et al.* 2004, 2006). Moreover, the fact that the second-order schedule we have principally used is of a fixed-interval type encourages this view, since interval schedules are known to result in the more rapid development of S–R habits through the weaker relationship between response and outcome

that obtains, when compared with ratio schedules (Dickinson 1985). In addition, studies with orally self-administered cocaine and alcohol have shown the more rapid development of habitual drug seeking compared with the seeking of a natural sweet reward (Dickinson *et al.* 2002, Miles *et al.* 2003).

These observations supporting the increasing importance of the dorsal striatum in well-established, or habitual, drug seeking raise the issue of how, in neural terms, such a shift in the locus of control from ventral to dorsal striatum might occur. The observations of Haber *et al.* (2000) in primates and, more recently, Ikemoto (2007) in rats provide a possible neuroanatomical basis. They showed that ventral tiers of the striatal complex regulate the dopaminergic innervation of more dorsal tiers through so-called ‘spiralling’ connections with the midbrain (figure 1a). Thus, the AcbS projects to dopamine neurons in the ventral tegmental area that innervate not only the shell but also the more dorsally situated AcbC. Neurons in the AcbC innervate dopamine neurons projecting to both the AcbC and the immediately dorsal regions of the dorsomedial caudate–putamen and so on, in a serially cascading pattern ultimately to encompass

more lateral parts of the dorsal striatum—the site at which dopamine release is increased during habitual drug seeking and where dopamine receptor antagonist infusions impair this behaviour.

We tested the hypothesis that the serial cascade of striato-nigro-striatal connectivity underlies this progressively greater control over well-established cocaine-seeking behaviour by the DL striatum using a novel ventral–dorsal striatal ‘disconnection’ (figure 1*b*). Thus, the AcbC was selectively lesioned on one side of the brain and combined with dopamine receptor blockade in the contralateral DL striatum, thereby functionally disconnecting serial interactions between these ventral and dorsal striatal domains bilaterally (Belin & Everitt 2008). This disconnection greatly and selectively decreased cocaine seeking in rats tested some weeks after stable responding had been attained under a second-order schedule of reinforcement (figure 1*c*). Two important additional observations underlined the specificity of the ventral–dorsal striatal disconnection effect. (i) The same animals were trained to perform a novel chain-pulling response for sucrose under a fixed ratio 1 schedule of reinforcement and again underwent the disconnection manipulation, or bilateral dorsal striatal dopamine receptor antagonist infusions (in non-lesioned rats), immediately after acquisition when the behaviour was under response–outcome control; neither manipulation had any effect (Belin & Everitt 2008). (ii) In separate groups of animals, either bilateral dorsal striatal dopamine receptor blockade or AcbC–dorsal striatal disconnection was performed at a much earlier stage of acquisition of the cocaine seeking, second-order schedule when responding was under ratio, rather than interval, control and when response–outcome mechanisms dominate performance. Again, neither manipulation had any effect on cocaine seeking (D. Belin & B. J. Everitt 2008, unpublished observations).

Taken together, the above results clearly indicate the devolution of control over drug seeking from ventral to dorsal striatum and also strongly suggest that this shift is progressive. Other data also support the notion of this shift. For example, using autoradiographic methods, Porrino and colleagues showed the development of neuroadaptations in $D_{2/3}$ dopamine receptors and other neurochemical, or metabolic, markers in the dorsal striatum following chronic, but not acute, cocaine self-administration by monkeys (Letchworth *et al.* 2001; Porrino *et al.* 2004). At earlier stages of training, these adaptations were largely restricted to the more ventral, nucleus accumbens region. The DL striatum has also been shown to be involved in ‘relapse’ to a cocaine-seeking habit, since neural inhibition induced by γ -aminobutyric acid receptor agonist infusion into this area, but not into the AcbS or AcbC, prevented the reinstatement of cocaine seeking after protracted withdrawal (Fuchs *et al.* 2006; See *et al.* 2007). Moreover, the presentation of drug cues to human cocaine addicts both induced drug craving (that has been shown to be correlated with the activation of the amygdala and limbic prefrontal cortical areas; Grant *et al.* 1996; Childress *et al.* 1999; Garavan *et al.* 2000) and also marked the activation of the dorsal striatum (Garavan *et al.* 2000; Volkow *et al.* 2006).

These observations therefore strongly indicate a link between limbic cortical mechanisms and the engagement of the dorsal striatum in long-term drug abusers exposed to drug cues, whereas the results of our ventral/dorsal striatal disconnection experiments reveal that this recruitment is mediated by antecedent limbic cortex-dependent activity in the AcbC and its regulation of dorsal striatal dopaminergic projections.

It seems likely that the ventral to dorsal striatum shift is not specific to drug seeking, but would apply equally to the control over instrumental responding for natural reinforcers under appropriate conditions. Indeed, lesion or inactivation of the AcbC, dorsomedial or DL striatum in rats responding for ingestive reinforcers does not globally impair instrumental behaviour, but instead has major effects that depend upon the response–outcome or S–R associative structure underlying the behaviour. Lesions or *N*-methyl-D-aspartate receptor blockade of the AcbC (Kelley *et al.* 1997; Corbit *et al.* 2001) or dorsomedial striatum (Yin *et al.* 2004, 2005), impair instrumental behaviour under response–outcome control, but actually enhance the development of S–R habits in which responding persists after reinforcer devaluation (Yin *et al.* 2004). By contrast, DL striatal lesions, inactivation or dopamine denervation return previously habitual responding to response–outcome control, reinstating sensitivity to reinforcer devaluation (Yin *et al.* 2004; Faure *et al.* 2005) or action–outcome contingency degradation (Yin *et al.* 2006). These observations emphasize that response–outcome and S–R learning mechanisms are probably engaged not serially, but in parallel, with DL striatum-dependent S–R mechanisms eventually dominating the control over behaviour.

However, it is possible that the shift from ventral to dorsal striatal control occurs more rapidly in animals seeking drugs owing to the effects of these agents themselves on the plasticity mechanisms involved, particularly in the case of psychomotor stimulants which so powerfully increase dopamine transmission. Thus, an amphetamine sensitization treatment regimen leads to the more rapid instantiation of habit learning in animals responding for food (Nelson & Killcross 2006). Moreover, animals that had escalated their cocaine intake showed, when subsequently challenged with cocaine, a marked enhancement of stereotyped behavioural responses that have long been known to depend upon the dorsal, rather than the ventral, striatal dopaminergic innervation (Ferrario *et al.* 2005). Finally, Schoenbaum and colleagues have demonstrated a shift in the balance of associative encoding from ventral to dorsal striatum correlated with concomitant enhancement of cue-evoked neuronal firing in the dorsal striatum (Takahashi *et al.* 2007). This finding resonates with the observation of drug-associated CS-induced activation of the dorsal striatum in human cocaine abusers (Garavan *et al.* 2000; Volkow *et al.* 2006). Therefore, the unique properties of drugs as reinforcers, especially stimulant drugs, might accelerate, or more effectively *consolidate*, the development of drug seeking as a S–R habit (Everitt *et al.* 2001).

But it should also be appreciated that while all animals responding for drugs or natural reinforcers will engage the dorsal striatal habit mechanism under

appropriate reinforcement contingencies, not all individuals that take addictive drugs will become addicted. That is, not all individuals self-administering drugs will escalate their intake and go on to develop compulsive drug seeking that persists in the face of negative or aversive outcomes, the key characteristics of addiction, or substance dependence, in DSM-IV. Thus, some individuals are vulnerable in terms of these characteristics, a proportion that is often estimated to be approximately 20% or less of those initially exposed to addictive drugs (Anthony *et al.* 1994). We have shown recently that impulsivity is a behavioural characteristic that both predicts the escalation of cocaine intake and the progression to compulsive drug seeking, as well as an increased propensity to relapse after abstinence.

3. IMPULSIVITY, VENTRAL STRIATAL DOPAMINE RECEPTORS AND VULNERABILITY TO ADDICTION

Studies of human addicts have implicated individual differences in impulsivity, or other traits, such as 'sensation seeking', in the vulnerability to drug use and abuse (Verdejo-Garcia & Perez-Garcias 2007); although it has never been clear whether the impulsivity observed in drug addicts predates the onset of addictive behaviour or is a consequence of protracted exposure to drugs (Dom *et al.* 2006; Zilberman *et al.* 2007). We have investigated this issue experimentally by defining in rats an operational measure of the human trait of impulsivity as premature responses in a five-choice serial reaction-time task (5-CSRTT; Dalley *et al.* 2007). A proportion (less than 10%) of the outbred Lister-hooded strain of rats in our study were impulsive on this task in that they showed high levels of anticipatory responses made before the presentation of a food-predictive, brief light stimulus—especially under conditions when the stimulus presentation was delayed after trial onset (Dalley *et al.* 2007). This impulsivity we term 'waiting impulsivity', as it is related to impulsivity measured as an inability to tolerate delays of reinforcement (Robinson *et al.* 2008), but appears different from the response impulsivity measured by the stop-signal reaction time in STOP-signal reaction-time tasks (Eagle *et al.* 2008; E. S. J. Robinson, D. M. Eagle, D. Economidou, D. E. H. Theobald, A. C. Mar, E. R. Murphy, T. W. Robbins & J. W. Dalley 2008, unpublished observations).

When allowed to self-administer cocaine intravenously, the impulsive rats showed a marked escalation of their cocaine intake compared with non-impulsive controls. They did not acquire cocaine self-administration more rapidly, but responded at a much higher rate for their cocaine infusions than did non-impulsive subjects (Dalley *et al.* 2007). This is in marked contrast to rats showing high locomotor responses to novelty (a 'sensation-seeking' phenotype), which self-administer cocaine at low doses that do not sustain self-administration in those rats that show low locomotor responses to novelty (Piazza *et al.* 1989). Thus, high impulsivity predicts the tendency to escalate cocaine intake, which is reminiscent of one of the diagnostic characteristics of substance dependence in DSM-IV which describes the taking of drugs in larger quantities than intended as a core symptom.

Highly impulsive animals were investigated neurobiologically using positron emission tomography to measure binding in the striatum of the selective, high affinity D_{2/3} dopamine receptor antagonist [¹⁸F]fallypride (Mukherjee *et al.* 1999). Impulsive animals showed markedly reduced fallypride binding within the ventral, but not the dorsal striatum. Moreover, the reduced D_{2/3} dopamine receptor availability in the ventral striatum was correlated with impulsivity on the 5-CSRTT (Dalley *et al.* 2007). Thus, low D_{2/3} dopamine receptors in the ventral striatum, encompassing the nucleus AcbC and shell regions, are correlated both with impulsivity and the marked propensity to escalate cocaine intake. This observation, taken together with the role of the AcbS in the reinforcing and stimulant effects of cocaine, as well as involvement of the AcbC in the acquisition of drug-seeking behaviour (Ito *et al.* 2004) and the ability to tolerate delays to reinforcement (Cardinal *et al.* 2001), indicates the importance of the ventral striatum in the neural mechanisms underlying the propensity to seek and work for cocaine over extended periods of time. Nader and colleagues, in studies of socially housed monkeys, have also demonstrated the significant relationship between D₂ dopamine receptors in the striatum and cocaine self-administration (Czoty *et al.* 2004; Nader *et al.* 2006); these studies are reviewed comprehensively by Nader *et al.* (2008).

Impulsive rats having self-administered cocaine subsequently showed reduced levels of impulsivity (Dalley *et al.* 2007)—perhaps related to the reduction in impulsive behaviour following treatment with the stimulant methylphenidate seen in humans with attention-deficit hyperactivity disorder. Moreover, during a subsequent period in which animals had no access to cocaine self-administration ('enforced abstinence'), impulsivity returned to near pre-cocaine self-administration levels (figure 2a). Since impulsivity predicts the escalation of drug intake, we investigated whether the return of impulsivity during abstinence might also be associated with a greater propensity to relapse, as suggested by de Wit & Richards (2004). In these experiments (Economidou *et al.* 2007; D. Economidou, J. W. Dalley & B. J. Everitt 2007, unpublished observations), rats were trained in a cocaine seeking-taking task. Once responding had stabilized, a punishment contingency was introduced whereby on a random basis 50% of the seeking responses were followed by the presentation of the cocaine taking link in the chain, but 50% were followed by mild footshock (Pelloux *et al.* 2007). This schedule of unpredictable cocaine taking and aversive footshock outcomes results in the suppression of drug-seeking responses, especially after a limited, or non-escalated, history of cocaine self-administration; it can therefore be viewed as the development of 'abstinence', in that animals voluntarily withhold their drug-seeking responses when the punishment contingency is present. After having attained abstinence in this way and some two weeks after their last cocaine infusion, groups of impulsive and non-impulsive rats were reintroduced to the test boxes and again allowed to respond on the seeking lever, which always resulted in access to the




Figure 2. (a) The return of impulsive behaviour in highly impulsive rats following withdrawal and abstinence from cocaine self-administration. The data shown are premature responses in a 5-CSRTT. (i) Highly impulsive rats respond prematurely before any cocaine experience and (ii) their impulsivity is reduced following sessions of cocaine self-administration; (iii) but premature responding returns to pre-cocaine levels following an extended period of withdrawal (J. W. Dalley 2007, unpublished observations). Filled circles, high; open circles, low. (b) Impulsive rats show an increased propensity to relapse after abstinence. (i) Impulsive and non-impulsive rats were trained to seek and take cocaine under a chained schedule. (ii) Subsequently, 50% of the seeking responses were followed unpredictably by punishment and 50% by access to the taking lever; this results in the suppression of drug seeking. (iii) Following a further period of abstinence, when no cocaine was available rats were returned to the self-administration setting in which seeking responses resulted in the presentation of a cocaine-associated conditioned reinforcer, but no drug. (iv) Impulsive rats showed much higher numbers of seeking responses than low impulsive rats and hence showed a greater propensity to 'relapse'. Filled circles, high; open circles, low.

taking lever, responses upon which resulted in the presentation of the cocaine-associated CS, but no drug. While both groups of animals reinstated their drug-seeking responses, i.e. 'relapsed', responding by impulsive rats was markedly and significantly greater than that of non-impulsive subjects (figure 2b). Thus, impulsivity not only confers an increased predisposition to escalate cocaine self-administration but also an increased propensity to relapse to a drug-seeking habit after abstinence. The neurobiological mechanisms underlying this persisting vulnerability to seek and take drugs after abstinence are an important area for future research.

4. FROM IMPULSIVITY TO COMPULSIVE DRUG SEEKING IN ADDICTION

Demonstrating that impulsivity is a factor underlying the tendency to escalate drug intake and to relapse after abstinence leaves open the important issue of whether impulsivity is also a vulnerability marker for drug addiction and the compulsive drug seeking this entails. There are relatively few accepted models of compulsive drug seeking, or indeed compulsive behaviour in general, in animals. Perseverative responding in reversal learning tasks may provide one interesting example because this form of compulsion is persistently enhanced following even relatively brief periods of

cocaine treatment (Jentsch *et al.* 2002; Calu *et al.* 2007). In theoretical terms, we have suggested (Everitt & Robbins 2005) that compulsive drug seeking can be characterized as a maladaptive S-R habit in which the ultimate goal of the behaviour has been devalued, perhaps through tolerance to the rewarding effects of the drug. Instead, drug seeking is increasingly controlled by a succession of drug-associated discriminative stimuli, which also function as conditioned reinforcers when presented as a consequence of instrumental responses, as in the second-order drug-seeking schedule described above. Central to drug addiction, then, is the persisting quality of these habits, which we have suggested (Everitt & Robbins 2005) may correspond to the subjective state of 'must do!'—the persistent reinitiation of habitual acts—not least to distinguish it from the subjective state of excessive 'wanting' embodied in the incentive salience sensitization view of addiction (Robinson & Berridge 1993; see Robinson & Berridge 2008).

In attempting to model drug addiction in animals, we have tried to capture the *compulsive* quality of drug seeking by measuring its persistence despite negative or aversive outcomes, as in the DSM-IV. In developing such behavioural procedures, we have shown that compulsive drug seeking only emerges following an extended, or chronic, history of cocaine taking

(Deroche-Gamonet *et al.* 2004; Vanderschuren & Everitt 2004; Pelloux *et al.* 2007). In the study by Deroche-Gamonet *et al.* (2004), three addiction-like behavioural criteria were measured in rats, namely (i) increased motivation to take the drug, (ii) inability to refrain from drug seeking, and (iii) maintained drug use despite aversive consequences. After approximately 40 days of cocaine self-administration, but not at earlier times, some 17% of subjects developed these addiction-like criteria showing increased break points under a progressive ratio of cocaine reinforcement, persistent responding during signalled periods of drug unavailability and, perhaps most importantly, persistence of the instrumental nose-poke response for cocaine even when it was punished by mild footshock. In the study by Pelloux *et al.* (2007), rats were trained on the seeking-taking chained schedule with intermittent punishment of the seeking response (i.e. on 50% of the seeking bouts, randomly occurring) to achieve suppression of drug seeking, or abstinence, as described previously. In this study too, whereas all rats suppressed their cocaine seeking after a limited history of cocaine self-administration, after an extended history 17–20% of subjects were completely resistant to punishment, continuing to seek and take drugs despite the ongoing, daily experience of the negative outcome. The proportion of rats compulsively seeking drugs, then, was similar both to that in the Belin study and to the addiction-vulnerable subgroup of human subjects, often estimated to be less than 20% of the population that initially use drugs (Anthony *et al.* 1994).

However, the origins of this propensity to seek cocaine *compulsively* have not been established. We hypothesized that impulsivity, which we have shown to be associated with low $D_{2/3}$ dopamine receptor availability in the ventral striatum and to predict the escalation of cocaine intake (Dalley *et al.* 2007), might also confer a vulnerability to develop compulsive drug seeking and addiction following extended access to cocaine. Rats were screened both for impulsivity in the 5-CSSRT and also for the sensation-seeking phenotype of high locomotor responsiveness to novelty (HR rats) which has earlier been suggested to be an addiction vulnerability phenotype (Piazza *et al.* 1989). The resultant groups were then studied in the acquisition of cocaine self-administration, and for the emergence of the three addiction-like behavioural criteria identified by Deroche-Gamonet *et al.* (2004), but particularly persistent responding for the drug in the face of punishment. As expected, and as reported previously (Piazza *et al.* 1989), the HR rats more readily acquired cocaine self-administration and showed an upward shift in the cocaine dose-response curve as compared both with rats with low responses to novelty and also high impulsivity (Belin *et al.* 2008). We also confirmed our earlier finding that impulsivity is not associated with more rapid acquisition of cocaine self-administration, but instead with the escalation of cocaine intake. However, and in marked contrast, it was high impulsivity, but not high reactivity to novelty, that predicted the switch (Leshner 1997) from controlled to compulsive cocaine taking (Belin *et al.* 2008). Highly impulsive rats displayed higher addiction scores and much greater resistance to punishment than rats with

high or low responses to novelty or low impulsivity (electronic supplementary material, figure S1). In fact, highly impulsive rats did not differ from rats showing the three addiction-like behavioural criteria in any of their addiction-like behaviours after the extended period of cocaine self-administration. Therefore, it seems, perhaps counter-intuitively, that the propensity to acquire cocaine self-administration when first encountering the drug and the vulnerability to develop compulsive cocaine intake (addiction), depend upon distinct and seemingly orthogonal behavioural characteristics—novelty/sensation-seeking versus impulsivity, respectively—each of which might have a genetic or environmental basis. The results of this study also provide experimental evidence that high levels of impulsivity can antedate the onset of compulsive drug use, thereby emphasizing the importance of pre-existing impulsivity seen in individuals addicted to drugs (Jentsch & Taylor 1999; Dom *et al.* 2006).

Now that we have established models of compulsive drug seeking and addiction, it will be possible to investigate not only predisposing factors, such as impulsivity, but also the underlying neurobiological mechanisms. There are several current views about the origins of compulsion within the brain, which are often thought of as being, but in reality are not, mutually exclusive. The neuroadaptations occurring during behavioural sensitization to stimulant drugs have been argued to underlie an extreme incentive motivational state of drug ‘wanting’ (Robinson & Berridge 1993). According to this view, addicts experience this state especially when exposed to drug-associated cues, which leads to over-activation of the sensitized dopaminergic innervation of the nucleus accumbens, in which plasticity-associated structural changes in dendritic spines have also been observed (Ferrario *et al.* 2005). This hypothesis is discussed in detail by Robinson & Berridge (2008). One interpretation of compulsive drug seeking, then, is that it is a behavioural manifestation of this potentiated motivational state, which has been demonstrated in some studies as an increased break point under progressive ratio schedules of reinforcement (for a review, see Vezina 2004). However, as noted above, a sensitization treatment regimen with amphetamine also leads to the more rapid instantiation of S–R habits (Nelson & Killcross 2006) and it is not easy at the behavioural level to differentiate between an increased tendency to repeat drug-seeking responses elicited and maintained by drug-associated stimuli—the ‘must do!’ of *compulsive* habits discussed above—from an increased desire for a drug, which might also seem counter-intuitive given the development of tolerance to its rewarding or reinforcing effects.

Perhaps more directly related to the notion of drug seeking as a compulsive habit, however, is the observation of reductions in dopamine D_2 receptors in the dorsal striatum in abstinent alcoholics, cocaine, heroin and methamphetamine addicts (Volkow & Wise 2005) and also following chronic, but not acute, cocaine self-administration in monkeys (Moore *et al.* 1998; Nader *et al.* 2002). The consequences of this change in striatal dopamine D_2 receptors for the plasticity underlying instrumental learning and performance is unclear, but an intriguing putative sequential mechanism

within the striatum and involving its dopaminergic innervation might be considered. Thus, the early vulnerability to escalate cocaine intake seen in impulsive rats is predicted by low $D_{2/3}$ dopamine receptor levels in the *ventral*, but not the dorsal striatum (Dalley *et al.* 2007). However, this escalated intake may lead to more rapid neuroadaptations, including downregulated D_2 dopamine receptors, in the *dorsal* striatum and mediated in part by aberrant engagement of the spiralling striato-nigro-striatal circuitry. This would lead to more rapid consolidation of drug-seeking habits that are difficult to relinquish, despite negative outcomes and are more readily reinstated after abstinence following exposure to response-eliciting drug-associated stimuli.

An alternative account may be provided by the impact of negative reinforcement, as has been suggested to underlie obsessive-compulsive disorder, whereby drug-seeking habits are maintained by the motivation to alleviate or avoid (self-medicate) the negative emotional state and dysregulation resulting from tolerance to, and withdrawal from, drugs taken in increasing amounts over time (Koob & Le Moal 2001). These counter-adaptations are prevalent in the central and extended amygdala and their motivational impact on drug self-administration is described in detail by Koob & Le Moal (2008). These mechanisms are not of course mutually exclusive. Addiction to drugs may reflect a combination of increased incentive motivation mediated by the upregulation of ventral striatal dopamine transmission, by 'hyper-consolidated' habit learning mediated by upregulated dorsal striatum, dopamine-dependent mechanisms and the drive engendered by negative emotional states in extra-striatal networks.

However, we and others have also hypothesized an additional neurobiological mechanism perhaps arising in part as the direct or indirect consequence of toxic drug effects. This mechanism may be implicated in a shift in balance of behavioural control processes from the prefrontal cortex to the striatum, thereby promoting *compulsive* habitual behaviour. There are abundant data suggesting prefrontal cortical, especially orbitofrontal (OFC), dysfunction in addicts, which are also increasingly supported by experimental studies in animals (Schoenbaum *et al.* 2006; Everitt *et al.* 2007; Olausson *et al.* 2007) and humans (see Garavan *et al.* 2008). Thus, in cocaine and methamphetamine abusers, reduced activity of the OFC that correlates with reduced $D_{2/3}$ dopamine receptors in the striatum (Volkow *et al.* 2001), and reduced grey matter volume in this region (Matochik *et al.* 2003) have been reported. There are also growing numbers of reports of impaired behavioural and cognitive functions, including poor behavioural adjustment (Bechara 2005) and impaired probabilistic reversal learning in cocaine abusers (Ersche *et al.* 2008), possibly due to reduced inhibitory control. Deficits have been reported in decision-making cognition on computerized versions of a gambling task, when stimulant abusers chose the most favourable option less frequently than control subjects and chose significantly against the odds in risky conditions, suggesting difficulties in estimating outcome probabilities (Rogers *et al.* 1999; Ersche *et al.* 2005). Similar changes in behaviour are seen in

individuals with OFC damage (Rogers *et al.* 1999) and this has encouraged the view that chronic drug taking may actually be a causal factor in inducing such prefrontal cortex-dependent deficits. But suboptimal prefrontal cortical, including OFC and anterior cingulate cortex, function (Volkow & Fowler 2000; Kaufman *et al.* 2003; Hester & Garavan 2004) may also represent a pre-existing vulnerability trait that results in poor decisions and/or a lack of sensitivity to the consequences of such decisions, and hence drug abuse leading to addiction.

Experimental studies primarily involving psychostimulant treatment of rats and monkeys even after brief periods of exposure have supported the view that disrupted OFC function may indeed be a consequence of toxic drug effects during an addict's history of drug abuse (Jentsch & Taylor 1999; Schoenbaum *et al.* 2006). Short term, usually experimenter-, and not self-, administered cocaine or amphetamine enhanced the development of impulsivity (Jentsch & Taylor 1999; Roesch *et al.* 2007). Reversal learning was impaired by cocaine treatment in monkeys (Jentsch *et al.* 2002) and rats (Schoenbaum *et al.* 2004). Rats having self-administered and then been withdrawn from cocaine exhibited both increased extinction responding and a marked deficit in reversal learning during withdrawal (Calu *et al.* 2007). Schoenbaum and colleagues have emphasized the similarity between OFC lesions and these apparently long-lasting effects of relatively short-term treatment with cocaine, and also showed that the deficit in reversal learning is reflected in a change in the properties of OFC neurons, which do not develop appropriate responses to cues predicting outcomes (Stalnaker *et al.* 2006).

It is remarkable that even brief periods of drug exposure, whether experimenter administered or self-administered, can result in enduring changes in behaviour indicative of OFC dysfunction. However, in the great majority of imaging and neuropsychological investigations of addicts, there has been an exceptionally long history of drug abuse and often poly-drug abuse. These drug-addicted individuals must represent, therefore, a relatively small proportion of the much larger number of individuals in a population who have abused drugs over varying periods of time, but who have not made the transition to an addicted state as characterized by compulsive drug use. Thus, it would seem unlikely that experimental groups of rats receiving the fairly modest exposure to stimulant drugs in the experiments described above would in any sense fulfil the criteria for addiction, yet the changes in behaviour indicative of OFC dysfunction are seen in the entire population of treated experimental animals. It will be important, therefore, to investigate neurobiologically those models that capture chronic drug self-administration (Dalley *et al.* 2005a,b) and the compulsive drug seeking that develops in vulnerable sub-populations of rats if we are to understand the mechanisms underlying the interaction between predisposing behavioural traits and chronic drug exposure in the development of drug addiction.

The majority of the theorizing and evidence summarized above comes from studies of the neural and psychological basis of the seeking and taking

of stimulant drugs such as cocaine. There is clearly a major need for studies of other drugs, especially opiates and alcohol, at both the psychological and neurobiological levels before the generalizability of the mechanisms so defined becomes clear and the gaps in our understanding are filled.

This research was supported by grants from the Medical Research Council and Wellcome Trust and was conducted within the Behavioural and Clinical Neuroscience Institute in the University of Cambridge.

REFERENCES

- Anthony, J. C., Warner, L. A. & Kessler, R. C. 1994 Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants. *Exp. Clin. Psychopharmacol.* **2**, 244–268. (doi:10.1037/1064-1297.2.3.244)
- Bechara, A. 2005 Decision making, impulse control and loss of willpower to resist drugs: a neurocognitive perspective. *Nat. Neurosci.* **8**, 1458–1463. (doi:10.1038/nn1584)
- Belin, D. & Everitt, B. J. 2008 Cocaine seeking habits depend upon dopamine-dependent serial connectivity linking the ventral with the dorsal striatum. *Neuron* **57**, 432–441. (doi:10.1016/j.neuron.2007.12.019)
- Belin, D., Mar, A. C., Dalley, J. W., Robbins, T. W. & Everitt, B. J. 2008 High impulsivity predicts the switch to compulsive cocaine-taking. *Science* **320**, 1352–1355. (doi:10.1126/science.1158136)
- Cador, M., Robbins, T. W. & Everitt, B. J. 1989 Involvement of the amygdala in stimulus-reward associations: interaction with the ventral striatum. *Neuroscience* **30**, 77–86. (doi:10.1016/0306-4522(89)90354-0)
- Calu, D. J., Stalnaker, T. A., Franz, T. F., Singh, T., Shaham, Y. & Schoenbaum, G. 2007 Withdrawal from cocaine self-administration produces long-lasting deficits in orbitofrontal-dependent reversal learning in rats. *Learn. Mem.* **14**, 325–328. (doi:10.1101/lm.534807)
- Cardinal, R. N. & Everitt, B. J. 2004 Neural and psychological mechanisms underlying appetitive learning: links to drug addiction. *Curr. Opin. Neurobiol.* **14**, 156–162. (doi:10.1016/j.conb.2004.03.004)
- Cardinal, R. N., Pennicott, D. R., Sugathapala, C. L., Robbins, T. W. & Everitt, B. J. 2001 Impulsive choice induced in rats by lesions of the nucleus accumbens core. *Science* **292**, 2499–2501. (doi:10.1126/science.1060818)
- Cardinal, R. N., Parkinson, J. A., Hall, J. & Everitt, B. J. 2002 Emotion and motivation: the role of the amygdala, ventral striatum, and prefrontal cortex. *Neurosci. Biobehav. Rev.* **26**, 321–352. (doi:10.1016/S0149-7634(02)00007-6)
- Childress, A. R., Mozley, P. D., McElgin, W., Fitzgerald, J., Reivich, M. & O'Brien, C. P. 1999 Limbic activation during cue-induced cocaine craving. *Am. J. Psychiatry* **156**, 11–18.
- Corbit, L. H., Muir, J. L. & Balleine, B. W. 2001 The role of the nucleus accumbens in instrumental conditioning: evidence of a functional dissociation between accumbens core and shell. *J. Neurosci.* **21**, 3251–3260.
- Czoty, P. W., Morgan, D., Shannon, E. E., Gage, H. D. & Nader, M. A. 2004 Characterization of dopamine D-1 and D-2 receptor function in socially housed cynomolgus monkeys self-administering cocaine. *Psychopharmacology* **174**, 381–388. (doi:10.1007/s00213-003-1752-z)
- Dalley, J. W., Laane, K., Pena, Y., Theobald, D. E. H., Everitt, B. J. & Robbins, T. W. 2005a Attentional and motivational deficits in rats withdrawn from intravenous self-administration of cocaine or heroin. *Psychopharmacology* **182**, 579–587. (doi:10.1007/s00213-005-0107-3)
- Dalley, J. W., Theobald, D. E. H., Berry, D., Milstein, J. A., Laane, K., Everitt, B. J. & Robbins, T. W. 2005b Cognitive sequelae of intravenous amphetamine self-administration in rats: evidence for selective effects on attentional performance. *Neuropsychopharmacology* **30**, 525–537. (doi:10.1038/sj.npp.1300590)
- Dalley, J. W. et al. 2007 Nucleus accumbens D2/3 receptors predict trait impulsivity and cocaine reinforcement. *Science* **315**, 1267–1270. (doi:10.1126/science.1137073)
- Deroche-Gamonet, V., Belin, D. & Piazza, P. V. 2004 Evidence for addiction-like behavior in the rat. *Science* **305**, 1014–1017. (doi:10.1126/science.1099020)
- de Wit, H. & Richards, J. B. 2004 Dual determinants of drug use in humans: reward and impulsivity. In *Motivational factors in the etiology of drug abuse*, vol. 50 (eds R. A. Bevins & M. T. Bardo), pp. 19–56. Lincoln, Nebraska: University of Nebraska Press.
- Di Chiara, G. et al. 2004 Dopamine and drug addiction: the nucleus accumbens shell connection. *Neuropharmacology* **47**, 227–241. (doi:10.1016/j.neuropharm.2004.06.032)
- Di Ciano, P. & Everitt, B. J. 2004 Direct interactions between the basolateral amygdala and nucleus accumbens core underlie cocaine-seeking behavior by rats. *J. Neurosci.* **24**, 7167–7173. (doi:10.1523/JNEUROSCI.1581-04.2004)
- Dickinson, A. 1985 Actions and habits: the development of behavioural autonomy. *Phil. Trans. R. Soc. B* **308**, 67–78. (doi:10.1098/rstb.1985.0010)
- Dickinson, A., Wood, N. & Smith, J. W. 2002 Alcohol seeking by rats: action or habit? *Q. J. Exp. Psychol. B Comp. Physiol. Psychol.* **55**, 331–348. (doi:10.1080/0272499024400016)
- Dom, G., D'Haene, P., Hulstijn, W. & Sabbe, B. 2006 Impulsivity in abstinent early- and late-onset alcoholics: differences in self-report measures and a discounting task. *Addiction* **101**, 50–59. (doi:10.1111/j.1360-0443.2005.01270.x)
- DSM-IV 1994 *Diagnostic and statistical manual of mental disorders*. Washington, DC: American Psychiatric Association.
- Eagle, D. M., Baunez, C., Hutcheson, D. M., Lehmann, O., Shah, A. P. & Robbins, T. W. 2008 Stop-signal reaction-time task performance: role of prefrontal cortex and subthalamic nucleus. *Cereb. Cortex* **18**, 178–188. (doi:10.1093/cercor/bhm044)
- Economidou, D., Pelloux, Y., Dalley, J. W., Robbins, T. W. & Everitt, B. J. 2007 Trait impulsivity as a predictive factor for relapse to drug-seeking behavior in rats. *Behav. Pharmacol.* **18**, S82–S83.
- Ersche, K. D., Fletcher, P. C., Lewis, S. J. G., Clark, L., Stocks-Gee, G., London, M., Deakin, J. B., Robbins, T. W. & Sahakian, B. J. 2005 Abnormal frontal activations related to decision-making in current and former amphetamine and opiate dependent individuals. *Psychopharmacology* **180**, 612–623. (doi:10.1007/s00213-005-2205-7)
- Ersche, K. D., Roiser, J. P., Robbins, T. W. & Sahakian, B. J. 2008 Chronic cocaine, but not chronic amphetamine use is associated with perseverative responding in humans. *Psychopharmacology* **197**, 421–431. (doi:10.1007/s00213-007-1051-1)
- Everitt, B. J. & Robbins, T. W. 2000 Second-order schedules of drug reinforcement in rats and monkeys: measurement of reinforcing efficacy and drug-seeking behaviour. *Psychopharmacology* **153**, 17–30. (doi:10.1007/s002130000566)
- Everitt, B. J. & Robbins, T. W. 2005 Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nat. Neurosci.* **8**, 1481–1489. (doi:10.1038/nn1579)
- Everitt, B. J., Dickinson, A. & Robbins, T. W. 2001 The neuropsychological basis of addictive behaviour. *Brain Res. Rev.* **36**, 129–138. (doi:10.1016/S0165-0173(01)00088-1)

- Everitt, B. J., Hutcheson, D. M., Ersche, K. D., Pelloux, Y., Dalley, J. W. & Robbins, T. W. 2007 The orbital prefrontal cortex and drug addiction in laboratory animals and humans. *Ann. NY Acad. Sci.* **1121**, 576–597. (doi:10.1196/annals.1401.022)
- Faure, A., Haberland, U., Conde, F. & El Massioui, N. 2005 Lesion to the nigrostriatal dopamine system disrupts stimulus–response habit formation. *J. Neurosci.* **25**, 2771–2780. (doi:10.1523/JNEUROSCI.3894-04.2005)
- Ferrario, C. R., Gorny, G., Crombag, H. S., Li, Y. L., Kolb, B. & Robinson, T. E. 2005 Neural and behavioral plasticity associated with the transition from controlled to escalated cocaine use. *Biol. Psychiatry* **58**, 751–759. (doi:10.1016/j.biopsych.2005.04.046)
- Fuchs, R. A., Branham, R. K. & See, R. E. 2006 Different neural substrates mediate cocaine seeking after abstinence versus extinction training: a critical role for the dorso-lateral caudate–putamen. *J. Neurosci.* **26**, 3584–3588. (doi:10.1523/JNEUROSCI.5146-05.2006)
- Garavan, H. *et al.* 2000 Cue-induced cocaine craving: neuroanatomical specificity for drug users and drug stimuli. *Am. J. Psychiatry* **157**, 1789–1798. (doi:10.1176/appi.ajp.157.11.1789)
- Garavan, H., Kaufman, J. N. & Hester, R. 2008 Acute effects of cocaine on the neurobiology of cognitive control. *Phil. Trans. R. Soc. B* **363**, 3267–3276. (doi:10.1098/rstb.2008.0106)
- Grant, S., London, E. D., Newlin, D. B., Villemagne, V. L., Xiang, L., Contoreggi, C., Phillips, R. L., Kimes, A. S. & Margolin, A. 1996 Activation of memory circuits during cue-elicited cocaine craving. *Proc. Natl Acad. Sci. USA* **93**, 12 040–12 045. (doi:10.1073/pnas.93.21.12040)
- Haber, S. N., Fudge, J. L. & McFarland, N. R. 2000 Striatonigral pathways in primates form an ascending spiral from the shell to the dorsolateral striatum. *J. Neurosci.* **20**, 2369–2382.
- Hester, R. & Garavan, H. 2004 Executive dysfunction in cocaine addiction: evidence for discordant frontal, cingulate, and cerebellar activity. *J. Neurosci.* **24**, 11 017–11 022. (doi:10.1523/JNEUROSCI.3321-04.2004)
- Ikemoto, S. 2007 Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens–olfactory tubercle complex. *Brain Res. Rev.* **56**, 27–78. (doi:10.1016/j.brainresrev.2007.05.004)
- Ikemoto, S., Qin, M. & Liu, Z. H. 2005 The functional divide for primary reinforcement of D-amphetamine lies between the medial and lateral ventral striatum: is the division of the accumbens core, shell, and olfactory tubercle valid? *J. Neurosci.* **25**, 5061–5065. (doi:10.1523/JNEUROSCI.0892-05.2005)
- Ito, R., Dalley, J. W., Howes, S. R., Robbins, T. W. & Everitt, B. J. 2000 Dissociation in conditioned dopamine release in the nucleus accumbens core and shell in response to cocaine cues and during cocaine-seeking behavior in rats. *J. Neurosci.* **20**, 7489–7495.
- Ito, R., Dalley, J. W., Robbins, T. W. & Everitt, B. J. 2002 Dopamine release in the dorsal striatum during cocaine-seeking behavior under the control of a drug-associated cue. *J. Neurosci.* **22**, 6247–6253.
- Ito, R., Robbins, T. W. & Everitt, B. J. 2004 Differential control over cocaine-seeking behavior by nucleus accumbens core and shell. *Nat. Neurosci.* **7**, 389–397. (doi:10.1038/nn1217)
- Jentsch, J. D. & Taylor, J. R. 1999 Impulsivity resulting from frontostriatal dysfunction in drug abuse: implications for the control of behavior by reward-related stimuli. *Psychopharmacology* **146**, 373–390. (doi:10.1007/PL00005483)
- Jentsch, J. D., Olsson, P., De la Garza, R. & Taylor, J. R. 2002 Impairments of reversal learning and response perseveration after repeated, intermittent cocaine administrations to monkeys. *Neuropsychopharmacology* **26**, 183–190. (doi:10.1016/S0893-133X(01)00355-4)
- Kaufman, J. N., Ross, T. J., Stein, E. A. & Garavan, H. 2003 Cingulate hypoactivity in cocaine users during a GO–NOGO task as revealed by event-related functional magnetic resonance imaging. *J. Neurosci.* **23**, 7839–7843.
- Kelley, A. E., SmithRoe, S. L. & Holahan, M. R. 1997 Response-reinforcement learning is dependent on N-methyl-D-aspartate receptor activation in the nucleus accumbens core. *Proc. Natl Acad. Sci. USA* **94**, 12 174–12 179. (doi:10.1073/pnas.94.22.12174)
- Koob, G. F. & Le Moal, M. 2001 Drug addiction, dysregulation of reward, and allostasis. *Neuropsychopharmacology* **24**, 97–129. (doi:10.1016/S0893-133X(00)00195-0)
- Koob, G. F. & Le Moal, M. 2005 *Neurobiology of addiction*. San Diego, CA: Academic Press.
- Koob, G. F. & Le Moal, M. 2008 Neurobiological mechanisms for opponent motivational processes in addiction. *Phil. Trans. R. Soc. B* **363**, 3113–3123. (doi:10.1098/rstb.2008.0094)
- Leshner, A. I. 1997 Addiction is a brain disease, and it matters. *Science* **278**, 45–47. (doi:10.1126/science.278.5335.45)
- Letchworth, S. R., Nader, M. A., Smith, H. R., Friedman, D. P. & Porrino, L. J. 2001 Progression of changes in dopamine transporter binding site density as a result of cocaine self-administration in rhesus monkeys. *J. Neurosci.* **21**, 2799–2807.
- Matochik, J. A., London, E. D., Eldreth, D. A., Cadet, J. L. & Bolla, K. I. 2003 Frontal cortical tissue composition in abstinent cocaine abusers: a magnetic resonance imaging study. *Neuroimage* **19**, 1095–1102. (doi:10.1016/S1053-8119(03)00244-1)
- Miles, F. J., Everitt, B. J. & Dickinson, A. 2003 Oral cocaine seeking by rats: action or habit? *Behav. Neurosci.* **117**, 927–938. (doi:10.1037/0735-7044.117.5.927)
- Moore, R. J., Vinsant, S. L., Nader, M. A., Porrino, L. J. & Friedman, D. P. 1998 Effect of cocaine self-administration on dopamine D-2 receptors in rhesus monkeys. *Synapse* **30**, 88–96. (doi:10.1002/(SICI)1098-2396(199809)30:1<88::AID-SYN11>3.0.CO;2-L)
- Mukherjee, J. *et al.* 1999 Preliminary assessment of extrastriatal dopamine D-2 receptor binding in the rodent and nonhuman primate brains using the high affinity radioligand, ¹⁸F-fallypride. *Nucl. Med. Biol.* **26**, 519–527. (doi:10.1016/S0969-8051(99)00012-8)
- Nader, M. A., Daunais, J. B., Moore, T., Nader, S. H., Moore, R. J., Smith, H. R., Friedman, D. P. & Porrino, L. J. 2002 Effects of cocaine self-administration on striatal dopamine systems in rhesus monkeys: initial and chronic exposure. *Neuropsychopharmacology* **27**, 35–46. (doi:10.1016/S0893-133X(01)00427-4)
- Nader, M. A., Morgan, D., Gage, H. D., Nader, S. H., Calhoun, T. L., Buchheimer, N., Ehrenkauf, R. & Mach, R. H. 2006 PET imaging of dopamine D₂ receptors during chronic cocaine self-administration in monkeys. *Nat. Neurosci.* **9**, 1050–1056. (doi:10.1038/nn1737)
- Nader, M. A., Czoty, P. W., Gould, R. W. & Riddick, N. V. 2008 Positron emission tomography imaging studies of dopamine receptors in primate models of addiction. *Phil. Trans. R. Soc. B* **363**, 3223–3232. (doi:10.1098/rstb.2008.0092)
- Nelson, A. & Killcross, S. 2006 Amphetamine exposure enhances habit formation. *J. Neurosci.* **26**, 3805–3812. (doi:10.1523/JNEUROSCI.4305-05.2006)
- Nestler, E. J. 2004 Molecular mechanisms of drug addiction. *Neuropharmacology* **47**, 24–32. (doi:10.1016/j.neuropharm.2004.06.031)

- O'Brien, C. P. & McLellan, A. T. 1996 Myths about the treatment of addiction. *Lancet* **347**, 237–240. (doi:10.1016/S0140-6736(96)90409-2)
- Olausson, P., Jentsch, J. D., Krueger, D. D., Tronson, N. C., Nairn, A. C. & Taylor, J. R. 2007 Orbitofrontal cortex and cognitive-motivational impairments in psychostimulant addiction. Linking affect to action: critical contributions of the orbitofrontal cortex. *Ann. N. Y. Acad. Sci.* **1121**, 610–638. (doi:10.1196/annals.1401.016)
- Olmstead, M. C., Lafond, M. V., Everitt, B. J. & Dickinson, A. 2001 Cocaine seeking by rats is a goal-directed action. *Behav. Neurosci.* **115**, 394–402. (doi:10.1037/0735-7044.115.2.394)
- Pelloux, Y., Everitt, B. J. & Dickinson, A. 2007 Compulsive drug seeking by rats under punishment: effects of drug taking history. *Psychopharmacology* **194**, 127–137. (doi:10.1007/s00213-007-0805-0)
- Piazza, P. V., Deminière, J. M., Le Moal, M. & Simon, H. 1989 Factors that predict individual vulnerability to amphetamine self-administration. *Science* **245**, 1511–1513. (doi:10.1126/science.2781295)
- Porrino, L. J., Daunais, J. B., Smith, H. R. & Nader, M. A. 2004 The expanding effects of cocaine: studies in a nonhuman primate model of cocaine self-administration. *Neurosci. Biobehav. Rev.* **27**, 813–820. (doi:10.1016/j.neubiorev.2003.11.013)
- Robbins, T. W. & Everitt, B. J. 1999 Drug addiction: bad habits add up. *Nature* **398**, 567–570. (doi:10.1038/19208)
- Robinson, T. E. & Berridge, K. C. 1993 The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res. Rev.* **18**, 247–291. (doi:10.1016/0165-0173(93)90013-P)
- Robinson, T. E. & Berridge, K. C. 2008 The incentive sensitization theory of addiction: some current issues. *Phil. Trans. R. Soc. B* **363**, 3137–3146. (doi:10.1098/rstb.2008.0093)
- Robinson, E. S. J., Eagle, D. M., Mar, A. C., Bari, A., Banerjee, G., Jiang, X. S., Dalley, J. W. & Robbins, T. W. 2008 Similar effects of the selective noradrenaline reuptake inhibitor atomoxetine on three distinct forms of impulsivity in the rat. *Neuropsychopharmacology* **33**, 1028–1037. (doi:10.1038/sj.npp.1301487)
- Roesch, M. R., Takahashi, Y., Gugs, N., Bissonette, G. B. & Schoenbaum, G. 2007 Previous cocaine exposure makes rats hypersensitive to both delay and reward magnitude. *J. Neurosci.* **27**, 245–250. (doi:10.1523/JNEUROSCI.4080-06.2007)
- Rogers, R. D. et al. 1999 Dissociable deficits in the decision-making cognition of chronic amphetamine abusers, opiate abusers, patients with focal damage to prefrontal cortex, and tryptophan-depleted normal volunteers: evidence for monoaminergic mechanisms. *Neuropsychopharmacology* **20**, 322–339. (doi:10.1016/S0893-133X(98)00091-8)
- Schoenbaum, G., Saddoris, M. P., Ramus, S. J., Shaham, Y. & Setlow, B. 2004 Cocaine-experienced rats exhibit learning deficits in a task sensitive to orbitofrontal cortex lesions. *Eur. J. Neurosci.* **19**, 1997–2002. (doi:10.1111/j.1460-9568.2004.03274.x)
- Schoenbaum, G., Roesch, M. R. & Stalnaker, T. A. 2006 Orbitofrontal cortex, decision-making and drug addiction. *Trends Neurosci.* **29**, 116–124. (doi:10.1016/j.tins.2005.12.006)
- See, R. E., Elliott, J. C. & Feltenstein, M. W. 2007 The role of dorsal vs ventral striatal pathways in cocaine-seeking behavior following prolonged abstinence in rats. *Psychopharmacology* **195**, 321–331. (doi:10.1007/s00213-007-0850-8)
- Stalnaker, T. A., Roesch, M. R., Franz, T. M., Burke, K. A. & Schoenbaum, G. 2006 Abnormal associative encoding in orbitofrontal neurons in cocaine-experienced rats during decision-making. *Eur. J. Neurosci.* **24**, 2643–2653. (doi:10.1111/j.1460-9568.2006.05128.x)
- Takahashi, Y., Roesch, M. R., Stalnaker, T. A. & Schoenbaum, G. 2007 Cocaine shifts the balance of cue-evoked firing from ventral to dorsal striatum. See <http://frontiersin.org/neuroscience/abstract/10.3389/neuro.07/011.2007>.
- Tiffany, S. T. 1990 A cognitive model of drug urges and drug-use behavior: role of automatic and non-automatic processes. *Psychol. Rev.* **97**, 146–168. (doi:10.1037/0033-295X.97.2.147)
- Vanderschuren, L. J. M. J. & Everitt, B. J. 2004 Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science* **305**, 1017–1019. (doi:10.1126/science.1098975)
- Vanderschuren, L., Di Ciano, P. & Everitt, B. J. 2005 Involvement of the dorsal striatum in cue-controlled cocaine seeking. *J. Neurosci.* **25**, 8665–8670. (doi:10.1523/JNEUROSCI.0925-05.2005)
- Verdejo-Garcia, A. & Perez-Garcia, M. 2007 Ecological assessment of executive functions in substance dependent individuals. *Drug Alcohol Depend.* **90**, 48–55. (doi:10.1016/j.drugalcdep.2007.02.010)
- Vezina, P. 2004 Sensitization of midbrain dopamine neuron reactivity and the self-administration of psychomotor stimulant drugs. *Neurosci. Biobehav. Rev.* **27**, 827–839. (doi:10.1016/j.neubiorev.2003.11.001)
- Volkow, N. D. & Fowler, J. S. 2000 Addiction, a disease of compulsion and drive: involvement of the orbitofrontal cortex. *Cereb. Cortex* **10**, 318–325. (doi:10.1093/cercor/10.3.318)
- Volkow, N. D. & Wise, R. A. 2005 How can drug addiction help us understand obesity? *Nat. Neurosci.* **8**, 555–560. (doi:10.1038/nn1452)
- Volkow, N. D. et al. 2001 Low level of brain dopamine D-2 receptors in methamphetamine abusers: association with metabolism in the orbitofrontal cortex. *Am. J. Psychiatry* **158**, 2015–2021. (doi:10.1176/appi.ajp.158.12.2015)
- Volkow, N. D., Wang, G. J., Telang, F., Fowler, J. S., Logan, J., Childress, A. R., Jayne, M., Ma, Y. M. & Wong, C. 2006 Cocaine cues and dopamine in dorsal striatum: mechanism of craving in cocaine addiction. *J. Neurosci.* **26**, 6583–6588. (doi:10.1523/JNEUROSCI.1544-06.2006)
- Whitelaw, R. B., Markou, A., Robbins, T. W. & Everitt, B. J. 1996 Excitotoxic lesions of the basolateral amygdala impair the acquisition of cocaine-seeking behaviour under a second-order schedule of reinforcement. *Psychopharmacology* **127**, 213–224.
- Wise, R. A. 2004 Dopamine, learning and motivation. *Nat. Rev. Neurosci.* **5**, 483–494. (doi:10.1038/nrn1406)
- Yin, H. H., Knowlton, B. J. & Balleine, B. W. 2004 Lesions of dorsolateral striatum preserve outcome expectancy but disrupt habit formation in instrumental learning. *Eur. J. Neurosci.* **19**, 181–189. (doi:10.1111/j.1460-9568.2004.03095.x)
- Yin, H. H., Ostlund, S. B., Knowlton, B. J. & Balleine, B. W. 2005 The role of the dorsomedial striatum in instrumental conditioning. *Eur. J. Neurosci.* **22**, 513–523. (doi:10.1111/j.1460-9568.2005.04218.x)
- Yin, H. H., Knowlton, B. J. & Balleine, B. W. 2006 Inactivation of dorsolateral striatum enhances sensitivity to changes in the action–outcome contingency in instrumental conditioning. *Behav. Brain Res.* **166**, 189–196. (doi:10.1016/j.bbr.2005.07.012)
- Zilberman, M. L., Tavares, H., Hodgins, D. C. & El-Guebaly, N. 2007 The impact of gender, depression, and personality on craving. *J. Addict. Dis.* **26**, 79–84. (doi:10.1300/J069v26n01_10)