

Type-1 metabotropic glutamate receptor in cerebellar Purkinje cells: a key molecule responsible for long-term depression, endocannabinoid signalling and synapse elimination

Masanobu Kano*, Kouichi Hashimoto† and Toshihide Tabata‡

Department of Cellular Neuroscience, Graduate School of Medicine, Osaka University, 2-2 Yamada-oka, Suita, Osaka 565-0871, Japan

The cerebellum is a brain structure involved in the coordination, control and learning of movements, and elucidation of its function is an important issue. Japanese scholars have made seminal contributions in this field of neuroscience. Electrophysiological studies of the cerebellum have a long history in Japan since the pioneering works by Ito and Sasaki. Elucidation of the basic circuit diagram of the cerebellum in the 1960s was followed by the construction of cerebellar network theories and finding of their neural correlates in the 1970s. A theoretically predicted synaptic plasticity, long-term depression (LTD) at parallel fibre to Purkinje cell synapse, was demonstrated experimentally in 1982 by Ito and co-workers. Since then, Japanese neuroscientists from various disciplines participated in this field and have made major contributions to elucidate molecular mechanisms underlying LTD. An important pathway for LTD induction is type-1 metabotropic glutamate receptor (mGluR1) and its downstream signal transduction in Purkinje cells. Sugiyama and co-workers demonstrated the presence of mGluRs and Nakanishi and his pupils identified the molecular structures and functions of the mGluR family. Moreover, the authors contributed to the discovery and elucidation of several novel functions of mGluR1 in cerebellar Purkinje cells. mGluR1 turned out to be crucial for the release of endocannabinoid from Purkinje cells and the resultant retrograde suppression of transmitter release. It was also found that mGluR1 and its downstream signal transduction in Purkinje cells are indispensable for the elimination of redundant synapses during post-natal cerebellar development. This article overviews the seminal works by Japanese neuroscientists, focusing on mGluR1 signalling in cerebellar Purkinje cells.

Keywords: metabotropic glutamate receptor; long-term depression; endocannabinoid; synapse elimination; Purkinje cell; cerebellum

1. INTRODUCTION

The cerebellum is a brain structure involved in the coordination, control and learning of movements, and elucidation of its function is an important issue in neuroscience. Japan has a long history of electrophysiological study of the cerebellum particularly since the pioneering work by two pupils of Sir John Eccles, Masao Ito and Kazuo Sasaki. After return from Eccles' laboratory in Canberra, Australia, Masao Ito discovered that the Purkinje cell, the principal cell type solely constituting the output of the cerebellar cortex, is an inhibitory neuron

(Ito & Yoshida 1964, 1966; Ito *et al.* 1964, 1966) and uses γ -aminobutyric acid (GABA) as its neurotransmitter (Obata *et al.* 1967). Since it was believed in the early 1960s that all principal neurons were excitatory and only local interneurons were inhibitory, Ito's discovery that the principal neuron of the cerebellar cortex is inhibitory was a great surprise. By using a microelectrode technique, Sasaki studied the neuronal circuit of the cerebellar cortex very intensively with Eccles, Llinas and Strata in Canberra. They clarified the synaptic actions of four types of interneurons in the cerebellar cortex (i.e. granule, basket, stellate and Golgi cells (Eccles *et al.* 1966*b,c,d,e,f*, 1967*b*)), as well as those of mossy and climbing fibres, the two major inputs to the cerebellar cortex (Eccles *et al.* 1966*a,d*, 1967*c*; Sasaki & Strata 1967). By the end of the 1960s, the basic neuronal circuit of the cerebellum was dissected in detail both electrophysiologically and morphologically, as documented in the monographs (Eccles *et al.* 1967*a*; Palay & Chan-Palay 1974; Ito 1984). Purkinje cells in the cerebellar cortex receive two distinct excitatory inputs, namely parallel and climbing fibres. Parallel fibres are the axons of granule cells in the

* Author and address for correspondence: Department of Neurophysiology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. (mkano-tky@m.u-tokyo.ac.jp).

† Present Address: Department of Neurophysiology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

‡ Present Address: Faculty of Engineering, The University of Toyama, 3190 Gofuku, Toyama 930-8555, Japan.

One contribution of 17 to a Theme Issue 'Japan: its tradition and hot topics in biological sciences'.

cerebellar cortex and form synapses on the spines of the Purkinje cell's dendrites. Synaptic inputs from individual parallel fibres are weak, but the number of parallel fibres innervating a single Purkinje cell counts as many as 100 000–200 000. Granule cells are driven by the excitatory inputs from mossy fibres originating from various precerebellar nuclei such as the pontine nuclei. On the other hand, climbing fibres originate from the inferior olive in the contralateral medulla and form contacts directly on Purkinje cells. In contrast to parallel fibres, only one climbing fibre innervates a single Purkinje cell in the adult cerebellum but each climbing fibre makes strong synaptic contacts on the Purkinje cell's proximal dendrites. Purkinje cells in turn form inhibitory synaptic contacts on neurons in the deep cerebellar and vestibular nuclei. After the dissection of the basic neuronal network of the cerebellum, the main interest of researchers shifted to uncovering the elaborate neuronal mechanisms of the cerebellum and defining its roles in the entire nervous system function. Ito proceeded in the former direction and discovered long-term depression (LTD) as detailed below, while Sasaki advanced in the latter direction and studied the roles of the cerebro-cerebellar communication loop.

2. LONG-TERM DEPRESSION

(a) *Discovery of LTD*

Marr proposed in his epoch-making network theory of the cerebellar cortex that conjunctive activation of certain parallel and climbing fibre synapses on the same Purkinje cell leads to long-term potentiation of these parallel fibre synapses (Marr 1969). Later, Albus suggested that depression rather than potentiation is expected to occur following conjunctive parallel and climbing fibre activation, in view of the stable operation of the system (Albus 1971). Ito and co-workers found that in the flocculus, an evolutionally old part of the cerebellum, mossy fibres arising from vestibular organs and climbing fibres conveying visual information from the retina converge on Purkinje cells (Fukuda *et al.* 1972; Ito 1972; Maekawa & Simpson 1973). Inspired by the theoretical works by Marr (1969) and Albus (1971), Ito proposed a hypothesis that the cerebellar flocculus adaptively controls the gain of the vestibulo-ocular reflex depending on the retinal error signals conveyed by the climbing fibre inputs to the flocculus (Ito 1970, 1972, 1982). This 'flocculus hypothesis' is based on plasticity assumption, i.e. the strength of parallel fibre to Purkinje cell synapse is modified when these synapses are activated conjunctively with climbing fibre synaptic inputs. It has been shown that, during the adaptation of vestibulo-ocular reflex, Purkinje cell firing frequency in the flocculus undergoes changes in the manner predicted by Albus' hypothesis (Ghelarducci *et al.* 1975; Dufosse *et al.* 1978). As well, Gilbert and Thach showed that, in monkeys adapting to compensate for a sudden change in arm load, Purkinje cells changed their firings in the manner predicted by Albus' hypothesis (Gilbert & Thach 1977). Despite these lines of circumstantial evidence, the direct demonstration of the plasticity at parallel fibre to Purkinje cell synapse did not occur until the demonstration of LTD by Ito *et al.* (1982). In decerebrate rabbits, they showed

that the responses of flocculus Purkinje cells to vestibular nerve stimulation undergoes LTD after the vestibular nerve stimulation is applied conjunctively with the stimulation of the contralateral inferior olive at 4 Hz for 25 s (Ito *et al.* 1982). Moreover, it turned out that LTD can be induced by the repeated stimulation of parallel fibres in conjunction with climbing fibres in decerebrate rabbits (Ito & Kano 1982; Ekerot & Kano 1985, 1989). These observations confirmed the validity of the Marr–Albus–Ito's synaptic plasticity assumption. Subsequently, Sakurai demonstrated LTD in cerebellar slice preparations from rat (Sakurai 1987) and that LTD requires an elevation of intracellular Ca^{2+} level in the Purkinje cell (Sakurai 1990). Hirano demonstrated LTD in cultured Purkinje cells (Hirano 1990). The demonstration of LTD in such reduced preparations paved the way for the researches seeking for molecular mechanisms of LTD in the 1990s.

Because parallel fibres were thought to use glutamate as their neurotransmitter, Kano and Kato explored the subtype of glutamate receptors involved in LTD (Kano & Kato 1987). In the middle of the 1980s, it was thought that glutamate receptors were classified into at least three subtypes: *N*-methyl-D-aspartate (NMDA), quisqualate, and kainate receptors (Foster & Fagg 1988). The clear concept of metabotropic glutamate receptor was not yet established at that time while it had already been shown that glutamate stimulation can enhance inositol phosphate turnover in nervous tissues (Sladeczek *et al.* 1985; Nicoletti *et al.* 1986). In addition, many subtype-specific antagonists that are widely used today were not developed. Therefore, we examined which of the glutamate agonists available at that time could induce LTD of parallel fibre to Purkinje cell synaptic transmission after an agonist was iontophoretically applied to parallel fibre synapses in conjunction with climbing fibre stimulation. The order of potency to induce LTD turned out to be quisqualate > glutamate >>> kainite and aspartate (Kano & Kato 1987). This order of preference did not match that of any glutamate receptor subtype known at that time (Foster & Fagg 1988). Therefore, this result indicated that LTD induction requires a novel subtype of glutamate receptor that is highly sensitive to quisqualate.

(b) *Identification of metabotropic glutamate receptor*

Sugiyama *et al.* (1987) demonstrated the presence of metabotropic glutamate receptor that is coupled to G protein, which activates inositol phospholipid metabolism and mobilizes Ca^{2+} from internal stores. When injected with mRNA from rat brain, *Xenopus* oocytes expressed a novel type of glutamate receptor that activated Ca^{2+} -dependent Cl currents due to inositol trisphosphate (IP_3)-induced Ca^{2+} release from internal stores. Masu *et al.* (1991) reported the cloning and characterization of cDNA for type-1 metabotropic glutamate receptor (mGluR1) that is coupled to IP_3 / Ca^{2+} signal transduction. Importantly, mGluR1 is richly expressed in cerebellar Purkinje cells and has an agonist preference of the order of quisqualate > glutamate >>> kainite (Masu *et al.* 1991; Aramori & Nakanishi 1992). Therefore, mGluR1 appeared to correspond to the 'quisqualate receptor' demonstrated

to be involved in cerebellar LTD (Kano & Kato 1987). After the identification of mGluR1, seven other members of mGluRs have been identified, and they are classified into three groups (Conn & Pin 1997). Group I consists of mGluR1 and mGluR5 and is coupled to the G protein G_q family (G_q and G_{11}) that mediates IP_3 -induced Ca^{2+} mobilization and the activation of protein kinase C (PKC). Group II consists of mGluR2 and mGluR3 and is coupled to the $G_{i/o}$ protein. Group III consists of mGluR4, mGluR6, mGluR7 and mGluR8 and is also coupled to the $G_{i/o}$ protein.

(c) Requirement of mGluR1 in LTD induction

In the meantime, Linden *et al.* (1991) showed pharmacologically that quisqualate-preferring glutamate receptors were involved in the LTD of the glutamate responsiveness of cultured cerebellar Purkinje cells. Crepel *et al.* (1991) also reported that an mGluR agonist induced LTD-like persistent reduction of parallel fibre to Purkinje cell synaptic transmission in cerebellar slice preparation. Shigemoto *et al.* (1994) raised antibodies against mGluR1, which functionally block mGluR1-mediated Ca^{2+} mobilization. They showed that these antibodies blocked LTD of the glutamate responsiveness of cultured cerebellar Purkinje cells. Meanwhile, Atsu Aiba produced an mGluR1 knockout mouse in Susumu Tonegawa's laboratory. In collaboration with Kano, Aiba observed that LTD is deficient in the mGluR1 knockout mice, yet the morphology of the cerebellum and the basic properties of excitatory synaptic transmission were apparently normal (Aiba *et al.* 1994). In addition, Aiba *et al.* (1994) observed that the mGluR1 knockout mice were ataxic, displayed motor discoordination, and were impaired in delay eyeblink conditioning, a paradigm known to require intact cerebellum and thought to represent learning of an elementary movement (discrete motor learning). Therefore, the mGluR1 knockout mouse is the first mouse model whose phenotypes support the notion that LTD at parallel fibre to Purkinje cell synapse is a cellular basis of cerebellum-dependent discrete motor learning. Conquet *et al.* (1994) also reported in the same year that cerebellar LTD was deficient in mice lacking mGluR1. After he returned to Japan, Aiba and co-workers created a mouse strain in which mGluR1 is expressed only in cerebellar Purkinje cells (Ichise *et al.* 2000). They introduced rat mGluR1 α (full-length mGluR1 splice variant) into the mGluR1 knockout mouse by using a Purkinje cell-specific promoter, L7. This mGluR1-rescue mouse exhibits apparently normal motor coordination, although the mGluR1 expression level is approximately 1/40 of that of the wild-type (Ichise *et al.* 2000). Importantly, both cerebellar LTD (Ichise *et al.* 2000) and delay eyeblink conditioning (Kishimoto *et al.* 2002) are restored in the mGluR1-rescue mouse, indicating that mGluR1 in Purkinje cell is responsible for cerebellar LTD and discrete motor learning.

Then, how is mGluR1 activated by synaptic activity? Immunoelectron microscopic examinations demonstrate that mGluR1 is localized in the perisynaptic annuli of the Purkinje cell dendritic spines facing parallel fibre synaptic terminals (Baude *et al.* 1993; Nusser *et al.* 1994; Petralia *et al.* 1998; Lopez-Bendito *et al.* 2001).

Therefore, glutamate released from parallel fibre synaptic terminals is expected to activate mGluR1 at the spine annuli. Consistent with this expectation, strong repetitive stimulation of parallel fibres was shown to evoke an mGluR1-mediated excitatory postsynaptic potential (EPSP) in the Purkinje cells (Batchelor *et al.* 1994; Batchelor & Garthwaite 1997). In 1998, two groups independently demonstrated that a brief train of parallel fibre stimulation induces mGluR1-mediated local Ca^{2+} release from the internal stores that is confined to the dendritic region receiving activated parallel fibre inputs (Finch & Augustine 1998; Takechi *et al.* 1998). These results clearly indicate that mGluR1 can be activated by parallel fibre activity at Purkinje cell dendritic spines. Therefore, mGluR1 is thought to be activated at parallel fibre synapses during conjunctive activation of parallel and climbing fibre inputs for inducing LTD.

(d) Modulation of mGluR1 signalling may influence LTD induction

Glutamate is thought to be the primary activator of mGluRs *in vivo*. However, it was clarified that mGluR1 activity could be modulated by extracellular factors other than glutamate. Kubo *et al.* (1998) found an unexpected phenomenon that mGluRs can be activated by extracellular Ca^{2+} in the absence of glutamate. Tabata *et al.* (2002) have demonstrated that extracellular Ca^{2+} significantly broadens the dynamic range of mGluR1-mediated responses to its agonist in cultured Purkinje cells. Hirono *et al.* (2001b) reported that the mGluR1-mediated responses of Purkinje cells in cerebellar slices were enhanced by the activation of type-B γ -aminobutyric acid receptor ($GABA_B$ R). This effect was mediated by the $G_{i/o}$ protein (Hirono *et al.* 2001b). $GABA_B$ R is localized perisynaptically at excitatory parallel fibre synapses on Purkinje cell dendritic spines where mGluR1 is concentrated (Kulik *et al.* 2002). In addition to the $G_{i/o}$ protein-mediated enhancement, Tabata *et al.* (2004) have shown that $GABA_B$ R activation enhances the mGluR1-mediated responses of cultured Purkinje cells through the $G_{i/o}$ protein-independent direct interaction between $GABA_B$ R and mGluR1. Surprisingly, this $GABA_B$ R–mGluR1 interaction occurs in the absence of GABA but is caused by extracellular Ca^{2+} (Tabata *et al.* 2004), indicating that $GABA_B$ R can act as Ca^{2+} -dependent cofactor of mGluR1 signalling in the Purkinje cells. These data suggest that LTD induction may be influenced by the $GABA_B$ R activity through $G_{i/o}$ protein-dependent and/or -independent pathway. In support of this hypothesis, Tabata *et al.* (2004) showed that blockade of $GABA_B$ R reduced the magnitude of LTD in cultured Purkinje cells. Furthermore, more direct evidence has been recently provided by Kamikubo *et al.* (2007) that in cultured Purkinje cells, $GABA_B$ R activation enhanced LTD in a $G_{i/o}$ protein-dependent manner.

Besides the interplay with $GABA_B$ R, mGluR1 has been reported to interact with $G_{i/o}$ protein-coupled A_1 -subtype adenosine receptor (A_1 R; Tabata *et al.* 2007). Tabata *et al.* reported that the agonists for A_1 R induced continuous depression of an mGluR1-coupled inward current. This inhibitory effect from A_1 R to mGluR1 was independent of $G_{i/o}$ protein, suggesting a

direct interaction between the two receptors. This AIR-mGluR1 interaction might influence the induction of LTD (Tabata *et al.* 2007).

Jin *et al.* (2007) recently reported that mGluR1-mediated slow excitatory postsynaptic current (EPSC) and its coincident Ca^{2+} transient after a brief burst of parallel fibres were selectively and persistently depressed by repeated climbing fibre-evoked depolarization of Purkinje cells. This 'LTD' of mGluR1 responses (LTD(mGluR1)) blocked subsequent induction of LTD of AMPA receptor-mediated EPSC at parallel fibre-Purkinje cell synapses (Jin *et al.* 2007). Thus, LTD(mGluR1) may have a metaplastic function, blocking induction of AMPA receptor LTD.

(e) Signalling molecules downstream of mGluR1 required for LTD induction

The mGluR1 downstream signalling cascade involved in LTD induction was studied by using mutant mice lacking the candidate molecules. mGluR1 is coupled to the G protein G_q family (G_q and G_{11}), which activates the β isoforms of phospholipase C (PLC), produces diacylglycerol (DG) and IP_3 , and leads to DG-mediated activation of PKC and IP_3 -mediated Ca^{2+} release from internal stores (Conn & Pin 1997). Immunohistochemical studies indicate that $G\alpha_q$ and $G\alpha_{11}$ are concentrated in the perisynaptic region of the Purkinje cell dendritic spines where mGluR1 is localized (Tanaka *et al.* 2000). Hartmann *et al.* (2004) have demonstrated that a $G\alpha_q$ -knockout mouse exhibits clear ataxia, deficient mGluR1-mediated Ca^{2+} mobilization, deficient LTD and impaired motor learning. By contrast, a $G\alpha_{11}$ knockout mouse has no apparent ataxia and normal mGluR1-mediated Ca^{2+} mobilization but exhibits mild impairment of LTD and mild motor deficit (Hartmann *et al.* 2004). These results can be attributed to the fact that the expression level of $G\alpha_q$ is more than 10-fold higher than that of $G\alpha_{11}$ in the Purkinje cells (Hartmann *et al.* 2004) despite their overlapping subcellular localization (Tanaka *et al.* 2000). These results unequivocally indicate that both $G\alpha_q$ and $G\alpha_{11}$ mediate mGluR1 signalling and LTD induction within Purkinje cells.

In situ hybridization and immunohistochemical studies indicate that Purkinje cells in the rostral cerebellum mainly express PLC β_4 , whereas those in the caudal cerebellum mainly have PLC β_3 (Kano *et al.* 1998; Nakamura *et al.* 2004). Miyata *et al.* (2001) analysed the PLC β_4 knockout mice and found that both mGluR1-mediated Ca^{2+} mobilization and LTD were deficient in the rostral cerebellum, whereas these responses were intact in the caudal cerebellum. Hirono *et al.* (2001a) confirmed the observation by Miyata *et al.* by analysing the other strain of PLC β_4 knockout mice. Importantly, Miyata *et al.* (2001) showed that delay eyeblink conditioning was severely impaired in the PLC β_4 knockout mice. Since lobule simplex or lobule HVI of Larsell, a part of the cerebellar cortex with rich PLC β_4 expression in the Purkinje cells, is crucial for delay eyeblink conditioning (Hesslow & Yeo 1998), these results indicate that PLC β_4 is crucial for LTD and discrete motor learning in the rostral cerebellum.

Several pharmacological data in the 1990s suggest that PKC is involved in LTD induction (Crepel & Jaillard 1990; Linden & Connor 1991; Hartell 1994; Freeman *et al.* 1998). Importantly, a transgenic mouse expressing a PKC inhibitor peptide, PKC-[19–31] in the Purkinje cells displays no LTD (De Zeeuw *et al.* 1998). Among the four classical PKC isoforms (α , β I, β II and γ) whose activation requires Ca^{2+} elevation, PKC α , PKC β I and PKC γ are strongly expressed in the Purkinje cells of the entire cerebellum (Hirono *et al.* 2001a). Although PKC γ knockout mice display ataxia and a defect in developmental synapse elimination (see below; Kano *et al.* 1995), LTD was intact in their Purkinje cells (Chen *et al.* 1995). Therefore, PKC α and/or PKC β I may be the probable candidate that plays a role in LTD induction.

IP_3 -induced Ca^{2+} release is also shown to be required for LTD induction in cerebellar slices (Hemart *et al.* 1995; Khodakhah & Armstrong 1997; Inoue *et al.* 1998; Daniel *et al.* 1999; Miyata *et al.* 2000), although it is not required for LTD of glutamate responsiveness in certain cultured Purkinje cell preparations (Narasimhan *et al.* 1998). Several studies indicate that pharmacological blockade of IP_3 receptors or depletion of Ca^{2+} stores blocks LTD or LTD-like synaptic depression in cerebellar slices (Hemart *et al.* 1995; Khodakhah & Armstrong 1997; Inoue *et al.* 1998). Mikoshiba and co-workers created a type-1 IP_3 receptor knockout mouse that exhibits seizure and ataxia and mostly dies during the first three post-natal weeks. LTD is clearly deficient in the Purkinje cells in cerebellar slices from the type-1 IP_3 receptor knockout mice (Inoue *et al.* 1998). Furthermore, Miyata *et al.* (2000) demonstrated that IP_3 -mediated Ca^{2+} signalling in the Purkinje cell dendritic spines and concomitantly LTD were absent in those mice and rats with mutations in myosin Va, in which the endoplasmic reticulum did not enter the dendritic spines. LTD was restored in these animals by photolysis of caged Ca^{2+} compound in the Purkinje cells (Miyata *et al.* 2000). Daniel *et al.* (1999) showed that photolytic release of IP_3 restored LTD in the mGluR1 knockout mice and this LTD was sensitive to a PKC inhibitor. These results unequivocally indicate that IP_3 -induced local Ca^{2+} release within Purkinje cell dendritic spines is crucial for LTD induction.

Figure 1 summarizes and schematically illustrates how mGluR1 and its downstream signal transduction cascade are involved in LTD induction. Other receptors and signalling molecules such as glutamate receptor δ_2 subunit (GluR δ_2) and nitric oxide known to be involved in LTD are not illustrated here for simplicity. The details of signal transduction underlying LTD are described in recent reviews (Ito 2001, 2002). However, the putative final common path to LTD is included in figure 1. It has been demonstrated that LTD results from the internalization of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA)-type glutamate receptors from the postsynaptic membrane of the parallel fibre to Purkinje cell synapses (Matsuda *et al.* 2000; Wang & Linden 2000; Xia *et al.* 2000). This internalization is caused by the PKC-dependent phosphorylation of Ser880 of AMPA receptor GluR2 subunit. This leads to the unbinding of GluR2 from glutamate receptor-interacting protein (GRIP), which usually anchors AMPA receptors




Figure 1. *Contribution of mGluR1 to LTD induction.* LTD is induced following conjunctive activation of parallel and climbing fibre synapses. During LTD induction, glutamate released from parallel fibre terminals binds to mGluR1 at the perisynaptic region of Purkinje cell dendritic spines. The molecular cascade involving G_q and PLCβ4 is activated in the spines, and DG and IP₃ are produced. IP₃ triggers Ca²⁺ release from internal stores in the spines. Climbing fibre-induced depolarization of Purkinje cell dendrites opens voltage-gated Ca²⁺ channels (VGCC) and induces Ca²⁺ influx. As a result of these two events, local Ca²⁺ levels in the spines are elevated such that they can activate PKC together with DG. PKC phosphorylates Ser880 of the GluR2 subunit of AMPA receptor, and then the phosphorylated AMPA receptors are internalized. GABA_BR activated presumably by extracellular Ca²⁺ and/or ambient GABA spilt over from neighbouring inhibitory synapses enhances mGluR1 signalling through its direct interaction with mGluR1 and/or through the action of G_{i/o} protein. AMPAR, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor.

to cytoskeleton lining the postsynaptic membrane (Matsuda *et al.* 2000). As described above, Japanese neuroscientists have made major contributions to the elucidation of the LTD signal transduction cascade as illustrated in figure 1.

3. ENDOCANNABINOID SIGNALLING

(a) *Retrograde synaptic suppression triggered by postsynaptic mGluR1 activation*

It has been shown that bath application of mGluR1 agonists to a cerebellar slice induces presynaptic inhibition of excitatory inputs to Purkinje cells (Conquet *et al.* 1994; Maejima *et al.* 2001). However, since mGluR1 is not detected at excitatory presynaptic terminals in immunoelectron microscopic studies (Baude *et al.* 1993; Nusser *et al.* 1994; Petralia *et al.* 1998; Lopez-Bendito *et al.* 2001), how mGluR1 agonists affect glutamate release from presynaptic terminals was a puzzling problem. Maejima *et al.* (2001) at our laboratory injected a non-hydrolyzable analogue of GTP or GDP (GTP γ -S or GDP β -S) into the recorded Purkinje cell through a patch-clamp recording pipette and inactivated the G-protein signalling. This manipulation completely eliminated the presynaptic inhibition of climbing fibre-mediated EPSCs induced by the mGluR1 agonist 3,5-dihydroxyphenylglycine (DHPG; Maejima *et al.* 2001). This result unequivocally indicates that mGluR1 signalling in the postsynaptic Purkinje cell is responsible for the presynaptic inhibition. Therefore,

some retrograde messenger must exist that transmits signals from Purkinje cells to climbing fibre terminals. After a 3-year struggle, we have finally revealed that antagonists of cannabinoid CB1 receptor abolish the DHPG-induced presynaptic suppression (Maejima *et al.* 2001), indicating that endogenous cannabinoids (endocannabinoids) mediate the retrograde signal.

(b) *An endocannabinoid, 2-arachidonoylglycerol, as a retrograde messenger*

Five months prior to Maejima's report, the three groups including the authors' laboratory published independently and simultaneously that strong depolarization and the resultant elevation of intracellular Ca²⁺ concentration in hippocampal neurons (Ohno-Shosaku *et al.* 2001; Wilson & Nicoll 2001) and cerebellar Purkinje cells (Kretzner & Regehr 2001) produce endocannabinoids, which in turn retrogradely suppress transmitter release from inhibitory and excitatory presynaptic terminals. These phenomena, termed depolarization-induced suppression of inhibition (DSI) and excitation (DSE), can be triggered by Ca²⁺ elevation alone without the activation of mGluRs. By contrast, mGluR1-mediated endocannabinoid release occurs even in the Purkinje cells dialysed with the Ca²⁺ chelator BAPTA (Maejima *et al.* 2001), indicating that mGluR1 activation produces endocannabinoids without the elevation of Ca²⁺. These results indicate that there are two distinct modes of endocannabinoid release from




Figure 2. *Contribution of mGluR1 to endocannabinoid signalling.* A brief burst of parallel fibre stimulation causes the release of a sufficient amount of glutamate to activate mGluR1 and induce Ca^{2+} influx through voltage-gated Ca^{2+} channels opened by AMPA receptor-mediated local depolarization. Then, G_q and Ca^{2+} synergistically activate PLC β 4. When climbing fibre stimulation is paired with parallel fibre stimulation, a shorter burst (e.g. two pulses) is sufficient to activate PLC β 4 because the Ca^{2+} flowing into the dendrites by climbing fibre-induced depolarization further sensitizes PLC β 4. Activated PLC β 4 produces DG, which in turn is converted to the endocannabinoid 2-AG by diacylglycerol lipase (DGL). 2-AG is released from the postsynaptic membrane and retrogradely binds to CB1 receptors at the parallel fibre terminals. Activated CB1 suppresses the release of glutamate presumably inhibiting voltage-gated Ca^{2+} channels and release machinery in the presynaptic terminals. AMPAR, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor.

neurons. Later, it was shown that Ca^{2+} elevation and activation of $\text{G}_{q/11}$ -coupled receptors cooperatively produce endocannabinoids in hippocampal neurons (Ohno-Shosaku *et al.* 2002, 2003; Varma *et al.* 2002). Hashimoto *et al.* (2005) at our laboratory have revealed that this cooperativity is attributable to PLC β activity that depends on both intracellular Ca^{2+} and $\text{G}_{q/11}$ protein. In cerebellar Purkinje cells, a brief burst of parallel fibre stimulation causes mGluR1 activation and local Ca^{2+} elevation due to AMPA receptor-mediated depolarization, and induces endocannabinoid-mediated retrograde suppression of glutamate release from parallel fibres (Maejima *et al.* 2001; Brown *et al.* 2003; Marcaggi & Attwell 2005, 2007). We have demonstrated that this suppression results from the Ca^{2+} dependency of PLC β 4 (Maejima *et al.* 2005). We also show that mGluR1 activation combined with depolarization leads to the production of an endocannabinoid, 2-arachidonoyl glycerol (2-AG) (Maejima *et al.* 2005), which is known to be produced from DG by diacylglycerol lipase (Sugiura *et al.* 2006). Brenowitz and Regehr observed that coactivation of parallel and climbing fibres markedly facilitated endocannabinoid-mediated suppression of the parallel fibre synapses, presumably owing to the cooperativity of parallel fibre-induced mGluR1 activation and climbing fibre-induced Ca^{2+} elevation (Brenowitz & Regehr 2005). Thus, endocannabinoid-mediated retrograde suppression, dependent on both mGluR1 and Ca^{2+} , may constitute a local negative feedback loop to prevent the hyper-excitation of Purkinje cells. Figure 2

schematically illustrates how this putative negative feedback loop works at parallel fibre to Purkinje cell synapses.

(c) *Endocannabinoid signalling involved in cerebellar LTD and motor learning*

Endocannabinoid signalling is also involved in the LTD of parallel fibre to Purkinje cell synapses. Safo and Regehr observed that LTD was blocked by a CB1 receptor antagonist and inhibiting diacylglycerol lipase, and LTD was deficient in a CB1 knockout mouse (Safo & Regehr 2005). Although the CB1 knockout mouse exhibits no apparent deficit in motor coordination, Kishimoto & Kano (2006) found that delay eyeblink conditioning was severely impaired in these mice and in wild-type mice injected with a CB1 antagonist. These results suggest that mGluR1-dependent endocannabinoid signalling is required for LTD induction and cerebellum-dependent discrete motor learning.

4. SYNAPSE ELIMINATION

(a) *Maturation of climbing fibre synapses during post-natal development*

A widely accepted scheme of functional neural circuit formation during post-natal development is that immature neurons initially make synaptic connections not only to their final targets, but also to other neurons. Then, functionally important synapses are strengthened, and less important synapses are weakened relative to the important ones. The weakened synapses are finally

eliminated morphologically (Purves & Lichtman 1980; Lichtman & Colman 2000). The climbing fibre to Purkinje cell synapse is a typical example in the CNS, which undergoes such changes during post-natal development. In early post-natal days, all Purkinje cells are innervated by multiple climbing fibres (multiple innervation) (Crepel *et al.* 1976; Crepel 1982; Lohof *et al.* 1996). These surplus climbing fibres are eliminated eventually with the progress of post-natal development, and most Purkinje cells become innervated by single climbing fibres (mono-innervation) by the end of the third post-natal week in mice (Kano *et al.* 1995, 1997, 1998; Offermanns *et al.* 1997).

In the mature cerebellum, a single climbing fibre forms synapses on the proximal dendrites of a Purkinje cell. Stimulation of such a mono-innervating climbing fibre elicits very large EPSPs that are enough to activate voltage-gated Ca^{2+} channels over whole dendrites (Konnerth *et al.* 1992; Miyakawa *et al.* 1992; Hashimoto *et al.* 2001a). By contrast, multiple climbing fibres initially form synapses around the somata of Purkinje cells in newborn mice (Chedotal & Sotelo 1993; Morando *et al.* 2001). EPSCs elicited by stimulating such multiply innervating climbing fibres are much smaller than those of mature climbing fibres (Kano *et al.* 1995, 1997, 1998; Hashimoto & Kano 2003). Therefore climbing fibre inputs become stronger during post-natal development, while redundant climbing fibres are eliminated during the same period. Mariani and Changeux showed that some Purkinje cells had two climbing fibre-mediated EPSPs, the amplitudes of which were quite different around P10–P13 (Mariani & Changeux 1981), suggesting that only one climbing fibre is strengthened relative to others prior to the completion of synapse elimination.

We analysed changes in the relative synaptic strengths of multiple climbing fibres innervating the same Purkinje cell in mice (Hashimoto & Kano 2003). In the Purkinje cells from newborn mice, the amplitudes of climbing fibre-mediated EPSCs (CF-EPSCs) are similar, indicating that multiple climbing fibres have similar strengths. In the Purkinje cells from mice of approximately two weeks of age, one CF-EPSC is much larger than other few CF-EPSCs. Our detailed quantitative analysis of the developmental course indicates that the disparity among multiply innervating CF-EPSC amplitudes progressively becomes larger from P3 to P6 and reaches a plateau after P7 (Hashimoto & Kano 2003). A morphological study by Sugihara has shown that the innervation pattern of climbing fibres over Purkinje cells drastically changes during this post-natal period in rats. At P4, climbing fibres have many creeping terminals in the Purkinje cell layer and their swellings do not aggregate at particular Purkinje cell somata (creeper type). Then, from P4 to P7, climbing fibres surround several specific Purkinje cell somata and form aggregated terminals on them (nest type; Sugihara 2005). This result indicates that one of the multiple climbing fibres innervating the same Purkinje cell is strengthened during the first post-natal week (Hashimoto & Kano 2003).

By contrast, Scelfo and Strata reported a conflicting result that the disparity among multiply innervating CF-EPSC amplitudes becomes smaller from

P4 to P7, and then progressively increases from P7 to P10 (Scelfo & Strata 2005). The reason for the apparent discrepancy between the two reports is not clear. One possibility is a difference between preparations and/or techniques used in these studies. Scelfo and Strata reported that the average number of climbing fibres innervating individual Purkinje cells at P4–P7 was approximately 3.3 in CD1 mice. This value is much smaller than that of our measurements in C57BL/6 mice (approx. 5.9; Hashimoto & Kano 2005). This may be attributed to the difference in mouse strain (Scelfo & Strata 2005). Alternately, Scelfo and Strata might have missed more climbing fibres than we might have when searching for multiple climbing fibres innervating individual Purkinje cells.

We examined the physiological parameters of CF-EPSCs among the strongest and weaker climbing fibre inputs in individual multiply innervated Purkinje cells as well as those of the single climbing fibre inputs in mono-innervated Purkinje cells. The size of glutamate transients at the synaptic clefts from weaker climbing fibres were consistently smaller than those for the strongest climbing fibres (Hashimoto & Kano 2003). This difference is attributable to the fact that the number of simultaneously released vesicles is smaller (i.e. the probability of multivesicular release is lower in weaker climbing fibres) when compared with the strongest climbing fibre innervating the same Purkinje cell or climbing fibres of mono-innervated Purkinje cells. The occurrence of multivesicular release is affected by the release probability and the number of release sites. We show that the paired-pulse ratio, which reflects release probability, is not different among the three types of climbing fibres in a condition in which postsynaptic AMPA receptors should not be saturated (i.e. with a low external Ca^{2+} concentration; Hashimoto & Kano 2003). Therefore, it is concluded that the number of functional release sites facing a narrow postsynaptic region of Purkinje cell is smaller in the weaker climbing fibres than in the strongest climbing fibre.

In addition to functional differences, we show that the innervation territories over Purkinje cell dendrites are different between the strongest and weaker climbing fibres. We also show that Ca^{2+} signals elicited by stimulating the strongest climbing fibres spread over the whole dendritic trees, indicating that these climbing fibres form synaptic contacts in a similar manner to mono-innervating mature climbing fibres in the adult (Konnerth *et al.* 1992; Miyakawa *et al.* 1992; Hashimoto *et al.* 2001a). By contrast, Ca^{2+} signals elicited by stimulating weaker climbing fibres were confined to the most proximal parts of the dendritic trees (Hashimoto & Kano 2003). This result indicates that synapses formed by weaker climbing fibres are confined to the proximal part of dendrites near the soma.

(b) Elimination of redundant climbing fibre synapses

Earlier studies in the 1970s and 1980s clarified that multiple innervation persists into adulthood in several spontaneously occurring mutant mice that are devoid of granule cells or have a defect in parallel fibre synaptogenesis. These mice include *reeler* (Mariani *et al.* 1977), *weaver* (Crepel & Mariani 1976;

Puro & Woodward 1977) and *staggerer* (Crepel *et al.* 1980; Mariani & Changeux 1980). The multiple innervation also persists in the animals whose migrating granule cells are destroyed by X-ray irradiation (Woodward *et al.* 1974; Crepel *et al.* 1981) or by infection of a mink enteritis virus (Benoit *et al.* 1987). These results suggest that the regression of surplus climbing fibres is critically dependent on the normal formation of granule cell to Purkinje cell connections (Crepel 1982; Lohof *et al.* 1996). Moreover, an analysis of the developmental course of multiple innervation in X-ray irradiated rats revealed that there are at least two distinct phases in climbing fibre elimination process (Crepel *et al.* 1981). In these rats, initial climbing fibre innervation and subsequent elimination until P8 are normal, but further regression of climbing fibres does not progress in the following period (Crepel *et al.* 1981). This indicates that the early phase of synapse elimination is independent of the formation of granule cell to Purkinje cell connection, but the late phase is critically dependent on it. This notion is confirmed by the analysis of a GluR δ 2 knockout mouse. GluR δ 2 belongs to a family of ionotropic glutamate receptors, which is expressed selectively in Purkinje cells (Araki *et al.* 1993; Lomeli *et al.* 1993; Takayama *et al.* 1996; Landsend *et al.* 1997). In the GluR δ 2 null mutant mice, the density of parallel fibre synapses is less than half of that of the wild-type mice, whereas the abnormalities in cerebellar structures, Purkinje cell dendrites and granule cell density are mild (Kashiwabuchi *et al.* 1995; Kurihara *et al.* 1997). Kashiwabuchi *et al.* (1995) showed that the climbing fibre synapse elimination was impaired, supporting that parallel fibre synapse formation is essential for the climbing fibre synapse elimination. In addition, Hashimoto and co-workers revealed that the spatial pattern of climbing fibre innervation over Purkinje cell dendrites was abnormal in the GluR δ 2 knockout mice. In the adult cerebellum, parallel fibres and a climbing fibre normally make synapses on distal and proximal dendrites, respectively. In the GluR δ 2 knockout mice, climbing fibres invade into the distal dendrites and form ectopic synapses there (Hashimoto *et al.* 2001a; Ichikawa *et al.* 2002). These ectopic climbing fibre synapses appear around P10 when parallel fibre synapse formation and Purkinje cell dendritic arborization occur most vigorously. A similar type of multiple climbing fibre innervation was also found in a mutant mouse deficient in cerebellin in which parallel fibre to Purkinje cell synapse formation is severely impaired (Hirai *et al.* 2005). These results suggest that parallel fibres compete with climbing fibres for the innervation territory during development and play a role in restricting climbing fibre innervation to proximal dendrites.

(c) Parallel fibres activate mGluR1 signalling cascades for driving the regression of surplus climbing factor synapses

Another role of parallel fibre synapse is to activate mGluR1 and its downstream signalling cascades in the Purkinje cell to drive the process of climbing fibre synapse elimination. Kano, Hashimoto and co-workers show that the mutant mice deficient in mGluR1, G α _q, PLC β 4 or PKC γ are all impaired in climbing fibre

synapse elimination (Kano *et al.* 1995, 1997, 1998; Offermanns *et al.* 1997; Hashimoto *et al.* 2000, 2001b; Ichise *et al.* 2000). Levenes *et al.* (1997) also reported incomplete synapse elimination in the mGluR1 knockout mice. In these mice, the regression of climbing fibre synapse normally occurs until around P10, indicating that the early phase of climbing fibre synapse elimination (Crepel *et al.* 1981) is independent of these signalling molecules. However, these mice display an abnormality in synapse elimination thereafter. These results suggest that the signalling cascade from mGluR1 to PKC γ is essential for the late phase of climbing fibre synapse elimination. Importantly, the formation and function of parallel fibre to Purkinje cell synapses are normal in these mutant mice, indicating that the multiple climbing fibre innervation is not caused secondarily by the defect in parallel fibre synaptogenesis.

The defect of the climbing fibre synapse elimination in the mGluR1 knockout mice is restored in mGluR1-rescue mice in which mGluR1a has been introduced specifically into Purkinje cells (Ichise *et al.* 2000). Regression of climbing fibre synapses is impaired in mice by Purkinje cell-specific expression of a PKC inhibitor peptide (De Zeeuw *et al.* 1998). Furthermore, the distribution of multiply innervated Purkinje cells in the PLC β 4 knockout mouse cerebellum exactly matches that of the Purkinje cells predominantly expressing PLC β 4 in the wild-type mouse cerebellum (Kano *et al.* 1998; Hashimoto *et al.* 2000). These lines of evidence clearly indicate that the signalling from mGluR1 to PKC γ in Purkinje cells but not other cell types plays a central role in mediating climbing fibre synapse elimination.

Although mGluR1 is concentrated at perisynaptic annuli of the postsynaptic Purkinje cell spines facing parallel and climbing fibre terminals (Baude *et al.* 1993; Nusser *et al.* 1994; Petralia *et al.* 1998; Lopez-Bendito *et al.* 2001), mGluR1 activation at parallel fibre synapses appears to be important for the late phase of climbing fibre synapse elimination. This could be correlated to the fact that mGluR1 can readily be activated by parallel fibre inputs (Finch & Augustine 1998; Takechi *et al.* 1998) but hardly by climbing fibre inputs without blockade of glutamate transporters (Dzubay & Otis 2002). Furthermore, we demonstrate that chronic blockade of NMDA receptors within the cerebellum specifically impairs the late phase of climbing fibre synapse elimination (Kakizawa *et al.* 2000). Because NMDA receptors are not present at excitatory synapses of Purkinje cells but are abundantly expressed at mossy fibre to granule cell synapses, the chronic blockade of NMDA receptors within the cerebellum should affect mossy fibre to granule cell transmission. These results suggest that neural activity along mossy fibre (granule cell) and parallel fibre (Purkinje cell) pathway, and subsequent activation of mGluR1 is a prerequisite for the late phase of climbing fibre synapse elimination (Kakizawa *et al.* 2000).

Figure 3 illustrates the current model for mechanisms underlying the late phase of climbing fibre synapse elimination. First, parallel fibre synapses confine the climbing fibre innervation sites to the proximal dendrites of Purkinje cells. Second, parallel fibre




Figure 3. *Contribution of mGluR1 to synapse elimination during cerebellar development.* Elimination of redundant climbing fibre synapses on Purkinje cells consists of two phases, the early phase (until P10) and the late phase (P10–P20) in mice. The mGluR1 signal is required for the late phase. Neural activity along mossy fibre (granule cell) and parallel fibre (Purkinje cell) pathway, which is gated by NMDA receptor at the mossy fibre to granule cell synapses, drives the mGluR1 signalling cascades in Purkinje cells. Subsequently, the downstream signalling cascade involving G_q , PLC β_4 and PKC γ is activated and, through unknown mechanisms, it facilitates the elimination of weaker climbing fibre synapses while leaving stronger climbing fibre synapses intact. AMPAR, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; NMDAR, *N*-methyl-D-aspartate receptor.

activity involving NMDA receptor at mossy fibre to granule cell synapses drives mGluR1 to PKC γ signalling cascades in the Purkinje cells to eliminate weaker climbing fibres. The signalling molecules downstream of PKC γ are currently unknown. Furthermore, we know little about the molecular mechanisms underlying the morphological elimination of the weaker climbing fibres. Some mechanisms must convey trans-synaptic retrograde signalling from Purkinje cells to weaker climbing fibres. One possibility would be that some cytokines such as insulin-like growth factor 1 (IGF-1) supports climbing fibre synapses (Kakizawa *et al.* 2003); the climbing fibres that do not receive a large enough amount of such cytokines may undergo regression. Another possibility would be that climbing fibre synapses may be maintained by protein–protein interaction between surface-expressed proteins such as cell adhesion molecules; disruption of such interaction may lead to the regression of climbing fibres.

5. CONCLUSION

In this article, we have made an overview of the three functional roles of mGluR1 in cerebellar Purkinje cells: LTD at parallel fibre to Purkinje cell synapses; endocannabinoid-mediated retrograde suppression of transmitter release; and climbing fibre synapse elimination during post-natal development. Japanese neuroscientists have made seminal contributions to the discovery and the elucidation of the mechanisms of these phenomena. Based on the theory that LTD

underlies cerebellar motor learning, Ito demonstrated LTD experimentally and paved a way to the subsequent researches on LTD. The presence of metabotropic glutamate receptor was first demonstrated by Sugiyama and co-workers, and the molecular identity and functions of mGluR family receptors were clarified by Nakanishi and his pupils. In collaboration with Aiba who created the mGluR1 knockout mouse and the mGluR1-rescue mouse, we elucidated that mGluR1 signal transduction in the Purkinje cell is crucial for LTD induction. We also discovered the mGluR1-mediated endocannabinoid release and resultant retrograde suppression of transmitter release. Finally, we show that mGluR1 signal transduction in the Purkinje cell is indispensable for the elimination of redundant climbing fibre synapses that occurs during second and third post-natal weeks in mice. The massive outcome in this field is due to the efforts of Japanese scientists from various disciplines including neurophysiology, neuroanatomy, neuropharmacology, molecular biology, computational neuroscience and genetic engineering.

We thank D. Kawakami for preparing the figures. This work was supported by grants-in-aid for scientific research 17023021 and 17100004 (M.K.), 17700305 and 18019022 (T.T.) and 16680014 (K.H.) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

REFERENCES

Aiba, A., Kano, M., Chen, C., Stanton, M. E., Fox, G. D., Herrup, K., Zwingman, T. A. & Tonegawa, S. 1994

- Deficient cerebellar long-term depression and impaired motor learning in mGluR1 mutant mice. *Cell* **79**, 377–388. (doi:10.1016/0092-8674(94)90204-6)
- Albus, J. S. 1971 A theory of cerebellar function. *Math. Biosci.* **10**, 25–61. (doi:10.1016/0025-5564(71)90051-4)
- Araki, K., Meguro, H., Kushiya, E., Takayama, C., Inoue, Y. & Mishina, M. 1993 Selective expression of the glutamate receptor channel delta 2 subunit in cerebellar Purkinje cells. *Biochem. Biophys. Res. Commun.* **197**, 1267–1276. (doi:10.1006/bbrc.1993.2614)
- Aramori, I. & Nakanishi, S. 1992 Signal transduction and pharmacological characteristics of a metabotropic glutamate receptor, mGluR1, in transfected CHO cells. *Neuron* **8**, 757–765. (doi:10.1016/0896-6273(92)90096-V)
- Batchelor, A. M. & Garthwaite, J. 1997 Frequency detection and temporally dispersed synaptic signal association through a metabotropic glutamate receptor pathway. *Nature* **385**, 74–77. (doi:10.1038/385074a0)
- Batchelor, A. M., Madge, D. J. & Garthwaite, J. 1994 Synaptic activation of metabotropic glutamate receptors in the parallel fibre-Purkinje cell pathway in rat cerebellar slices. *Neuroscience* **63**, 911–915. (doi:10.1016/0306-4522(94)90558-4)
- Baude, A., Nusser, Z., Roberts, J. D., Mulvihill, E., McIlhinney, R. A. & Somogyi, P. 1993 The metabotropic glutamate receptor (mGluR1 alpha) is concentrated at perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction. *Neuron* **11**, 771–787. (doi:10.1016/0896-6273(93)90086-7)
- Benoit, P., Mariani, J., Delhay-Bouchaud, N. & Chappuis, G. 1987 Evidence for a multiple innervation of cerebellar Purkinje cells by climbing fibers in adult ferrets infected at birth by a mink enteritis virus. *Brain Res.* **431**, 51–57.
- Brenowitz, S. D. & Regehr, W. G. 2005 Associative short-term synaptic plasticity mediated by endocannabinoids. *Neuron* **45**, 419–431. (doi:10.1016/j.neuron.2004.12.045)
- Brown, S. P., Brenowitz, S. D. & Regehr, W. G. 2003 Brief presynaptic bursts evoke synapse-specific retrograde inhibition mediated by endogenous cannabinoids. *Nat. Neurosci.* **6**, 1048–1057. (doi:10.1038/nn1126)
- Chedotal, A. & Sotelo, C. 1993 The 'creeper stage' in cerebellar climbing fiber synaptogenesis precedes the 'pericellular nest'-ultrastructural evidence with parvalbumin immunocytochemistry. *Brain Res. Dev. Brain Res.* **76**, 207–220. (doi:10.1016/0165-3806(93)90209-S)
- Chen, C., Kano, M., Abeliovich, A., Chen, L., Bao, S., Kim, J. J., Hashimoto, K., Thompson, R. F. & Tonegawa, S. 1995 Impaired motor coordination correlates with persistent multiple climbing fiber innervation in PKC gamma mutant mice. *Cell* **83**, 1233–1242. (doi:10.1016/0092-8674(95)90148-5)
- Conn, P. J. & Pin, J. P. 1997 Pharmacology and functions of metabotropic glutamate receptors. *Annu. Rev. Pharmacol. Toxicol.* **37**, 205–237. (doi:10.1146/annurev.pharmtox.37.1.205)
- Conquet, F. et al. 1994 Motor deficit and impairment of synaptic plasticity in mice lacking mGluR1. *Nature* **372**, 237–243. (doi:10.1038/372237a0)
- Crepel, F. 1982 Regression of functional synapses in the immature mammalian cerebellum. *Trends Neurosci.* **5**, 266–269. (doi:10.1016/0166-2236(82)90168-0)
- Crepel, F. & Jaillard, D. 1990 Protein kinases, nitric oxide and long-term depression of synapses in the cerebellum. *Neuroreport* **1**, 133–136. (doi:10.1097/00001756-199010000-00013)
- Crepel, F. & Mariani, J. 1976 Multiple innervation of Purkinje cells by climbing fibers in the cerebellum of the Weaver Mutant Mouse. *J. Neurobiol.* **7**, 579–582. (doi:10.1002/neu.480070610)
- Crepel, F., Mariani, J. & Delhay-Bouchaud, N. 1976 Evidence for a multiple innervation of Purkinje cells by climbing fibers in the immature rat cerebellum. *J. Neurobiol.* **7**, 567–578. (doi:10.1002/neu.480070609)
- Crepel, F., Delhay-Bouchaud, N., Guastavino, J. M. & Sampaio, I. 1980 Multiple innervation of cerebellar Purkinje cells by climbing fibres in staggerer mutant mouse. *Nature* **283**, 483–484. (doi:10.1038/283483a0)
- Crepel, F., Delhay-Bouchaud, N. & Dupont, J. L. 1981 Fate of the multiple innervation of cerebellar Purkinje cells by climbing fibers in immature control, X-irradiated and hypothyroid rats. *Brain Res.* **227**, 59–71.
- Crepel, F., Daniel, H., Hemart, N. & Jaillard, D. 1991 Effects of ACPD and AP3 on parallel-fibre-mediated EPSPs of Purkinje cells in cerebellar slices *in vitro*. *Exp. Brain Res.* **86**, 402–406. (doi:10.1007/BF00228964)
- Daniel, H., Levenes, C., Fagni, L., Conquet, F., Bockaert, J. & Crepel, F. 1999 Inositol-1,4,5-trisphosphate-mediated rescue of cerebellar long-term depression in subtype 1 metabotropic glutamate receptor mutant mouse. *Neuroscience* **92**, 1–6. (doi:10.1016/S0306-4522(99)00136-0)
- De Zeeuw, C. I., Hansel, C., Bian, F., Koekkoek, S. K., van Alphen, A. M., Linden, D. J. & Oberdick, J. 1998 Expression of a protein kinase C inhibitor in Purkinje cells blocks cerebellar LTD and adaptation of the vestibulo-ocular reflex. *Neuron* **20**, 495–508. (doi:10.1016/S0896-6273(00)80990-3)
- Dufosse, M., Ito, M., Jastreboff, P. J. & Miyashita, Y. 1978 A neuronal correlate in rabbit's cerebellum to adaptive modification of the vestibulo-ocular reflex. *Brain Res.* **150**, 611–616. (doi:10.1016/0006-8993(78)90825-9)
- Dzubay, J. A. & Otis, T. S. 2002 Climbing fiber activation of metabotropic glutamate receptors on cerebellar purkinje neurons. *Neuron* **36**, 1159–1167. (doi:10.1016/S0896-6273(02)01052-8)
- Eccles, J. C., Llinas, R. & Sasaki, K. 1966a The excitatory synaptic action of climbing fibres on the Purkinje cells of the cerebellum. *J. Physiol.* **182**, 268–296.
- Eccles, J. C., Llinas, R. & Sasaki, K. 1966b The inhibitory interneurons within the cerebellar cortex. *Exp. Brain Res.* **1**, 1–16.
- Eccles, J. C., Llinas, R. & Sasaki, K. 1966c Intracellularly recorded responses of the cerebellar Purkinje cells. *Exp. Brain Res.* **1**, 161–183.
- Eccles, J. C., Llinas, R. & Sasaki, K. 1966d The mossy fibre-granule cell relay of the cerebellum and its inhibitory control by Golgi cells. *Exp. Brain Res.* **1**, 82–101.
- Eccles, J. C., Llinas, R. & Sasaki, K. 1966e Parallel fibre stimulation and the responses induced thereby in the Purkinje cells of the cerebellum. *Exp. Brain Res.* **1**, 17–39.
- Eccles, J. C., Sasaki, K. & Strata, P. 1966f The profiles of physiological events produced by a parallel fibre volley in the cerebellar cortex. *Exp. Brain Res.* **2**, 18–34. (doi:10.1007/BF00234358)
- Eccles, J. C., Ito, M. & Szentagothai, J. 1967a *The cerebellum as a neuronal machine*. Berlin, Germany; New York, NY: Springer.
- Eccles, J. C., Sasaki, K. & Strata, P. 1967b A comparison of the inhibitory actions of Golgi cells and of basket cells. *Exp. Brain Res.* **3**, 81–94.
- Eccles, J. C., Sasaki, K. & Strata, P. 1967c Interpretation of the potential fields generated in the cerebellar cortex by a mossy fibre volley. *Exp. Brain Res.* **3**, 58–80.
- Ekerot, C. F. & Kano, M. 1985 Long-term depression of parallel fibre synapses following stimulation of climbing fibres. *Brain Res.* **342**, 357–360. (doi:10.1016/0006-8993(85)91136-9)
- Ekerot, C. F. & Kano, M. 1989 Stimulation parameters influencing climbing fibre induced long-term depression of parallel fibre synapses. *Neurosci. Res.* **6**, 264–268. (doi:10.1016/0168-0102(89)90065-5)

- Finch, E. A. & Augustine, G. J. 1998 Local calcium signalling by inositol-1,4,5-trisphosphate in Purkinje cell dendrites. *Nature* **396**, 753–756. (doi:10.1038/25541)
- Foster, A. C. & Fagg, G. E. 1988 Acidic amino acid receptor nomenclature: time for change. *Trends Neurosci.* **11**, 17–18. (doi:10.1016/0166-2236(88)90043-4)
- Freeman Jr, J. H., Shi, T. & Schreurs, B. G. 1998 Pairing-specific long-term depression prevented by blockade of PKC or intracellular Ca²⁺. *Neuroreport* **9**, 2237–2241.
- Fukuda, J., Highstein, S. M. & Ito, M. 1972 Cerebellar inhibitory control of the vestibulo-ocular reflex investigated in rabbit 3rd nucleus. *Exp. Brain Res.* **14**, 511–526. (doi:10.1007/BF00236593)
- Ghelarducci, B., Ito, M. & Yagi, N. 1975 Impulse discharges from flocculus Purkinje cells of alert rabbits during visual stimulation combined with horizontal head rotation. *Brain Res.* **87**, 66–72. (doi:10.1016/0006-8993(75)90780-5)
- Gilbert, P. F. & Thach, W. T. 1977 Purkinje cell activity during motor learning. *Brain Res.* **128**, 309–328. (doi:10.1016/0006-8993(77)90997-0)
- Hartell, N. A. 1994 cGMP acts within cerebellar Purkinje cells to produce long term depression via mechanisms involving PKC and PKG. *Neuroreport* **5**, 833–836.
- Hartmann, J. et al. 2004 Distinct roles of Galpha(q) and Galpha11 for Purkinje cell signaling and motor behavior. *J. Neurosci.* **24**, 5119–5130. (doi:10.1523/JNEUROSCI.4193-03.2004)
- Hashimoto, K. & Kano, M. 2003 Functional differentiation of multiple climbing fiber inputs during synapse elimination in the developing cerebellum. *Neuron* **38**, 785–796. (doi:10.1016/S0896-6273(03)00298-8)
- Hashimoto, K. & Kano, M. 2005 Postnatal development and synapse elimination of climbing fiber to Purkinje cell projection in the cerebellum. *Neurosci. Res.* **53**, 221–228. (doi:10.1016/j.neures.2005.07.007)
- Hashimoto, K. et al. 2000 Climbing fiber synapse elimination during postnatal cerebellar development requires signal transduction involving G alpha q and phospholipase C beta 4. *Prog. Brain Res.* **124**, 31–48.
- Hashimoto, K. et al. 2001a Roles of glutamate receptor delta 2 subunit (GluRdelta 2) and metabotropic glutamate receptor subtype 1 (mGluR1) in climbing fiber synapse elimination during postnatal cerebellar development. *J. Neurosci.* **21**, 9701–9712.
- Hashimoto, K., Miyata, M., Watanabe, M. & Kano, M. 2001b Roles of phospholipase Cbeta4 in synapse elimination and plasticity in developing and mature cerebellum. *Mol. Neurobiol.* **23**, 69–82. (doi:10.1385/MN:23:1:69)
- Hashimoto, Y., Ohno-Shosaku, T., Tsubokawa, H., Ogata, H., Emoto, K., Maejima, T., Araishi, K., Shin, H. S. & Kano, M. 2005 Phospholipase Cb serves as a coincidence detector through its Ca²⁺ dependency for triggering retrograde endocannabinoid signal. *Neuron* **45**, 257–268. (doi:10.1016/j.neuron.2005.01.004)
- Hemart, N., Daniel, H., Jaillard, D. & Crepel, F. 1995 Receptors and second messengers involved in long-term depression in rat cerebellar slices *in vitro*: a reappraisal. *Eur. J. Neurosci.* **7**, 45–53. (doi:10.1111/j.1460-9568.1995.tb01019.x)
- Hesslow, G. & Yeo, C. 1998 Cerebellum and learning: a complex problem. *Science* **280**, 1817–1819. (doi:10.1126/science.280.5371.1815e)
- Hirai, H. et al. 2005 Cbln1 is essential for synaptic integrity and plasticity in the cerebellum. *Nat. Neurosci.* **8**, 1534–1541. (doi:10.1038/nn1576)
- Hirano, T. 1990 Depression and potentiation of the synaptic transmission between a granule cell and a Purkinje cell in rat cerebellar culture. *Neurosci. Lett.* **119**, 141–144. (doi:10.1016/0304-3940(90)90818-T)
- Hirono, M. et al. 2001a Phospholipase Cbeta4 and protein kinase Calpha and/or protein kinase CbetaI are involved in the induction of long term depression in cerebellar Purkinje cells. *J. Biol. Chem.* **276**, 45 236–45 242. (doi:10.1074/jbc.M105413200)
- Hirono, M., Yoshioka, T. & Konishi, S. 2001b GABA(B) receptor activation enhances mGluR-mediated responses at cerebellar excitatory synapses. *Nat. Neurosci.* **4**, 1207–1216. (doi:10.1038/nn764)
- Ichikawa, R., Miyazaki, T., Kano, M., Hashikawa, T., Tatsumi, H., Sakimura, K., Mishina, M., Inoue, Y. & Watanabe, M. 2002 Distal extension of climbing fiber territory and multiple innervation caused by aberrant wiring to adjacent spiny branchlets in cerebellar Purkinje cells lacking glutamate receptor delta 2. *J. Neurosci.* **22**, 8487–8503.
- Ichise, T., Kano, M., Hashimoto, K., Yanagihara, D., Nakao, K., Shigemoto, R., Katsuki, M. & Aiba, A. 2000 mGluR1 in cerebellar Purkinje cells essential for long-term depression, synapse elimination, and motor coordination. *Science* **288**, 1832–1835. (doi:10.1126/science.288.5472.1832)
- Inoue, T., Kato, K., Kohda, K. & Mikoshiba, K. 1998 Type 1 inositol 1,4,5-trisphosphate receptor is required for induction of long-term depression in cerebellar Purkinje neurons. *J. Neurosci.* **18**, 5366–5373.
- Ito, M. 1970 Neurophysiological aspects of the cerebellar motor control system. *Int. J. Neurol.* **7**, 162–176.
- Ito, M. 1972 Neural design of the cerebellar motor control system. *Brain Res.* **40**, 81–84. (doi:10.1016/0006-8993(72)90110-2)
- Ito, M. 1982 Cerebellar control of the vestibulo-ocular reflex-around the flocculus hypothesis. *Annu. Rev. Neurosci.* **5**, 275–296. (doi:10.1146/annurev.ne.05.030182.001423)
- Ito, M. 1984 *The cerebellum and neural control*. New York, NY: Raven Press.
- Ito, M. 2001 Cerebellar long-term depression: characterization, signal transduction, and functional roles. *Physiol. Rev.* **81**, 1143–1195.
- Ito, M. 2002 The molecular organization of cerebellar long-term depression. *Nat. Rev. Neurosci.* **3**, 896–902. (doi:10.1038/nrn962)
- Ito, M. & Kano, M. 1982 Long-lasting depression of parallel fiber-Purkinje cell transmission induced by conjunctive stimulation of parallel fibers and climbing fibers in the cerebellar cortex. *Neurosci. Lett.* **33**, 253–258. (doi:10.1016/0304-3940(82)90380-9)
- Ito, M. & Yoshida, M. 1964 The cerebellar-evoked monosynaptic inhibition of Deiters' neurones. *Experientia* **20**, 515–516. (doi:10.1007/BF02154085)
- Ito, M. & Yoshida, M. 1966 The origin of cerebral-induced inhibition of Deiters neurones. I. Monosynaptic initiation of the inhibitory postsynaptic potentials. *Exp. Brain Res.* **2**, 330–349.
- Ito, M., Obata, K. & Ochi, R. 1966 The origin of cerebellar-induced inhibition of Deiters neurones. II. Temporal correlation between the trans-synaptic activation of Purkinje cells and the inhibition of Deiters neurones. *Exp. Brain Res.* **2**, 350–364.
- Ito, M., Yoshida, M. & Obata, K. 1964 Monosynaptic inhibition of the intracerebellar nuclei induced from the cerebellar cortex. *Experientia* **20**, 575–576. (doi:10.1007/BF02150304)
- Ito, M., Sakurai, M. & Tongroach, P. 1982 Climbing fibre induced depression of both mossy fibre responsiveness and glutamate sensitivity of cerebellar Purkinje cells. *J. Physiol.* **324**, 113–134.
- Jin, Y., Kim, S. J., Kim, J., Worley, P. F. & Linden, D. J. 2007 Long-term depression of mGluR1 signaling. *Neuron* **55**, 277–287. (doi:10.1016/j.neuron.2007.06.035)

- Kakizawa, S., Yamasaki, M., Watanabe, M. & Kano, M. 2000 Critical period for activity-dependent synapse elimination in developing cerebellum. *J. Neurosci.* **20**, 4954–4961.
- Kakizawa, S., Yamada, K., Iino, M., Watanabe, M. & Kano, M. 2003 Effects of insulin-like growth factor I on climbing fibre synapse elimination during cerebellar development. *Eur. J. Neurosci.* **17**, 545–554. (doi:10.1046/j.1460-9568.2003.02486.x)
- Kamikubo, Y., Tabata, T., Kakizawa, S., Kawakami, D., Watanabe, M., Ogura, A., Iino, M. & Kano, M. 2007 Postsynaptic GABA_B receptor signalling enhances LTD in mouse cerebellar Purkinje cells. *J. Physiol.* **585**, 549–563. (doi:10.1113/jphysiol.2007.141010)
- Kano, M. & Kato, M. 1987 Quisqualate receptors are specifically involved in cerebellar synaptic plasticity. *Nature* **325**, 276–279. (doi:10.1038/325276a0)
- Kano, M., Hashimoto, K., Chen, C., Abeliovich, A., Aiba, A., Kurihara, H., Watanabe, M., Inoue, Y. & Tonegawa, S. 1995 Impaired synapse elimination during cerebellar development in PKC gamma mutant mice. *Cell* **83**, 1223–1231. (doi:10.1016/0092-8674(95)90147-7)
- Kano, M., Hashimoto, K., Kurihara, H., Watanabe, M., Inoue, Y., Aiba, A. & Tonegawa, S. 1997 Persistent multiple climbing fiber innervation of cerebellar Purkinje cells in mice lacking mGluR1. *Neuron* **18**, 71–79. (doi:10.1016/S0896-6273(01)80047-7)
- Kano, M. *et al.* 1998 Phospholipase cbeta4 is specifically involved in climbing fiber synapse elimination in the developing cerebellum. *Proc. Natl Acad. Sci. USA* **95**, 15 724–15 729. (doi:10.1073/pnas.95.26.15724)
- Kashiwabuchi, N. *et al.* 1995 Impairment of motor coordination, Purkinje cell synapse formation, and cerebellar long-term depression in GluR delta 2 mutant mice. *Cell* **81**, 245–252. (doi:10.1016/0092-8674(95)90334-8)
- Khodakhah, K. & Armstrong, C. M. 1997 Induction of long-term depression and rebound potentiation by inositol trisphosphate in cerebellar Purkinje neurons. *Proc. Natl Acad. Sci. USA* **94**, 14 009–14 014. (doi:10.1073/pnas.94.25.14009)
- Kishimoto, Y. & Kano, M. 2006 Endogenous cannabinoid signaling through the CB1 receptor is essential for cerebellum-dependent discrete motor learning. *J. Neurosci.* **26**, 8829–8837. (doi:10.1523/JNEUROSCI.1236-06.2006)
- Kishimoto, Y., Fujimichi, R., Araishi, K., Kawahara, S., Kano, M., Aiba, A. & Kirino, Y. 2002 mGluR1 in cerebellar Purkinje cells is required for normal association of temporally contiguous stimuli in classical conditioning. *Eur. J. Neurosci.* **16**, 2416–2424. (doi:10.1046/j.1460-9568.2002.02407.x)
- Konnerth, A., Dreesen, J. & Augustine, G. J. 1992 Brief dendritic calcium signals initiate long-lasting synaptic depression in cerebellar Purkinje cells. *Proc. Natl Acad. Sci. USA* **89**, 7051–7055. (doi:10.1073/pnas.89.15.7051)
- Kreitzer, A. C. & Regehr, W. G. 2001 Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* **29**, 717–727. (doi:10.1016/S0896-6273(01)00246-X)
- Kubo, Y., Miyashita, T. & Murata, Y. 1998 Structural basis for a Ca²⁺-sensing function of the metabotropic glutamate receptors. *Science* **279**, 1722–1725. (doi:10.1126/science.279.5357.1722)
- Kulik, A., Nakadate, K., Nyiri, G., Notomi, T., Malitschek, B., Bettler, B. & Shigemoto, R. 2002 Distinct localization of GABA(B) receptors relative to synaptic sites in the rat cerebellum and ventrobasal thalamus. *Eur. J. Neurosci.* **15**, 291–307. (doi:10.1046/j.0953-816x.2001.01855.x)
- Kurihara, H., Hashimoto, K., Kano, M., Takayama, C., Sakimura, K., Mishina, M., Inoue, Y. & Watanabe, M. 1997 Impaired parallel fiber->Purkinje cell synapse stabilization during cerebellar development of mutant mice lacking the glutamate receptor delta2 subunit. *J. Neurosci.* **17**, 9613–9623.
- Landsend, A. S., Amiry-Moghaddam, M., Matsubara, A., Bergersen, L., Usami, S., Wenthold, R. J. & Ottersen, O. P. 1997 Differential localization of delta glutamate receptors in the rat cerebellum: coexpression with AMPA receptors in parallel fiber-spine synapses and absence from climbing fiber-spine synapses. *J. Neurosci.* **17**, 834–842.
- Levenes, C., Daniel, H., Jaillard, D., Conquet, F. & Crepel, F. 1997 Incomplete regression of multiple climbing fibre innervation of cerebellar Purkinje cells in mGluR1 mutant mice. *Neuroreport* **8**, 571–574. (doi:10.1097/00001756-199701200-00038)
- Lichtman, J. W. & Colman, H. 2000 Synapse elimination and indelible memory. *Neuron* **25**, 269–278. (doi:10.1016/S0896-6273(00)80893-4)
- Linden, D. J. & Connor, J. A. 1991 Participation of postsynaptic PKC in cerebellar long-term depression in culture. *Science* **254**, 1656–1659. (doi:10.1126/science.1721243)
- Linden, D. J., Dickinson, M. H., Smeyne, M. & Connor, J. A. 1991 A long-term depression of AMPA currents in cultured cerebellar Purkinje neurons. *Neuron* **7**, 81–89. (doi:10.1016/0896-6273(91)90076-C)
- Lohof, A. M., Delhaye-Bouchaud, N. & Mariani, J. 1996 Synapse elimination in the central nervous system: functional significance and cellular mechanisms. *Rev. Neurosci.* **7**, 85–101.
- Lomeli, H., Sprengel, R., Laurie, D. J., Kohr, G., Herb, A., Seeburg, P. H. & Wisden, W. 1993 The rat delta-1 and delta-2 subunits extend the excitatory amino acid receptor family. *FEBS Lett.* **315**, 318–322. (doi:10.1016/0014-5793(93)81186-4)
- Lopez-Bendito, G., Shigemoto, R., Lujan, R. & Juiz, J. M. 2001 Developmental changes in the localisation of the mGluR1alpha subtype of metabotropic glutamate receptors in Purkinje cells. *Neuroscience* **105**, 413–429. (doi:10.1016/S0306-4522(01)00188-9)
- Maejima, T., Hashimoto, K., Yoshida, T., Aiba, A. & Kano, M. 2001 Presynaptic inhibition caused by retrograde signal from metabotropic glutamate to cannabinoid receptors. *Neuron* **31**, 463–475. (doi:10.1016/S0896-6273(01)00375-0)
- Maejima, T., Oka, S., Hashimoto, Y., Ohno-Shosaku, T., Aiba, A., Wu, D., Waku, K., Sugiura, T. & Kano, M. 2005 Synaptically driven endocannabinoid release requires Ca²⁺-assisted metabotropic glutamate receptor subtype 1 to phospholipase C b4 signaling cascade in the cerebellum. *J. Neurosci.* **25**, 6826–6835. (doi:10.1523/JNEUROSCI.0945-05.2005)
- Maekawa, K. & Simpson, J. L. 1973 Climbing fiber responses evoked in vestibulocerebellum of rabbit from visual system. *J. Neurophysiol.* **36**, 649–666.
- Marcaggi, P. & Attwell, D. 2005 Endocannabinoid signaling depends on the spatial pattern of synapse activation. *Nat. Neurosci.* **8**, 776–781. (doi:10.1038/nm1458)
- Marcaggi, P. & Attwell, D. 2007 Short- and long-term depression of rat cerebellar parallel fibre synaptic transmission mediated by synaptic crosstalk. *J. Physiol.* **578**, 545–550. (doi:10.1113/jphysiol.2006.115014)
- Mariani, J. & Changeux, J. P. 1980 Multiple innervation of Purkinje cells by climbing fibers in the cerebellum of the adult staggerer mutant mouse. *J. Neurobiol.* **11**, 41–50. (doi:10.1002/neu.480110106)
- Mariani, J. & Changeux, J. P. 1981 Ontogenesis of olivocerebellar relationships. I. Studies by intracellular recordings of the multiple innervation of Purkinje cells by climbing fibers in the developing rat cerebellum. *J. Neurosci.* **1**, 696–702.

- Mariani, J., Crepel, F., Mikoshiba, K., Changeux, J. P. & Sotelo, C. 1977 Anatomical, physiological and biochemical studies of the cerebellum from Reeler mutant mouse. *Phil. Trans. R. Soc. B* **281**, 1–28. (doi:10.1098/rstb.1977.0121)
- Marr, D. 1969 A theory of cerebellar cortex. *J. Physiol.* **202**, 437–470.
- Masu, M., Tanabe, Y., Tsuchida, K., Shigemoto, R. & Nakanishi, S. 1991 Sequence and expression of a metabotropic glutamate receptor. *Nature* **349**, 760–765. (doi:10.1038/349760a0)
- Matsuda, S., Launey, T., Mikawa, S. & Hirai, H. 2000 Disruption of AMPA receptor GluR2 clusters following long-term depression induction in cerebellar Purkinje neurons. *EMBO J.* **19**, 2765–2774. (doi:10.1093/emboj/19.12.2765)
- Miyakawa, H., Lev-Ram, V., Lasser-Ross, N. & Ross, W. N. 1992 Calcium transients evoked by climbing fiber and parallel fiber synaptic inputs in guinea pig cerebellar Purkinje neurons. *J. Neurophysiol.* **68**, 1178–1189.
- Miyata, M. et al. 2000 Local calcium release in dendritic spines required for long-term synaptic depression. *Neuron* **28**, 233–244. (doi:10.1016/S0896-6273(00)00099-4)
- Miyata, M. et al. 2001 Deficient long-term synaptic depression in the rostral cerebellum correlated with impaired motor learning in phospholipase C beta4 mutant mice. *Eur. J. Neurosci.* **13**, 1945–1954. (doi:10.1046/j.0953-816x.2001.01570.x)
- Morando, L., Cesa, R., Rasetti, R., Harvey, R. & Strata, P. 2001 Role of glutamate delta -2 receptors in activity-dependent competition between heterologous afferent fibers. *Proc. Natl Acad. Sci. USA* **98**, 9954–9959. (doi:10.1073/pnas.171098398)
- Nakamura, M., Sato, K., Fukaya, M., Araishi, K., Aiba, A., Kano, M. & Watanabe, M. 2004 Signaling complex formation of phospholipase Cbeta4 with metabotropic glutamate receptor type 1alpha and 1,4,5-trisphosphate receptor at the perisynapse and endoplasmic reticulum in the mouse brain. *Eur. J. Neurosci.* **20**, 2929–2944. (doi:10.1111/j.1460-9568.2004.03768.x)
- Narasimhan, K., Pessah, I. N. & Linden, D. J. 1998 Inositol-1,4,5-trisphosphate receptor-mediated Ca mobilization is not required for cerebellar long-term depression in reduced preparations. *J. Neurophysiol.* **80**, 2963–2974.
- Nicoletti, F., Meek, J. L., Iadarola, M. J., Chuang, D. M., Roth, B. L. & Costa, E. 1986 Coupling of inositol phospholipid metabolism with excitatory amino acid recognition sites in rat hippocampus. *J. Neurochem.* **46**, 40–46. (doi:10.1111/j.1471-4159.1986.tb12922.x)
- Nusser, Z., Mulvihill, E., Streit, P. & Somogyi, P. 1994 Subsynaptic segregation of metabotropic and ionotropic glutamate receptors as revealed by immunogold localization. *Neuroscience* **61**, 421–427. (doi:10.1016/0306-4522(94)90421-9)
- Obata, K., Ito, M., Ochi, R. & Sato, N. 1967 Pharmacological properties of the postsynaptic inhibition by Purkinje cell axons and the action of gamma-aminobutyric acid on deiters neurones. *Exp. Brain Res.* **4**, 43–57. (doi:10.1007/BF00235216)
- Offermanns, S., Hashimoto, K., Watanabe, M., Sun, W., Kurihara, H., Thompson, R. F., Inoue, Y., Kano, M. & Simon, M. I. 1997 Impaired motor coordination and persistent multiple climbing fiber innervation of cerebellar Purkinje cells in mice lacking Galphaq. *Proc. Natl Acad. Sci. USA* **94**, 14 089–14 094. (doi:10.1073/pnas.94.25.14089)
- Ohno-Shosaku, T., Maejima, T. & Kano, M. 2001 Endogenous cannabinoids mediate retrograde signals from depolarized postsynaptic neurons to presynaptic terminals. *Neuron* **29**, 729–738. (doi:10.1016/S0896-6273(01)00247-1)
- Ohno-Shosaku, T., Matsui, M., Fukudome, Y., Shosaku, J., Tsubokawa, H., Taketo, M. M., Manabe, T. & Kano, M. 2003 Postsynaptic M1 and M3 receptors are responsible for the muscarinic enhancement of retrograde endocannabinoid signalling in the hippocampus. *Eur. J. Neurosci.* **18**, 109–116. (doi:10.1046/j.1460-9568.2003.02732.x)
- Ohno-Shosaku, T., Shosaku, J., Tsubokawa, H. & Kano, M. 2002 Cooperative endocannabinoid production by neuronal depolarization and group I metabotropic glutamate receptor activation. *Eur. J. Neurosci.* **15**, 953–961. (doi:10.1046/j.1460-9568.2002.01929.x)
- Palay, S. L. & Chan-Palay, V. 1974 *Cerebellar Cortex*. New York, NY: Springer.
- Petralia, R. S., Zhao, H. M., Wang, Y. X. & Wenthold, R. J. 1998 Variations in the tangential distribution of postsynaptic glutamate receptors in Purkinje cell parallel and climbing fiber synapses during development. *Neuropharmacology* **37**, 1321–1334. (doi:10.1016/S0028-3908(98)00118-X)
- Puro, D. G. & Woodward, D. J. 1977 The climbing fiber system in the Weaver mutant. *Brain Res.* **129**, 141–146. (doi:10.1016/0006-8993(77)90976-3)
- Purves, D. & Lichtman, J. W. 1980 Elimination of synapses in the developing nervous system. *Science* **210**, 153–157. (doi:10.1126/science.7414326)
- Safo, P. K. & Regehr, W. G. 2005 Endocannabinoids control the induction of cerebellar LTD. *Neuron* **48**, 647–659. (doi:10.1016/j.neuron.2005.09.020)
- Sakurai, M. 1987 Synaptic modification of parallel fibre-Purkinje cell transmission in vitro guinea-pig cerebellar slices. *J. Physiol.* **394**, 463–480.
- Sakurai, M. 1990 Calcium is an intracellular mediator of the climbing fiber in induction of cerebellar long-term depression. *Proc. Natl Acad. Sci. USA* **87**, 3383–3385. (doi:10.1073/pnas.87.9.3383)
- Sasaki, K. & Strata, P. 1967 Responses evoked in the cerebellar cortex by stimulating mossy fibre pathways to the cerebellum. *Exp. Brain Res.* **3**, 95–110. (doi:10.1007/BF00233255)
- Scelfo, B. & Strata, P. 2005 Correlation between multiple climbing fibre regression and parallel fibre response development in the postnatal mouse cerebellum. *Eur. J. Neurosci.* **21**, 971–978. (doi:10.1111/j.1460-9568.2005.03933.x)
- Shigemoto, R., Abe, T., Nomura, S., Nakanishi, S. & Hirano, T. 1994 Antibodies inactivating mGluR1 metabotropic glutamate receptor block long-term depression in cultured Purkinje cells. *Neuron* **12**, 1245–1255. (doi:10.1016/0896-6273(94)90441-3)
- Sladeczek, F., Pin, J. P., Recasens, M., Bockaert, J. & Weiss, S. 1985 Glutamate stimulates inositol phosphate formation in striatal neurones. *Nature* **317**, 717–719. (doi:10.1038/317717a0)
- Sugihara, I. 2005 Microzonal projection and climbing fiber remodeling in single olivocerebellar axons of newborn rats at postnatal days 4–7. *J. Comp. Neurol.* **487**, 93–106. (doi:10.1002/cne.20531)
- Sugiura, T., Kishimoto, S., Oka, S. & Gokoh, M. 2006 Biochemistry, pharmacology and physiology of 2-arachidonoylglycerol, an endogenous cannabinoid receptor ligand. *Prog. Lipid Res.* **45**, 405–446. (doi:10.1016/j.plipres.2006.03.003)
- Sugiyama, H., Ito, I. & Hirono, C. 1987 A new type of glutamate receptor linked to inositol phospholipid metabolism. *Nature* **325**, 531–533. (doi:10.1038/325531a0)

- Tabata, T., Aiba, A. & Kano, M. 2002 Extracellular calcium controls the dynamic range of neuronal metabotropic glutamate receptor responses. *Mol. Cell Neurosci.* **20**, 56–68. (doi:10.1006/mcne.2002.1118)
- Tabata, T., Araishi, K., Hashimoto, K., Hashimotodani, Y., van der Putten, H., Bettler, B. & Kano, M. 2004 Ca²⁺ activity at GABA_B receptors constitutively promotes metabotropic glutamate signaling in the absence of GABA. *Proc. Natl Acad. Sci. USA* **101**, 16 952–16 957. (doi:10.1073/pnas.0405387101)
- Tabata, T. *et al.* 2007 G protein-independent neuromodulatory action of adenosine on metabotropic glutamate signaling in mouse cerebellar Purkinje cells. *J. Physiol.* **581**, 693–708. (doi:10.1113/jphysiol.2007.129866)
- Takayama, C., Nakagawa, S., Watanabe, M., Mishina, M. & Inoue, Y. 1996 Developmental changes in expression and distribution of the glutamate receptor channel delta 2 subunit according to the Purkinje cell maturation. *Brain Res. Dev. Brain Res.* **92**, 147–155. (doi:10.1016/0165-3806(95)00212-X)
- Takechi, H., Eilers, J. & Konnerth, A. 1998 A new class of synaptic response involving calcium release in dendritic spines. *Nature* **396**, 757–760. (doi:10.1038/25547)
- Tanaka, J. *et al.* 2000 Gq protein alpha subunits Galphaq and Galpha11 are localized at postsynaptic extra-junctional membrane of cerebellar Purkinje cells and hippocampal pyramidal cells. *Eur. J. Neurosci.* **12**, 781–792. (doi:10.1046/j.1460-9568.2000.00959.x)
- Varma, N., Brager, D., Morishita, W., Lenz, R. A., London, B. & Alger, B. 2002 Presynaptic factors in the regulation of DSI expression in hippocampus. *Neuropharmacology* **43**, 550–562. (doi:10.1016/S0028-3908(02)00168-5)
- Wang, Y. T. & Linden, D. J. 2000 Expression of cerebellar long-term depression requires postsynaptic clathrin-mediated endocytosis. *Neuron* **25**, 635–647. (doi:10.1016/S0896-6273(00)81066-1)
- Wilson, R. I. & Nicoll, R. A. 2001 Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. *Nature* **410**, 588–592. (doi:10.1038/35069076)
- Woodward, D. J., Hoffer, B. J. & Altman, J. 1974 Physiological and pharmacological properties of Purkinje cells in rat cerebellum degranulated by postnatal x-irradiation. *J. Neurobiol.* **5**, 283–304. (doi:10.1002/neu.480050402)
- Xia, J., Chung, H. J., Wihler, C., Haganir, R. L. & Linden, D. J. 2000 Cerebellar long-term depression requires PKC-regulated interactions between GluR2/3 and PDZ domain-containing proteins. *Neuron* **28**, 499–510. (doi:10.1016/S0896-6273(00)00128-8)