



Published in final edited form as:

Mol Microbiol. 2008 January ; 67(2): 336–349. doi:10.1111/j.1365-2958.2007.06030.x.

The disulfide isomerase DsbC cooperates with the oxidase DsbA in a DsbD-independent manner

Didier Vertommen¹, Matthieu Depuydt¹, Jonathan Pan², Pauline Leverrier¹, Laurent Knoops^{1,6}, Jean-Pierre Szikora¹, Joris Messens^{3,4,5}, James C.A. Bardwell², and Jean-Francois Collet^{1,5,*}

¹ de Duve Institute, Université catholique de Louvain, B-1200 Brussels, Belgium

² Program in Cellular and Molecular Biology, Department of Molecular, Cellular and Developmental Biology, University of Michigan, Ann Arbor, MI 48109-1048

³ Department of Molecular and Cellular Interactions, VIB

⁴ Ultrastructure Laboratory, Vrije Universiteit Brussel, B-1050 Brussels, Belgium

⁵ Brussels Center for Redox Biology, Belgium

⁶ Cliniques Universitaires Saint-Luc, B-1200 Brussels, Belgium

Summary

In *Escherichia coli*, DsbA introduces disulfide bonds into secreted proteins. DsbA is recycled by DsbB, which generates disulfides from quinone reduction. DsbA is not known to have any proofreading activity and can form incorrect disulfides in proteins with multiple cysteines. These incorrect disulfides are thought to be corrected by a protein disulfide isomerase, DsbC, which is kept in the reduced and active configuration by DsbD. The DsbC/DsbD isomerization pathway is considered to be isolated from the DsbA/DsbB pathway. We show that the DsbC and DsbA pathways are more intimately connected than previously thought. *dsbA⁻dsbC⁻* mutants have a number of phenotypes not exhibited by either *dsbA⁻*, *dsbC⁻*, or *dsbA⁻dsbD⁻* mutations: they exhibit an increased permeability of the outer membrane, are resistant to the lambdaoid phage $\Phi 80$, and are unable to assemble the maltoporin LamB. Using differential 2D-LC-MS/MS, we estimated the abundance of about 130 secreted proteins in various *dsb⁻* strains. *dsbA⁻dsbC⁻* mutants exhibit unique changes at the protein level that are not exhibited by *dsbA⁻dsbD⁻* mutants. Our data indicate that DsbC can assist DsbA in a DsbD-independent manner to oxidatively fold envelope proteins. The view that DsbC's function is limited to the disulfide isomerization pathway should therefore be reinterpreted.

Keywords

DsbC; envelope; SigmaE; outer membrane; disulfide; DsbA

Introduction

DsbA introduces disulfide bonds into secreted proteins in the *Escherichia coli* periplasm (Bardwell *et al.*, 1991). DsbA has a CXXC catalytic site motif present within a thioredoxin fold. The cysteine residues of this motif are found oxidized *in vivo*. The disulfide bond of

*For correspondence. Jean-Francois Collet, de Duve Institute, Université catholique de Louvain, 75-39 Avenue Hippocrate, B-1200 Brussels, Belgium. Tel. 32-2-764-7562; Fax 32-2-764-7598; E-mail jfcollet@uclouvain.be.

DsbA is very unstable and is rapidly transferred to secreted unfolded proteins. DsbA is then reoxidized by the inner-membrane protein DsbB (Bardwell *et al.*, 1993). DsbA forms a mixed disulfide complex with DsbB. Disulfide bond transfer occurs after conformational changes within the DsbB protein (Inaba *et al.*, 2006). DsbB has two pairs of cysteine residues and generates disulfide bonds *de novo* from quinone reduction. Electrons are then successively transferred from quinone to cytochrome oxidases and finally to molecular oxygen (Bader *et al.*, 1999).

DsbA is likely to be involved in the oxidative folding of many periplasmic and outer membrane proteins. A search in the sequence databases reveals that about 40% of the 700 secreted proteins in *E. coli* have at least two cysteine residues and are therefore potential DsbA substrates (Dana Boyd, personal communication). It is therefore not surprising that *dsbA*⁻ strains have a pleiotropic phenotype; they show an attenuated virulence, lack motility, form mucoidal colonies on minimal media and in the presence of some antibiotics such as tetracycline, and are more sensitive to dithiothreitol and cadmium (reviewed in Collet and Bardwell, 2002). About 15 DsbA substrates have been identified so far; these were obtained by trapping of mixed disulfides with DsbA mutants (Kadokura *et al.*, 2004), from differential thiol trapping experiments (Leichert and Jakob, 2004), and by 2D-gel analysis, which showed decreased levels of expression of several cysteine containing proteins (Hiniker and Bardwell, 2004).

Despite the significant number of proteins that are DsbA substrates, *dsbA*⁻ strains are surprisingly healthy, particularly when grown on rich media. This in part may be due to the fact that small molecule oxidants, like cysteine, are present in rich media (Bardwell *et al.*, 1993). Consistent with this, *dsbA*⁻ strains grow very poorly on minimal media lacking cysteine, particularly in some strain backgrounds. Although the rate of disulfide bond formation is about a 100-fold decreased in *dsbA*⁻ strains (Bardwell *et al.*, 1991), some slow residual disulfide bond formation does occur, and proteins that are stable in the absence of their disulfide accumulate in the oxidized form to near normal levels at steady state. This disulfide formation appears to be dependent on oxygen, as much lower levels of oxidized proteins are seen in *dsbA*⁻ strains when they are grown anaerobically.

DsbA is a powerful oxidant that apparently lacks proofreading activity. DsbA oxidizes cysteine residues on secreted proteins as they emerge into the periplasm. If the native disulfide bond pattern involves cysteine residues that are consecutive in the amino acid sequence, DsbA can form disulfides correctly. However, when secreted proteins have disulfides that need to be formed between nonconsecutive cysteines, DsbA can introduce non-native disulfides, leading to protein misfolding and degradation by proteases (Berkmen *et al.*, 2005). Noteworthy, our recent work on RNase I, a periplasmic protein with one non-consecutive disulfide, showed that DsbA is more specific than generally assumed (Messens *et al.*, 2007). The correction of non-native disulfides is thought to be the role of a disulfide isomerization system. This system is composed of two soluble periplasmic proteins, DsbC and DsbG, which are thought to function as disulfide isomerase proteins *in vivo* and *in vitro* (Zapun *et al.*, 1995; Bessette *et al.*, 1999). Like DsbA, DsbC and DsbG possess a thioredoxin fold and a CXXC catalytic site motif. In contrast to DsbA, whose CXXC active site motif is maintained in an oxidized form, the CXXC motif of DsbC and DsbG is kept reduced in the periplasm. This allows DsbC and DsbG to attack non-native disulfides, a necessary step in the isomerization reaction (reviewed in Messens and Collet, 2006). The protein that keeps DsbC and DsbG reduced is the inner-membrane protein, DsbD. DsbD transfers reducing equivalents from the cytoplasmic thioredoxin system to the periplasm via a succession of disulfide exchange reactions (Rietsch *et al.*, 1996; Katzen and Beckwith, 2000; Collet *et al.*, 2002; Rozhkova *et al.*, 2004). Residues that are important for this electron cascade have recently been identified (Cho and Beckwith, 2006; Hiniker *et al.*,

2006), but the precise mechanism used by DsbD to transport electrons from one side of the membrane to the other is still obscure.

In contrast to the central role played by DsbA in oxidizing a large number of periplasmic proteins, DsbC is thought to be required for the expression of a limited subset of proteins that contain nonconsecutive disulfides, including the penicillin insensitive endopeptidase MepA, the ribonuclease RNase I, and the acid phosphatase AppA (Hiniker and Bardwell, 2004; Berkmen *et al.*, 2005). In agreement with the relatively small number of DsbC substrates, *dsbC*⁻ strains have a milder phenotype than *dsbA*⁻ mutants (reviewed in Collet and Bardwell, 2002). DsbC seems to be particularly important under some oxidative stress conditions. For instance, DsbC is required for growth in the presence of high concentrations of copper, a redox metal that catalyzes the formation of non-native disulfide bonds (Hiniker *et al.*, 2005).

The role of DsbG is less clear. DsbG was originally reported to be an essential oxidase (Andersen *et al.*, 1997). However, subsequent work showed it to be nonessential for the growth of *E. coli* (Bessette *et al.*, 1999). *DsbG* null mutants have no defect in the folding of heterologous proteins containing multiple disulfide bonds, and are unable to catalyze disulfide bond rearrangement using either hirudin or the bovine pancreatic trypsin inhibitor intermediate as substrates (Bessette *et al.*, 1999; Hiniker *et al.*, 2007). However, DsbG overexpression is able to restore the ability of *dsbC*⁻ mutants to express some heterologous proteins containing multiple disulfide bonds (Bessette *et al.*, 1999). It is also possible to select mutations in DsbG that complement DsbC, and these mutations show increased isomerase activity (Hiniker *et al.*, 2007). These latter observations and DsbG's homology to DsbC (Heras *et al.*, 2004) have led to the conclusion that DsbG is a disulfide isomerase with restricted substrate specificity.

The situation in eukaryotic disulfide bond formation is more complex and controversial. Protein disulfide isomerase (PDI) is thought to function as both an oxidase and an isomerase *in vivo*. However, there has been quite a bit of controversy regarding the relative importance of the oxidative and isomerase activities to the cell (Sevier and Kaiser, 2006). In addition to PDI, yeast possesses several thioredoxin-like proteins that are localized to the ER. Those proteins are likely to play important roles in the isomerization and oxidation of proteins (reviewed in Gruber *et al.*, 2006).

In *E. coli*, the current view is that the distinct disulfide catalytic pathways have well defined roles: the DsbA/DsbB system is important in oxidizing disulfide bonds, and the DsbC-G/DsbD system is important in isomerizing them. The results presented in this paper show that this conclusion needs to be reinterpreted. We show that the simultaneous absence of DsbA and DsbC has severe consequences on *E. coli*'s viability and outer membrane integrity. *dsbA*⁻*dsbC*⁻ mutants are resistant to the Φ80 bacteriophage and seem unable to fold the trimeric porin LamB. Using a 2D-LC-MS/MS proteomics approach, we show that the absence of DsbA and DsbC affects the global protein content of the periplasm and leads to a decreased abundance of several cysteine-containing proteins. Our data indicate that the function of DsbC goes beyond the correction of the non-native disulfides formed by DsbA. On the basis of our results, we propose a new model for the oxidative protein folding pathways in *E. coli*.

Results and Discussion

***dsbA*⁻*dsbC*⁻ double mutants have a more severe phenotype than *dsbA*⁻ mutants**

To test whether DsbC can assist DsbA outside the framework of the isomerization pathway, we made *dsbA*⁻*dsbC*⁻ double mutants and compared their phenotype to that of *dsbA* null

strains. The logic behind these experiments is as follows: if DsbC's function is restricted to the correction of DsbA's mistakes, then *dsbA*⁻ and *dsbA*⁻*dsbC*⁻ mutants should have a similar phenotype. In contrast, if DsbC has functions in addition to the correction of DsbA's mistakes, then *dsbA*⁻*dsbC*⁻ double mutants should have a more severe phenotype than *dsbA*⁻ mutants. We observed that a *dsbA*⁻*dsbC*⁻ mutant has a more severe growth defect than a *dsbA*⁻ strain when cells are grown in minimal media (Fig. 1A and Table 2A). Both the *dsbA*⁻ and *dsbA*⁻*dsbC*⁻ strains are more sensitive to antibiotics and detergents than wild-type strains. However, the sensitivity of the *dsbA*⁻*dsbC*⁻ double mutant is more severe than that of a *dsbA*⁻ mutant. The *dsbA*⁻*dsbC*⁻ mutant is more sensitive to SDS (Fig. 1B), and, unlike *dsbA*⁻ strains, is unable to grow in the presence of 4 μg/ml rifampin, a large hydrophobic antibiotic (Table 2A). These phenotypes suggest that the permeability of the outer membrane is increased in the double mutant. A single *dsbC*⁻ mutant does not exhibit any growth defect or sensitivity toward antibiotics and detergents compared to isogenic wild-type strains.

dsbA⁻*dsbC*⁻ mutants form pink colonies on maltodextrin MacConkey agar, in contrast to wild-type, *dsbA*⁻, and *dsbC*⁻ colonies, which are red (Table 2). This pink phenotype is often indicative of a decreased abundance of LamB, the outer membrane component of the maltose transport system in *E. coli* (Duguay and Silhavy, 2002). LamB is a trimeric protein that possesses one disulfide bond per subunit. By Western blot analysis, we confirmed that the expression level of LamB is strongly decreased in a *dsbA*⁻*dsbC*⁻ mutant compared to wild-type, *dsbC*⁻, and *dsbA*⁻ strains (Fig. 1C). The *dsbA*⁻*dsbC*⁻ mutants are also significantly more resistant to the lambdoid Φ80 phage (Table 2B). The *E. coli* receptor for this phage is the ferrichrome iron receptor protein FhuA. Interestingly, FhuA contains four cysteine residues that form two consecutive disulfide bonds. These data suggest that the function of DsbC is not restricted to the correct folding of proteins that have disulfides formed between nonconsecutive cysteines. As the disulfides of LamB and FhuA do not seem to be important for the function of these proteins (Ferenci and Stretton, 1989; Bos *et al.*, 1998), it is tempting to speculate that in the absence of DsbA and DsbC intermolecular disulfides are formed, preventing these proteins from correctly folding in the outer membrane.

Our results clearly show that *dsbA*⁻*dsbC*⁻ mutants have a more severe phenotype than *dsbA*⁻ mutants and suggest that DsbC may be involved in the folding of LamB and FhuA, two proteins that have only consecutive disulfides. Our data thus support the hypothesis that DsbC's function is not restricted to the correction of non-native disulfides. DsbC also has chaperone activity that is independent of its active site cysteine residues (Liu and Wang, 2001). Thus we needed to consider the possibility that the defects observed in *dsbC*⁻ strains is due to the lack of DsbC's chaperone activity. We however observed that expression of a mutant of DsbC in which both catalytic site cysteines are replaced by serine failed to improve the growth rate of *dsbA*⁻*dsbC*⁻ strains, in contrast to expression of the wild-type DsbC protein. This mutant of DsbC is expected to lose its thiol-disulfide oxidoreductase activity, while retaining its chaperone activity (Liu and Wang, 2001). Thus the severe phenotype of the *dsbA*⁻*dsbC*⁻ mutant is unlikely to be due to a chaperone deficiency in *dsbC*⁻ strains, but rather a thiol-disulfide oxidoreductase defect.

DsbC is able to assist DsbA in a DsbD-independent manner

To test whether the activity of DsbC always depends on the presence of DsbD, we generated a *dsbA*⁻*dsbD*⁻ double mutant. This mutant is phenotypically similar to a *dsbA*⁻ strain in terms of detergent and antibiotics sensitivity. It is even slightly more resistant to detergent than a *dsbA*⁻ strain. As such, its phenotype is less severe than that of a *dsbA*⁻*dsbC*⁻ strain. DsbD functions to keep DsbC reduced in the periplasm so that DsbC can react with non-native disulfides to correct them. The lack of equivalence between a *dsbA*⁻*dsbC*⁻ mutant

and a *dsbA*⁻*dsbD*⁻ mutant indicates that DsbC does not always require DsbD to function in the periplasm.

DsbC is reduced in a *dsbA*⁻*dsbD*⁻ mutant

DsbC is found reduced in wild-type cells, but is oxidized in strains lacking DsbD. We determined the *in vivo* redox state of DsbC in a *dsbA*⁻*dsbD*⁻ mutant using AMS trapping. DsbC is found reduced in this genetic background, even when cells are grown in LB, a media that contains small molecule oxidants (Fig. 2). Addition of diamide, a disulfide generating compound, to growing *dsbA*⁻*dsbD*⁻ cells leads to immediate oxidation of DsbC (not shown). However, when this disulfide stress is over, DsbC goes back to its reduced state even though DsbD is absent. We propose that DsbC accumulates in the reduced state in the periplasm of *dsbA*⁻*dsbD*⁻ strains by donating its disulfide bond to folding proteins.

The deletion of DsbG has no effect

The *E. coli* periplasm contains another protein disulfide isomerase, DsbG. DsbG has been proposed to function as an isomerase for essentially two reasons: first, like DsbC, DsbG is found reduced in the periplasm and second, DsbG can assist the folding of eukaryotic proteins with multiple cysteine residues when it is overexpressed (Bessette *et al.*, 1999). However, no physiological substrate has been identified so far for DsbG, and the exact function of this protein remains unclear. To see whether DsbG is also able to function in the periplasm of *dsbA*⁻ strains in a manner similar to DsbC, we constructed a *dsbA*⁻*dsbG*⁻ double mutant.

We found that deletion of DsbG does not affect the phenotype of a *dsbA*⁻ mutant (data not shown). Similarly, a triple *dsbA*⁻*dsbC*⁻*dsbG*⁻ mutant is phenotypically similar to a *dsbA*⁻*dsbC*⁻ double mutant. However, DsbG may be able to donate its disulfides in ways similar to DsbC as we found the protein to be mostly reduced in *dsbA*⁻*dsbD*⁻ double mutants.

2D-LC-MS/MS analysis of *dsb*⁻ strains

The data presented above suggest that DsbC cooperates with DsbA in a DsbD-independent manner. These observations prompted us to characterize the periplasmic proteome of various *dsb*⁻ strains by two-dimensional liquid chromatographic mass-spec/mass spec analysis (2D-LC-MS/MS) to see whether deletion of both *dsbA* and *dsbC* has specific consequences on the protein content of the periplasm. 2D-LC-MS/MS allows a global and semi-quantitative analysis of protein expression ratios. The periplasmic proteomes of wild-type, *dsbA*⁻, *dsbC*⁻, *dsbA*⁻*dsbC*⁻, and *dsbA*⁻*dsbD*⁻ strains were compared.

Two proteins with multiple cysteine were not identified in the *dsbC*⁻ strains

To examine the consequences of the absence of DsbC at the protein level, wild-type and *dsbC*⁻ strains were grown in minimal media, and periplasmic extracts were prepared. Periplasmic proteins were then digested by trypsin, and the generated peptides were analyzed by 2D-LC-MS/MS. The experiments were repeated three times for both strains. Each run allowed us to identify up to 175 secreted proteins, but only 115 proteins that could be reproducibly identified were kept for further analysis. To our knowledge, this is the first time that a proteomic approach allowed the identification of such a large number of secreted proteins, representing about 18% of all the proteins present in the cell envelope. A number of outer-membrane proteins, probably present in outer-membrane vesicles that did not pellet during the centrifugation, were reproducibly identified. Since they also represent potential targets for DsbA and DsbC, they were kept for further analysis.

The same proteins were identified as being present in both the wild-type and *dsbC*⁻ strains with the exception of three proteins that were absent in the latter strain. We discovered that these three proteins include DsbC itself and two proteins with multiple cysteine residues: a penicillin insensitive murine endopeptidase (MepA) and an endonuclease (End1). MepA had previously been found to depend on DsbC for expression (Hiniker and Bardwell, 2004), but End1 had not previously been reported to be a DsbC substrate. The structure of *Vibrio cholera* End1 shows four disulfide bonds, one of which is formed between nonconsecutive cysteines (Altermark *et al.*, 2006). The eight cysteines that form these four disulfide bonds are conserved in the *E. coli* End1 protein. The *E. coli* and *V. cholera* proteins are 66% identical at the amino acid sequence level; this suggests that they have similar structures and makes it almost certain that the *E. coli* protein shares the *V. cholera* disulfide bond pattern. Since DsbC is required for the formation of nonconsecutive disulfides, the folding of End1 is likely to require the presence of DsbC.

We then searched for proteins that, although present in both strains, vary substantially in their abundance. For quantification of abundance, we used the number of spectral counts (SC) reported for every protein. The number of SC for a protein is the total number of MS/MS spectra taken on peptides from this protein in a given 2D-LC-MS/MS analysis. This value is linearly correlated with the protein abundance over a dynamic range of two orders of magnitude (Liu *et al.*, 2004). Protein ratios determined by spectral counting agree well with those determined from peak area intensity measurements and are consistent with independent measurements based on gel staining intensities (Old *et al.*, 2005). To validate this quantification method, we added varying amounts (2–60 pmoles) of two eukaryotic proteins, ovalbumin and carbonic anhydrase, to 300 µg of periplasmic proteins. Linear regression based on different sampling statistics was performed for each of the 2D-LC-MS/MS runs; the R² values obtained for the spectral counts were 0.94 and 0.92 for ovalbumin and carbonic anhydrase, respectively (Fig. 3). We concluded that this method is quantitative and that the ratio of spectral counts reliably reflects changes in protein expression levels.

We selected proteins whose abundance was decreased or increased by at least two-fold. To test the significance of the data, we used the unpaired Student's *t* test and defined significance as a *P* < 0.05 (2-tail 2-sample equal variance test). No protein was more abundant in the *dsbC*⁻ strain (Table S1), and only two were decreased. YebF, a small protein with an unknown function, was the protein most decreased by the absence of DsbC (6 SC instead of 36), whereas Ivy, an inhibitor of lysozyme, was about two-fold less abundant. Both of these proteins have two cysteine residues, and our results suggest that they may partially depend on DsbC for correct folding.

Deletion of both *dsbA* and *dsbC* affects the global protein content of the periplasm

The results of the analysis of the periplasmic proteome of *dsbC*⁻ and wild-type strains by 2D-LC-MS/MS agree well with those obtained using 2D gels (Hiniker and Bardwell, 2004). This indicates that our 2D-LC-MS/MS method is reliable and should allow for the detection of changes in the periplasmic protein content of other *dsb*⁻ strains. Periplasmic extracts were prepared from *dsbA*⁻, *dsbA*⁻*dsbC*⁻, and *dsbA*⁻*dsbD*⁻ mutants, proteins were digested, and peptides were separated by HPLC followed by LC-MS/MS analysis. The results from the MS/MS analysis were then compared to those obtained previously for the wild-type strain. First, the expression levels of several proteins were dramatically modified in all the strains that lack DsbA. These differences, which are described below, are consistent with the previously reported role of DsbA in oxidative protein folding. Second, we found that a *dsbA*⁻*dsbC*⁻ double mutant has a significantly altered periplasmic proteome when compared to *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ strains. This is reflected by the graphs shown in Fig. 4. The SC values from the *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ strains are linearly distributed, reflecting similar protein content. Comparison of the *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ mutants shows indeed that there

is only one protein (Spy) whose abundance is significantly different between these two strains (see supplementary table 2). In contrast, when the SC values reported for proteins from the *dsbA*⁻*dsbC*⁻ mutant are plotted against those from the *dsbA*⁻ strain, the distribution is much more dispersed. This indicates a distinct overall protein content of the *dsbA*⁻*dsbC*⁻ mutant and confirms the deleterious effect of the simultaneous absence of DsbA and DsbC on the periplasm. One possibility is that the broader dispersion of the SC values observed in the *dsbA*⁻*dsbC*⁻ mutant reflects an increased sensitivity of this strain to the osmotic shock procedure. However, this would probably be reflected by an increased overall yield of proteins in periplasmic extracts, not decreased amounts of specific proteins in these extracts. The distinct protein content of *dsbA*⁻*dsbC*⁻ strains is a new finding and suggests that DsbC and DsbA cooperate in the folding of proteins. Specifically our finding suggests that DsbC can take over the function of DsbA. This is not consistent with the assigned roles of DsbA as an oxidase and DsbC as an isomerase. It suggests that a revision of the current model of disulfide bond formation in *E. coli* is called for. We note that a more likely interpretation is that DsbC acts to compensate for the absence of DsbA, and DsbA and DsbC cooperate in helping to fold proteins.

Several cysteine-containing proteins are less abundant in a *dsbA*⁻*dsbC*⁻ strain

We searched for proteins with a significantly decreased abundance in the *dsbA*⁻*dsbC*⁻ strain, compared to the *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ strains. Ten proteins were not detected in the double *dsbA*⁻*dsbC*⁻ mutant (Cn16, DsbC, FepB, GltI, Slp, SubI, YcfS, YggN, YjhT, and YnjE), and 10 were at least two-fold less abundant in the *dsbA*⁻*dsbC*⁻ than in the *dsbA*⁻ and the *dsbA*⁻*dsbD*⁻ mutants ($P < 0.05$) (Table 3). Interestingly, six of these proteins (Cn16, GltI, YggN, OppA, TreA, and YhjJ) have two cysteine residues. Noteworthy, GltI, YggN, and OppA are known DsbA substrates (Hiniker and Bardwell, 2004; Kadokura *et al.*, 2004). Two other cysteine-containing DsbA substrates, PhoA and DppA, were also more than 2 fold less abundant in the *dsbA*⁻*dsbC*⁻ mutant than in the *dsbA*⁻ or the *dsbA*⁻*dsbD*⁻ mutant. However, the decrease was not statistically significant when compared to the *dsbA*⁻*dsbD*⁻ mutant (see Supplementary Table 2) and these proteins were not included in Table 3. Altogether, our data show that the simultaneous absence of both DsbA and DsbC decreases the level of several proteins, including several that contain cysteine residues. In contrast, the absence of DsbD has no effect on *dsbA*⁻ strains. Determination of RNA expression levels (Table 3) showed that, for most of the decreased proteins, their lower abundance is not due to a decreased transcription and is therefore likely to represent a direct consequence of the absence of DsbC. Altogether, our results further support the hypothesis that DsbC can assist DsbA in a DsbD-independent manner.

Proteins are decreased in all strains impaired in disulfide bond formation

In all strains lacking DsbA, the relative abundance of several proteins is dramatically modified compared to wild-type (Table 4). In 125 proteins that were identified, we observed that the abundance of about 50 proteins was modified by at least two-fold.

In addition to DsbA, eight proteins with at least two cysteine residues were missing in *dsbA*⁻ strains. Two of these proteins (MepA and End1) were also missing in the *dsbC*⁻ mutant, which suggests they require the presence of both DsbA and DsbC. The other cysteine containing proteins are a periplasmic ribonuclease (RNase I), the outer membrane colicin 1 receptor protein (CirA), a protein that is exported to the periplasm according to Psort (CreA), and three proteins with an unknown function (YebY, YtfQ, and YfhM). We also found nine proteins with two or more cysteine residues whose abundance was decreased by at least two-fold ($P < 0.05$) (Table 4). HisJ, GltI, DppA, PhoA and YggN are known DsbA substrates (Hiniker and Bardwell, 2004; Kadokura *et al.*, 2004), but the other proteins (ProX, ArgT, ArtJ, and YebF) had not previously been identified to depend on DsbA for correct

folding. Sequence analysis revealed that the cysteine residues of these proteins are conserved in homologous sequences, which suggests that they are structurally important and probably form disulfide bonds. Our results allow us to add these proteins to the list of the potential DsbA substrates. The formation of a disulfide in YtfQ, YebF, and ArtJ was confirmed by differential thiol trapping (see below).

The absence of DsbA also leads to decreased levels of several proteins that do not contain cysteine residues (Table 4). This can either be a direct consequence of a misfolding problem caused by the absence of DsbA or can result from a decreased transcription of their genes. To discriminate between these two possibilities, we determined the RNA expression levels of the genes coding for these proteins (Table 4). Our data show that the absence of DsbA has no consequence on the transcription of most of these genes and that some genes, such as *artJ*, *phoA*, and *yebF* are even induced. This indicates that the decreased protein abundance is due to an impaired folding of these proteins. Because DsbA does not have a chaperone activity, we propose that envelope perturbations in *dsbA*⁻ strains prevent these proteins to correctly fold in the periplasm. In contrast, we found that the RNA expression levels of *ompF* and *flgG* were significantly decreased, which indicates that the diminution in the corresponding protein abundance is due to a decreased transcription of their genes. Regarding the outer membrane protein OmpF, our data agree with previous results (Pugsley, 1993). The other protein, FlgG, is a protein of the bacterial flagellum. Previous reports have shown that in a *dsbA*⁻ strain, the flagellar P-ring protein FlgI is not properly folded and is degraded (Dailey and Berg, 1993). The degradation of FlgI prevents the assembly of a functional flagellum, which leads to the repression of the transcription of other flagellum genes (Chilcott and Hughes, 2000). The misfolding of FlgI is therefore the reason why the transcription of *flgG* is repressed and the corresponding protein is not identified in the *dsbA*⁻ strain. Noteworthy, we found that the transcription of *flgH*, the gene coding for the other flagellum protein that was decreased in our proteomics analysis, was also diminished, but by less than 2 fold.

Several stress-related proteins are more abundant in strains lacking DsbA

The absence of DsbA leads to increased levels of 25 proteins, including nine proteins that were not identified in the wild-type. Similar data were obtained for the *dsbA*⁻*dsbC*⁻ and *dsbA*⁻*dsbD*⁻ mutants (see supplementary tables). Determination of the RNA expression levels allowed us to show that the transcription rates of 14 of the genes coding for these proteins are increased (Table 3).

Seven of these 25 proteins are part of the SigmaE regulon. SigmaE is a transcriptional activator that controls the expression of a variety of genes involved in maintaining the integrity of the cell envelope (reviewed in Ruiz and Silhavy, 2005). SigmaE is induced under conditions of stress in the cell envelope, including accumulation of misfolded outer membrane proteins in the periplasm, aberrant lipopolysaccharides, and lack of periplasmic folding agent. Gross and coworkers already showed that the absence of DsbA leads to an increased transcription of the gene coding for SigmaE (Mecsas *et al.*, 1993), but this is the first time that the induction of SigmaE was confirmed at the protein level. The SigmaE regulon members whose abundance is increased in the *dsbA*⁻ strain include the outer membrane proteins OmpA and OmpX, the periplasmic chaperone FkbA, the periplasmic proteases DegP and YhjJ, a protein involved in the biosynthesis of osmoregulated glycans (OpgG) and a negative regulator of SigmaE activity (RseB). The induction of these proteins further indicates that lack of disulfide bond formation leads to a global stress in the cell envelope.

In addition to the induction of SigmaE regulated proteins, we observed the induction of proteins that are known to be induced under high osmotic pressure: OsmE, an osmotically-

inducible lipoprotein (Bordes *et al.*, 2002), OsmY, a small protein of unknown function that has been proposed to interact with phospholipids on both sides of the periplasm (Lange *et al.*, 1993), and the periplasmic trehalase TreA (Repoila and Gutierrez, 1991). Determination of the expression levels of the genes coding for these proteins allowed us to confirm that the increased protein abundance of OsmE and OsmY can be attributed to an increased RNA synthesis. Similar changes in the abundance of these three proteins, as well as a decreased transcription of *ompF* (see above), have been observed in strains grown under high osmotic pressure, suggesting that *dsbA*⁻ strains mimic the effects of increased osmotic pressure. The induction of Spy, a protein that is specifically induced in spheroplasts (Hagenmaier *et al.*, 1997), suggests that the induction of these osmo-related proteins in *dsbA*⁻ strains may be the consequence of an altered peptidoglycan layer.

Determination of the in vivo redox state of periplasmic proteins by differential thiol trapping

To confirm the presence of disulfide bonds in the newly identified DsbA substrates, we adapted the differential thiol trapping technique developed by (Leichert and Jakob, 2004) to determine the redox state of the cysteine residues present in the periplasm of *dsbA*⁻ and wild-type strains.

As expected, the majority of the cysteine residues identified in peptides from the wild-type strain were oxidized (Table 5). In contrast, more reduced cysteine residues were found in proteins from the *dsbA*⁻ strain. In particular, cysteine residues from known DsbA substrates including OmpA, PhoA, DppA, and GltI were found oxidized in the wild-type and, when detected, reduced in the *dsbA*⁻ strain. The differential thiol trapping technique allowed us to confirm the formation of a disulfide bond in three newly identified DsbA substrates, ArtJ, YebF and YtfQ.

We also determined the redox state of the cysteine residues in the periplasm of the *dsbA*⁻*dsbC*⁻ and *dsbA*⁻*dsbD*⁻ double mutants and we found that they are similar to those observed in the *dsbA*⁻ strain.

A revised model for the oxidative protein folding pathways in *E. coli*

In conclusion, our results show that the simultaneous absence of DsbA and DsbC leads to a decreased integrity of the cell envelope and affects the global protein content of the periplasm. In contrast, strains lacking both DsbA and DsbD, the protein that is responsible for keeping DsbC active as an isomerase, do not share these characteristics. Our results suggest therefore that DsbC cooperates with DsbA in a DsbD-independent manner to ensure the correct folding of *E. coli* envelope proteins.

Kinetic, structural and genetic data showed that DsbB is unable to oxidize DsbC at physiological rates, unless the dimerization domain is removed and DsbC is expressed as a monomeric protein (Bader *et al.*, 2001). Similarly, DsbD is unable to reduce DsbA (Rozhkova *et al.*, 2004). This led to the assumption that the DsbA/DsbB oxidation pathway was isolated from the DsbC/DsbD isomerization pathway. Our results show that one can open a door in the barrier separating the oxidative and isomerization pathways. Our results indicate that, in contrast to the current view, DsbC can function independently of DsbD and is therefore able to function in both the oxidation and isomerization pathways. When DsbC gets oxidized upon reduction of a non-native disulfide, it is either reduced by DsbD or by transferring its disulfide to a reduced protein. DsbC may possibly be acting as a stand-alone protein folding catalyst that is able to cycle from the reduced to the oxidized state upon substrate oxidation and substrate reduction, respectively. This activity of DsbC seems important to maintain the integrity of the cell envelope and is not restricted to the correction

of non-consecutive disulfides. Our results extend those from Bader and co-workers who showed that monomeric mutants of DsbC are substrates for DsbB and can catalyze disulfide bond formation (Bader *et al.*, 2001). On the basis of our results, we have adapted the model of disulfide bond formation in the *E. coli* periplasm (Fig. 5).

Experimental procedures

Bacterial strains and growth conditions

The bacterial strains used in this study are described in Table 1. Strains JP114, JP220, JP539, JP557, JP649 and C600 were used for the titration experiment with phage *F80*. All the other experiments were performed with strains AH50, JFC383, AH396, MD1 and MD3 in the MC1000 background. Strains JFC383, MD1, and MD3 were constructed by P1 transduction. Cells were grown aerobically in either LB or M63 minimal media, at 37°C. Unless otherwise indicated, M63 minimal medium was supplemented with 0.2% glucose, vitamins (Thiamine 10 µg/ml, Biotine 1 µg/ml, Riboflavine 10 µg/ml, and Nicotinamide 10 µg/ml), 1 mM MgSO₄, leucine (20 µg/ml) and isoleucine (20 µg/ml). Sensitivity to antibiotics was assayed by streaking the strains on LB plates containing 4 µg/ml rifampin. To test the sensitivity to SDS, strains were grown in LB at 37°C to an A₆₀₀ of 0.5. The cultures were then serially diluted 10⁷-fold in 10-fold increments. 10 µl of each dilution were then spotted on LB plates containing 2.5% SDS and grown overnight. To study the ability of *dsb*⁻ strains to assemble a functional LamB protein, strains were streaked on MacConkey agar indicator plates containing 1% maltodextrin.

Φ80 phage titration

Cells were grown overnight to late logarithmic phase at 30°C. 100 µL of cells was used to inoculate 3 mL of LB top agar (0.7% LB agar). The suspension was vortexed and plated onto a pre-warmed LB plate. Serial dilutions of *F80* stock (> 10¹¹ pfu/mL) were made at 10⁻³, 10⁻⁶, and 10⁻⁹. 5 µL of these serial dilutions was spotted onto the LB agar plate containing cells, and allowed to incubate at 30°C overnight (16–18 h). The number of plaques and plaque sizes were tabulated.

Expression of a DsbC SXXS mutant

The catalytic site cysteine residues of DsbC were replaced by serine residues using the QuickChange Mutagenesis Protocol (Stratagene). Both the wild-type and mutated DNA sequence were then inserted in the pBAD33 expression plasmid. The plasmids were transferred into the MD3 strain and DsbC expression was induced by adding L-arabinose (0.2 %).

Periplasmic extracts preparation

Cells (100 ml) were grown aerobically at 37°C in M63 minimal media to an A₆₀₀ of 0.8, and periplasmic extracts were prepared as in (Hiniker and Bardwell, 2004). Protein concentration was determined using the Bradford assay.

Differential thiol trapping and digestion

300 µg of periplasmic proteins were precipitated by adding trichloroacetic acid (TCA) to a final concentration of 10% w/v, followed by incubation on ice for 30 min. Samples were then centrifuged at 14,000 rpm for 20 min and the resulting pellets washed with 5% ice cold TCA. The pellets were then resuspended in 100 µl denaturing buffer (6M urea, 200 mM Tris-HCl pH 8.5, 10 mM EDTA) supplemented with 100 mM iodoacetamide. At this stage, various amounts ranging from 2 to 60 pmoles of carbonic anhydrase and ovalbumin were added to the samples as internal standards. After a 20 min incubation at 25°C, the reaction

was stopped by adding 10 μ l of ice cold 100% TCA and left on ice for 20 min. The alkylated proteins were centrifuged and the pellet washed as described above. The proteins were then dissolved in 100 μ l of 10 mM DTT in denaturing buffer. After a 1 h incubation at 25°C, 100 μ l of denaturing buffer supplemented with 100 mM N-ethylmaleimide was added to titrate out the remaining DTT and alkylate all newly reduced cysteines. The reaction was stopped by addition of 10% TCA and the proteins collected by centrifugation. The resulting pellet was successively washed with TCA and ice cold acetone, dried in a Speedvac, resuspended in 0.1 M NH_4HCO_3 pH 8.0 with 3 μ g sequencing grade trypsin, and digested overnight at 30°C. Peptide samples were then acidified to pH 3.0 with formic acid and stored at -20°C.

Differential analysis of periplasmic proteins by label-free 2D-LC-MS/MS

Peptides were loaded onto a strong cation exchange column GROM-SIL 100 SCX (100 \times 2 mm, GROM, Rottenburg, Germany) equilibrated with solvent A (5% acetonitrile v/v, 0.05% v/v formic acid pH 2.5 in water) and connected to an Agilent 1100 HPLC system. Peptides were separated using a 50 min elution gradient that consisted of 0%–50% solvent B (5% acetonitrile v/v, 1 M ammonium formate adjusted to pH 3.0 with formic acid in water) at a flow rate of 200 μ l/min. Fractions were collected at 2 min intervals (20 in total) and dried using a Speedvac. Peptides were resuspended in 10 μ l of solvent C (5% acetonitrile v/v, 0.01% v/v TFA in water) and analyzed by LC-MS/MS as described below.

The LC-MS/MS system consisted of an LCQ DECA XP Plus ion trap mass spectrometer (ThermoFinnigan, San José, CA, USA) equipped with a microflow electrospray ionization source and interfaced to an LCPackings Ultimate Plus Dual gradient pump, Switchos column switching device, and Famos Autosampler (Dionex, Amsterdam, Netherlands). Two reverse phase peptide traps C18 Pepmap 100 Dionex (300 μ m \times 5 mm) were used in parallel with two analytical BioBasic-C18 columns from ThermoElectron (0.18 mm \times 150 mm). Samples were injected and desalted on the peptide trap equilibrated with solvent C at a flow rate of 30 μ l/min. After valve switching, peptides were eluted in backflush mode from the trap onto the analytical column equilibrated in solvent D (5% acetonitrile v/v, 0.05% v/v formic acid in water) and separated using a 100 min gradient from 0% to 70% solvent E (80% acetonitrile v/v, 0.05% formic acid in water) at a flow rate of 1.5 μ l/min.

The mass spectrometer was set up to acquire one full MS scan in the mass range of 400–2000 m/z, followed by three MS/MS spectra of the three most intense peaks in the mass range 400–1500 m/z. The dynamic exclusion feature was enabled to obtain MS/MS spectra on co-eluting peptides, and the exclusion time was set at 2 min.

Protein identification

Raw data collection of approximately 54,000 MS/MS spectra per 2D-LC-MS/MS experiment was followed by protein identification using the TurboSequest algorithm in the Bioworks 3.2 software package (ThermoFinnigan) against an *E. coli* protein database (SwissProt) using the following constraints: only tryptic peptides up to one missed cleavage site were allowed; tolerances for MS and MS/MS fragment ions were set to 1.2 Da and 1.0 Da, respectively; and methionine oxidation (+ 16.0 Da), carboxamidomethyl cysteine or N-ethylmaleimide cysteine (+ 57.0 Da or + 125.0 Da, respectively) were specified as variable modifications. The identified peptides were further evaluated using charge state versus cross correlation number (Xcorr). The criteria for positive identification of peptides were Xcorr > 1.5 for singly charged ions, Xcorr > 2.0 for doubly charged ions, and Xcorr > 2.5 for triply charged ions. Protein scores (Su, Xcorr), peak areas, and spectral counts were calculated within Bioworks 3.2. The data were converted into Microsoft Excel spreadsheets by the export function contained in Bioworks and the output files were compared and processed by an in house software program. Relative quantification of protein abundance was estimated

by calculating the ratio of spectral counts determined within the BioWorks software package. This parameter was shown to follow a linear relationship over two orders of magnitude, as determined from spiked internal standard proteins, within a dynamic range of at least 10^3 .

Preparation of outer membrane proteins

Outer membrane proteins were prepared from strains grown in LB. Cultures were grown to an A_{600} of 0.8; cells were harvested by centrifugation at 6000 rpm for 10 min then resuspended in 25 mM Tris pH 8.0, 0.5 M sucrose, 1 mM EDTA, and 0.25 mg/ml lysozyme. After 15 min at room temperature, 20 mM $MgCl_2$ was added. The extracts were then centrifuged for 5 min at 12,000 rpm. The pellets were discarded and supernatants were centrifuged at 45,000 rpm for 1 h at 4°C. The pellets were resuspended in Laemli buffer and loaded on SDS-PAGE.

In vivo redox state of DsbC

Cells were grown in LB at 37°C to an A_{600} of 0.8, and 1 ml samples were taken. Proteins were precipitated with 5% ice cold TCA and centrifuged at $16,000 \times g$ for 15 min. The pellets were washed with acetone, dried, and resuspended in 50 mM Tris-HCl, pH 7.5, 0.1% SDS, 10 mM EDTA, and 10 mM AMS. AMS is a reagent that covalently reacts with free thiol groups, adding a 490 Da group. This leads to a major mobility shift respective of modified protein in SDS-PAGE gels. Samples were analyzed by SDS-PAGE under nonreducing conditions.

Antibodies

Antibodies against LamB were kindly provided by Natividad Ruiz and Tom Silhavy (Princeton), and antibodies against DsbC were provided by Jon Beckwith (Harvard).

Microarray analysis

RNA from WT, *dsbA*- and *dsbA-dsbC*- strains was extracted using the Tripure reagent and the RNeasy purification kit (Qiagen). Microarray analysis were performed in triplicates by using “GeneChip® *E. Coli* Genome 2.0 Array” and the protocol provided by Affymetrix for prokaryotic expression analysis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Geneviève Connerotte for technical help, Annie Hiniker and Maria Veiga da Cunha for helpful advice, and Emile Van Schaftingen for criticism of the manuscript. JFC is Chercheur Qualifié, LK is Chargé de Recherche, PL is Collaborateur Scientifique, and DV is Collaborateur logistique of the Belgian FRS-FNRS. MD is a research fellow of the FRIA and JM is a project leader of the VIB. This work was supported by the Interuniversity Attraction Pole Programme-Belgian Science Policy to JFC and MD (network P6/05) and to DV (network P6/28). This research was supported in part by grants from the FRS-FNRS to JFC and from the National Institutes of Health to J.C.A.B., an investigator of the Howard Hughes Medical Institute.

References

Altermark B, Smalas AO, Willassen NP, Helland R. The structure of *Vibrio cholerae* extracellular endonuclease I reveals the presence of a buried chloride ion. *Acta Crystallogr D Biol Crystallogr*. 2006; 62:1387–1391. [PubMed: 17057343]

- Andersen CL, Matthey-Dupraz A, Missiakas D, Raina S. A new *Escherichia coli* gene, *dsbG*, encodes a periplasmic protein involved in disulphide bond formation, required for recycling DsbA/DsbB and DsbC redox proteins. *Mol Microbiol.* 1997; 26:121–132. [PubMed: 9383195]
- Bader M, Muse W, Ballou DP, Gassner C, Bardwell JC. Oxidative protein folding is driven by the electron transport system. *Cell.* 1999; 98:217–227. [PubMed: 10428033]
- Bader MW, Hiniker A, Regeimbal J, Goldstone D, Haebel PW, Riemer J, Metcalf P, Bardwell JC. Turning a disulfide isomerase into an oxidase: DsbC mutants that imitate DsbA. *Embo J.* 2001; 20:1555–1562. [PubMed: 11285220]
- Bardwell JC, McGovern K, Beckwith J. Identification of a protein required for disulfide bond formation in vivo. *Cell.* 1991; 67:581–589. [PubMed: 1934062]
- Bardwell JC, Lee JO, Jander G, Martin N, Belin D, Beckwith J. A pathway for disulfide bond formation in vivo. *Proc Natl Acad Sci U S A.* 1993; 90:1038–1042. [PubMed: 8430071]
- Berkmen M, Boyd D, Beckwith J. The nonconsecutive disulfide bond of *Escherichia coli* phytase (AppA) renders it dependent on the protein-disulfide isomerase, DsbC. *J Biol Chem.* 2005; 280:11387–11394. [PubMed: 15642731]
- Bessette PH, Cotto JJ, Gilbert HF, Georgiou G. In vivo and in vitro function of the *Escherichia coli* periplasmic cysteine oxidoreductase DsbG. *J Biol Chem.* 1999; 274:7784–7792. [PubMed: 10075670]
- Bordes P, Bouvier J, Conter A, Kolb A, Gutierrez C. Transient repressor effect of Fis on the growth phase-regulated *osmE* promoter of *Escherichia coli* K12. *Mol Genet Genomics.* 2002; 268:206–213. [PubMed: 12395194]
- Bos C, Lorenzen D, Braun V. Specific in vivo labeling of cell surface-exposed protein loops: reactive cysteines in the predicted gating loop mark a ferrichrome binding site and a ligand-induced conformational change of the *Escherichia coli* FhuA protein. *J Bacteriol.* 1998; 180:605–613. [PubMed: 9457864]
- Chilcott GS, Hughes KT. Coupling of flagellar gene expression to flagellar assembly in *Salmonella enterica* serovar typhimurium and *Escherichia coli*. *Microbiol Mol Biol Rev.* 2000; 64:694–708. [PubMed: 11104815]
- Cho SH, Beckwith J. Mutations of the membrane-bound disulfide reductase DsbD that block electron transfer steps from cytoplasm to periplasm in *Escherichia coli*. *J Bacteriol.* 2006; 188:5066–5076. [PubMed: 16816179]
- Collet JF, Bardwell JC. Oxidative protein folding in bacteria. *Mol Microbiol.* 2002; 44:1–8. [PubMed: 11967064]
- Collet JF, Riemer J, Bader MW, Bardwell JC. Reconstitution of a disulfide isomerization system. *J Biol Chem.* 2002; 277:26886–26892. [PubMed: 12004064]
- Dailey FE, Berg HC. Mutants in disulfide bond formation that disrupt flagellar assembly in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 1993; 90:1043–1047. [PubMed: 8503954]
- Duguay AR, Silhavy TJ. Signal sequence mutations as tools for the characterization of LamB folding intermediates. *J Bacteriol.* 2002; 184:6918–6928. [PubMed: 12446642]
- Ferenci T, Stretton S. Cysteine-22 and cysteine-38 are not essential for the functions of maltoporin (LamB protein). *FEMS Microbiol Lett.* 1989; 52:335–339. [PubMed: 2693195]
- Gruber CW, Cemazar M, Heras B, Martin JL, Craik DJ. Protein disulfide isomerase: the structure of oxidative folding. *Trends Biochem Sci.* 2006; 31:455–464. [PubMed: 16815710]
- Hagenmaier S, Stierhof YD, Henning U. A new periplasmic protein of *Escherichia coli* which is synthesized in spheroplasts but not in intact cells. *J Bacteriol.* 1997; 179:2073–2076. [PubMed: 9068658]
- Heras B, Edeling MA, Schirra HJ, Raina S, Martin JL. Crystal structures of the DsbG disulfide isomerase reveal an unstable disulfide. *Proc Natl Acad Sci U S A.* 2004; 101:8876–8881. [PubMed: 15184683]
- Hiniker A, Bardwell JC. In vivo substrate specificity of periplasmic disulfide oxidoreductases. *J Biol Chem.* 2004; 279:12967–12973. [PubMed: 14726535]
- Hiniker A, Collet JF, Bardwell JC. Copper stress causes an in vivo requirement for the *Escherichia coli* disulfide isomerase DsbC. *J Biol Chem.* 2005; 280:33785–33791. [PubMed: 16087673]

- Hiniker A, Vertommen D, Bardwell JC, Collet JF. Evidence for conformational changes within DsbD: possible role for membrane-embedded proline residues. *J Bacteriol.* 2006; 188:7317–7320. [PubMed: 17015672]
- Hiniker A, Ren G, Heras B, Zheng Y, Laurinec S, Jobson RW, Stuckey JA, Martin JL, Bardwell JC. Laboratory evolution of one disulfide isomerase to resemble another. *Proc Natl Acad Sci U S A.* 2007; 104:11670–11675. [PubMed: 17609373]
- Inaba K, Murakami S, Suzuki M, Nakagawa A, Yamashita E, Okada K, Ito K. Crystal structure of the DsbB-DsbA complex reveals a mechanism of disulfide bond generation. *Cell.* 2006; 127:789–801. [PubMed: 17110337]
- Kadokura H, Tian H, Zander T, Bardwell JC, Beckwith J. Snapshots of DsbA in action: detection of proteins in the process of oxidative folding. *Science.* 2004; 303:534–537. [PubMed: 14739460]
- Katzen F, Beckwith J. Transmembrane electron transfer by the membrane protein DsbD occurs via a disulfide bond cascade. *Cell.* 2000; 103:769–779. [PubMed: 11114333]
- Lange R, Barth M, Hengge-Aronis R. Complex transcriptional control of the sigma s-dependent stationary-phase-induced and osmotically regulated *osmY* (*csi-5*) gene suggests novel roles for Lrp, cyclic AMP (cAMP) receptor protein-cAMP complex, and integration host factor in the stationary-phase response of *Escherichia coli*. *J Bacteriol.* 1993; 175:7910–7917. [PubMed: 8253679]
- Leichert LI, Jakob U. Protein thiol modifications visualized in vivo. *PLoS Biol.* 2004; 2:e333. [PubMed: 15502869]
- Liu H, Sadygov RG, Yates JR 3rd. A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal Chem.* 2004; 76:4193–4201. [PubMed: 15253663]
- Liu X, Wang CC. Disulfide-dependent folding and export of *Escherichia coli* DsbC. *J Biol Chem.* 2001; 276:1146–1151. [PubMed: 11042167]
- Mecenas J, Rouviere PE, Erickson JW, Donohue TJ, Gross CA. The activity of sigma E, an *Escherichia coli* heat-inducible sigma-factor, is modulated by expression of outer membrane proteins. *Genes Dev.* 1993; 7:2618–2628. [PubMed: 8276244]
- Messens J, Collet JF. Pathways of disulfide bond formation in *Escherichia coli*. *Int J Biochem Cell Biol.* 2006; 38:1050–1062. [PubMed: 16446111]
- Messens J, Collet JF, Van Belle K, Brosens E, Loris R, Wyns L. The oxidase DsbA folds a protein with a nonconsecutive disulfide. *J Biol Chem.* 2007
- Old WM, Meyer-Arendt K, Aveline-Wolf L, Pierce KG, Mendoza A, Sevinsky JR, Resing KA, Ahn NG. Comparison of label-free methods for quantifying human proteins by shotgun proteomics. *Mol Cell Proteomics.* 2005; 4:1487–1502. [PubMed: 15979981]
- Pugsley AP. A mutation in the *dsbA* gene coding for periplasmic disulfide oxidoreductase reduces transcription of the *Escherichia coli* *ompF* gene. *Mol Gen Genet.* 1993; 237:407–411. [PubMed: 8483456]
- Repoila F, Gutierrez C. Osmotic induction of the periplasmic trehalase in *Escherichia coli* K12: characterization of the *treA* gene promoter. *Mol Microbiol.* 1991; 5:747–755. [PubMed: 1710760]
- Rietsch A, Belin D, Martin N, Beckwith J. An in vivo pathway for disulfide bond isomerization in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 1996; 93:13048–13053. [PubMed: 8917542]
- Rozhkova A, Stirnimann CU, Frei P, Grauschopf U, Brunisholz R, Grutter MG, Capitani G, Glockshuber R. Structural basis and kinetics of inter- and intramolecular disulfide exchange in the redox catalyst DsbD. *Embo J.* 2004; 23:1709–1719. [PubMed: 15057279]
- Ruiz N, Silhavy TJ. Sensing external stress: watchdogs of the *Escherichia coli* cell envelope. *Curr Opin Microbiol.* 2005; 8:122–126. [PubMed: 15802241]
- Sevier CS, Kaiser CA. Conservation and diversity of the cellular disulfide bond formation pathways. *Antioxid Redox Signal.* 2006; 8:797–811. [PubMed: 16771671]
- Zapun A, Missiakas D, Raina S, Creighton TE. Structural and functional characterization of DsbC, a protein involved in disulfide bond formation in *Escherichia coli*. *Biochemistry.* 1995; 34:5075–5089. [PubMed: 7536035]




Fig. 1. The absence of *dsbA* and *dsbC* has phenotypical consequences

A. Growth curves of wild-type (□), *dsbC*⁻ (▲), *dsbA*⁻ (*), *dsbA*⁻*dsbD*⁻ (■) and *dsbA*⁻*dsbC*⁻ (●) strains in M63 minimal media at 37°C. Growth was monitored at A₆₀₀.

B. SDS sensitivity of wild-type (lane 1), *dsbA*⁻ (lane 2), *dsbA*⁻*dsbC*⁻ (lane 3) and *dsbA*⁻*dsbD*⁻ (lane 4) strains. Strains were grown in LB at 37°C to an A₆₀₀ of 0.5. The cultures were then serially diluted 10⁷-fold in 10-fold increments. 10 μl of each dilution were then spotted on LB plates containing 2.5% SDS and grown overnight.

C. Western blot showing protein expression levels. The upper bands correspond to the LamB protein. Outer membrane proteins prepared from wild-type (lane 1), *dsbA*⁻ (lane 2), *dsbC*⁻ (lane 3), *dsbA*⁻*dsbC*⁻ (lane 4), and *LamB*⁻ (lane 5) strains. The lower bands correspond to an unknown protein recognized by the anti-LamB antibody, which was used as an internal standard.




Fig. 2. *In vivo* redox state of DsbC. Exponentially growing cells (in LB) were TCA-precipitated, free cysteines were modified by AMS, and DsbC was detected by Western blot analysis. Lanes: 1, wild-type; 2, *dsbA*⁻; 3, *dsbA*⁻*dsbD*⁻; 4, *dsbD*⁻.




Fig. 3. The number of spectral counts correlates with the abundance of a protein. Varying amounts (2–60 pmoles) of two eukaryotic proteins, ovalbumin and carbonic anhydrase, were added to 300 μg of periplasmic proteins. The SC values obtained for these two proteins in the various samples were then plotted against the corresponding protein amounts. After linear regression, we found that the R^2 values obtained for the spectral counts were 0.94 and 0.92 for ovalbumin and carbonic anhydrase, respectively. This indicates that the number of spectral counts reliably reflects protein abundance in the sample.




Fig. 4. The overall protein content of a $dsbA^-dsbC^-$ mutant is different compared to $dsbA^-$ and $dsbA^-dsbD^-$ strains

A. The logarithms of the SC values reported for the $dsbA^-$ strain were plotted against those reported for the $dsbA^-dsbD^-$ mutant. Most of the SC values are similar in both strains, which is reflected by a quasi-linear distribution.

B. The logarithms of the SC values reported for the $dsbA^-$ strain were plotted against those reported for the $dsbA^-dsbC^-$ mutant. The distribution is more dispersed, which indicates that the overall protein content of this double mutant is different.




Fig. 5. A revised model for the formation of disulfide bonds in the *E. coli* periplasm

Disulfide bonds are introduced by the DsbA/DsbB pathway. Non-native disulfides are corrected by DsbC, which is recycled by DsbD. Both pathways are kinetically isolated. Our results indicate that DsbC is also able to function on the other side of the barrier where it assists DsbA in a DsbD-independent manner. DsbC may be acting as a stand-alone protein folding catalyst that cycles from the reduced to the oxidized state upon substrate oxidation and substrate reduction, respectively. Although kinetics data showed that DsbC is not a good substrate for DsbB, we cannot exclude that a slow oxidation of DsbC by DsbB may play a more significant role in the absence of DsbA. The Western blot data presented in Figure 2 also suggest that in the absence of DsbD, DsbA may be responsible for the oxidation of DsbC. The redox potentials of DsbA and DsbC are -125 mV and -130 mV, respectively.

Table 1

Strains used in this study and their relevant genotypes.

Strain	Relevant genotype	Source
AH50	MC1000 <i>phoR</i> Δ <i>ara714</i> <i>leu</i> ⁺ <i>phoA68</i>	Hiniker <i>et al.</i> , 2005
JFC383	AH50 <i>dsbC::kan</i>	This study
AH396	AH50 <i>dsbD::cm</i> , <i>dsbA::kan1</i>	Hiniker <i>et al.</i> , 2005
MD1	AH50 <i>dsbA::kan1</i>	This study
MD3	AH50 <i>dsbC::cm</i> , <i>dsbA::kan</i>	This study
JP114	ER1821	New England Biolabs
JP220	JP114 Δ <i>dsbA::kan</i>	This study
JP539	JP114 Δ <i>dsbC::cm</i>	This study
JP557	JP220 Δ <i>dsbC::cm</i>	This study
JP649	JP114 Δ <i>dsbA::kan dsbD</i> ⁻ <i>::cm</i>	This study
C600	JP114 <i>fluA::kan</i>	This study

Table 2Phenotypic characterization of *dsbA*⁻*dsbC*⁻ mutants.

Table 2A.				
Strain	Genotype	MacConkey Maltodextrin (1%)	Rifampin (4 µg/ml)	Growth rate (h⁻¹) *
JFC209	Wild-type	Mal+ (red)	+++	1.15
MD1	<i>dsbA</i> ⁻	Mal+ (red)	++	0.79
JFC383	<i>dsbC</i> ⁻	Mal+ (red)	+++	1.15
MD3	<i>dsbA</i> ⁻ <i>dsbC</i> ⁻	Mal+/- (pink)	-	0.69
AH396	<i>dsbA</i> ⁻ <i>dsbD</i> ⁻	Mal+ (red)	++	0.87

Table 2B.		
Strain	Genotype	Pfu with Φ80
JP114	Wild-type	5.5 x 10 ¹⁰
JP220	<i>dsbA</i> ⁻	1.0 x 10 ⁹
JP539	<i>dsbC</i> ⁻	5.1 x 10 ¹⁰
JP557	<i>dsbA</i> ⁻ <i>dsbC</i> ⁻	2.0 x 10 ⁴
JP649	<i>dsbA</i> ⁻ <i>dsbD</i> ⁻	7.0 x 10 ⁹
C600	<i>fhua</i> ⁻	0

* Growth rate have been calculated using the following formula: $\mu = \ln(A_{600})/\text{time}$

Table 3

Proteins more than two-fold less abundant in a *dsbA*⁻*dsbC*⁻ mutant than in *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ strains ($P < 0.05$)¹. The levels of expression of the genes coding for these proteins in the *dsbA*⁻*dsbC*⁻ double mutant relative to the *dsbA*⁻ mutant are shown in grey.

Protein	# Cysteines	Spectral counts			RNA ² (<i>dsbA</i> ⁻ <i>dsbC</i> ⁻ vs <i>dsbA</i> ⁻)
		<i>dsbA</i> ⁻	<i>dsbA</i> ⁻ <i>dsbC</i> ⁻	<i>dsbA</i> ⁻ <i>dsbD</i> ⁻	
Proteins with at least two cysteine residues					
Cn16	2	2	0	3	NSD
DsbC	4	12	0	8	↓ (110)
GitI	2	5	0	6	NSD
OppA ³	2	305	137	259	NSD
YggN	2	2	0	2	NSD
TreA ³	2	8	1	11	nd
YhjJ	2	10	2	9	NSD
Proteins with one or no cysteine residue					
FepB	0	5	0	3	NSD
FliY	0	151	65	132	NSD
SubI	0	9	0	6	NSD
YcfS	1	2	0	2	NSD
GgT	0	8	3	8	NSD
MalE	0	41	1	24	nd
MppA	0	14	3	15	NSD
PhnD	0	159	79	168	NSD
Slp	1	5	0	3	↓ (2.9)
YthB	1	10	4	11	NSD
YjhT	0	4	0	2	NSD
YnjE	1	14	0	6	NSD
YncE	0	15	4	14	NSD

¹ All proteins that were identified in the *dsbA*⁻ and *dsbA*⁻*dsbD*⁻ strains but were absent in all three independent analysis of the *dsbA*⁻*dsbC*⁻ were also considered as significantly less abundant in the latter strain.

² Genes whose expression ratios were >2 fold different ($P < 0.05$) in the $dsbA^- dsbC^-$ double mutant relative to the $dsbA^-$ mutant were considered as induced (↑) or repressed (↓). All other genes were considered as not significantly different (NSD). The ratio indicated next to the arrows is the fold increase or decrease in transcript level in the $dsbA^- dsbC^-$ strain compared to the $dsbA^-$ strain. "nd" means that no data were obtained for the corresponding gene.

³ $P = 0.05$ when SC values from the $dsbA^- dsbC^-$ are compared to those from the $dsbA^- dsbD^-$.

A complete list with all the identified proteins is available in Table S2. Microarray data for the genes corresponding to the identified proteins are available in Table S3.

Table 4

Proteins whose abundance is significantly changed in *dsbA*⁻ strains

The levels of expression of the genes coding for these proteins in the *dsbA*⁻ mutant relative to the wild-type are shown in grey.

Protein	Proteins > two-fold less abundant in <i>dsbA</i> ⁻ strains ($P < 0.05$) ^d		RNA ² <i>dsbA</i> ⁻ vs wild-type		Proteins > two-fold more abundant in <i>dsbA</i> ⁻ strains ($P < 0.05$)		RNA ² (<i>dsbA</i> ⁻ vs wild-type)
	Spectral counts		NSD	↓ (2.6)	Spectral counts		
	wild-type	<i>dsbA</i> ⁻			Wild-type	<i>dsbA</i> ⁻	
<u>ArgT</u> (2 cys) ^d	54	9	NSD		1	27	↑ (2.0)
<u>ArtJ</u> (2 cys) ^d	301	150	↑ (3.5)		24	62	NSD
<u>DppA</u> (4 cys)	99	34	NSD		0	8	↑ (2.3)
<u>GljI</u> (2 cys)	49	5	NSD		26	224	↑ (7.9)
<u>HslI</u> (2 cys)	170	84	↓ (2.6)		131	401	NSD
<u>PhoA</u> (4 cys)	212	29	↑ (19.4)		27	152	NSD
<u>ProX</u> (2 cys) ^d	46	8	NSD		8	32	NSD [#]
<u>YfhM</u> (2 cys)	1	0	NSD		16	39	NSD [#]
<u>YebF</u> (2 cys) ^d	36	1	↑ (3.3)		104	285	↑ (6.0)
<u>YggN</u> (2 cys)	5	2	NSD		9	31	↑ (9.6)
<u>YtfQ</u> (2 cys) ^d	8	0	↑ (27.7)		0	3	nd
<u>CirA</u> (2 cys) ^d	8	0	NSD		0	10	↑ (5.3)
<u>CreA</u> (2 cys) ^d	2	0	NSD		1	8	nd
<u>EndI</u> (8 cys) ^d	1	0	nd		0	8	↑ (9.3)
<u>MepA</u> (6 cys)	4	0	NSD		8	30	↑ (4.5)
<u>RNase I</u> (8 cys)	4	0	NSD		11	108	↑ (9.7)
<u>YehY</u> (2 cys) ^d	8	0	NSD		7	41	↑ (2.8)
<u>DsbA</u>	131	0	↓ (3.4)		1	14	↑ (3.7)
<u>FlgH</u>	5	0	NSD		6	20	NSD
<u>FltC</u>	219	0	↑ (92.5)		3	10	NSD
<u>OmpF</u>	136	13	↓ (10.7)		0	6	↑ (5.0)
<u>PhoE</u>	86	9	↑ (3.9)		0	5	↑ (5.9)

Protein	Proteins > two-fold less abundant in <i>dsbA</i> ⁻ strains (<i>P</i> < 0.05) ¹		RNA ² <i>dsbA</i> ⁻ vs wild-type		Proteins > two-fold more abundant in <i>dsbA</i> ⁻ strains (<i>P</i> < 0.05)			RNA ² (<i>dsbA</i> ⁻ vs wild-type)
	Spectral counts		↓ (*) NSD	↓ (*) NSD	Spectral counts			
	wild-type	<i>dsbA</i> ⁻			Wild-type	<i>dsbA</i> ⁻		
FlgG	5	0	↓ (*)	NSD	0	2		nd
OmpN	2	0	NSD	NSD	0	3		↑ (4,5)
					1	3		NSD

DsbA substrates are underlined.

¹ Newly identified DsbA substrates.

* *flgG* transcripts were not detected in the *dsbA*⁻

osmE and *opgG* transcription rates were increased by 1.82 (*P*=0.026) and 1.7 (*P*=0.0039), respectively.

¹ All proteins that were identified in the wild-type strain but were absent in all three independent analysis of the *dsbA*⁻ were also considered as significantly less abundant in the latter strain.

² Genes whose expression ratios were >2 fold different (*P*<0.05) in the *dsbA*⁻ mutant relative to the wild-type were considered as induced (↑) or repressed (↓). All other genes were considered as not significantly different (NSD). The ratio indicated next to the arrows is the fold increase or decrease in transcript level in the *dsbA*⁻ strain compared to the wild-type strain. "nd" means that no data were obtained for the corresponding gene.

A complete list with all the identified proteins is available in Table S1. Microarray data for the genes corresponding to the identified proteins are available in Table S4.

Table 5

Redox state of cysteine-containing peptides.

Protein	cysteine-containing peptides found	wild-type (% red)	<i>dsbA</i> ⁻ (% red)	<i>dsbA</i> ⁻ <i>dsbD</i> ⁻ (% red)	<i>dsbA</i> ⁻ <i>dsbC</i> ⁻ (% red)
PhoA	ATYHGNDKPAVTCTPNPQR	2	45	66	66
	CYGPSATSEK	0	-	-	-
	QDHAANPCGQIGETVLDLDEAVQR	0	50	33	50
OmpA	AALIDCLAPDRR	0	47	32	52
	GMGESNPVTGNTCDNVK	2	85	83	90
AtJ	QMQAECTFTNHFDSLPSLK	11	100	100	80
YebF	SADIHYSVSDCK	15	-	-	-
	CEDLDAAGIAASVK	0	-	-	-
DppA	NEQVMPYPNPADIAR	0	-	-	-
GitI	PQSQEA YGCMLR	0	-	-	-
MepA	TPPPLPPSQALLDEHVI	0	-	-	-
RNaseI	AVKLTCQGNPAYLTEIQSIK	0	-	-	-
YneA	VLTWDSDTKPECR	0	-	-	-
YtfQ	KPCNVVELQGTVGASVAIDR	0	-	-	-
	VSSPVSTMMACPDGK	-	100	100	100
EcoT	VELLIGQTLEVDCNLHR	-	100	100	100
YedD	VDRPTAEC AAALDK	-	75	100	60
YegG	TLSDQACQEMDSK	-	66	66	50