

## Association of Fibril Structure Formation with Cell Surface Properties of *Yersinia enterocolitica*

R. V. LACHICA,<sup>1</sup>\* D. L. ZINK,<sup>1,2</sup>† AND W. R. FERRIS<sup>3</sup>

Departments of Nutrition and Food Science,<sup>1</sup> Microbiology,<sup>2</sup> and Cellular Biology,<sup>3</sup> The University of Arizona, Tucson, Arizona 85721

Received 5 March 1984/Accepted 25 June 1984

**Electron microscopic examination of virulence plasmid-bearing cells of *Yersinia enterocolitica* revealed the formation of a fibril structure when grown at 37°C but not at 22°C. Plasmidless cells did not exhibit any surface matrix. The formation of this surface component was associated with increased cell surface charge and hydrophobicity, formation of long chains of cells, and autoagglutination. This structure is distinct from the rigid appendages (fimbriae) which are elaborated by certain *Yersinia* strains.**

Adherence of various pathogenic bacteria to eucaryotic cells has been demonstrated to play a central role in their ability to colonize epithelial surfaces (24). Both specific and nonspecific interactions have been shown to mediate the adherence of certain pathogens. Specific interaction involves the binding of a specific surface structure on the bacterial cell surface with receptors on the epithelial cells. Nonspecific interaction depends on physicochemical forces between the surfaces of the epithelial tissue and bacterial cells. Formation of fimbriae has been associated with cell surface hydrophobicity among various pathogens (7, 10).

There is considerable interest in the pathogenicity of *Yersinia enterocolitica* since a resident plasmid contributes to its virulence (9, 21-23, 27). Although various properties of *Y. enterocolitica* are associated with the virulence plasmid (9, 21-23, 28), attachment to the mucosal surface is thought to be chromosomally mediated (12, 17, 23). Certain strains have been reported to elaborate fimbriae (19) which mediate increased cell surface hydrophobicity (6). However, low growth temperatures (<30°C) are required for their synthesis. Moreover, no correlation has been observed between the formation of fimbriae and the attachment to epithelial cells in vitro (19). We have recently reported the association of mouse gastrointestinal tract colonization with the plasmid-mediated cell surface properties of *Y. enterocolitica* (13, 14). In the present study, the formation of a fibrillar material associated with the virulence plasmid and the cell surface properties of *Y. enterocolitica* are described.

We investigated in greater detail the autoagglutination (AA) reaction, which is one of the properties associated with the virulence of *Y. enterocolitica* (15, 22, 28). Plasmid-bearing *Y. enterocolitica* MCH56A2 (p<sup>+</sup>; serotype O:3) was AA<sup>+</sup> at 37°C but not at 22°C; its plasmidless derivative (p<sup>-</sup>) was AA<sup>-</sup>, regardless of the growth temperature, as described in a previous report (14). During growth at 37°C in Eagle minimal essential medium plus 10% fetal calf serum, p<sup>+</sup> cell aggregates adhered to the wall of the polystyrene test tube and eventually sank to the bottom of the tube. Some

remained attached to the tube wall; these were best observed by staining with crystal violet. The AA reaction of p<sup>+</sup> cells is in marked contrast to the autoaggregation of fimbriated bacteria which form thick pellicles at the broth-air interphases of undisturbed still cultures (6, 20).

Under phase-contrast light microscopy, the p<sup>+</sup> cells grown at 37°C were shown (Fig. 1) to form long chains (some chains consisted of 12 cells) which were attached to each other and to the glass surface. None of these were observed when p<sup>+</sup> cells were grown in brain heart infusion or tryptic soy broth (Difco Laboratories, Detroit, Mich.). The p<sup>-</sup> cells grew at 37°C as single, unattached pleomorphic cells. At 22°C, both strains were motile, short rods.

Transmission electron microscopy was employed to determine the formation of the surface structure on p<sup>+</sup> cells. From our previous study (13), p<sup>+</sup> cells grown at 37°C were shown to exhibit a highly anionic surface. This information suggested to us the efficacy of ruthenium red (RR; Sigma Chemical Co., St. Louis, Mo.) for staining the putative surface structure. This electron-dense cationic stain has been useful in visualizing anionic polysaccharide fibers (glycocalyx) on the surface of various bacteria (5). Pellets of cells or mixtures of cells and latex particles (0.5 μm in diameter; E. F. Fullum, Inc., Schenectady, N.Y.) were fixed in 1.5% glutaraldehyde in 0.2 M cacodylate buffer (pH 7.2) containing 0.05% RR at 22°C for 2 h. After a brief rinse in cacodylate buffer, the pellets were postfixed in 2% OsO<sub>4</sub> in cacodylate buffer containing RR for 1 h at 0°C. After washing the pellets with cacodylate buffer containing RR, they were dehydrated with graded series of ethanol and embedded with Epon 812 by the method of Luft (16). Sections were cut and examined with a Philips model 200 electron microscope at 60 kV.

The plasmid-bearing cells grown at 37°C elaborated a matrix of fibers distal to the outer membrane (Fig. 2A). This fibrillar material appeared to facilitate the attachment of the cells to each other. Recently, we reported this cell-cell interaction (AA) to be accelerated by the addition of divalent cations such as Ca<sup>2+</sup> or Zn<sup>2+</sup>, which are thought to form a bridge between the anionic surfaces of the cells (14). The cells also interacted with the hydrophobic latex particles. Interestingly, the latter were coated with what appeared to be fibrillar material that may have been dislodged from the cell surface. No fibrillar component was observed on plasmid-bearing cells previously grown at 22°C (Fig. 2B) or on plasmidless cells, regardless of the growth temperature.

\* Corresponding author.

† Present address: Eastern Regional Research Center, Agricultural Research Service, U.S. Department of Agriculture, Philadelphia, PA 19118.

‡ Present address: Campbell Institute for Research and Technology, Campbell Place, Camden, NJ 08101.




FIG. 1. Phase-contrast micrograph showing chain formation and clumping of *Y. enterocolitica* MCH56A2 ( $p^+$ ) cells grown in Eagle minimal essential medium plus 10% fetal calf serum. Magnification,  $\times 1,500$ .

The fact that the fibrous matrix was demonstrated despite the absence of any stabilizing influence suggests that the matrix is not as highly hydrated as the extracellular polysaccharide matrix exhibited by other bacteria (5). Yet, it is possible that a better image of the surface matrix could be attained with the use of stabilizing influences such as lectins or antiserum (5) or the employment of different dehydration techniques (25). The flexible nature of the fibril structure is similar to the capsular antigen of *Yersinia pestis* described by Chen et al. (4). The flexible fibrillar material on the cell surface of *Y. enterocolitica* and the rigid gonococcal and *Escherichia coli* type 1 fimbriae share the property of attachment to latex particles (2, 3), indicating their hydrophobic nature.

Our results suggest that the fibrillar material of *Y. enterocolitica* mediates the surface properties of the cell which remained undiminished after treatment with sodium meta-periodate (10 mM, 30 min) and actually increased when heated at 60°C for 60 min (data not shown). These observations indicate that the fibrillar material is flexible, amphipathic protein similar to  $\beta$ -casein (18). It is possible that some of the plasmid-mediated outer membrane proteins elaborated by *Y. enterocolitica* (1, 17, 23) may comprise components of the fibril structure.

Our results indicate that the AA reaction is based on the formation of long chains of cells and the alteration of cell surface properties likely to be mediated by the fibrillar material. The recently described mannose-resistant aggluti-




FIG. 2. Electron micrographs of RR-stained  $p^+$  cells grown at 37°C (A) and 22°C (B). Note the presence of fibrous material on the surface of the cells grown at 37°C and on latex particles (L) but not on cells grown at 22°C.

nation of guinea pig erythrocytes by virulent cells of *Y. enterocolitica* (11) is also likely to be mediated by the fibril structure. It is envisaged that this surface structure promotes the attachment of p<sup>+</sup> cells to the intestinal mucosa. The increased susceptibility to phagocytosis as a result of the increased surface hydrophobicity (26) may be counteracted by the formation of long chains and microcolonies (5). The fibril structure may act also as a molecular barrier against the killing action of serum (21).

It should be emphasized that the surface structure described in this report is distinct from the rigid appendages (fimbriae) which are elaborated by certain yersiniae after several subculturings at low (<35°C) but not high (>35°C) temperatures (6, 19). Presumably, the expression of the fimbriae is chromosomally encoded among the environmental group of yersiniae (11). Whether virulent strains of *Y. enterocolitica* are capable of elaborating fimbriae is not known.

Studies are in progress as to the determination of whether the fibril structure or the plasmid-mediated outer membrane proteins or both play the major role in mediating the altered cell surface properties of virulent strains of *Y. enterocolitica*.

This work was supported in part by Arizona Agriculture Experiment Station, project no. 173804.

#### LITERATURE CITED

- Bölin, I., L. Norlander, and H. Wolf-Watz. 1982. Temperature-inducible outer membrane protein of *Yersinia pseudotuberculosis* and *Yersinia enterocolitica* is associated with the virulence plasmid. *Infect. Immun.* **37**:506-512.
- Brinton, C. C. 1965. The structure, function, synthesis, and genetic control of bacterial pili and a molecular model for DNA and RNA transport in gram negative bacteria. *Trans. N.Y. Acad. Sci.* **27**:1003-1054.
- Buchanan, T. M. 1976. Attachment of purified gonococcal pili to latex spheres, p. 491-493. *In* D. Schlessinger (ed.), *Microbiology—1976*. American Society for Microbiology, Washington, D.C.
- Chen, T. H., S. S. Elberg, J. Boyles, and M. A. Velez. 1975. *Yersinia pestis*: correlation of ultrastructure and immunological status. *Infect. Immun.* **11**:1382-1390.
- Costerton, J. W., and R. T. Irvin. 1981. The bacterial glycocalyx in nature and disease. *Annu. Rev. Microbiol.* **35**:299-324.
- Faris, A., M. Lindahl, Å. Ljungh, D. C. Old, and T. Wadström. 1983. Autoaggregating *Yersinia enterocolitica* express surface fimbriae with high surface hydrophobicity. *J. Appl. Bacteriol.* **55**:97-100.
- Faris, A., M. Lindahl, and T. Wadström. 1982. High surface hydrophobicity of hemagglutinating *Vibrio cholerae* and other vibrios. *Curr. Microbiol.* **7**:357-362.
- Faris, A., T. Wadström, and J. H. Freer. 1981. Hydrophobic adsorptive and hemagglutinating properties of *Escherichia coli* possessing colonization factor antigens (CFA/I or CFA/II), type 1 pili, and other pili. *Curr. Microbiol.* **5**:67-72.
- Gemski, P., J. R. Lazere, and T. Casey. 1980. Plasmid associated with pathogenicity and calcium dependency of *Yersinia enterocolitica*. *Infect. Immun.* **27**:682-685.
- Gibbons, R. J., I. Etherden, and Z. Skobe. 1983. Association of fimbriae with the hydrophobicity of *Streptococcus sanguis* FC-1 and adherence to salivary pellicles. *Infect. Immun.* **41**:414-417.
- Kapperud, G., and J. Lassen. 1983. Relationship of virulence-associated autoagglutination to hemagglutinin production in *Yersinia enterocolitica*-like bacteria. *Infect. Immun.* **42**:163-169.
- Kilström, E., and K.-E. Magnusson. 1983. Haemagglutinating, adhesive, and physico-chemical surface properties of different *Yersinia enterocolitica* and *Yersinia enterocolitica*-like bacteria. *Acta Pathol. Microbiol. Scand. Sect. B* **91**:113-119.
- Lachica, R. V., and D. L. Zink. 1984. Plasmid-associated cell surface charge and hydrophobicity of *Yersinia enterocolitica*. *Infect. Immun.* **44**:540-543.
- Lachica, R. V., and D. L. Zink. 1984. Determination of plasmid-associated hydrophobicity of *Yersinia enterocolitica* by a latex particle agglutination test. *J. Clin. Microbiol.* **19**:660-663.
- Laird, W. J., and D. C. Cavanaugh. 1980. Correlation of autoagglutination and virulence of yersiniae. *J. Clin. Microbiol.* **11**:430-432.
- Luft, J. H. 1971. Ruthenium red and ruthenium violet. I. Chemistry purification, methods of use for electron microscopy, and mechanism of action. *Anat. Rec.* **171**:347-368.
- Martinez, R. J. 1983. Plasmid-mediated and temperature-regulated surface properties of *Yersinia enterocolitica*. *Infect. Immun.* **41**:921-930.
- Morr, C. V. 1979. Conformation and functionality of milk proteins, p. 65-79. *In* A. Pour-El (ed.), *Functionality and protein structure*. American Chemical Society, Washington, D.C.
- Old, D. C., and J. Robertson. 1981. Adherence of fimbriate and nonfimbriate strains of *Yersinia enterocolitica* to human epithelial cells. *Microbiol. Immunol.* **25**:993-998.
- Ottow, J. C. G. 1975. Ecology, physiology, and genetics of fimbriae and pili. *Annu. Rev. Microbiol.* **29**:79-108.
- Pai, C. H., and L. DeStephano. 1982. Serum resistance associated with virulence in *Yersinia enterocolitica*. *Infect. Immun.* **35**:605-611.
- Perry, R. D., and R. R. Brubaker. 1983. Vwa<sup>+</sup> phenotype of *Yersinia enterocolitica*. *Infect. Immun.* **40**:166-171.
- Portnoy, D. A., S. L. Moseley, and S. Falkow. 1981. Characterization of plasmids and plasmid-associated determinants of *Yersinia enterocolitica* pathogenesis. *Infect. Immun.* **31**:775-782.
- Smith, H. 1977. Microbial surfaces in relation to pathogenicity. *Bacteriol. Rev.* **41**:475-500.
- Springer, E. L., and I. L. Roth. 1973. The ultrastructure of the capsules of *Diplococcus pneumoniae* and *Klebsiella pneumoniae* strains with ruthenium red. *J. Gen. Microbiol.* **74**:21-31.
- Van Oss, C. J. 1978. Phagocytosis as a surface phenomenon. *Annu. Rev. Microbiol.* **32**:19-39.
- Zink, D. L., J. C. Feeley, J. G. Wells, C. Vanderzant, J. C. Vickery, W. C. Roof, and G. A. O'Donovan. 1980. Plasmid-mediated tissue invasiveness in *Yersinia enterocolitica*. *Nature (London)* **283**:224-226.
- Zink, D. L., R. V. Lachica, and J. R. Dubel. 1982. *Yersinia enterocolitica* and *Yersinia enterocolitica*-like species: their pathogenicity and significance in foods. *J. Food Safety* **4**:223-241.